



HAL
open science

A high resolution authigenic $^{10}\text{Be}/^{9}\text{Be}$ record of geomagnetic moment variations over the last 300 ka from sedimentary cores of the Portuguese margin.

Julien Carcaillet, Didier Bourlès, Nicolas Thouveny, Maurice Arnold

► **To cite this version:**

Julien Carcaillet, Didier Bourlès, Nicolas Thouveny, Maurice Arnold. A high resolution authigenic $^{10}\text{Be}/^{9}\text{Be}$ record of geomagnetic moment variations over the last 300 ka from sedimentary cores of the Portuguese margin.. Earth and Planetary Science Letters, 2004, 219, pp.397-412. hal-00099482

HAL Id: hal-00099482

<https://hal.science/hal-00099482>

Submitted on 26 Sep 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A HIGH RESOLUTION AUTHIGENIC $^{10}\text{Be}/^9\text{Be}$ RECORD OF GEOMAGNETIC MOMENT VARIATIONS OVER THE LAST 300 KA FROM SEDIMENTARY CORES OF THE PORTUGUESE MARGIN.

Julien Carcaillet¹, Didier L. Bourlès¹, Nicolas Thouveny¹ and Maurice Arnold².

1. *CEREGE, Europôle Méditerranéen de l'Arbois, BP 80, F-13545 Aix en Provence cedex 04, France.*

2. *UMS2004 (CNRS-CEA), Bâtiment 30, Avenue de la Terrasse, F-91198 Gif-sur-Yvette Cedex France*

Corresponding author : carcaillet@cerege.fr

Abstract

A high resolution study of authigenic Be isotopes (^{10}Be and ^9Be) combined with continuous relative paleointensity records has been performed along the same marine sedimentary sequences from the Portuguese margin (N.E. Atlantic) covering the past 300 ka in order to assess relationships between geomagnetic moment variations and ^{10}Be production rate variations. A careful examination of the various ways of taking into account environmental disturbing effects on the authigenic ^{10}Be concentration leads to the conclusion that the most reliable proxy of cosmonuclide production rates is presently the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio. Eight intervals of significant authigenic $^{10}\text{Be}/^9\text{Be}$ enhancement evidence geomagnetic moment drops related to global paleomagnetic excursions, some being already admitted, others being proposed as new geomagnetic features. Since, contrarily to sedimentary magnetic remanence, the authigenic $^{10}\text{Be}/^9\text{Be}$ records dipole moment variations without significant acquisition delay, it provides better constraints on their timing. Comparison of $^{10}\text{Be}/^9\text{Be}$ and benthic $\delta^{18}\text{O}$ records from the same cores suggests that dipole moment lows preferentially occurred during or at the end of interglacial episodes, with a quasi-period of 100 ka.

Keywords : authigenic Be isotopes, cosmonuclides, geomagnetic dipole lows, geomagnetic excursions.

1. Introduction

Production of cosmogenic nuclides that varies with space (altitude and latitude), also varies with time. Composed primarily of high energy protons and α -particles, the cosmic-ray flux entering the magnetosphere is deflected by Lorentz forces. Cosmogenic production rate is thus proportional to the galactic cosmic ray flux and inversely proportional to the solar activity and the Earth's magnetic field intensity [1-4]. Measurements of cosmogenic radionuclides with different half-lives in lunar samples and in meteorites having different irradiation histories [5], and in marine sediments [6] and manganese crusts [7] yield to the conclusion that the average galactic cosmic ray flux was fairly constant on time-scales of millions of years. Solar activity varies on annual to centennial time-scales around a mean that remained roughly constant over the observed long-term (from thousands to millions of years) variations in cosmogenic production rate. Therefore, such long-term variations are most likely controlled by variations of the magnetic cutoff rigidity, directly related to the intensity of the magnetic field [8].

Since production rates of cosmogenic nuclides play a key role in their geophysical applications, mainly dating and tracing of environmental processes, the knowledge of past variations in cosmogenic nuclide production rates is a subject of considerable importance. Previous studies of the coincidence between low geomagnetic field intensity and high cosmogenic nuclide production rates were restricted to few geomagnetic events [9-12]. We present here, after the introduction of a paleomagnetic study [13], high resolution authigenic $^{10}\text{Be}/^9\text{Be}$ and continuous relative paleointensity records along the same marine sedimentary sequences from the Portuguese margin (N.E. Atlantic). Covering the past 300 ka, this work allows to assess relationships between geomagnetic moment and ^{10}Be production rate variations.

2. Description of sedimentary sequences and methodology

2.1. Sampling context

Authigenic ^{10}Be and ^9Be concentrations have been measured along three sedimentary sequences collected on the Portuguese Margin by the French vessel R. V. Marion Dufresne (Institut Français pour la Recherche et la Technologie Polaires) during IMAGES cruises using the Calypso piston corer: MD95-2042 (37°48N., 10°10W., 3146 m water depth); MD95-2040 (40°35N., 9°52W., 2465 m water depth) and MD01-2440G, a duplicate of core MD95-2042.

The top of cores MD95-2040 and -2042 are affected by coring deformation leading to unreliable paleomagnetic records [13]. Our sampling therefore only concerned the intervals

yielding reliable paleomagnetic data: 12-30 m of MD95-2040 and 9.5-28.5 m of MD95-2042. Few complementary samples were collected from the undisturbed core MD01-2440G, in order to document the youngest part of the record. Subsampling of ~0.5 g every 50 cm was reduced to every 10 cm at and close to the intervals of low relative paleointensity assignable to geomagnetic excursions.

2.2. Sediment description

The sampling area is under the influence of continental contributions arising mainly from both the Tagus and Douro rivers that flow over 800 km in the Sierra Albaraçin and over 600 km in the Sierra de la Virgen, respectively. These rivers drain sedimentary, plutonic and metamorphic formations. Deep currents and Mediterranean Outflow Waters (MOW) also contributed to the sedimentation.

The sediments mainly consist of mid-gray clayey-mud deposited during cold periods, with intercalation of beige carbonate ooze deposited during temperate periods. The clayey fraction (40 to 70%) and the “sortable silt” fraction vary in phase opposition; the later being more concentrated in carbonate layers, due to bottom current transport [14]. Carbonate contents range from 10 to 65% and from 14 to 40% in MD95-2040 [15] and in MD95-2042 [16], respectively. With a mean grain size of 7-8 μm , the sediment matrix contains two independent coarse fractions (63 μm – 1mm) : the carbonate biogenic fraction, and the lithogenic fraction composed of detrital carbonates, quartz, feldpars and magnetite grains, identified as ice rafted detritus linked to Heinrich events [17]. Like in most marine sediments, pyrite patches and concretions are frequent along the core.

2.3. Chronostratigraphy

Chronostratigraphic data are presented in details in Thouveny et al. [13] and Moreno et al. [18]. They are based on radiocarbon data and correlation of marine isotope stages (MIS) identified in the benthic $\delta^{18}\text{O}$ curves of cores MD95-2042 [19] and MD95-2039 [15] with the SPECMAP record [20]. Depth to time transform functions were constructed for each core using high degree polynomial best fit functions over contiguous windows, and used to transfer our data records on a time scale (Figure 1).

The studied cores cover the last climatic cycle in core MD95-2042, and the last three climatic cycles (i.e. since MIS9) in core MD95-2040. The time interval covered by our sampling thus ranges from 0 to 300 ka BP and concerns the succession of 3 glacial-interglacial cycles.

2.4. Chemical procedures

^{10}Be concentrations measured in marine sediments depend on ^{10}Be production rates as well as on environmental parameters (chemical and granulometric composition) [21-24]. We thus extracted the authigenic (i.e. adsorbed onto particles from the water column) $^{10}\text{Be}/^9\text{Be}$ from the studied sediments, since only the soluble form of both beryllium isotopes have been homogenized in the water column before deposition in the sediment [21, 23].

Authigenic beryllium was extracted using a 0.04M $\text{NH}_2\text{OH}\cdot\text{HCl}$ in 25% acetic acid leaching solution at $95^\circ\text{C} \pm 5$ during 7 hours from ~ 0.5 g samples previously crushed and oven-dried. The method is similar to that fully described by Bourlès et al. [21]. An aliquot of each sample solution was taken for furnace atomic absorption determination of ^9Be using the method of standard additions and a Zeeman effect background correction (Hitachi Z-8200). Uncertainties in ^9Be concentrations are based on the reproducibility of measurements when more than one has been made, or estimated from the fit of the standard addition lines in the case of a single measurement. The remainder of the sample was spiked for isotope dilution with 0.3 mg of ^9Be and then purified for ^{10}Be measurements by a series (usually two) of solvent extractions of Be acetylacetonate in the presence of EDTA, followed by precipitation of $\text{Be}(\text{OH})_2$ at pH 8.5. The beryllium precipitate, rinsed twice with deionized water, was finally dissolved in a few drops of HNO_3 . This small amount of solution being dried in a quartz crucible, BeO was at the end obtained by heating at 1000°C .

$^{10}\text{Be}/^9\text{Be}$ ratio measurements performed at the Tandetron AMS facility in Gif-sur-Yvette [25, 26] were calibrated directly against $^{10}\text{Be}/^9\text{Be}$ of the National Institute of Standards and Technology (NIST) standard reference material SRM 4325. The ^{10}Be uncertainties have been estimated using a conservative 5% instrumental uncertainty together with one standard deviation statistics of the number of ^{10}Be events counted (less than $\pm 3\%$).

3. Beryllium isotope results

3.1. Authigenic ^9Be

The ^9Be concentrations measured along MD95-2042 (Figure 2a), ranging from 3.1 to $8.8 \cdot 10^{-7}$ g/g, are systematically higher than those measured along MD95-2040 that range from 1.4 to $5.9 \cdot 10^{-7}$ g/g (Figure 3a). The lowest (resp. highest) ^9Be concentrations measured in these two cores

yield a ratio of ~2.1 (resp. ~1.5), that may most likely be entirely attributed to the difference in sedimentation rates between these two cores, since the sedimentation rate of MD95-2042 (~13 cm/kyr) is 1.7-1.9 times higher than that of MD95-2040 (~7.5 cm/ka). Linked to the continental origin of ^9Be , its higher concentrations occur in intervals of higher continental input, i.e. higher sedimentation rate. This direct linkage is confirmed by the observed increases in ^9Be concentration (Figure 2a and 3a) generally coinciding with high values of magnetic susceptibility.

3.2. Authigenic ^{10}Be

The ^{10}Be concentrations measured along MD95-2042 and MD95-2040 are by contrast bounded within similar range, that is between 5.1 to 10.8 10^{-15} g/g (mean value : ~7.3 10^{-15} g/g) in MD95-2042 (Figure 2b) and between 3.5 and 11.1 10^{-15} g/g (mean value : ~7.35 10^{-15} g/g) in MD95-2040 (Figure 3b).

Compared to the mean values, both ^{10}Be records show significant increases (above mean value $+1\sigma$) of the ^{10}Be concentration at the following depth intervals : 1325-1475 cm, 1800-1850 cm, 2000-2105 cm, 2245-2450 cm, 2480-2500 cm, 2590-2605 cm in MD95-2042, and 1340-1513 cm, 2094-2148 cm in MD95-2040. Compared to the base levels defined by the lowest ^{10}Be concentrations (~5 10^{-15} g/g for MD 95-2042 and ~4 10^{-15} g/g for MD95-2040), sharp increases of the ^{10}Be concentration appear at the following depth intervals : 1750-1950 cm, 2560-2645 cm in MD95-2042, and 2244-2305 cm, 2375-2450 cm, 2502-2600 cm and 2800-3006 cm in MD95-2040.

As stated in section 2.4, in order to interpret the previously discussed ^{10}Be concentration variations in terms of production rate variations, the environmental disturbing effects have to be taken in account by i) normalizing the authigenic ^{10}Be concentrations to the authigenic ^9Be concentrations and ii) computing ^{10}Be fluxes that are not affected by ^9Be input variations mainly controlled by the eolian flux to the ocean [27, 28].

3.3. Authigenic $^{10}\text{Be}/^9\text{Be}$ ratio

Since authigenic ^{10}Be concentrations measured along the two cores are within the same range while authigenic ^9Be concentrations along MD95-2042 are ~1.8 higher than along MD95-2040, the authigenic $^{10}\text{Be}/^9\text{Be}$ ratios measured along MD95-2042 (0.7 to 2.4 10^{-8}) (Figure 2c), are roughly a factor of two lower than those measured along MD95-2040 (1.17 to 5.2 10^{-8}) (Figure 3c).

Along MD95-2042, increases of at least a factor of two relative to the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio base line (1.1 10^{-8}) are evidenced at the following depth intervals : 1405-1425 cm (Interval I),

2265-2330 cm (Interval II) and 2420-2440 cm and 2490-2530 cm (Interval III). In addition, minor but significant increases are observed at the following depths : 1850 cm, 2105 cm and 2210 cm.

Along MD95-2040, numerous major significant increases corresponding at least to a doubling of the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio base line ($\sim 1.5 \cdot 10^{-8}$) are also clearly evidenced. Because, based on susceptibility records, the top of core MD95-2040 overlaps with the bottom of core MD95-2042, depth intervals from 1372 to 1414 cm and from 1473 to 1513 cm in MD95-2040 correspond to previously noticed intervals II and III. Additional major increases appear at depth intervals from 2094 to 2118 cm (interval IV) and from 2216 to 2244 cm (interval V). Less constrained but major significant increases are recorded from 2375 to 2424 cm (interval VI), and at ~ 2550 cm (interval VII). Finally, the last major increase is observed between 2891 and 2931 cm (interval VIII). Like in core MD95-2042, minor but significant increases are also observed at 1255-1274 cm and 1996-2024 cm.

3.4. ^{10}Be flux

Another way of reconstructing ^{10}Be production rate variations is to compute ^{10}Be flux variations along the sedimentary cores. ^{10}Be concentrations in marine sediments being inversely proportional to the carbonate content [21, 29, 30], ^{10}Be flux values were calculated using the following equation: ^{10}Be flux = ^{10}Be concentration * bulk density * sedimentation rate * (100/100-CaCO₃%). As shown in Figures 2c, d and 3c, d, the ^{10}Be flux significantly increases coincidentally with authigenic $^{10}\text{Be}/^9\text{Be}$ ratios. This confirms that the normalization of authigenic ^{10}Be to authigenic ^9Be takes into account -as expected- all biases that may result from differences in sources and pathways of both isotopes, and therefore implies that the observed increases should not result from environmental effects. However, given uncertainties about density and sedimentation rates estimation, flux determinations are less accurate than authigenic $^{10}\text{Be}/^9\text{Be}$ ratios.

Since a recently published study [29] of the influence of particle composition on sorption of ^{230}Th and ^{10}Be from sea water shows that the $^{10}\text{Be}/^{230}\text{Th}$ ratio depends on the silica/ carbonate ratio, the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio presently remains the only reliable proxy for reconstructing cosmonuclide production rates.

3.5. Relation with $\delta^{18}\text{O}$ variations ?

Authigenic $^{10}\text{Be}/^9\text{Be}$ ratios generally present significant increases at levels corresponding to interglacial conditions, as documented by $\delta^{18}\text{O}$ records measured on the same cores [15, 19] (Figures 2e and 3e). Such a correlation was previously reported [30-32] and attributed to a

dependence of ^{10}Be fluxes to glacial /interglacial alternation. However, close examination of Figures 2 and 3 reveals that : i) interval IV and VII, fall at the transition between interglacial MIS7 and glacial MIS6 and within glacial MIS 8, respectively ; ii) as evidenced by intervals I, II and VIII, falling in MIS 3, MIS 5.3, and MIS 8.2, respectively, the amplitudes of authigenic $^{10}\text{Be}/^9\text{Be}$ ratio variations are not comparable with the amplitudes of $\delta^{18}\text{O}$ signals. These observations lead to the conclusion that climatic variability is probably not the main forcing parameter.

4. Relations with relative paleointensity variations

As previously stated, variations of the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio thus most likely reflect variations of the cosmonuclide production rate induced by variations of the geomagnetic moment. In this section, we recall major observations and conclusions drawn from the paleomagnetic study, and introduce the comparison with the beryllium isotopes record.

In the paleomagnetic stack [13], large amplitude deviations of the inclination occur during periods of low relative paleointensity (RPI) : some are excursions, while the others remain within the limits of paleosecular variation and may represent excursions smoothed out by post-depositional processes.

Significant phases of low RPI, defined in Figure 4 by values significantly lower than the average value minus 1σ , are observed at successive depths in the two studied cores (Figures 4b and 4d). These are related to the authigenic $^{10}\text{Be}/^9\text{Be}$ significant increases as indicated in the following table.

Low RPI Interval	Depth (cm) (in corresponding core)	Corresponding high authigenic $^{10}\text{Be}/^9\text{Be}$ interval
A	1415 – 1530 (MD95-2042)	I
B (B1 and B2)	2284 - 2613 (MD95-2042) 1391 – 1522 (MD95-2040)	II and III
C	2100 – 2169	IV
D	2230 - 2296	V
E	2384 - 2408	VI
F	2539 – 2558	VII
G	2867 – 2982	VIII

A careful analysis of the relative depth position of RPI lows and authigenic $^{10}\text{Be}/^9\text{Be}$ peaks (Figure 4) reveals either that they occur at the same depth, or that RPI minima are significantly shifted downwards relatively to authigenic $^{10}\text{Be}/^9\text{Be}$ maxima. Namely, the mid-point of the

minimum RPI of interval A leads by ~30 cm the authigenic $^{10}\text{Be}/^9\text{Be}$ maximum of interval I ; the mid-point of interval B1 leads the mid-point of interval II by ~16 cm in core MD95-2042 ; the mid-point of interval B2 leads by ~26 cm the mid-point of interval III in core MD95-2040. The minimum RPI of interval C leads by ~24 cm the authigenic $^{10}\text{Be}/^9\text{Be}$ maximum of interval IV.

Given the sedimentation rates of these sequences, beryllium isotopes residence time in the water column in continental margin environments on the order of 200 to 300 years [33] may induce maximum shifts of 6 cm for interval A in core MD95-2042 and 2 cm for other intervals. Therefore, lags on the order of 14 to 24 cm remain to be explained by other effects.

Former studies suggested that post-depositional processes not only impose a smoothing of the geomagnetic input signal [34 - 36] but also lead to a delayed lock-in of the magnetization beneath a critical thickness which depends on grain size distribution, clay concentration, organic matter concentration and bioturbation. More recent studies [e.g. 37] tend to consider that such effects are negligible in quiet and undisturbed sedimentation environments. However, the lags evidenced by our results confirm that lock-in depth are not negligible, and agree with lock-in depth of ~25 cm revealed in a core of the Blake Outer Ridge [38].

5. Discussion

Measurements of the relative paleointensity and authigenic $^{10}\text{Be}/^9\text{Be}$ ratios along the same cores offer the unique opportunity to construct stacked records of both measured parameters. This allows to directly compare the proxy record of the cosmonuclide production rates to the proxy record of the geomagnetic forcing. The composite record of authigenic $^{10}\text{Be}/^9\text{Be}$ ratio has been reconstructed using all data measured along MD95-2042 and MD95-2040; 8 values from another short core MD01-2440G complete the record in its upper part (electronic annex Figure). In order to account for sedimentation rate differences between MD95-2042 and MD95-2040, authigenic $^{10}\text{Be}/^9\text{Be}$ ratio were readjusted by dividing the MD95-2040 data serie by a constant factor 1.8 (cf section 3.1.). Then, wheighted averages have been computed within 1ka time windows.

Using these reconstructions we first discuss the relationship between the two proxies. As shown in Figure 5, the averaged $^{10}\text{Be}/^9\text{Be}$ ratios plotted against averaged paleointensity values yields a simple exponential best-fit relationship, analogous to the Elsasser algorithm [2]:

$$P/P_0 = \sqrt{(M_0/M)}.$$

When compared to the global stack formerly established for the interval 0-200 ka BP by Frank et al. [39] (Figure 6), both records show similar trends. However, our authigenic $^{10}\text{Be}/^9\text{Be}$ variations have significantly stronger amplitudes than the stacked ^{10}Be deposition rates, equivalent

to a ^{10}Be flux, presented in [39] as the ^{10}Be production rate. Namely, the equivalent of Interval I has a restricted amplitude ($\sim 60\%$), between 30 and 42 ka BP, instead of a more significant spike ($\sim 100\%$) centered at 41 ka BP in our record. Equivalent of Interval II and III appears as a unique massive phase of relatively enhanced ^{10}Be flux ($\sim 30\%$) between 85 and 110 ka BP instead of two separate phases of enhancement ($\sim 140\%$ and $\sim 120\%$) lasting from 90 to 105 ka BP and from 112 to 120 ka BP, respectively. The equivalent of Interval IV is expressed by a $\sim 70\%$ increase of ^{10}Be flux, instead of a $\sim 130\%$ increase in our record, centered at 185 ka BP in both cases.

A major difference between the two records is the occurrence of a $\sim 40\%$ increase located between 60 and 75 ka BP in the global stack expressed in our record by a non-significant peak at 75 ka BP.

These discrepancies likely result from the fact that the global curve was constructed by stacking mostly data sets from central areas of oceanic basins, i.e. relatively low sedimentation rate areas, while our records result from a high resolution sampling focused on depth intervals recording low RPI features, in high sedimentation rate sequences.

A record of the relative virtual dipole moment (VDM) has been derived from the cosmonuclide record using the algorithm of Elsasser et al. [2] with a P_0 value = 1. As introduced in section 4 and evidenced by the comparison of Figures 7a, b and c, the authigenic $^{10}\text{Be}/^9\text{Be}$ oscillations are translated in terms of relative VDM variations tightly connected to relative paleointensity variations.

The interval 0-25 ka BP, documented in core MD01-2440G, exhibits high relative VDM values for the Holocene period, reaching a maximum at 20 ka BP in coincidence with high RPI values (17-22 ka BP). Although the relative VDM trend for this time interval reasonably agrees with the RPI records, one single relative VDM at 20 ka BP reaches a value contrasting with absolute values obtained from volcanic rocks (e.g. Teanby et al. [40]).

Beyond 20 ka BP, the evolution of relative VDM is coherent with trends revealed by RPI sedimentary records (Figures 7c and d) and with available volcanic VDM data [42]. Low VDM phases derived from the authigenic $^{10}\text{Be}/^9\text{Be}$ record (Figures 7a, b) are confronted with those reported from the RPI record (Figures 7c, d) and with paleomagnetic excursions documented in the studied time interval [13].

The minimum VDM at 41 ka BP of interval I well corresponds to interval A identified as the Laschamps excursion recorded in lavas flows from France [43] and Iceland [44], as well as in lacustrine [45] and marine sediments [46].

Intervals II (90-105 ka BP) and III (110-120 ka BP) confirm the long period of low RPI with two minima centered at ~95 ka (B1) and ~117 ka BP (B2) also documented in the Sint-record. While interval B2 can easily be ascribed to the well-known Blake event, interval B1 may be related to an excursionsal feature previously described as a post-Blake excursion [36]. The Blake event, first reported in sediments [47, 48], was described in details by Tric et al. [49] as a full reversal, but also as two successive anomalous inclination features in the soil of the last interglacial of a loess sequence of the China plateau and accurately dated at 114 and 120 ka BP by thermoluminescence [50].

Interval IV centered at 190 ka BP coincides with the low RPI interval C (~190 ka BP), which is accompanied by an excursion ascribed to the “Icelandic basin event” at ODP sites 983 and 984 [51] and particularly well expressed in the Sint-record.

Intervals V (205-208 ka BP), VI (~230 ka BP) and VII (250 ka BP), corresponding respectively to intervals D, E and F, can be ascribed to the Jamaica and/or Pringle falls [52, 53], the Mamaku excursion [54], and the Calabrian ridge excursion [53], respectively.

Interval VIII centered at 290 ka BP, corresponds to the low RPI phase of interval G (285-295 ka BP) which correlate to a low RPI phase of small amplitude lasting from 265 to 295 ka BP in the Sint-curve. The excursionsal directions linked to interval G were named “Portuguese margin excursion”, because, the Levantine excursion, formerly dated at ~290 ka BP, has been reassessed to an excursion occurring in older sediments in the reference location [53].

Although the extremely low RPI phases associated to excursions are well constrained along the Portuguese sequence, the VDM record derived from authigenic $^{10}\text{Be}/^9\text{Be}$ ratios, in contrast to the relative equilibrium of “oscillations” along the RPI stacks, appears to be dominated by long lasting phases of medium or low VDM (25-125 ka BP, 170-240 ka BP, 275-300 ka BP). This may be an artefact of the used simple algorithm while translating extreme authigenic $^{10}\text{Be}/^9\text{Be}$ ratios into VDM values, which emphasizes the need to perform a calibration based on the assignment of authigenic $^{10}\text{Be}/^9\text{Be}$ ratios to absolute VDM values along this time interval (Carcaillet et al. in prep.).

Spectral analyses of the 300 ka authigenic $^{10}\text{Be}/^9\text{Be}$ composite record (Figure 7a) using the Maximum Entropy and Blackman-Tukey methods [Analyseries program of Paillard et al. [55]] (Figure 8) evidence one single significant periodicity at 100 ka that also dominates paleointensity and inclination variations [13]. Major authigenic $^{10}\text{Be}/^9\text{Be}$ increases of the studied time serie are thus most likely essentially related to overproductions of cosmogenic ^{10}Be induced by geomagnetic moment drops associated with excursions or events.

Due to the lack of lock-in delays in the authigenic $^{10}\text{Be}/^9\text{Be}$ record, the apparent coincidence of most authigenic $^{10}\text{Be}/^9\text{Be}$ increases with interglacial (section 3.5) can be critically examined. We note that for 7 out of 8 cases, the exception being interval VII, cosmonuclide overproductions occur during or at the end of interglacial periods, supporting the hypothesis of a preferential occurrence of geomagnetic excursions during interglacials [51] or during interglacial/glacial transitions.

6. Conclusions

The records along the same sedimentary sequences of relative virtual dipole moment variations, authigenic ^{10}Be and ^9Be concentrations allow to demonstrate that over the last 300 ka, the geomagnetic moment variations exert the main control on ^{10}Be production variations. Phases of overproduction of ^{10}Be confirm the global occurrence of well-recognized and well-dated phases of low geomagnetic moment associated to the Laschamps excursion, the Blake event and the Jamaica/Pringle Falls excursion. They strengthen the validity of the recently identified Icelandic basin and Calabrian Ridge 0 excursions, and distinguish the Mamaku excursion (~240 ka BP), formerly assigned to the Jamaica/Pringle falls excursion. Moreover, they confirm two new excursions evidenced by [13] : the Post-Blake excursion (~95 ka BP) and the Portuguese margin excursion (290 ka BP).

Because they are not affected by post-depositional lock-in depths of ~15-24 cm, the authigenic $^{10}\text{Be}/^9\text{Be}$ results allow an accurate determination of ages of VDM minimas associated with excursions, which strongly suggest that most of them occurred during, or at the end of, interglacials or interstadials.

Acknowledgements

Cores were collected by the technical crew of the R.V. Marion Dufresne under the direction of Y. Balut. IMAGES campaign were funded by "I.F.T.R.P." (presently I.P.R.E.V. for "Institut de Recherche Paul-Emile Victor"). We thank the chiefs Scientist of MD IMAGES. We thank Nick J. Shackelton and J. Schönfeld for providing $\delta^{18}\text{O}$ data of cores MD95-2042 and MD95-2040. We thank E. Bard for usefull suggestions and discussions, R. Braucher for assistance in ^{10}Be measurements and M. Frank for providing the $^{10}\text{Be}/^{230}\text{Th}$ stack data set . This study was partly funded by the INSU program "Intérieur de la Terre" to N.T. and D. B. The UMS Tandetron is supported by the CNRS, CEA and IN2P3. We thank Mélanie Héroguel for her help in the English writing. This manuscript benefited from valuable comments of D. Lal and L. Mc Hargue.

References

- [1] J.C. Gosse and F.M. Phillips, Terrestrial in situ cosmogenic nuclides : theory and application, *Quat. Sci. Rev.* 20 (2001) 1475-1560.
- [2] W. Elsasser, E.P. Ney and J.R. Winckler, Cosmic-ray intensity and geomagnetism, *Nature* 178 (1956) 1226-1227.
- [3] D. Lal, Expected secular variations in the global terrestrial production rate of radiocarbon, in : E. Bard and W.S. Broecker (eds.), *The last Deglaciation : Absolute and Radiocarbon Chronologies, Series I : Global Environmental Change 2*, (1992) pp. 114-126.
- [4] J. Masarik and J. Beer, Simulation of particule fluxes and cosmogenic nuclide production in the Earth's atmosphere, *J. Geophys. Res.* 104 (D10) (1999) 12099-12111.
- [5] S. Vogt, G.F. Herzog and R.C. Reedy, Cosmogenic nuclides in extraterrestrial materials, *Rev. Geophys.* 28 (1990) 253-275.
- [6] S. Tanaka and T. Inoue, ^{10}Be dating of North Pacific sediment cores up to 2.5 million years B.P., *Earth Planet. Sci. Lett.* 45 (1979) 181-187.
- [7] T.L. Ku, M. Kusakabe, D.E. Nelson, J.R. Southon, R.G. Korteling, J. Vogel and I. Nokikow, Constancy of oceanic deposition of ^{10}Be as recorded in manganese crusts, *Nature* 299 (1982) 240-242.
- [8] T.J. Dunai, Influence of the secular variation of the geomagnetic field on the production rates of in situ produced cosmogenic nuclides, *Earth Planet. Sci. Lett.* 193 (2001) 197-212.
- [9] D. Aylmer, M. Herzog, G.F. Guyton, B. Clement, J. Klein and R. Middleton, Beryllium-10 contents of sediments associated with Matuyama-Brunhes geomagnetic reversal., *EOS* 69 (16) (1988) 341.
- [10] G. Cini Castagnoli, A. Albrecht, J. Beer, G. Bonino, C. Shen, E. Callegari, C. Taricco, B. Dittrich-Hannen, P. Kubik, M. Suter and G.M. Zhu, Evidence for enhanced ^{10}Be deposition in Mediterranean sediments 35 Kyr BP, *Geophys. Res. Lett.* 22 (6) (1995) 707-710.
- [11] W.U. Henken-Mellies, J. Beer, F. Heller, K.J. Hsü, C. Shen, G. Bonani, H.J. Hofmann, M. Suter and W. Wölfli, ^{10}Be and ^9Be in South Atlantic DSPD Site 519 : Relation to geomagnetic reversals and to sediment composition, *Earth Planet. Sci. Lett.* 98 (1990) 267-276.
- [12] G.M. Raisbeck, F. Yiou, D. Bourlès and D.V. Kent, Evidence for an increase in cosmogenic ^{10}Be during a geomagnetic reversal, *Nature* 315 (1985) 315-317.

- [13] N. Thouveny, J. Carcaillet, E. Moreno, G. Leduc and D. Nérini, A 400 ka record of geomagnetic paleosecular variation and excursions from sedimentary cores of the Portuguese Margin, *Earth Planet. Sci. Lett.* This issue.
- [14] I.R. Hall and I.N. Mc Cave, Palaeocurrent reconstruction, sediment and thorium focusing on the Iberian margin over the last 140 ka, *Earth Planet. Sci. Lett.* 178 (2000) 151-164.
- [15] J. Thomson, S. Nixon, C.P. Summerhayes, J. Schönfeld, R. Zahn and P. Grootes, Implications for sedimentation changes on the Iberian margin over the last two glacial/interglacial transitions from $(^{230}\text{Th}_{\text{excess}})_o$ systematics, *Earth Planet. Sci. Lett.* 165 (1999) 255-270.
- [16] D. Pailler and E. Bard, High frequency paleoceanographic changes during the past 140.000 years recorded by the organic matter in sediments of the Iberian Margin, *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 181 (2002) 431-452.
- [17] N. Thouveny, E. Moreno, D. Delanghe, L. Candon, Y. Lancelot and N.J. Shackleton, Rock magnetic detection of distal ice-rafted debris : clue for the identification of Heinrich layers on the Portuguese margin, *Earth Planet. Sci. Lett.* 180 (2000) 61-75.
- [18] E. Moreno, N. Thouveny, D. Delanghe, I.N. Mc Cave and N.J. Shackleton, Climatic and oceanographic changes in the Northeast Atlantic reflected by magnetic properties of sediments deposited on the Portuguese Margin during the last 340 ka, *Earth Planet. Sci. Lett.* 202 (2002) 465-480.
- [19] N.J. Shackleton, M. Hall and E. Vincent, Phase relationship between millennial-scales events 64.000-24.000 years ago, *Paleoceanography* 15 (6) (2000) 565-569.
- [20] D.G. Martinson, N. Pisias, J.D. Hays, J. Imbrie, T.C. Moore and N.J. Shackleton, Age dating and the orbital theory of the ice ages : development of a high-resolution 0 to 300.000-years chronostratigraphy, *Quat. Res.* 27 (1987) 1-29.
- [21] D. Bourlès, G.M. Raisbeck and F. Yiou, ^{10}Be and ^9Be in marine sediments and their potential for dating, *Geochim. Cosmochim. Acta* 53 (1989) 443-452.
- [22] R.F. Anderson, Y. Lao, W.S. Broecker, S.E. Trumbore, H.J. Hofmann and W. Wolfli, Boundary scavenging in the Pacific Ocean : a comparison of ^{10}Be and ^{231}Pa , *Earth Planet. Sci. Lett.* 96 (1990) 287-304.
- [23] C. Robinson, G.M. Raisbeck, F. Yiou, B. Lehman and C. Laj, The relationship between ^{10}Be and geomagnetic field strength records in central North Atlantic sediments during the last 80 ka, *Earth Planet. Sci. Lett.* 136 (1995) 551-557.
- [24] M. Frank, Comparison of cosmogenic radionuclide production and geomagnetic field intensity over the last 200 000 years, *Phil. Trans. R. Soc. Lond.* 358 (A) (2000) 1089-1107.

- [25] G.M. Raisbeck, F. Yiou, D.L. Bourles, J. Lestringuez and D. Deboffle, Measurements of ^{10}Be and ^{26}Al with a Tandetron AMS facility., Nucl. Instr. and Meth. B29 (1987) 22-26.
- [26] G.M. Raisbeck, F. Yiou, D. Bourles, E. Brown, D. Deboffle, P. Jouhannau, J. Lestringuez and Z.Q. Zhou, The AMS facility at Gif-sur-Yvette : progress, perturbations and projects., Nucl. Instr. and Meth. B92 (1994) 43-46.
- [27] E.T. Brown, C.I. Measures, J.M. Edmond, D.L. Bourlès, G.M. Raisbeck and F. Yiou, Continental input of beryllium to the oceans. Earth Planet. Sci. Lett. 114 (1992) 101-111.
- [28] W. Dong, D. Lal, B. Ransom, W. Berger and M.W. Caffee, Marine biogeochemistries of Be and Al: A study based on cosmogenic ^{10}Be , Be and Al in marine calcite, aragonite, and opal. Proc. Indian Acad. Sci. (Earth Planet. Sci.) 110 (2001) 95-102.
- [29] Z. Chase, R.F. Anderson, M.Q. Fleisher and P.W. Kubik, The influence of particle composition and particle flux on scavenging of Th, Pa and Be in the ocean, Earth Planet. Sci. Lett. 204 (2002) 215-229.
- [30] J.R. Southon, T.L. Ku, D.E. Nelson, J.L. Reyss, J.C. Duplessy and J.S. Vogel, ^{10}Be in a deep-sea core : implications regarding ^{10}Be production changes over the past 420 ka, Earth Planet. Sci. Lett. 85 (1987) 356-364.
- [31] A. Eisenhauer, R.F. Spielhagen, M. Frank, G. Hentschel, A. Mangini, P.W. Kubik, B. Dittrich-Hannen and T. Billen, ^{10}Be records of sediments cores from high northern latitudes : Implications for environmental and climatic changes, Earth Planet. Sci. Lett. 124 (1994) 171-184.
- [32] A.A. Aldahan, S. Ning, G. Possnert, J. Backman and B. K., ^{10}Be records from sediments of the Arctic Ocean covering the past 350 ka, Mar. Geol. 144 (1-3) (1997) 147-162.
- [33] Y. Lao, R.F. Anderson, W.S. Broecker, S.E. Trumbore, H.J. Hofmann and W. Wolfli, Transport and burial rates of ^{10}Be and ^{231}Pa in the Pacific Ocean during the Holocene period, Earth Planet. Sci. Lett. 113 (1992) 173-189.
- [34] M. Hyodo, Possibility of reconstruction of the past geomagnetic field from homogeneous sediments, J. Geomagn. Geoelectr. 36 (1984) 45-62.
- [35] P. Vlag, N. Thouveny and P. Rochette, Synthetic and sedimentary records of geomagnetic excursions, Geophys. Res. Lett. 24 (6) (1997) 723-726.
- [36] N. Thouveny, K.M. Creer, I. Blunk, Extension of the Lac du Bouchet paleomagnetic record over the last 120, 000 years. Earth Planet. Sci. Lett. 97 (1990) 140-161.
- [37] L. Tauxe, Sedimentary records of relative paleointensity of the geomagnetic field : theory and practice, Rev. Geophys. 31 (3) (1993) 319-354.

- [38] L.R. Mc Hargue, D. Donahue, P.E. Damon, C.P. Sonett, D. Biddulph and G. Burr, Geomagnetic modulation of the late Pleistocene cosmic-ray flux as determined by ^{10}Be from Blake Outer Ridge marine sediments, *Nucl. Instr. and Meth.* 172 (B) (2000) 555-561.
- [39] M. Frank, B. Schwarz, S. Baumann, P.W. Kubik, M. Suter and A. Mangini, A 200 kyr record of cosmogenic radionuclide production rate and geomagnetic field intensity from ^{10}Be in globally stacked deep-sea sediments, *Earth Planet. Sci. Lett.* 149 (1997) 121-129.
- [40] N. Teanby, Laj, C., Gubbins, D., Pringles, M., A detailed palaeointensity and inclination record from drill core SOH1 on Hawaii, *Phys. Earth Planet. Inter.* 131 (2002) 101-140.
- [41] Y. Guyodo and J.P. Valet, Global changes in intensity of the Earth's magnetic field during the past 800 kyr, *Nature* 399 (1999) 249-252.
- [42] M. Perrin and V. Shcherbakov, Paleointensity database updated, *Eos Trans. AGU.* 79 (1998) 198.
- [43] N. Bonhommet and J. Babkine, Sur la présence d'aimantations inversées dans la Chaîne des Puys, *C. R. Acad. Sci. Paris* 264 (1967) 92-94.
- [44] S. Levi, H. Audunsson, R.A. Duncan, L. Kristjansson, P.-Y. Gillot and S. Jakobsson, Late Pleistocene geomagnetic excursion in Icelandic lavas : confirmation of the Late Laschamp excursion, *Earth Planet. Sci. Lett.* 96 (1990) 443-457.
- [45] P. Vlag, N. Thouveny, D. Williamson, P. Rochette and F. Ben-Atig, Evidence for a geomagnetic excursion recorded in the sediments of lac St. Front, France : A link with the Laschamp excursion? *J. Geophys. Res.* 101 (B12) (1996) 28211-28230.
- [46] C. Laj, C. Kissel, A. Mazaud, J.E.T. Channell and J. Beer, North Atlantic paleointensity stack since 75 ka (NAPIS-75) and the duration of the Laschamp event, *Phil. Trans. R. Soc. Lond. A* 358 (2000) 1009-1025.
- [47] J.D. Smith and J.H. Foster, Geomagnetic reversal in Brunhes normal polarity epoch, *Science* 163 (1969) 565.
- [48] W.B.F. Ryan, Stratigraphy of the late Quaternary sediments in the Eastern Mediterranean, in : J. Stanley (ed.), *The mediterranean Sea : a natural sedimentation laboratory* (1972) pp. 149-169.
- [49] E. Tric, C. Laj, J. Valet, P. Tucholka, M. Paterne and F. Guichard, The Blake geomagnetic event : transition geometry, dynamical characteristics and geomagnetic significance., *Earth Planet. Sci. Lett.* 102 (1991) 1-13.
- [50] X.M. Fang, J.J. Li, R. Van der Voo, C. Mac Niocail, X.R. Dai, R.A. Kemp, E. Derbyshire, J.X. Cao, J.M. Wang and G. Wang, A record of the Blake Event during the last interglacial paleosol in the western Loess Plateau of China, *Earth Planet. Sci. Lett.* 146 (1997) 73-82.

- [51] J.E.T. Channell, Geomagnetic paleointensity and directional secular variation at Ocean Drilling Program (ODP) Site 984 (Bjorn Drift) since 500 ka : Comparisons with ODP Site 983 (Gardar Drift), *J. Geophys. Res.* 104 (B10) (1999) 22937-22951.
- [52] E. Herrero-Bervera, C.E. Helsley, A.M. Sarma-Wojcicki, K.R. Lajoie, C.E. Meyer, M.O. Mc Williams, R.M. Negrini, B.D. Turin, J.M. Donnelly-Nolan and J.C. Liddicoat, Age and correlation of a paleomagnetic episode in the western united states by Ar-40/Ar-39 dating and tephrochronology : The Jamaica, Blake or a new polarity episode?, *J. Geophys. Res.* 99 (1994) 24091-24103.
- [53] C.G. Langereis, M.J. Dekkers, G.J. De Lange, M. Paterne and P.J.M. Van Santvoort, Magnetostratigraphy and astronomical calibration of the last 1.1 Myr from an eastern Mediterranean piston core and dating of short events in the Brunhes, *Geophys. J. Int* 129 (1997) 75-94.
- [54] P. Shane, T. Black and J. Westgate, Isothermal plateau fission-track age for a paleomagnetic excursion in the Mamaku ignimbrite, New Zealand and implication for late Quaternary stratigraphy, *Geophys. Res. Lett.* 21 (16) (1994) 1695-1698.
- [55] D. Paillard, L. Labeyrie and P. Yiou, Macintosh program performs time-series analysis, *EOS Transcr. AGU* 77 (1996) 379.

Figure 1: Chronological reconstruction of the sequences. Ages listed in [13] are plotted along the depth of each core (a: MD95-2042; b: MD95-2039; c: MD95-2040): calibrated ^{14}C ages (dots), Heinrich events 3 and 4 (H), and dated paleoclimatic markers (circles). Best-fit polynomial curves were computed in successive windows to obtain age/depth function used to transfer the data from each core depth scale to the time scale. The grey vertical band locate the transition between the anomalous (left) and normal (right) sedimentary fabrics.

Figure 2 : Core MD95-2042. (a) ^9Be concentrations (10^{-7} g/g) ; errors express reproducibility or statistical error of the fit of the standard addition lines (see text). (b) ^{10}Be concentrations (10^{-15} g/g) ; errors include counting statistics and instrumental uncertainty (see text). (c) $^{10}\text{Be}/^9\text{Be}$ (10^{-8}) ratios ; errors result from the propagation of the ^9Be and ^{10}Be concentration errors ; bold line corresponds to the mean value, thin line to mean value+ 1σ , grey bars underline $^{10}\text{Be}/^9\text{Be}$ values higher than mean value+ 1σ . (d) ^{10}Be carbonate-free flux (10^{10} at/cm²/ka). Note the flux scale change at 1900 cm. (e) Benthic $\delta^{18}\text{O}$ (‰) [19].

Figure 3 : Core MD95-2040. (a) ^9Be concentrations (10^{-7} g/g) ; errors express reproducibility or statistical error of the fit of the standard addition lines (see text). (b) ^{10}Be concentrations (10^{-15} g/g) ; errors include counting statistics and instrumental uncertainty (see text). (c) $^{10}\text{Be}/^9\text{Be}$ (10^{-8}) ratios ; errors result from the propagation of the ^9Be and ^{10}Be concentration errors ; bold line corresponds to the mean value, thin line to mean value+ 1σ , grey bars underline $^{10}\text{Be}/^9\text{Be}$ values higher than mean value+ 1σ . (d) ^{10}Be carbonate-free flux (10^{10} at/cm²/ka). (e) Benthic $\delta^{18}\text{O}$ (‰) [15].

Figure 4 : Comparison of authigenic $^{10}\text{Be}/^9\text{Be}$ and relative paleointensity records. (a) MD95-2042 $^{10}\text{Be}/^9\text{Be}$ ratios (10^{-8}). (b) MD95-2042 relative paleointensity, bold line corresponds the mean value, thin line to mean value- 1σ , grey bars underline relative paleointensities lower than mean- 1σ . (c) MD95-2040 $^{10}\text{Be}/^9\text{Be}$ ratios (10^{-8}). (d) MD95-2040 relative paleointensities : open circles correspond to discrete specimens, open triangles to U-channels. Bold line is the mean value, thin line the mean values- 1σ , grey bars underline relative paleointensities values lower than mean- 1σ . Vertical lines connecting $^{10}\text{Be}/^9\text{Be}$ peaks to relative paleointensity features point out several shifts due to lock-in depths of the remanence.

Figure 5 : Authigenic $^{10}\text{Be}/^9\text{Be}$ ratios as a function of relative paleointensities. The black curve represents the best exponential fit of the data. The grey curve represents the $^{10}\text{Be}/^9\text{Be}$ values

derived from the Elsasser algorithm, with P_0 equal to the $^{10}\text{Be}/^9\text{Be}$ ratios average value and M_0 equal to the relative paleointensities average value.

Figure 6 : Comparison of the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio record from the Portuguese margin stack with the global $^{10}\text{Be}/^{230}\text{Th}$ stack of Frank et al. [39].

Figure 7 : Comparison of the authigenic $^{10}\text{Be}/^9\text{Be}$ stack and the dipole moments reconstructed from authigenic $^{10}\text{Be}/^9\text{Be}$ with relative paleointensity and VDM curves. (a) Relative Authigenic $^{10}\text{Be}/^9\text{Be}$ ratio stack ; (b) M/M_0 reconstructed from authigenic $^{10}\text{Be}/^9\text{Be}$ ratios ; errors are propagated from $^{10}\text{Be}/^9\text{Be}$ errors (see text) (c) Portuguese paleointensity stack [13]. New excursions are distinguished from well-known by bold italic characters (d). Sint-800 VDM stack [41]

Figure 8 : Spectral analysis of the authigenic $^{10}\text{Be}/^9\text{Be}$ record using analysis series [56] (a) Maximum Entropy Method and (b) Blackman Tukey Method. Periods are expressed in ka.

Electronic annex

Figure EA : Results from core MD01-2440G. (a) ^9Be concentrations (10^{-7} g/g) ; errors express reproducibility or statistical error of the fit of the standard addition lines (see text). (b) ^{10}Be concentrations (10^{-15} g/g) ; errors include counting statistics and instrumental uncertainty (see text). (c) Authigenic $^{10}\text{Be}/^9\text{Be}$ (10^{-8}) ratios.

Comment : While ^9Be data of the 23-30 ka interval are comparable to those of the upper part of core MD95-2042, a doubling of the ^9Be concentrations occurs at 20 ka BP with a maximum value at 17 ka BP. Similarly, ^{10}Be concentrations are comparable to those obtained at the upper part of MD95-2042, but with a decreasing trend to minimum values at 20 and 17 ka BP. Therefore, authigenic $^{10}\text{Be}/^9\text{Be}$ ratios of this latter interval are reaching anomalously low values. At least for the 17 ka BP data, this can be explained by the accidental sampling of a layer of Ice rafted debris linked to Heinrich event n°1 ; indeed it appears that this sample corresponds to a high susceptibility layer.