



HAL
open science

Multiple Qubits as Symplectic Polar Spaces of Order Two

Metod Saniga, Michel R. P. Planat

► **To cite this version:**

Metod Saniga, Michel R. P. Planat. Multiple Qubits as Symplectic Polar Spaces of Order Two. Advanced Studies in Theoretical Physics, 2007, 1, pp.1 - 4. hal-00121565

HAL Id: hal-00121565

<https://hal.science/hal-00121565>

Submitted on 21 Dec 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multiple Qubits as Symplectic Polar Spaces of Order Two

Metod Saniga¹ and Michel Planat²

¹Astronomical Institute, Slovak Academy of Sciences
SK-05960 Tatranská Lomnica, Slovak Republic
(msaniga@astro.sk)

and

²Institut FEMTO-ST, CNRS, Département LPMO, 32 Avenue de l'Observatoire
F-25044 Besançon Cedex, France
(michel.planat@femto-st.fr)

Abstract

It is surmised that the algebra of the Pauli operators on the Hilbert space of N -qubits is embodied in the geometry of the symplectic polar space of rank N and order two, $W_{2N-1}(2)$. The operators (discarding the identity) answer to the points of $W_{2N-1}(2)$, their partitionings into maximally commuting subsets correspond to spreads of the space, a maximally commuting subset has its representative in a maximal totally isotropic subspace of $W_{2N-1}(2)$ and, finally, “commuting” translates into “collinear” (or “perpendicular”).

MSC Codes: 51Exx, 81R99

PACS Numbers: 02.10.Ox, 02.40.Dr, 03.65.Ca

Keywords: Symplectic Polar Spaces of Order Two – N-Qubits

It is well known that a complete basis of operators in the Hilbert space of N -qubits, $N \geq 2$, can be given in terms of the Pauli operators — tensor products of classical 2×2 Pauli matrices. Although the Hilbert space in question is 2^N -dimensional, the operators' space is of dimension 4^N . Excluding the identity matrix, the set of $4^N - 1$ Pauli operators can be partitioned into $2^N + 1$ subsets, each comprising $2^N - 1$ mutually commuting elements [1]. The purpose of this note is to put together several important facts supporting the view that this operators' space can be identified with $W_{2N-1}(q = 2)$, the symplectic polar space of rank N and order two.

A (finite-dimensional) classical polar space (see [2–6] for more details) describes the geometry of a d -dimensional vector space over the Galois field $GF(q)$, $V(d, q)$, carrying a non-degenerate reflexive sesquilinear form σ . The polar space is called symplectic, and usually denoted as $W_{d-1}(q)$, if this form is bilinear and alternating, i.e., if $\sigma(x, x) = 0$ for all $x \in V(d, q)$; such a space exists only if $d = 2N$, where N is called its rank. A subspace of $V(d, q)$ is called totally isotropic if σ vanishes identically on it. $W_{2N-1}(q)$ can then be regarded as the space of totally isotropic subspaces of $PG(2N - 1, q)$, the ordinary $(2N - 1)$ -dimensional projective space over $GF(q)$, with respect to a symplectic form (also known as a null polarity), with its maximal totally isotropic subspaces, also called generators G , having dimension $N - 1$. For $q = 2$ this polar space contains

$$|W_{2N-1}(2)| = |PG(2N - 1, 2)| = 2^{2N} - 1 = 4^N - 1 \quad (1)$$

points and

$$|\Sigma(W_{2N-1}(2))| = (2 + 1)(2^2 + 1) \dots (2^N + 1) \quad (2)$$

generators [2–4]. An important object associated with any polar space is its *spread*, i.e., a set of generators partitioning its points. A spread S of $W_{2N-1}(q)$ is an $(N - 1)$ -spread of its ambient projective space $PG(2N - 1, q)$ [4, 5, 7], i.e., a set of $(N - 1)$ -dimensional subspaces of $PG(2N - 1, q)$ partitioning its points. The cardinalities of a spread and a generator of $W_{2N-1}(2)$ thus read

$$|S| = 2^N + 1 \quad (3)$$

and

$$|G| = 2^N - 1, \tag{4}$$

respectively [2, 3]. Finally, it needs to be mentioned that two distinct points of $W_{2N-1}(q)$ are called perpendicular if they are “isotropically” collinear, i. e., joined by a totally isotropic line of $W_{2N-1}(q)$; for $q = 2$ there are

$$\#\Delta = 2^{2N-1} \tag{5}$$

points that are *not* perpendicular to a given point of $W_{2N-1}(2)$ [2, 3].

Now, in light of Eq. (1), we can identify the Pauli operators with the points of $W_{2N-1}(2)$. If, further, we identify the operational concept “commuting” with the geometrical one “perpendicular,” from Eqs. (3) and (4) we readily see that the points lying on generators of $W_{2N-1}(2)$ correspond to maximally commuting subsets (MCSs) of operators and a spread of $W_{2N-1}(2)$ is nothing but a partitioning of the whole set of operators into MCSs. From Eq. (2) we then infer that the operators’ space possesses $(2+1)(2^2+1)\dots(2^N+1)$ MCSs and, finally, Eq. (5) tells us that there are 2^{2N-1} operators that do *not* commute with a given operator; the last two statements are, for $N > 2$, still conjectures to be rigorously proven. However, the case of two-qubits ($N = 2$) is recovered in full generality [1, 8, 9], with the geometry behind being that of the *generalized quadrangle of order two* [9] — the simplest nontrivial symplectic polar space.¹

Acknowledgements

This work was partially supported by the Science and Technology Assistance Agency under the contract # APVT-51-012704, the VEGA project # 2/6070/26 (both from Slovak Republic) and the trans-national ECO-NET project # 12651NJ “Geometries Over Finite Rings and the Properties of Mutually Unbiased Bases” (France).

References

- [1] Lawrence, J., Brukner, Č., and Zeilinger, A., “Mutually unbiased binary observable sets on N qubits,” *Physical Review A* 65, 032320 (2002).
- [2] Ball, S., “The geometry of finite fields,” *Quaderni Elettronici del Seminario di Geometria Combinatoria 2E* (2001); available on-line from <http://www.mat.uniroma1.it/~combinat/quaderni/>.
- [3] Cameron, P. J., “Projective and polar spaces,” available on-line from <http://www.maths.qmw.ac.uk/~pjc/pps/>.
- [4] De Clerck, F., and Van Maldeghem, H., “Ovoids and spreads of polar spaces and generalized polygons,” a lecture given at an intensive course on “Galois Geometry and Generalized Polygons,” University of Ghent, April 14–25, 1998; available from <http://cage.rug.ac.be/~fdc/intensivecourse/ovspr.ps>.
- [5] Hirschfeld, J. W. P., and Thas, J. A., “General Galois Geometries” (Oxford University Press, Oxford, 1991).
- [6] Payne, S. E., and Thas, J. A., “Finite Generalized Quadrangles” (Pitman, London, 1984).
- [7] Thas, J. A., “Ovoids and spreads in classical polar spaces,” *Geom. Dedicata* 10, 135–144 (1981).
- [8] Wootters, W. K., “Picturing qubits in phase space,” *IBM J. Res. Dev.* 48, 99–110 (2004).
- [9] Saniga, M., Planat, M., and Pracna, P., “Projective ring line encompassing two-qubits,” *Theoretical and Mathematical Physics*, submitted; quant-ph/0611063-v4.

¹This object can also be recognized as the projective line over the Jordan system of the full 2×2 matrix ring with coefficients in $GF(2)$ [9].