



# A Remark on Almost Umbilical Hypersurfaces

Julien Roth

► **To cite this version:**

Julien Roth. A Remark on Almost Umbilical Hypersurfaces. *Archivum Mathematicum*, Masarykova  
Universita, 2013, 49 (1), pp.1-7. hal-00182518v2

**HAL Id: hal-00182518**

**<https://hal.archives-ouvertes.fr/hal-00182518v2>**

Submitted on 9 Oct 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A Remark on Almost Umbilical Hypersurfaces

Julien Roth

October 9, 2012

Université Paris-Est, LAMA (UMR 8050), UPEMLV, UPEC, CNRS,  
F-77454, Marne-la-Vallée, France

julien.roth@univ-mlv.fr

## Abstract

In this article, we prove new stability results for almost-Einstein hypersurfaces of the Euclidean space, based on previous eigenvalue pinching results. Then, we deduce some comparable results for almost umbilical hypersurfaces.

**Keywords:** Hypersurfaces, Rigidity, Pinching, Ricci Curvature, Umbilicity Tensor, Higher Order Mean Curvatures.

**Mathematical Subject Classification:** 53A07, 53A10, 53C20, 53C24.

## 1 Introduction

It is a well-known fact that a totally umbilical hypersurface of the Euclidean space  $\mathbb{R}^{n+1}$  which is not totally geodesic is a round sphere. An other classical rigidity result states that an Einstein (with positive scalar curvature) hypersurface of the Euclidean space  $\mathbb{R}^{n+1}$  is a round sphere. This was proved by Thomas [11] and independently by Fialkow [3] in the 30's. Recently, Grosjean [4] gave a new proof based on the equality case of an estimate of the first eigenvalue of the Laplacian involving the scalar curvature.

It is then natural to consider the case of almost umbilical and almost Einstein hypersurfaces and ask if they are close to round spheres (in a sense to be precised). Shiohama and Xu [9, 10] proved that under a condition on Betti numbers, almost umbilical hypersurfaces of Euclidean space are homeomorphic to the sphere. For almost Einstein hypersurfaces, Vlachos obtained a comparable result in [12].

By studying the stability problem of Grosjean's proof, we showed in [8] another promity result for almost Einstein, namely these hypersurfaces are

diffeomorphic and quasi-isometric to round sphere. We got stronger proximity but in counterpart, with stronger assumption that in the result of Vlachos.

The aim of the present paper is to relax the assumptions of this result in one hand (Theorem 1) and on the other hand to get stability results for almost umbilical hypersurfaces to be compared with the results of Shiohama and Xu (see Theorems 2 and 3).

## 2 Preliminaries

Let  $(M^n, g)$  be a  $n$ -dimensional compact, connected, oriented Riemannian manifold without boundary, isometrically immersed into the  $(n+1)$ -dimensional Euclidean space  $(\mathbb{R}^{n+1}, can)$ . The second fundamental form  $B$  of the immersion is the bilinear symmetric form defined by

$$B(Y, Z) = -g(\bar{\nabla}_Y \nu, Z),$$

where  $\bar{\nabla}$  is the Riemannian connection on  $\mathbb{R}^{n+1}$  and  $\nu$  the outward normal unit vector field on  $M$ .

From  $B$ , we can define the mean curvature,

$$H = \frac{1}{n} \text{tr}(B),$$

and, more generally, the higher order mean curvatures,

$$H_r = \frac{1}{\binom{n}{r}} \sigma_r(\kappa_1, \dots, \kappa_n),$$

where  $\sigma_r$  is the  $r$ -th symmetric polynomial and  $\kappa_1, \dots, \kappa_n$  are the principal curvatures of the immersion. By convention, we set  $H_0 = 1$ .

Note that  $H_1 = H$  and from the Gauss equation,  $H_2$  is, up to a multiplicative constant, the scalar curvature. Namely, we have  $H_2 = \frac{1}{n(n-1)} \text{Scal}$ .

These extrinsic curvatures satisfy the well-known Hsiung-Minkowski formula, for  $1 \leq r \leq n$ ,

$$(1) \quad \int_M (H_{r-1} + H_r \langle X, \nu \rangle) dv_g = 0,$$

where  $X$  is the position vector and  $\nu$  the normal vector of the immersion. They also satisfy the following inequalities if  $H_r$  is a positive function:

$$(2) \quad H_r^{\frac{1}{r}} \leq H_{r-1}^{\frac{1}{r-1}} \leq \dots \leq H_2^{\frac{1}{2}} \leq H.$$

Moreover, Reilly [7] proved some upper bounds for the first eigenvalue of the Laplacian for hypersurfaces of  $\mathbb{R}^{n+1}$  in terms of higher order mean

curvatures. Precisely, he showed

$$(3) \quad \lambda_1(M) \left( \int_M H_{r-1} dv_g \right)^2 \leq \frac{n}{\text{Vol}(M)} \int_M H_r^2 dv_g,$$

with equality if and only if  $M$  is a geodesic hyperspheres of  $\mathbb{R}^{n+1}$ .

By the Hölder inequality, we obtain for  $p \geq 2$ ,

$$\lambda_1(M) \leq n \frac{\|H_r\|_{2p}^2 \text{Vol}(M)^2}{\left( \int_M H_{r-1} dv_g \right)^2}.$$

Now, for  $p \geq 2$  and  $1 \leq r \leq n$ , we define  $k_{p,r} = \frac{\|H_r\|_{2p}^2 \text{Vol}(M)^2}{\left( \int_M H_{r-1} dv_g \right)^2}$ , which are the constants involved in Theorem 1.

Note that we use the following convention, for any smooth function  $f$  and any  $p \geq 1$ ,

$$\|f\|_p = \frac{1}{\text{Vol}(M)^{\frac{1}{p}}} \left( \int_M |f|^p dv_g \right)^{\frac{1}{p}}.$$

The main tool that we will use in the present paper is the following pinching result, associated with these inequalities, that we proved in [8].

**Theorem A** (Roth [8]). *Let  $(M^n, g)$  be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in  $\mathbb{R}^{n+1}$ ,  $n \geq 2$ . Let  $r \in \{1, \dots, n\}$  and assume that  $H_r > 0$  if  $r > 1$ . Then for any  $q \geq 2$  and any  $\theta \in ]0, 1[$ , there exists a constant  $K_\theta$  depending only on  $n$ ,  $\|H\|_\infty$ ,  $\text{Vol}(M)$ ,  $\|H_r\|_{2q}$  and  $\theta$  such that if the pinching condition*

$$(P_{K_\theta}) \quad 0 \geq \lambda_1(M) \left( \int_M H_{r-1} dv_g \right)^2 - n \text{Vol}(M)^2 \|H_r\|_{2q}^2 > -K_\theta$$

*is satisfied, then  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n \left( \sqrt{\frac{n}{\lambda_1}} \right)$ .*

**Remark 1.** *Note that if  $q \geq \frac{n}{2r}$ ; then  $K_\theta$  does not depend on  $\|H_r\|_{2q}$ .*

**Remark 2.** *The case  $r = 1$  was proved by Colbois and Grosjean [2]. In that case we do not need to assume that  $H > 0$ .*

### 3 Almost Einstein hypersurfaces

We showed in [8] that almost-Einstein hypersurfaces of  $\mathbb{R}^{n+1}$  are close to round spheres. Namely,

**Theorem B** (Roth [8]). *Let  $(M^n, g)$  be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in  $\mathbb{R}^{n+1}$ ,  $n \geq 2$ . Let  $\theta \in ]0, 1[$ . If  $(M^n, g)$  is almost-Einstein, that is,  $\|\text{Ric} - (n-1)kg\|_\infty \leq \varepsilon$  for a positive constant  $k$ , with  $\varepsilon$  small enough depending on  $n$ ,  $k$ ,  $\|H\|_\infty$  and  $\theta$ , then  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n \left( \sqrt{\frac{1}{k}} \right)$*

By  $\theta$ -quasi-isometric, we understand that there exists a diffeomorphism  $F$  from  $M$  into  $\mathbb{S}^n\left(\sqrt{\frac{1}{k}}\right)$  such that, for any  $x \in M$  and for any unitary vector  $u \in T_x M$ , we have

$$\left| |d_x F(u)|^2 - 1 \right| \leq \theta.$$

This theorem is a corollary of our pinching result for the first eigenvalue of the Laplacian (Theorem A).

In this article, we consider almost-Einstein hypersurfaces of  $\mathbb{R}^{n+1}$  in a weaker sense, namely for the  $L^q$ -norm, that is,  $\|\text{Ric} - (n-1)kg\|_q \leq \varepsilon$  for some positive constant  $k$  and a sufficiently small  $\varepsilon$ . We prove that for some suitable constants  $k$ , such manifolds are close to round spheres. Precisely, we prove the following

**Theorem 1.** *Let  $(M^n, g)$  be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in  $\mathbb{R}^{n+1}$ . Let  $q > \frac{n}{2}$ ,  $r \in \{1, \dots, n\}$  and if  $r > 1$ , assume that  $H_r > 0$ . Let  $\theta \in ]0, 1[$ , if  $(M^n, g)$  satisfies  $\|\text{Ric} - (n-1)k_{p,r}g\|_q \leq \varepsilon$  for some sufficiently small  $\varepsilon$  depending on  $n, q, \|H\|_\infty, \text{Vol}(M)$  and  $\theta$ , then  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n\left(\sqrt{\frac{1}{k_{p,r}}}\right)$ .*

The constants  $k_{p,r}$  in the theorem are defined from the higher order mean curvature (see Sect. 2).

After giving the proof of this theorem, we will give some applications to almost-umbilical hypersurfaces.

*Proof:* The proof is based on the above pinching result combined with a lower bound for the first eigenvalue of the Laplacian due to Aubry [1]. Assume that  $\|\text{Ric} - (n-1)kg\|_q \leq \varepsilon(n, q, k)$  for a positive constant  $k$ ,  $q > \frac{n}{2}$  and  $\varepsilon$  small enough, then from Theorem 1.1 of [1], we deduce that  $\lambda_1(\Delta)$  satisfies

$$(4) \quad \lambda_1(\Delta) \geq nk(1 - C_\varepsilon),$$

where  $C_\varepsilon$  is an explicit constant such that  $C_\varepsilon \rightarrow 0$  when  $\varepsilon \rightarrow 0$ .

Now, with the particular choice of  $k = k_{p,r}$ , we get:

$$\lambda_1(M) \left( \int_M H_{r-1} dv_g \right)^2 - n \text{Vol}(M)^2 \|H_r\|_{2p}^2 > -K_\varepsilon$$

for some constant  $K_\varepsilon$  such that  $K_\varepsilon \rightarrow 0$  when  $\varepsilon \rightarrow 0$ .

Let  $\theta \in ]0, 1[$ , we choose  $\varepsilon(n, q, k, \theta)$  small enough such that  $K_\varepsilon$  is small enough in Theorem A to obtain a diffeomorphism and  $\theta$ -quasi-isometry between  $M$  and  $\mathbb{S}^n\left(\sqrt{\frac{1}{k_{p,r}}}\right)$ .  $\square$

**Remark 3.** Note that in this Theorem,  $\varepsilon$  depends on  $\text{Vol}(M)$  contrary to Theorem B. We can remove this dependence on the volume and replace it by a dependence on  $\|H_{r-1}\|_1$ . Indeed, using an upper bound on the volume under the  $L^p$  condition on Ricci (result by Aubry [1]) we have that the volume is bounded from above by a constant depending on  $n, q, \|H\|_\infty$  and  $k_{r,p}$  that is in fact on  $n, q, \|H\|_\infty$  and  $\|H_{r-1}\|_1$  because of the definition of  $k_{r,p}$  and (2). On the other hand, by the classical extrinsic Sobolev inequality of Hoffman and Spruck [6], we can bound the volume from below by a constant depending only on  $n$  and  $\|H\|_\infty$ .

Now, we will deduce from Theorem 1 some Corollaries for almost-umbilical hypersurfaces of  $\mathbb{R}^{n+1}$ .

## 4 Almost umbilical hypersurfaces

First, we give the following theorem, which is a direct application of Theorem B.

**Theorem 2.** Let  $(M^n, g)$  be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in  $\mathbb{R}^{n+1}$ . Let  $\theta \in ]0, 1[$ . If  $(M^n, g)$  is almost-umbilical, that is,  $\|B - kg\|_\infty \leq \varepsilon$  for a positive constant  $k$ , with  $\varepsilon$  small enough depending on  $n, k$  and  $\theta$  then  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n(\frac{1}{k})$ .

*Proof :* From the Gauss formula, we have

$$(5) \quad \text{Ric}(Y, Y) = nH \langle B(Y), Y \rangle - \langle B(Y), B(Y) \rangle,$$

for a tangent vector field  $Y$ . From (5) and  $\|B - kg\|_\infty \leq \varepsilon$ , we deduce

$$\begin{aligned} \text{Ric}(Y, Y) &\geq nk^2\|Y\|^2(1 - \varepsilon)^2 - k^2\|Y\|^2(1 + \varepsilon)^2 \\ &\geq (n - 1)k^2\|Y\|^2 - \alpha_n(\varepsilon)\|Y\|^2, \end{aligned}$$

where  $\alpha_n$  is a positive function such that  $\alpha_n(\varepsilon) \rightarrow 0$  when  $\varepsilon \rightarrow 0$ . Similarly, we get

$$\text{Ric}(Y, Y) \leq (n - 1)k^2\|Y\|^2 + \alpha_n(\varepsilon)\|Y\|^2.$$

Finally, we have

$$\|\text{Ric} - (n - 1)k^2g\|_\infty \leq \alpha_n(\varepsilon),$$

which implies, by Theorem B, that for  $\varepsilon$  small enough,  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n(\frac{1}{k})$ .  $\square$

Now, from Theorem 1, it is possible to obtain, in some particular cases,

comparable results for almost umbilical hypersurfaces in an  $L^q$ -sense. We recall that the umbilicity tensor is defined by

$$\tau = B - H\text{Id}.$$

As we mentioned above, if  $M$  is umbilical, *i.e.*,  $\tau = 0$ , and if  $M$  is compact, then  $M$  is a geodesic sphere. Here, we prove the following stability result for almost umbilical hypersurfaces.

**Theorem 3.** *Let  $(M^n, g)$  be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in  $\mathbb{R}^{n+1}$ . Let  $q > \frac{n}{2}$ ,  $r \in \{1, \dots, n\}$  and if  $r > 1$ , assume that  $H_r > 0$ . For any  $\theta \in ]0, 1[$ , there exists two constants  $\varepsilon_i(\theta, n, \|H\|_\infty, \text{Vol}(M))$ ,  $i = 1, 2$ , such that if*

1.  $\|\tau\|_{2q} \leq \varepsilon_1$ ,
2.  $\|H^2 - k_{p,r}\|_q \leq \varepsilon_2$ , for  $p \geq 4$  and  $1 \leq r \leq n$ ,

*then  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n \left( \frac{1}{\sqrt{k_{p,r}}} \right)$*

**Remark 4.** *Note that for  $r = 1$ , the result is due to Grosjean-Roth (see [5]), using a pinching result which involves only  $H_1$  (see [2]).*

*Proof.* Here again, from the Gauss formula, we have

$$\text{Ric} = nHB - B^2.$$

From this, we deduce that

$$\begin{aligned} \text{Ric} - (n-1)H^2g &= nHB - B^2 - (n-1)H^2g \\ &= (n-2)H\tau - \tau^2, \end{aligned}$$

which implies

$$\begin{aligned} \|\text{Ric} - (n-1)k_g\|_q &\leq \|\text{Ric} - (n-1)H^2g\|_q + (n-1)\sqrt{n}\|(H^2 - k)\|_q \\ &\leq (n-2)\|H\|_\infty\|\tau\|_{2q} + \|\tau\|_{2q}^2 + (n-1)\sqrt{n}\|(H^2 - k)\|_q \\ &\leq (n-2)\|H\|_\infty\varepsilon_1 + \varepsilon_1^2 + (n-1)\sqrt{n}\varepsilon_2 \end{aligned}$$

Now, we conclude by taking  $\varepsilon_1$  and  $\varepsilon_2$  small enough depending on  $n$ ,  $\|H\|_\infty$  and  $\theta$  in order to apply Theorem 1 and obtain the  $\theta$ -quasi-isometry.  $\square$

Then, we deduce the following corollary which is to compare with Theorem 2.

**Corollary 1.** *Let  $(M^n, g)$  be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in  $\mathbb{R}^{n+1}$ . Let  $\theta \in ]0, 1[$ . If  $(M^n, g)$  is almost-umbilic, that is,  $\|B - \sqrt{k_{p,r}}g\|_{2q} \leq \varepsilon$ , for  $q > \frac{n}{2}$ , with  $\varepsilon$  small enough depending on  $n$ ,  $\|H\|_\infty$  and  $\theta$  then  $M$  is diffeomorphic and  $\theta$ -quasi-isometric to  $\mathbb{S}^n \left( \frac{1}{\sqrt{k_{p,r}}} \right)$ .*

*Proof :* A simple computation shows that

$$\|H^2 - k_{p,r}\|_{2q} \leq \alpha_1 \|B - \sqrt{k_{p,r}}g\|_{2q}, \text{ and}$$

$$\|\tau\|_{2q} \leq \alpha_2 \|B - \sqrt{k_{p,r}}g\|_{2q},$$

for two constants  $\alpha_1$  and  $\alpha_2$  depending on  $n$  and  $\|H\|_\infty$ . Since we assume that  $\|B - \sqrt{k_{p,r}}g\|_{2q} \leq \varepsilon$ , we get

1.  $\|H^2 - k_{p,r}\|_{2q} \leq \alpha_1 \varepsilon$ ,
2.  $\|\tau\|_{2q} \leq \alpha_2 \varepsilon$ .

For  $\varepsilon$  small enough, the assumptions of Theorem 3 are satisfied and we can conclude that  $M$  is diffeomorphic and quasi-isometric to  $\mathbb{S}^n \left( \frac{1}{\sqrt{k_{p,r}}} \right)$ .  $\square$

**Remark 5.** *We want to point out that this corollary is an improvement of Theorem 2 only in some sense. Indeed, we improve the  $L^\infty$ -proximity to an  $L^{2q}$ -proximity, but this corollary is valid only for some special constants  $k_{p,r}$  and not for any positive constant as in Theorem 2.*

## References

- [1] E. Aubry, *Finiteness of  $\pi_1$  and geometric inequalities in almost positive Ricci curvature*, Ann. Sci. Ecole Norm. Sup. (4) **40** (2007), no 4, 675-695.
- [2] B. Colbois and J.F. Grosjean, *A pinching theorem for the first eigenvalue of the Laplacian on hypersurfaces of the Euclidean space*, Comment. Math. Helv. **82** (2007), 175-195.
- [3] A. Fialkow, *Hypersurfaces of a space of constant curvature*, Ann. of Math. **39** (1938), 762-783.
- [4] J.F. Grosjean, *Upper bounds for the first eigenvalue of the Laplacian on compact manifolds*, Pac. J. Math. **206** (2002), no 1, 93-111.
- [5] J.F. Grosjean and J. Roth, *Eigenvalue pinching and application to the stability and the almost umbilicity of hypersurfaces*, Math. Z. **271** (2012), no 1, 469-488.



- [6] D. Hoffman and D. Spruck, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, Comm. Pure Appl. Math. **27** (1974), 715-727
- [7] R.C. Reilly, *On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space*, Comment. Math. Helv. **52** (1977), 525-533.
- [8] J. Roth, *Pinching of the first eigenvalue of the Laplacian and almost-Einstein hypersurfaces of Euclidean space*, Ann. Glob. Anal. Geom. **33** (2008), no 3, 293-306.
- [9] K. Shiohama and H. Xu *Rigidity and sphere theorems for submanifolds*, Kyushu J. Math. **48**, (1994), no 2, 291-306.
- [10] K. Shiohama and H. Xu *Rigidity and sphere theorems for submanifolds II*, Kyushu J. Math. **54**, (2000), no 1, 103-109.
- [11] T.Y. Thomas, *On closed space of constant mean curvature*, Amer. J. Math. **58** (1936), 702-704.
- [12] T. Vlachos, *Almost-einstein hypersurfaces in the euclidean space*, Illinois J. Math. **53** (2009), no 4, 1221-1235.