Synthesis, Structural Characterization of the first N-heterocyclic carbene ligand fused to a porphyrin

Sébastien Richeter, Aurélie Hadj-Aïssa, Céline Taffin, Arie van Der Lee,
Dominique Leclercq

► To cite this version:

Sébastien Richeter, Aurélie Hadj-Aïssa, Céline Taffin, Arie van Der Lee, Dominique Leclercq. Synthesis, Structural Characterization of the first N-heterocyclic carbene ligand fused to a porphyrin. Chemical Communications, 2007, pp.2148-2150. 10.1039/b704681b . hal-00183367

HAL Id: hal-00183367

<https://hal.science/hal-00183367>

Submitted on 30 Oct 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis and Structural Characterization of the First N-Heterocyclic Carbene Fused to a Porphyrin

Sébastien Richeter^{*a}, Aurélie Hadj-Aïssa^a, Céline Taffin^a, Arie van der Lee^b and Dominique Leclercq^a

Receipt/Acceptance Data [DO NOT ALTER/DELETE THIS TEXT]

Publication data [DO NOT ALTER/DELETE THIS TEXT]

DOI: 10.1039/b000000x [DO NOT ALTER/DELETE THIS TEXT]

The functionalization of two neighboring β -pyrrolic positions of a porphyrin by a fused N-heterocyclic carbene, the subsequent metallation of this external coordination site by palladium(II) and the structural characterization of the compounds obtained are presented.

The synthesis of multiporphyrin systems is of great interest because of their wide potential applications in catalysis,¹ molecular materials² and medicine.³ A lot of synthetic strategies have been developed to build multiporphyrin systems through covalent⁴ and non-covalent bonds⁵. The use of coordination bonds to link porphyrins together opened exciting opportunities to build systems with different shapes, such as linear,⁶ cyclic,⁷ 2D⁸ and 3D⁹ geometries. The classical building blocks employed for this strategy are porphyrins bearing one or more peripheral coordination sites, as it is well illustrated by the examples of porphyrins linked to pyridyl derivatives.^{8,10}

Surprisingly, the number of examples of porphyrins bearing N-heterocyclic carbene (NHC) ligands as external coordination site remains limited¹¹ and, to our knowledge, there is no example of NHC conjugated with the aromatic core of the porphyrin. Stable N-heterocyclic carbenes (NHCs), first isolated by Arduengo *et al.*¹² are versatile ligands for transition metal complexes and remain very important in the field of organometallic chemistry

Fig. 1 Structures of the complexes 1-7.

^a Institut Charles Gerhardt, UM2, CNRS, équipe CMOS, CC 007 Université Montpellier 2, Place Eugène Bataillon, 34095 Montpellier Cedex 05, France. E-mail: Sebastien.Richeter@univ-montp2.fr

^b Institut Européen des Membranes, UM2, CNRS, ENSCM, CC 045, Université Montpellier 2, Place Eugène Bataillon, 34095 Montpellier Cedex 05, France.

† Electronic Supplementary Information (ESI) available: Synthesis and characterization of the compounds 1, 2, 4-6 and 10 and the cif file of the crystal structure of 10.

See <http://dx.doi.org/10.1039/b000000x/>

Conditions:

(a) LiNO₃, CHCl₃/AcOH/Ac₂O, 45°C, 2h.

(b) 4-amino-4H-1,2,4-triazole, NaOH, toluene/EtOH, reflux, 1h.

(c) (i) NaBH₄, Pd/C, CH₂Cl₂/MeOH, 25°C, 1h.; (ii) HCO₂H, toluene, 110°C, 10 mn.; (iii) TFA, toluene, 110°C, 3h.

(d) MeI, K₂CO₃, 25°C, 24h.

Overall yields are indicated.

Scheme 1 Synthesis of the imidazolium salt 2.

and homogeneous catalysis.¹³ Beyond catalysis, NHCs are emerging as relevant components in materials science.¹⁴ It is obvious that there is a considerable potential for NHCs containing functional or tunable groups in conjugation with the carbene moiety.¹⁵ Thus, we focused our attention in the design of the porphyrin 1 containing an imidazole ring fused across a β,β' -pyrrolic position and fully conjugated to the macrocycle. The formation of the imidazolium salt 2 gives a synthetic access to the porphyrin 3 fused to an NHC (Figure 1).

The peripheral functionalization of porphyrins is of great interest in order to access to new chromophores or to new multiporphyrin systems.¹⁶ Crossley and co. prepared fused porphyrin-imidazole systems by the condensation of porphyrin-2,3-diones with an arylaldehyde in the presence of ammonia.¹⁷ Following this procedure, the compounds obtained have a 2'-arylimidazole ring. Herein, we report another synthetic approach to access to the fused porphyrin-imidazole 1 with a non functionalized 2'-imidazole carbon (Scheme 1).

The synthetic porphyrin used as starting material is the complex 4.¹⁸ First, a nitration reaction was realized according to the literature procedure to yield the β -nitroporphyrin 5.¹⁹ The electron withdrawing group NO₂ switched the reactivity of the neighboring pyrrolic β -carbon to an electrophilic center. Callot and co. showed that the 4-amino-4H-1,2,4-triazole is a powerful amination reagent for α,β -unsaturated ketones and aldehydes conjugated with the porphyrin core.²⁰ This can also be applied to 5 and it was possible to get the compound 6 in good yield (81%). The β -NH₂ ¹H NMR signal of 6 could be detected as a broad singlet at δ = 6.55 ppm in CDCl₃. Its UV-visible absorption bands are bathochromically shifted compared to 5. The lowest energy absorption band of 6 shifts to 600 nm (587 nm for 5). The

Scheme 2

diaminoporphyrin **7** was obtained by the reduction of the NO₂ group of **6** with NaBH₄ in the presence of methanol and Pd/C.²¹ The porphyrin **7** was not isolated but used straightforward for the formation of the imidazole ring.

The cyclization reaction with formic acid was more challenging and realized in two steps compared to the classical procedure described for 2,3-diaminobenzene derivatives (one quantitative step).²² First, the alcohol **8** was obtained by the complete condensation reaction of **7** with formic acid in refluxing toluene (1:1 mixture) in less than 10 minutes. Then, two competitive reactions occur in these conditions: the formation of the porphyrin-imidazole **1** by the elimination of a molecule of water and the formation of the ester **9** (Scheme 2). To prevent the formation of the undesirable ester, it was preferable to change the reaction conditions after the formation of the alcohol **8**. The imidazole **1** was preferentially obtained by the slow addition of TFA to a refluxing solution of **8** in toluene. The overall yield from **6** to **1** was 48%. The ¹H NMR spectrum of **1** in CDCl₃ shows broad signals due to the slow imidazole tautomerism on the ¹H NMR timescale. The double N-alkylation of the imidazole ring of **1** with iodomethane yields the corresponding imidazolium salt **2** in 76% yield. A sharp singlet at δ = 11.50 ppm corresponding to the iminium C(2')-H is observed in the ¹H NMR spectrum (C₆D₆) of **2**. This downfield chemical shift shows the highly electron-withdrawing nature of the metalloporphyrin macrocycle.

Scheme 3 Synthesis of the complex **10**

Then, we investigated the possibility to generate the carbene fused to the porphyrin **3**. As a proof of its existence, we performed the direct metallation of **2** with the basic metal precursor Pd(OAc)₂ in refluxing toluene.²³ This reaction proceeded smoothly leading to the formation of the complex **10** which was purified by chromatography on silica gel (Scheme 3). No signal corresponding to the iminium proton was observed by ¹H NMR spectroscopy (CDCl₃) showing the formation of the NHC-Pd(II) bond. The evidence of the formation of the complex **10** was also provided by FAB⁺ mass spectrometry. The molecular pic corresponding to the dimer was observed at m/z = 2289. Two fragments were observed at m/z = 2162 and 1324 and correspond to the dimer whose one ligand was removed, respectively the iodine and the porphyrin-NHC. The signal observed at m/z = 963 corresponds to the cationic porphyrin **2**. The UV-visible

Fig. 2 Two views of the X-ray structure of the complex **10** (solvent molecules have been omitted for clarity). C in gray, N in blue, Ni in yellow, Pd in orange and I in purple. (a) *t*-butyl groups and H omitted for clarity. (b) *meso* aryl groups and H omitted for clarity.

absorption bands of **10** are slightly broadened and red-shifted compared to those of the imidazolium **2**. This result tends to show that there is negligible ground state electronic communication between the porphyrins through the palladium center.

The single-crystal X-ray analysis[†] unambiguously establish the structure of the complex **10** and the *trans* geometry around the palladium (II) (Figure 2). The two porphyrin moieties are not flat but are ruffled because the C(5)-Ni-C(15) and C(10)-Ni-C(20) angles are close to 165°. Despite this distorted geometry of the aromatic core of the porphyrin, the additional imidazoylidene ring and the adjacent pyrrole are coplanar. The nickel (II) was found in an slightly distorted square planar geometry: the four Ni-N distances are almost equivalent (1.900, 1.906, 1.924, and 1.960 Å) and the four N-Ni-N angles are close to 90° (88.45, 90.09, 91.16, and 91.29°). In this square-planar *trans* palladium (II) complex, the Pd-C distances which are consistent with Pd-C single bonds (2.037 Å) are shorter than the Pd-I distances (2.591 Å). The two imidazoylidene rings are close to coplanarity and are tilted ~82° from the square panar central plane PdI₂C₂. This tilted conformation minimizes the steric interactions between the methyl groups and the iodines linked to the palladium (II) but also the steric interactions between the *t*Bu groups close to the palladium center.

To conclude, we have presented here a new synthetic procedure to obtain a porphyrin with an additional imidazole ring fused across a β,β'-pyrrolic position. We have shown that it is possible to generate an NHC at the periphery of a porphyrin and to use it as a coordination site to built multiporphyrin assemblies. We are currently studying the tuning of the optical and redox properties of the NHC-porphyrin systems. The synthesis od bis(NHC)porphyrins are also under investigation.

Notes and references

[†] Crystallographic data were recorded with an Xcalibur CCD camera (Oxford Diffraction) using graphite monochromated Mo Kα radiation (λ = 0.71073 Å), 40 s per frame. The structure was solved using Patterson methods²⁴ and refined using full-matrix least squares methods²⁵.

- Hydrogen atoms (except one belonging to a chloroform molecule) were located from the Fourier difference map and not refined. Crystal data for **10**: $\text{C}_{130}\text{H}_{132}\text{Cl}_{12}\text{I}_2\text{N}_{12}\text{Ni}_2\text{Pd}$, $M_w = 2765.48$, monoclinic, space group $P 1 2_1/c 1$ (no. 14), $a = 22.1838(7) \text{ \AA}$, $b = 9.9174(4) \text{ \AA}$, $c = 31.4826(13) \text{ \AA}$, $\beta = 103.73(0)^\circ$, $V = 6728.43(50) \text{ \AA}^3$, $Z = 2$, $d_{\text{calc}} = 1.365 \text{ g/cm}^3$, $T = 173 \text{ K}$, $\lambda (\text{Mo K}\alpha) = 0.71073 \text{ \AA}$, $\mu (\text{Mo K}\alpha) = 0.128 \text{ mm}^{-1}$, 18567 reflns collected, 3351 unique reflns $I > 2\sigma(I)$, $R_F = 0.263$ and $wR_F = 0.0887$ for all reflections, $R_F = 0.0764$ and $wR_F = 0.0527$ for observed reflections.
- 1 D. Mansuy, *Coord. Chem. Rev.*, 1993, **125**, 129; L. M Slaughter, E. Galaron, P. Le Maux and G. Simonneaux, *Chem. Commun.*, 1997, 927; J. P. Collman, T. A. Eberspacher and J. I. Brauman, *Inorg. Chem.*, 2004, **43**, 5198; J. P. Collman, L. Zeng, H. J. H. Wang, A. Lei and J. I. Brauman, *Chem. Eur. J.*, 2006, **12**, 2707.
 - 2 M. G. H. Vicente, L. Jaquinod and K. M. Smith, *Chem. Commun.*, 1999, 1771; H. L. Anderson, *Chem. Commun.*, 1999, 2323.
 - 3 X. Chen, L. Hui, D. A. Foster and C. M. Drain, *Biochemistry*, 2004, **43**, 10918; C. E. Stilts, M. I. Nelen, D. G. Hilmey, S. R. Davies, S. O. Gollnick, A. R. Oseroff, S. L. Gibson, R. Hilf and M. R. Detty, *J. Med. Chem.*, 2000, **43**, 2403; H. Gu, K. Xu, Z. Yang, C. K. Chang and B. Xu, *Chem. Commun.*, 2005, 4270.
 - 4 A. K. Burrell, D. L. Officer, P. G. Plieger and D. C. W. Reid, *Chem. Rev.*, 2001, **101**, 2751.
 - 5 J. Wojaczynski and L. Latos-Grazynski, *Coord. Chem. Rev.*, 2000, **204**, 113.
 - 6 C. Ikeda, E. Fujiwara, A. Satake and Y. Kobuke, *Chem. Commun.*, 2003, 616; U. Michelsen and C. A. Hunter, *Angew. Chem., Int. Ed.*, 2000, **39**, 764.
 - 7 R. Takahashi and Y. Kobuke, *J. Am. Chem. Soc.*, 2003, **125**, 2372; R. A. Haycock, A. Hunter, D. A. James, U. Michelsen and L. R. Sutton, *Org. Lett.*, 2000, **2**, 2435.
 - 8 C. M. Drain, F. Nifiatis, A. Vasenko and J. D. Batteas, *Angew. Chem., Int. Ed.*, 1998, **37**, 2344.
 - 9 I. Goldberg, *Chem. Commun.*, 2005, 1243; I. Goldberg, *Chem. Eur. J.*, 2000, **6**, 3863; Y. Diskin-Posner, S. Dahal and I. Goldberg, *Angew. Chem., Int. Ed.*, 2000, **39**, 1288.
 - 10 A. Tsuda, T. Nakamura, S. Sakamoto, K. Yamaguchi and A. Osuka, *Angew. Chem., Int. Ed.*, 2002, **41**, 2817; N. Fujita, K. Biradha, M. Fujita, S. Sakamoto and K. Yamaguchi, *Angew. Chem., Int. Ed.*, 2001, **40**, 1718; A. Harriman, F. Odobel and J. -P. Sauvage, *J. Am. Chem. Soc.*, 1994, **116**, 5481.
 - 11 V. Seng, X. Wang, R. Ali, W. Fang, F. Charles-Pierre, S. Bhagan and A. G. Hyslop, *J. Heter. Chem.*, 2006, **43**, 1077.
 - 12 A. J. Arduengo III, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361; A. J. Arduengo III, H. V. Rasika Dias, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1992, **114**, 5530.
 - 13 W. A. Herrmann and C. Köcher, *Angew. Chem., Int. Ed.*, 1997, **36**, 2163; D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39; W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1290; M. C. Perry and K. Burgess, *Tetrahedron: Asymmetry*, 2003, **14**, 951; E. Peris and R. H. Crabtree, *Coord. Chem. Rev.*, 2004, **248**, 2239; K. J. Cavell and D. S. McGuinness, *Coord. Chem. Rev.*, 2004, **248**, 671; M. C. Crudden and D. P. Allen, *Coord. Chem. Rev.*, 2004, **248**, 2247; A. Zapf and M. Beller, *Chem. Commun.*, 2005, 910; N. M. Scott and S. P. Nolan, *Eur. J. Inorg. Chem.*, 2005, 1815.
 - 14 S. -W. Zhang, R. Ishii and S. Takahashi, *Organometallics*, 1997, **16**, 20; O. Guerret, S. Solé, H. Gornitzka, M. Teichert, G. Trinquier and G. Bertrand, *J. Am. Chem. Soc.*, 1997, **119**, 6668; C. K. Lee, J. C. C. Chen, K. M. Lee, C. W. Liu and I. J. B. Lin, *Chem. Mater.*, 1999, **11**, 1237; A. J. Boydston, K. A. Williams and C. W. Bielawski, *J. Am. Chem. Soc.*, 2005, **127**, 12496; A. J. Boydston and C. W. Bielawski, *Dalton Trans.*, 2006, 4073.
 - 15 M. D. Sanderson, J. W. Kamplain and C. W. Bielawski, *J. Am. Chem. Soc.*, 2006, **128**, 16514.
 - 16 L. Jaquinod, in *The Porphyrin Handbook*, ed. K. M. Kadish, K. M. Smith and R. Guilard, Academic Press, San Diego, 2000, vol. 6, pp 201–237; P. S. S. Lacerda, A. M. G. Silva, A. C. Tomé, M. G. P. M. S. Neves, A. M. S. Silva, J. A. S. Cavaleiro and A. L. Llamas-Saiz, *Angew. Chem., Int. Ed.*, 2006, **45**, 5487.
 - 17 M. J. Crossley, J. A. McDonald, *J. Chem. Soc., Perkin Trans. 1*, 1999, 2429.
 - 18 A. D. Adler, F. R. Longo, J. D. Finarelli, J. Goldmacher, J. Assour and L. Korsakoff, *J. Org. Chem.*, 1967, **32**, 476. The complex **4** was synthesized according to Adler's methodology. See Supporting Information.
 - 19 A. Giraudeau, H. J. Callot, J. Jordan, I. Ezahr and M. Gross, *J. Am. Chem. Soc.*, 1979, **101**, 3857; A. W. Johnson, D. J. Oldfield, *J. Chem. Soc.*, 1965, 4303.
 - 20 S. Richeter, C. Jeandon, R. Ruppert and H. J. Callot, *Chem. Commun.*, 2002, 266; S. Richeter, C. Jeandon, J. -P. Gisselbrecht, R. Ruppert and H. J. Callot, *J. Am. Chem. Soc.*, 2002, **124**, 6168; S. Richeter, C. Jeandon, J. -P. Gisselbrecht, R. Graff, R. Ruppert and H. J. Callot, *Inorg. Chem.*, 2004, **43**, 251.
 - 21 J. E. Baldwin, M. J. Crossley and J. DeBernardis, *Tetrahedron*, 1982, **38**, 685; V. Promarak and P. A. Burn, *J. Chem. Soc., Perkin Trans. 1*, 2001, 14.
 - 22 E. C. Wagner and W. H. Millett, *Org. Synth.*, 1939, **19**, 12.
 - 23 W. A. Herrmann, M. Ellison, J. Fischer, C. Kocher and G. R. J. Artus, *Angew. Chem., Int. Ed.* 1995, **34**, 2371.
 - 24 P. T. Beurskens, G. Beurskens, W. P. Bosman, R. de Gelder, S. Garcia Granda, R. O. Gould, R. Israel and J.M.M. Smits, in '*a Computer Program System for Crystallography Laboratory*,' Nijmegen, 1996.
 - 25 P. W. Betteridge, J. R. Carruthers, R. I. Cooper, K. Prout and D. J. Watkin, *J. Appl. Cryst.*, 2003, **36**, 1487.