



HAL
open science

An alternative expression for the Black-Scholes formula in terms of Brownian first and last passage times

D. Madan, Bernard Roynette, Marc Yor

► **To cite this version:**

D. Madan, Bernard Roynette, Marc Yor. An alternative expression for the Black-Scholes formula in terms of Brownian first and last passage times. 2008. hal-00257403

HAL Id: hal-00257403

<https://hal.science/hal-00257403>

Preprint submitted on 19 Feb 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An alternative expression for the Black-Scholes formula in terms of Brownian first and last passage times

D. Madan ⁽¹⁾, B. Roynette ⁽²⁾, M. Yor ⁽³⁾⁽⁴⁾

February 11, 2008

⁽¹⁾ Robert H. Smith School of Business, University of Maryland,
Van Munching Hall, College Park, MD 20742 - USA
E-mail:dmadan@rhsmith.umd.edu

⁽²⁾ Université Henri Poincaré, Institut Elie Cartan, BP239,
F-54506 Vandoeuvre-les-Nancy Cedex
E-mail:bernard.roynette@iecn.u-nancy.fr

⁽³⁾ Laboratoire de Probabilités et Modèles Aléatoires,
Universités Paris VI et VII, 4 Place Jussieu - Case 188,
F-75252 Paris Cedex 05
E-mail:deaproba@proba.jussieu.fr

⁽⁴⁾ Institut Universitaire de France

Abstract: The celebrated Black-Scholes formula which gives the price of a European option, may be expressed as the cumulative function of a last passage time of Brownian motion. A related result involving first passage times is also obtained.

Keywords: Black-Scholes formula, last passage times, time-inversion, Cameron-Martin relationship, reflection principle.

1 Introduction and main results

► **a)** Let $(B_t, t \geq 0)$ denote a one-dimensional Brownian motion starting from 0, and let

$$\mathcal{E}_t = \exp\left(B_t - \frac{t}{2}\right), \quad (t \geq 0)$$

A reduced form of the celebrated Black-Scholes formula is the following.

$$(1_{\mathbf{K}}) \quad E[(\mathcal{E}_t - K)^+] = \mathcal{N}\left(-\frac{\ln K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right) - K \mathcal{N}\left(-\frac{\ln K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right)$$

where $K \geq 0$, and as usual:

$$\mathcal{N}(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x dy \exp\left(-\frac{y^2}{2}\right)$$

In fact, formula $(1_{\mathbf{K}})$ may be split into two parts:

$$(1_{\mathbf{K}}^+) \quad E[\mathcal{E}_t 1_{\mathcal{E}_t > K}] = \mathcal{N}\left(-\frac{\ln K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right)$$

$$(1_{\mathbf{K}}^-) \quad K P(\mathcal{E}_t > K) = K \mathcal{N}\left(-\frac{\ln K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right)$$

As noted in Section 5 of [1], formula $(1_{\mathbf{K}}^\pm)$ is obtained in an elementary manner, after performing the change of probability:

$$P'_{|\mathcal{F}_t} = \mathcal{E}_t \bullet P_{|\mathcal{F}_t}$$

which transforms (B_t) in $(B_t + t)$ hence $(B_t - \frac{t}{2})$ in $(B_t + \frac{t}{2})$.

► **b)** In this Note, we give and discuss a different representation of $(1_{\mathbf{K}}^\pm)$.

Theorem 1. *For any $K \geq 0$, there are the representations:*

$$(2_{\mathbf{K}}^-) \quad E[\mathcal{E}_t 1_{(\mathcal{E}_t > K)}] - K P(\mathcal{E}_t > K) = P\left(G_{(\ln K)}^{(1/2)} \leq t\right) \quad (K \geq 0)$$

$$(2_{\mathbf{K} \geq 1}^+) \quad E[\mathcal{E}_t 1_{\mathcal{E}_t > K}] + K P(\mathcal{E}_t > K) = P\left(T_{(\ln K)}^{(1/2)} \leq t\right) \quad (K \geq 1)$$

$$(2_{\mathbf{K} \leq 1}^+) \quad E[\mathcal{E}_t 1_{\mathcal{E}_t < K}] + K P(\mathcal{E}_t < K) = P\left(T_{(\ln K)}^{(1/2)} \leq t\right) \quad (0 \leq K \leq 1)$$

or, equivalently:

$$(\mathbf{2}_{\mathbf{K} \leq 1}^{++}) \quad E[\mathcal{E}_t 1_{\mathcal{E}_t > K}] + K P(\mathcal{E}_t > K) = K + P\left(T_{(\ell n K)}^{(1/2)} > t\right) \quad (0 \leq K \leq 1)$$

where, for $\nu \in \mathbb{R}$, and $B_t^{(\nu)} \equiv B_t + \nu t$, we write:

$$T_a^{(\nu)} = \inf\{t : B_t^{(\nu)} = a\}; \quad G_a^{(\nu)} = \sup\{t : B_t^{(\nu)} = a\}$$

Comment and Complements about Theorem 1:

(i) Our motivation to prove formulae such as $(\mathbf{2}_{\mathbf{K}}^-)$ was our desire to obtain an expression on the RHS showing in a clear manner that the LHS ($= E((\mathcal{E}_t - K)_+)$) is an increasing function of t .

This is not clear from $(\mathbf{1}_{\mathbf{K}})$, although this property of increase is a consequence of the submartingale property of $(\mathcal{E}_t - K)^+$; see Section 4 for a more extended discussion.

(ii) Obviously, an equivalent presentation of the "system" $(\mathbf{2}_{\mathbf{K}}^\pm)$ is, for $K \geq 1$:

$$(\mathbf{3}_{\mathbf{K} \geq 1}^+) \quad E[\mathcal{E}_t 1_{(\mathcal{E}_t > K)}] = \frac{1}{2} \left\{ P\left(T_{(\ell n K)}^{(1/2)} \leq t\right) + P\left(G_{(\ell n K)}^{(1/2)} \leq t\right) \right\}$$

$$\begin{aligned} (\mathbf{3}_{\mathbf{K} \geq 1}^-) \quad K P(\mathcal{E}_t > K) &= \frac{1}{2} \left\{ P\left(T_{(\ell n K)}^{(1/2)} \leq t\right) - P\left(G_{(\ell n K)}^{(1/2)} \leq t\right) \right\} \\ &= \frac{1}{2} P\left(T_{(\ell n K)}^{(1/2)} \leq t \leq G_{(\ell n K)}^{(1/2)}\right) \end{aligned}$$

and, for $0 \leq K \leq 1$:

$$(\mathbf{3}_{\mathbf{K} \leq 1}^+) \quad E[\mathcal{E}_t 1_{(\mathcal{E}_t > K)}] = \frac{1}{2} \left\{ 1 + K - P\left(T_{(\ell n K)}^{(1/2)} \leq t \leq G_{(\ell n K)}^{(1/2)}\right) \right\}$$

$$(\mathbf{3}_{\mathbf{K} \leq 1}^-) \quad K P(\mathcal{E}_t > K) = \frac{1}{2} \left\{ 1 + K - \left[P\left(T_{(\ell n K)}^{(1/2)} \leq t\right) + P\left(G_{(\ell n K)}^{(1/2)} \leq t\right) \right] \right\}$$

(iii) In order to give formulae $(\mathbf{2}_{\mathbf{K}}^\pm)$ an "explicit" character, we now recall the distributions of

$$T_a^{(\nu)} = \inf\{t : B_t^{(\nu)} = a\} \text{ and } G_a^{(\nu)} = \sup\{t : B_t^{(\nu)} = a\},$$

for $\nu > 0$, and $a > 0$ (these formulae will then be used with $\nu = \frac{1}{2}$ and $a = \ell n K$): denoting by $p_t^{(\nu)}(a)$ the density of $B_t^{(\nu)}$, we have:

$$(4) \quad P(T_a^{(\nu)} \in dt) = \left(\frac{a}{t}\right) p_t^{(\nu)}(a) dt \equiv \frac{a}{\sqrt{2\pi t^3}} \exp\left(-\frac{1}{2t}(a - \nu t)^2\right) dt$$

whereas

$$(5) \quad P(G_a^{(\nu)} \in dt) = \nu p_t^{(\nu)}(a) dt \equiv \frac{\nu}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t}(a - \nu t)^2\right) dt$$

Formula (4) may be obtained from the combination of (4) for $\nu = 0$, which is very well known, followed by an application of the Cameron-Martin relationship between the laws of $B^{(\nu)}$ and B . Formula (5) is a particular case of a more general formula for last passage times of transient diffusions, obtained in Pitman-Yor [3].

(iv) Although this is not strictly necessary at this point (but will be useful in our proof of Theorem 1), we also present the distributions of $T_a^{(-\nu)}$ and $G_a^{(-\nu)}$, for $a \geq 0$.

In fact, they may be obtained easily from those of $T_a^{(\nu)}$ and $G_a^{(\nu)}$ thanks to the Cameron-Martin absolute continuity relationships:

$$(6) \quad \begin{aligned} W_{|\mathcal{F}_{T_a} \cap (T_a < \infty)}^{(-\nu)} &= \exp(-2\nu a) \cdot W_{|\mathcal{F}_{T_a}}^{(\nu)} \\ W_{|\mathcal{F}_{G_a} \cap (G_a > 0)}^{(-\nu)} &= \exp(-2\nu a) \cdot W_{|\mathcal{F}_{G_a}}^{(\nu)} \end{aligned}$$

where, here, $W^{(\mu)}$ denotes the law of $(B_t + \mu t, t \geq 0)$ on canonical space, and T_a , resp. G_a , is the first, resp. last, hitting time of a by the coordinate process.

Thus, we deduce from formulae (6), (4) and (5) that:

$$P(G_a^{(-\nu)} > 0) = P(T_a^{(-\nu)} < \infty) = \exp(-2\nu a)$$

whereas:

$$(7) \quad \begin{aligned} P(T_a^{(-\nu)} \in dt) &= \left(\frac{a}{t}\right) p_t^{(-\nu)}(a) dt = \frac{a}{\sqrt{2\pi t^3}} \exp\left(-\frac{1}{2t}(a + \nu t)^2\right) dt \\ P(G_a^{(-\nu)} \in dt) &= \nu p_t^{(-\nu)}(a) dt = \frac{\nu}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t}(a + \nu t)^2\right) dt \end{aligned}$$

► c) Organisation of the remainder of the paper:

- In Section 2, we prove Theorem 1, independently from formulae $(\mathbf{1}_K^\pm)$
- In Section 3, we give an elementary proof of the agreement between formulae $(\mathbf{1}_K^\pm)$ and $(\mathbf{3}_K^\pm)$
- Section 4 concludes, by setting the matter in a broader context.

2 Proof of Theorem 1

Clearly, in order to prove Theorem 1, it now suffices to prove $(\mathbf{2}_{\mathbf{K}}^-)$, $(\mathbf{3}_{\mathbf{K} \geq 1}^-)$ and $(\mathbf{2}_{\mathbf{K} \leq 1}^+)$.

► **a) Proof of $(\mathbf{2}_{\mathbf{K}}^-)$** (for any $K \geq 0$)

We shall show that

$$E[(\mathcal{E}_t - K)^+] \stackrel{(i)}{=} K P\left(0 < G_{(\ell n K)}^{(-1/2)} \leq t\right) \stackrel{(ii)}{=} P\left(G_{(\ell n K)}^{(+1/2)} \leq t\right)$$

The proof of (ii) follows from the relationship between the laws of $G_a^{(-\nu)}$ and $G_a^{(\nu)}$ as discussed in Section 1; namely:

$$P(G_a^{(-\nu)} > 0) = \exp(-2\nu a) \text{ and } P(G_a^{(-\nu)} \in dt | G_a^{(-\nu)} > 0) = P(G_a^{(\nu)} \in dt)$$

For the proof of (i) , we rely upon the following formula

$$(8) \quad P(G_a^{(\mu)} \geq t | \mathcal{F}_t) = \left(\frac{\exp(2\mu a)}{\exp(2\mu B_t^{(\mu)})} \right) \wedge 1$$

which is valid for all $\mu \in \mathbb{R}$; this is a particular case of the results for last passage times of a transient real-valued diffusion, as discussed in Pitman-Yor [3].

In particular, for $\mu = -\nu$, $\nu > 0$, we get:

$$\exp(2\nu a) P(0 < G_a^{(-\nu)} \leq t | \mathcal{F}_t) = (\exp(2\nu B_t^{(-\nu)}) - \exp(2\nu a))^+$$

This obviously proves (i) , by taking $a = (\ell n K)$, $\nu = 1/2$.

► **b) Proof of $(\mathbf{3}_{\mathbf{K} \geq 1}^-)$** (for $K \geq 1$)

Using again formula (8), we see that $(\mathbf{3}_{\mathbf{K} \geq 1}^-)$ is equivalent to:

$$(9_{\mathbf{K}}) \quad K P\left(B_t - \frac{t}{2} > (\ell n K)\right) = \frac{1}{2} E\left(1_{(T_{(\ell n K)}^{(1/2)} \leq t)} \cdot \left(\frac{K}{\exp(B_t + \frac{1}{2})} \wedge 1\right)\right)$$

We now use the Cameron-Martin relationship on both sides to reduce the statement of $(9_{\mathbf{K}})$ to a statement about standard Brownian motion (B_t) , for which we denote: $M_t = \sup_{s \leq t} B_s$. Thus, we find that $(9_{\mathbf{K}})$ is equivalent to:

$$(10_{\mathbf{K}}) \quad K E\left(1_{(B_t > (\ell n K))} \cdot \exp\left(-\frac{B_t}{2}\right)\right)$$

$$= \frac{1}{2} E \left(\mathbf{1}_{(M_t > (\ell n K))} \cdot \left(\frac{K}{\exp(B_t)} \wedge 1 \right) \cdot \exp \left(\frac{B_t}{2} \right) \right)$$

We now decompose the RHS of $(\mathbf{10}_K)$ in a sum of two quantities:

$$\frac{1}{2} \left\{ \begin{array}{l} E \left[\mathbf{1}_{(M_t > (\ell n K))} \cdot \mathbf{1}_{(B_t > \ell n K)} \cdot K \exp \left(-\frac{B_t}{2} \right) \right] \\ + E \left[\mathbf{1}_{(M_t > \ell n K)} \cdot \mathbf{1}_{(B_t < \ell n K)} \cdot \exp \left(+\frac{B_t}{2} \right) \right] \end{array} \right\}$$

Thus, $(\mathbf{10}_K)$ now gets simplified to the equivalent form:

$$\begin{aligned} (\mathbf{11}_K) \quad & \frac{K}{2} E \left(\mathbf{1}_{(B_t > (\ell n K))} \cdot \exp \left(-\frac{B_t}{2} \right) \right) \\ & = \frac{1}{2} E \left(\mathbf{1}_{(M_t > (\ell n K))} \cdot \mathbf{1}_{(B_t < \ell n K)} \cdot \exp \left(\frac{B_t}{2} \right) \right) \end{aligned}$$

which, taking $x = (\ell n K)$, may be written as:

$$(\mathbf{12}_x) \quad E \left(\mathbf{1}_{(B_t > x)} \cdot \exp \left(x - \frac{B_t}{2} \right) \right) = E \left(\mathbf{1}_{(M_t > x > B_t)} \cdot \exp \left(\frac{B_t}{2} \right) \right)$$

We now show $(\mathbf{12}_x)$, from the right to the left, as a consequence of the reflection principle:

conditionally on \mathcal{F}_{T_x} , and $T_x < t$, we have:

$$B_t - x = \widehat{B}_{(t-T_x)}, \text{ with } \widehat{B} \text{ independent from } \mathcal{F}_{T_x};$$

hence, under this condition, the reflection principle boils down to:

$$(\spadesuit) \quad B_t - x \stackrel{(\text{law})}{=} -(B_t - x)$$

Thus, the RHS of $(\mathbf{12}_x)$ is:

$$\begin{aligned} & E \left(\mathbf{1}_{(T_x < t)} \cdot \mathbf{1}_{(B_t - x < 0)} \cdot \exp \left(\frac{1}{2} \{x + (B_t - x)\} \right) \right) \\ & \stackrel{(\text{from } (\spadesuit))}{=} E \left(\mathbf{1}_{(T_x < t)} \cdot \mathbf{1}_{(B_t - x < 0)} \cdot \exp \left(\frac{1}{2} \{x - (B_t - x)\} \right) \right) \\ & = E \left(\mathbf{1}_{(B_t > x)} \cdot \exp \left(x - \frac{B_t}{2} \right) \right), \text{ which is the LHS of } (\mathbf{12}_x) \end{aligned}$$

This proves $(\mathbf{3}_{K \geq 1})$, and, with $(\mathbf{2}_K^-)$, $(\mathbf{2}_{K \geq 1}^+)$.

► **c)** We now prove that $(\mathbf{2}_{K \geq 1}^+)$ implies $(\mathbf{2}_{K \leq 1}^+)$ (for $0 \leq K \leq 1$):
We introduce the probability P' such that:

$$P'_{\mathcal{F}_t} = \mathcal{E}_t \bullet P_{\mathcal{F}_t}$$

We note that, under P' , $\frac{1}{\mathcal{E}_t} := \widehat{\mathcal{E}}_t = \exp\left(\widehat{B}_t - \frac{t}{2}\right)$, for a new Brownian motion $(\widehat{B}_t, t \geq 0)$. Thus, the LHS of $(\mathbf{2}_{K \leq 1}^+)$ writes:

$$\begin{aligned} & P'(\mathcal{E}_t < K) + K P'(\widehat{\mathcal{E}}_t - 1_{(\mathcal{E}_t < K)}) \\ = & P'\left(\widehat{\mathcal{E}}_t > \frac{1}{K}\right) + K E'\left(1_{(\widehat{\mathcal{E}}_t > \frac{1}{K})}\right) \\ = & K E'\left(1_{(\widehat{\mathcal{E}}_t > \frac{1}{K})}\right) + \frac{1}{K} P'\left(\widehat{\mathcal{E}}_t > \frac{1}{K}\right) \\ = & K P\left(T_{(\ln K)}^{(1/2)} \leq t\right) \quad (\text{from } (\mathbf{2}_{\frac{1}{K}}^+)) \\ = & K P\left(T_{(\ln K)}^{(-1/2)} \leq t\right) \quad (\text{by symmetry}) \\ = & P\left(T_{(\ln K)}^{(1/2)} \leq t\right) \quad (\text{from (6)}) \end{aligned}$$

► **d)** Finally, we observe that $(\mathbf{2}_{K \leq 1}^+)$ is equivalent to $(\mathbf{2}_{K \leq 1}^{++})$, since:

$$\begin{aligned} & E(\mathcal{E}_t 1_{(\mathcal{E}_t < K)}) + K P(\mathcal{E}_t < K) \\ = & 1 - E(\mathcal{E}_t 1_{(\mathcal{E}_t > K)}) + K(1 - P(\mathcal{E}_t > K)) \quad (\text{since } E(\mathcal{E}_t) = 1) \\ = & 1 + K - \{E(\mathcal{E}_t 1_{(\mathcal{E}_t > K)}) + K P(\mathcal{E}_t > K)\} \end{aligned}$$

3 On the agreement between the classical Black-Scholes formula $(\mathbf{1}_{\mathbf{K}}^\pm)$ and our main result

► **a)** We now check in an elementary manner formulae $(\mathbf{2}_{\mathbf{K}}^\pm)$ by comparing their LHS, as given from the "traditional" Black-Scholes formulae $(\mathbf{1}_{\mathbf{K}}^\pm)$, with their RHS, as given by (4) and (5).

► **b)** The case $K \geq 1$. Since both sides of $(\mathbf{2}_{\mathbf{K}}^-)$ and $(\mathbf{2}_{K \geq 1}^+)$ are equal to 0 for $t = 0$, we need only check that the derivatives in t are equal; thus, our task is to show:

$$(13_{\mathbf{K}}^-) \quad \frac{\partial}{\partial t} \left\{ \mathcal{N}\left(-\frac{\ln K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right) - K \mathcal{N}\left(-\frac{\ln K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right) \right\}$$

$$\begin{aligned}
&= \left(\frac{K}{2}\right) \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{1}{2t} \left((\ell n K) + \frac{t}{2}\right)^2\right) \\
(13_{\mathbf{K} \geq 1}^+) \quad &\frac{\partial}{\partial t} \left\{ \mathcal{N}\left(-\frac{\ell n K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right) + K \mathcal{N}\left(-\frac{\ell n K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right) \right\} \\
&= K(\ell n K) \frac{1}{\sqrt{2\pi t^3}} \exp\left(-\frac{1}{2t} \left((\ell n K) + \frac{t}{2}\right)^2\right)
\end{aligned}$$

(We also see on these expressions the relationships between $p_t^{(-1/2)}(x)$ and $p_t^{(+1/2)}(x)$).

To prove $(13_{\mathbf{K}}^-)$ and $(13_{\mathbf{K} \geq 1}^+)$, we compute:

$$\begin{aligned}
&\bullet \quad \frac{\partial}{\partial t} \left(\mathcal{N}\left(-\frac{\ell n K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right) \right) \\
&= \frac{\sqrt{K}}{\sqrt{2\pi}} \exp\left(-\frac{1}{2} \left(\frac{(\ell n K)^2}{t} + \frac{t}{4}\right)\right) \left(\frac{\partial}{\partial t} \left\{ -\frac{(\ell n K)}{\sqrt{t}} + \frac{\sqrt{t}}{2} \right\} \right) \\
&\bullet \quad K \frac{\partial}{\partial t} \left(\mathcal{N}\left(-\frac{\ell n K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right) \right) \\
&= \frac{\sqrt{K}}{\sqrt{2\pi}} \exp\left(-\frac{1}{2} \left(\frac{(\ell n K)^2}{t} + \frac{t}{4}\right)\right) \left(\frac{\partial}{\partial t} \left\{ -\frac{(\ell n K)}{\sqrt{t}} - \frac{\sqrt{t}}{2} \right\} \right)
\end{aligned}$$

and $(13_{\mathbf{K}}^-)$ and $(13_{\mathbf{K} \geq 1}^+)$ are then obtained by elementary algebraic manipulations. (In fact, it is these very manipulations which led us to believe in the truth of Theorem 1!!).

► **c)** The case $0 \leq K \leq 1$. Since both sides of $(2_{\mathbf{K} \leq 1}^{++})$ are equal to $1 + K$ for $t = 0$, it suffices to prove, for $K \leq 1$:

$$\begin{aligned}
&\frac{\partial}{\partial t} \left\{ \mathcal{N}\left(-\frac{\log K}{\sqrt{t}} + \frac{\sqrt{t}}{2}\right) + K \mathcal{N}\left(-\frac{\log K}{\sqrt{t}} - \frac{\sqrt{t}}{2}\right) \right\} \\
&= \frac{\partial}{\partial t} P\left(T_{(\ell n K)}^{(1/2)} > t\right)
\end{aligned}$$

and this latter relation follows immediately from the computations done in point **b)** above and from (7).

Note also that $(2_{\mathbf{K} \geq 1}^+)$ and $(2_{\mathbf{K} \leq 1}^+)$ coincide for $K = 1$.

4 Further remarks and conclusion

► **a)** Easy variants of formulae $(\mathbf{2}_{\mathbf{K}}^{\pm})$ may be written, e.g. by using the scaling property of Brownian motion, so that $T_a^{(\nu)}$ and $G_a^{(\nu)}$ appear on the RHS of $(\mathbf{2}_{\mathbf{K}}^{\pm})$; however, writing down these variants would only complicate unnecessarily these formulae.

► **b)** We recall that, as a consequence of the time inversion property of Brownian motion, there are the relations:

$$(T_a^{(\nu)}, G_a^{(\nu)}) \stackrel{(\text{law})}{=} \left(\frac{1}{G_\nu^{(a)}}, \frac{1}{T_\nu^{(a)}} \right)$$

(see e.g., Pitman-Yor [3] for a more general discussion).

In particular, for $a = 0$, one gets:

$$G_0^{(\nu)} \stackrel{(\text{law})}{=} \frac{1}{T_\nu^{(0)}} \stackrel{(\text{law})}{=} \frac{B_1^2}{\nu^2}$$

In particular, formula $(\mathbf{2}_{\mathbf{K}=1}^-)$ becomes:

$$(14_1) \quad E[(\mathcal{E}_t - 1)^+] = E[(\mathcal{E}_t - 1)^-] = P(4B_1^2 \leq t)$$

which allowed us to answer M. Qian's question [4]:

is there a simple formula for:

$$\int_0^\infty \theta(dt) E[(\mathcal{E}_t - 1)^\pm]$$

where $\theta(dt)$ is a probability on \mathbb{R}_+ ?

From (14_1) , we easily obtain:

$$(15) \quad \int_0^\infty \theta(dt) E[(\mathcal{E}_t - 1)^\pm] = E[\bar{\theta}(4B_1^2)]$$

where $\bar{\theta}(x) = \theta([x, \infty))$ is the tail of θ .

To particularise even more, we give the explicit form of the Laplace transform:

$$(16) \quad \int_0^\infty dt e^{-\lambda t} E[(\mathcal{E}_t - 1)^\pm] = \frac{1}{\lambda} E[\exp(-\lambda(4B_1^2))] \\ = \frac{1}{\lambda} \frac{1}{\sqrt{1 + 8\lambda}}$$

It is this question which set us on the general quest for a representation of

$$E[(\mathcal{E}_t - K)^+]$$

as a cumulative distribution function in t .

► **c)** We come back to the time inversion property of BM, in order to throw another light upon our main result $(\mathbf{2}_K^-)$, which relates the European call price with the cumulative function of last Brownian passage times. (This paragraph has been partly inspired by unpublished notes by Peter Carr [2].) Indeed, a variant of $(\mathbf{2}_K^-)$ is the following:

for every $t \geq 0$, $K \geq 0$, and $\phi : C([0, t]) \rightarrow \mathbb{R}_+$, measurable,

$$(17) \quad E[\phi(B_u, u \leq t)(K - \mathcal{E}_t)^+] = K E[\phi(B_u, u \leq t)1_{(\mathcal{G}_K \leq t)}]$$

where $\mathcal{G}_K = \sup\{u : \mathcal{E}_u = K\}$.

Writing (17) in terms of the Brownian motion $(\widehat{B}_v, v \geq 0)$ such that: $B_u = u\widehat{B}_{(1/u)}$, and setting $s = 1/t$, it is clearly seen that (17) is equivalent to:

$$K P(\widehat{T}_{1/2}^{(-\ell n K)} \geq s | \widehat{B}_s) = \left(K - \exp\left(\frac{1}{s}\widehat{B}_s - \frac{1}{2s}\right) \right)^+$$

where: $\widehat{T}_a^{(\nu)} = \inf\{u : \widehat{B}_u + \nu u = a\}$.

Since hats are no longer necessary for our purpose, we drop them, and we now look for an independent proof of:

$$(18) \quad P(T_{1/2}^{(-\ell n K)} \geq s | B_s = x) = \left(1 - \frac{1}{K} \exp\left(\frac{x}{s} - \frac{1}{2s}\right) \right)^+$$

On the LHS of (18), we may replace $(B_s = x)$ by $(B_s - s(\ell n K) = x - s(\ell n K))$. Now, as a consequence of the Cameron-Martin relationship, the conditional expectation:

$$E[F(B_u - \nu u, u \leq s) | B_s - \nu s = y]$$

does not depend on ν ; hence, (18) is equivalent to:

$$P(T_{1/2}^{(-\ell n K)} \geq s | B_s = x - s(\ell n K)) = \left(1 - \frac{1}{K} \exp\left(\frac{x - \frac{1}{2}}{s}\right) \right)^+$$

which simplifies to:

$$P(\sup_{u \leq s} B_u < \frac{1}{2} | B_s = y) = \left(1 - \exp\left(\frac{y - \frac{1}{2}}{s}\right) \right)^+$$

or, by scaling:

$$P(\sup_{u \leq 1} B_u < \frac{1}{2\sqrt{s}} | B_1 = \frac{y}{\sqrt{s}}) = \left(1 - \exp \left(\frac{1}{\sqrt{s}} \left(\frac{y}{\sqrt{s}} - \frac{1}{2\sqrt{s}} \right) \right) \right)^+$$

This is equivalent to:

$$(19) \quad P(\sup_{u \leq 1} B_u < \sigma | B_1 = y) = (1 - \exp(2\sigma(y - \sigma)))^+$$

for $\sigma \geq 0$, and $y \in \mathbb{R}$.

This formula is trivial for $\sigma < y$, and, for $\sigma \geq y$, it follows from the classical formula:

$$(20) \quad P(\sup_{u \leq 1} B_u \in d\sigma, B_1 \in da) = \frac{da d\sigma}{\sqrt{2\pi}} 2(2\sigma - a) e^{-\frac{(2\sigma - a)^2}{2}} 1_{\{a < \sigma; \sigma \geq 0\}}$$

► **d)** In a future work, we plan to study more generally how quantities such as the calls and puts:

$$E[(S_t - K)^+] \text{ and } E[(S_t - K)^-]$$

associated with a general \mathbb{R}_+ -valued continuous local martingale $(S_t, t \geq 0)$ may be written in terms of cumulative functions.

► **e)** In [1], the authors present eight different approaches to the Black-Scholes formula, among which the change of numéraire approach (Section 5 of [1]), and the local time approach (Section 6 of [1]). This local time approach, together with $(\mathbf{2}_{\mathbf{K}}^-)$ yields the relationship:

$$(21) \quad P \left(G_{(\ell_n K)}^{(1/2)} \leq t \right) = \left(\frac{1}{2} \right) E[\mathcal{L}_t^K]$$

where $(\mathcal{L}_t^K, t \geq 0)$ denotes the local time at level K of $(\mathcal{E}_t, t \geq 0)$. It is this kind of relationship $(\mathbf{21})$ which is central in the obtention in [3] of a general expression for the law of a last passage time of a transient diffusion. However, to our knowledge, despite the remarkable survey [1] of methods leading to the Black-Scholes formula, no interpretation of this formula seems to have been made in terms of last passage times distributions.

References

- [1] J. Andreasen, B. Jensen, R. Poulsen. Eight valuation methods in financial mathematics: the Black-Scholes formula as an example. *Math. Scientist*, 23, 18–40 (1998).

- [2] Peter Carr. Put No Touch-Inversion. Unpublished Notes. (July 2007).
- [3] J. Pitman and M. Yor. Bessel processes and infinitely divisible laws. In Stochastic integrals (Proc. Sympos., Univ. Durham, Durham, 1980), LNM 851, pages 285–370. Springer, Berlin, 1981.
- [4] M. Qian. Personal communication. August 2007.

Acknowledgments: We are grateful to M. Qian for raising the question mentioned in Section 4, and to T. Fujita for pointing out reference [1].