



HAL
open science

On Kazhdan-Lusztig cells in type B

Cédric Bonnafé

► **To cite this version:**

| Cédric Bonnafé. On Kazhdan-Lusztig cells in type B. 2009. hal-00283937v2

HAL Id: hal-00283937

<https://hal.science/hal-00283937v2>

Preprint submitted on 14 Jan 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON KAZHDAN-LUSZTIG CELLS IN TYPE B

CÉDRIC BONNAFÉ

ABSTRACT. We prove that, for any choice of parameters, the Kazhdan-Lusztig cells of a Weyl group of type B are unions of combinatorial cells (defined using the domino insertion algorithm).

Let (W_n, S_n) be the Weyl group of type B_n , where $S_n = \{t, s_1, \dots, s_{n-1}\}$ and where the Dynkin diagram is given by



Let $\ell : W_n \rightarrow \mathbb{N} = \{0, 1, 2, 3, \dots\}$ be the length function. Let Γ be a totally ordered abelian group and let $\varphi : W_n \rightarrow \Gamma$ be a weight function (in the sense of Lusztig [10, §3.1]). We set

$$\varphi(t) = b \quad \text{and} \quad \varphi(s_1) = \dots = \varphi(s_{n-1}) = a.$$

To this datum, the Kazhdan-Lusztig theory (with unequal parameters [10]) associates a partition of W_n into left, right or two-sided cells [10, Chapter 8].

In [3, Conjectures A and B], Geck, Iancu, Lam and the author have proposed several conjectures for describing these partitions (at least whenever $a, b > 0$, but this is not such a big restriction, as can be seen from [2, Corollary 5.8]): they involve a *domino insertion algorithm*. Roughly speaking, one can define a partition of W_n into *combinatorial (left, right or two-sided) (a, b) -cells* (which depend on a, b and which are defined combinatorially using the domino insertion algorithm): the combinatorial (left, right or two-sided) cells should coincide with the Kazhdan-Lusztig (left, right or two-sided) cells. The aim of this paper is to prove one of the two inclusions (see Theorem 1.24):

Theorem. *If two elements of W_n lie in the same combinatorial (left, right or two-sided) cell, then they lie in the same Kazhdan-Lusztig (left, right or two-sided) cell.*

In the case of the symmetric group, the partition into left cells (obtained by Kazhdan and Lusztig [7, Theorem 1.4]) uses the Robinson-Schensted correspondence, and the key tool is a description of this correspondence using plactic/coplactic relations

Date: January 14, 2009.

1991 Mathematics Subject Classification. According to the 2000 classification: Primary 20C08; Secondary 20C15.

The author is partly supported by the ANR (Project No JC07-192339).

(also called Knuth relations). For W_n , whenever $b > (n - 1)a$, the partition into left, right or two-sided cells was obtained by Iancu and the author (see [4, Theorem 7.7] and [1, Corollaries 3.6 and 5.2]) again by using the translation of a generalised Robinson-Schensted correspondence through plactic/coplactic relations.

Recently, M. Taskin [13] and T. Pietraho [12] have independently provided plactic/coplactic relations for the domino insertion algorithm. Our methods rely heavily on their results: we show that, if two elements of W_n are directly related by a plactic relation, then they are in the same Kazhdan-Lusztig cell. The key step will be the Propositions 2.14 and 2.15, where some multiplications between elements of the Kazhdan-Lusztig basis are computed by brute force. We then derive some consequences (see Propositions 4.1 and 6.1), where it is proved that some elements are in the same left cells. Then, the rest of the proof just uses the particular combinatoric of Weyl groups of type B , together with classical properties of Kazhdan-Lusztig cells.

Acknowledgements. Part of this work was done while the author stayed at the MSRI during the winter 2008. The author wishes to thank the Institute for its hospitality and the organizers of the two programs of that period for their invitation.

The author also wants to thank warmly L. Iancu and N. Jacon for their careful reading of a preliminary version of this paper and for their useful comments.


CONTENTS

1. Notation	3
1.A. Weyl group.	3
1.B. Decomposition of elements of W_n .	5
1.C. Subgroups W_m of W_n .	7
1.D. Hecke algebra.	7
1.E. Kazhdan-Lusztig basis.	8
1.F. Cells.	8
1.G. Domino insertion.	8
2. Kazhdan-Lusztig polynomials, structure constants	10
2.A. Recollection of general facts.	10
2.B. Special features in type B .	11
2.C. $*$ -operation.	14
2.D. Two relations \leftarrow^L .	15
3. Proof of Proposition 2.14	15
3.A. Easy reduction.	15
3.B. The Bruhat interval $[\mathbf{x}; \mathbf{y}]$.	15

3.C. The elements $z \in [x; y]$ such that $tz < z$.	16
3.D. Computation of the $\tilde{\mu}_I$.	17
4. Consequence of Proposition 2.14	19
5. Proof of Proposition 2.15	22
6. Consequences of Proposition 2.15	25
7. Proof of Theorem 1.24	27
7.A. Knuth relations.	27
7.B. Proof of Theorem 1.24.	27
7.C. Proof of Lemma 7.1.	28
7.D. Proof of Lemma 7.2.	28
7.E. Proof of Lemma 7.3.	29
References	30

1. NOTATION

1.A. Weyl group. Let (W_n, S_n) be the Weyl group of type B_n , where $S_n = \{t, s_1, \dots, s_{n-1}\}$ and where the Dynkin diagram is given by



Let $\ell : W_n \rightarrow \mathbb{N} = \{0, 1, 2, 3, \dots\}$ be the length function. Let $I_n = \{\pm 1, \dots, \pm n\}$: we shall identify W_n with the group of permutations w of I_n such that $w(-i) = -w(i)$ for all $w \in I_n$. The identification is through the following map

$$t \longmapsto (1, -1) \quad \text{and} \quad s_i \longmapsto (i, i + 1)(-i, -i - 1).$$

The next notation comes from [4, §4]: it is rather technical but will be used throughout this paper. We set $t_1 = r_1 = t$ and, for $1 \leq i \leq n - 1$, we set

$$r_{i+1} = s_i r_i \quad \text{and} \quad t_{i+1} = s_i t_i s_i.$$

We shall often use the following well-known lemma:

Lemma 1.1. *Let $w \in W_n$, $i \in \{1, 2, \dots, n - 1\}$ and $j \in \{1, 2, \dots, n\}$. Then:*

- (a) $\ell(ws_i) > \ell(w)$ (that is, $ws_i > w$) if and only if $w(i) < w(i + 1)$.
- (b) $\ell(wt_j) > \ell(w)$ if and only if $w(j) > 0$.

As a permutation of I_n , we have

(1.2)
$$t_i = (i, -i)$$

and

$$(1.3) \quad r_i(j) = \begin{cases} -i & \text{if } j = 1, \\ j - 1 & \text{if } 2 \leq j \leq i, \\ j & \text{if } i + 1 \leq j \leq n. \end{cases}$$

An easy computation shows that, if $j \in \{1, 2, \dots, n-1\}$ and $i \in \{1, 2, \dots, n\}$, then

$$(1.4) \quad s_j r_i = \begin{cases} r_i s_j & \text{if } j > i, \\ r_{i+1} & \text{if } j = i, \\ r_{i-1} & \text{if } j = i - 1, \\ r_i s_{j+1} & \text{if } 1 \leq j < i - 1. \end{cases}$$

Note also that, if $l \geq 2$, then

$$(1.5) \quad r_l r_l = r_{l-1} r_l s_1.$$

We set $a_0 = 1$ and, if $0 \leq l \leq n$, we set

$$a_l = r_1 r_2 \cdots r_l.$$

As a permutation of I_n , we have

$$(1.6) \quad a_l(i) = \begin{cases} i - 1 - l & \text{if } 1 \leq i \leq l, \\ i & \text{if } l + 1 \leq i \leq n. \end{cases}$$

In particular,

$$(1.7) \quad a_l^{-1} = a_l$$

and, if $i \in \{1, 2, \dots, n-1\} \setminus \{l\}$, then

$$(1.8) \quad a_l s_i a_l = \begin{cases} s_{l-i} & \text{if } i < l, \\ s_i & \text{if } i > l. \end{cases}$$

Note also that

$$(1.9) \quad \ell(a_l) = \frac{l(l+1)}{2}.$$

We shall identify the symmetric group \mathfrak{S}_n with the subgroup of W_n generated by s_1, \dots, s_{n-1} . We also set $I_n^+ = \{1, 2, \dots, n\}$. Then, as a group of permutations of I_n , we have

$$(1.10) \quad \mathfrak{S}_n = \{w \in W_n \mid w(I_n^+) = I_n^+\}.$$

If $1 \leq i \leq j \leq n$, we denote by $[i, j]$ the set $\{i, i+1, \dots, j\}$ and by $\mathfrak{S}_{[i, j]}$ the subgroup of W_n (or of \mathfrak{S}_n) generated by $s_i, s_{i+1}, \dots, s_{j-1}$. If $j < i$, then we set $[i, j] = \emptyset$ and $\sigma_{[i, j]} = 1$. As a group of permutations of I_n , we have

$$(1.11) \quad \mathfrak{S}_{[i, j]} = \{w \in \mathfrak{S}_n \mid \forall k \in I_n^+ \setminus [i, j], w(k) = k\}.$$

The longest element of W_n will be denoted by w_n (it is usually denoted by w_0 , but since we shall use induction on n , we need to emphasize its dependence on n).

We denote by σ_n the longest element of \mathfrak{S}_n . The longest element of $\mathfrak{S}_{[i,j]}$ will be denoted by $\sigma_{[i,j]}$. As a permutation of I_n , we have

$$(1.12) \quad w_n = (1, -1)(2, -2) \cdots (n, -n).$$

Note also that

$$(1.13) \quad \begin{cases} w_n &= t_1 t_2 \cdots t_n = t_n \cdots t_2 t_1 \\ w_n &= a_n \sigma_n = \sigma_n a_n, \\ \sigma_n &= \sigma_{[1,n]} \end{cases}$$

and that

$$(1.14) \quad w_n \text{ is central in } W_n.$$

1.B. Decomposition of elements of W_n . If $0 \leq l \leq n$, we denote by $\mathfrak{S}_{l,n-l}$ the subgroup of \mathfrak{S}_n generated by $\{s_1, \dots, s_{n-1}\} \setminus \{s_l\}$. Then $\mathfrak{S}_{l,n-l} = \mathfrak{S}_{[1,l]} \times \mathfrak{S}_{[l+1,n]} \simeq \mathfrak{S}_l \times \mathfrak{S}_{n-l}$. We denote by $Y_{l,n-l}$ the set of elements $w \in \mathfrak{S}_n$ which are of minimal length in $w\mathfrak{S}_{l,n-l}$. Note that a_l normalizes $\mathfrak{S}_{l,n-l}$ (this follows from 1.8).

If $w \in W_n$, we denote by $\ell_t(w)$ the number of occurrences of t in a reduced decomposition of w (this does not depend on the choice of the reduced decomposition). We set $\ell_s(w) = \ell(w) - \ell_t(w)$.

Lemma 1.15. *Let $w \in W_n$. Then there exists a unique quadruple $(l, \alpha, \beta, \sigma)$ where $0 \leq l \leq n$, $\alpha, \beta \in Y_{l,n-l}$ and $\sigma \in \mathfrak{S}_{l,n-l}$ are such that $w = \alpha a_l \sigma \beta^{-1}$. Moreover, there exists a unique sequence $1 \leq i_1 < i_2 < \cdots < i_l \leq n$ such that $\alpha a_l = r_{i_1} r_{i_2} \cdots r_{i_l}$. We have*

$$\begin{aligned} \ell(w) &= \ell(\alpha) + \ell(a_l) + \ell(\sigma) + \ell(\beta), \\ \ell_t(w) &= l \end{aligned}$$

and

$$\{i_1, \dots, i_l\} = \{i \in [1, n] \mid w^{-1}(i) < 0\}.$$

Note also that

$$\ell(\alpha) = \sum_{k=1}^l (i_k - k).$$

Proof. See [4, §4, and especially Proposition 4.10]. □

If $l \in [0, n]$ and if $1 \leq i_1 < \cdots < i_l \leq n$ and $1 \leq j_1 < \cdots < j_{n-l} \leq n$ are two sequences such that $[1, n] = \{i_1, \dots, i_l\} \cup \{j_1, \dots, j_{n-l}\}$, then it follows easily from 1.3 that

$$(1.16) \quad \begin{cases} (r_{i_1} \cdots r_{i_l})^{-1}(i_k) = k - l - 1 & \text{if } 1 \leq k \leq l, \\ (r_{i_1} \cdots r_{i_l})^{-1}(j_k) = l + k & \text{if } 1 \leq k \leq n - l. \end{cases}$$

The elements α , β and σ of the previous lemma will be denoted by α_w , β_w and σ_w respectively. We have

$$(1.17) \quad \ell_t(w^{-1}) = \ell_t(w), \quad \alpha_{w^{-1}} = \beta_w, \quad \beta_{w^{-1}} = \alpha_w \quad \text{and} \quad \sigma_{w^{-1}} = a_l(\sigma_w)^{-1}a_l.$$

We shall now describe how the multiplication by the longest element w_n acts on the decomposition given by Lemma 1.15. For this, we denote by $\sigma_{l,n-l}$ the longest element of $\mathfrak{S}_{l,n-l}$.

Proposition 1.18. *Let $w \in W_n$ and let $l = \ell_t(w)$. Then:*

- (a) $\ell_t(w_n w) = n - l$.
- (b) $\alpha_{w_n w} = \alpha_w \sigma_n \sigma_{n-l,l}$ and $\beta_{w_n w} = \beta_w \sigma_n \sigma_{n-l,l}$.
- (c) $\sigma_{w_n w} = \sigma_n \sigma_n^{-1} \sigma_{n-l,l}$.
- (d) *Let $1 \leq i_1 < \dots < i_l \leq n$ be the sequence such that $\alpha_w a_l = r_{i_1} \dots r_{i_l}$. Then $\alpha_{w_n w} = r_{j_1} \dots r_{j_{n-l}}$, where $1 \leq j_1 < \dots < j_{n-l} \leq n$ is the sequence such that $\{i_1, \dots, i_l\} \cup \{j_1, \dots, j_{n-l}\} = [1, n]$.*

Proof. (a) is clear. (d) follows from Lemma 1.15. We now prove (b) and (c) simultaneously. For this, let $\alpha' = \alpha_w \sigma_n \sigma_{n-l,l}$, $\beta' = \beta_w \sigma_n \sigma_{n-l,l}$ and $\sigma' = \sigma_n \sigma_w \sigma_n^{-1} \sigma_{n-l,l}$. By the unicity statement of the Lemma 1.15, we only need to show the following three properties:

- (1) $\alpha', \beta' \in Y_{n-l,l}$.
- (2) $\sigma' \in \mathfrak{S}_{n-l,l}$.
- (3) $w_n w = \alpha' a_{n-l} \sigma' \beta'^{-1}$.

For this, note first

$$\sigma_n \mathfrak{S}_{l,n-l} \sigma_n^{-1} = \mathfrak{S}_{n-l,l},$$

so that (2) follows immediately. This also implies that $\sigma_n \sigma_{n-l,l} \sigma_n^{-1} = \sigma_{l,n-l}$ because conjugacy by σ_n in \mathfrak{S}_n preserves the length.

Let us now show (1). Let $i \in \{1, 2, \dots, n\} \setminus \{n-l\}$. We want to show that $\ell(\alpha' s_i) > \ell(\alpha')$. By Lemma 1.1, this amounts to show that $\alpha'(i+1) > \alpha'(i)$. But $\alpha' = \alpha_w \sigma_{l,n-l} \sigma_n$. Also $\sigma_n(i) = n+1-i > \sigma_n(i+1) = n-i$ and $n+1-i$ and $n-i$ both belong to the same interval $[1, l]$ or $[l+1, n]$. Hence $\sigma_{l,n-l} \sigma_n(i) < \sigma_{l,n-l} \sigma_n(i+1)$ and $\alpha_w \sigma_{l,n-l} \sigma_n(i) < \alpha_w \sigma_{l,n-l} \sigma_n(i+1)$ since $\alpha_w \in Y_{l,n-l}$. This shows that $\alpha' \in Y_{n-l,l}$. Similarly, $\beta' \in Y_{n-l,l}$. So (1) is proved.

It remains to show (3). We have

$$\begin{aligned} \alpha' a_{n-l} \sigma' \beta'^{-1} &= (\alpha_w \sigma_n \sigma_{n-l,l}) \cdot a_{n-l} \cdot (\sigma_n \sigma_w \sigma_n^{-1} \sigma_{n-l,l}) \cdot (\sigma_{n-l,l}^{-1} \sigma_n^{-1} \beta_w^{-1}) \\ &= \alpha_w \sigma_n \sigma_{n-l,l} a_{n-l} \sigma_n \sigma_w \beta_w^{-1} \end{aligned}$$

But $\sigma_{n-l,l} = \sigma_{[n-l+1,n]}\sigma_{n-l}$ and $\sigma_n\sigma_{[n-l+1,n]}\sigma_n^{-1} = \sigma_{[1,l]} = \sigma_l$ so

$$\begin{aligned}\alpha' a_{n-l}\sigma'\beta'^{-1} &= \alpha_w\sigma_l\sigma_n\sigma_{n-l}a_{n-l}\sigma_n^{-1}\sigma_w\beta_w^{-1} \\ &= \alpha_w\sigma_l\sigma_n w_{n-l}\sigma_n^{-1}\sigma_w\beta_w^{-1},\end{aligned}$$

the last equality following from 1.13. Now, $\sigma_n w_{n-l}\sigma_n^{-1} = w_l w_n$ (see again 1.13) so

$$\begin{aligned}\alpha' a_{n-l}\sigma'\beta'^{-1} &= \alpha_w\sigma_l w_l w_n \sigma_w \beta_w^{-1} \\ &= \alpha_w a_l w_n \sigma_w \beta_w^{-1} \\ &= w_n \alpha_w a_l \sigma_w \beta_w^{-1} = w_n w,\end{aligned}$$

the second equality following from 1.13 and the third one from the fact that w_n is central (see 1.14). \square

1.C. Subgroups W_m of W_n . If $m \leq n$, we shall view W_m naturally as a subgroup of W_n (the pointwise stabilizer of $[m+1, n]$). It is the standard parabolic subgroup generated by $S_m = \{t, s_1, \dots, s_{m-1}\}$: we denote by $X_n^{(m)}$ the set of $w \in W_n$ which are of minimal length in wW_m . For simplification, we set $X_n = X_n^{(n-1)}$. It follows from Lemma 1.1 that:

Lemma 1.19. *Let w be an element of W_n . Then w belongs to $X_n^{(m)}$ if and only if $0 < w(1) < w(2) < \dots < w(m)$.*

If $I = \{i_1, \dots, i_l\} \subseteq [1, n-1]$ with $i_1 < \dots < i_l$, then we set

$$c_I = s_{i_1} s_{i_2} \cdots s_{i_l} \quad \text{and} \quad d_I = s_{i_l} \cdots s_{i_2} s_{i_1}.$$

By convention, $c_\emptyset = d_\emptyset = 1$. We have

$$(1.20) \quad X_n = \{c_{[i,n-1]} \mid 1 \leq i \leq n\} \dot{\cup} \{d_{[1,i]} t c_{[1,n-1]} \mid 0 \leq i \leq n-1\}.$$

1.D. Hecke algebra. We fix a totally ordered abelian group Γ (denoted additively) and a weight function $\varphi : W_n \rightarrow \Gamma$. We set

$$\varphi(t) = b \quad \text{and} \quad \varphi(s_1) = a \quad (= \varphi(s_2) = \dots = \varphi(s_{n-1})).$$

Note that

$$(1.21) \quad \varphi(w) = \ell_t(w)b + \ell_s(w)a$$

for all $w \in W_n$.

We denote by A the group algebra $\mathbb{Z}[\Gamma]$. We shall use an exponential notation: $A = \bigoplus_{\gamma \in \Gamma} \mathbb{Z}e^\gamma$, where $e^\gamma \cdot e^{\gamma'} = e^{\gamma+\gamma'}$ for all $\gamma, \gamma' \in \Gamma$. We set

$$Q = e^b \quad \text{and} \quad q = e^a.$$

Note that Q and q are not necessarily algebraically independent. We set

$$A_{<0} = \bigoplus_{\gamma < 0} \mathbb{Z}e^\gamma,$$

and we define similarly $A_{\leq 0}$, $A_{>0}$ and $A_{\geq 0}$.

We shall denote by \mathcal{H}_n the *Hecke algebra of W_n with parameter φ* : it is the free A -module with basis $(T_w)_{w \in W_n}$ and the multiplication is A -bilinear and is completely determined by the following rules:

$$\begin{cases} T_w T_{w'} = T_{ww'} & \text{if } \ell(ww') = \ell(w) + \ell(w'), \\ (T_t - Q)(T_t + Q^{-1}) = 0, \\ (T_{s_i} - q)(T_{s_i} + q^{-1}) = 0 & \text{if } 1 \leq i \leq n-1. \end{cases}$$

We also set

$$\mathcal{H}_n^{<0} = \bigoplus_{w \in W_n} A_{<0} T_w.$$

Finally, we denote by $\bar{\cdot} : \mathcal{H}_n \rightarrow \mathcal{H}_n$ the unique A -semilinear involution of \mathcal{H}_n such that $\overline{e^\gamma} = e^{-\gamma}$ and $\overline{T_w} = T_{w^{-1}}$ for all $\gamma \in \Gamma$ and $w \in W_n$.

1.E. Kazhdan-Lusztig basis. We shall recall here the basic definitions of Kazhdan-Lusztig theory. If $w \in W_n$, then [10, Theorem 5.2] there exists a unique $C_w \in \mathcal{H}_n$ such that

$$\begin{cases} \overline{C_w} = C_w \\ C_w \equiv T_w \pmod{\mathcal{H}_n^{<0}}. \end{cases}$$

Note that [10, §5.3]

$$(1.22) \quad C_w - T_w \in \bigoplus_{w' < w} A_{<0} T_{w'},$$

where \leq denotes the Bruhat order on W_n . In particular, $(C_w)_{w \in W_n}$ is an A -basis of \mathcal{H}_n , called the *Kazhdan-Lusztig basis* of \mathcal{H}_n .

1.F. Cells. If $x, y \in W_n$, then we shall write $x \xleftarrow{L} y$ (resp. $x \xleftarrow{R} y$, resp. $x \xleftarrow{LR} y$) if there exists $h \in \mathcal{H}_n$ such that the coefficient of C_x in the decomposition of hC_y (resp. $C_y h$, resp. hC_y or $C_y h$) is non-zero. We denote by \leq_L (resp. \leq_R , resp. \leq_{LR}) the transitive closure of \xleftarrow{L} (resp. \xleftarrow{R} , resp. \xleftarrow{LR}). Then \leq_L , \leq_R and \leq_{LR} are preorders on W_n and we denote respectively by \sim_L , \sim_R and \sim_{LR} the associated equivalence relations [10, Chapter 8]. An equivalence class for \sim_L (resp. \sim_R , resp. \sim_{LR}) is called a *left* (resp. *right*, resp. *two-sided*) *cell*. We recall the following result [10, §8.1]: if $x, y \in W_n$, then

$$(1.23) \quad x \sim_L y \iff x^{-1} \sim_R y^{-1}.$$

1.G. Domino insertion. If $r \geq 0$ and $w \in W_n$, then the domino insertion algorithm (see [8], [14], [15]) into the 2-core $\delta_r = (r, r-1, \dots, 2, 1)$ associates to w

a standard domino tableau $D_r(w)$ (with n dominoes, filled with $\{1, 2, \dots, n\}$). If D is a domino tableau, we denote by $\mathbf{sh}(D)$ its *shape*: we shall denote by $\mathbf{sh}_r(w)$ the shape of $D_r(w)$ (which is equal to the shape of $D_r(w^{-1})$, loc. cit.).

If x and $y \in W_n$ we shall write $x \sim_L^r y$ (resp. $x \sim_R^r y$, resp. $x \sim_{LR}^r y$) if $D_r(x^{-1}) = D_r(y^{-1})$ (resp. $D_r(x) = D_r(y)$, resp. $\mathbf{sh}_r(x) = \mathbf{sh}_r(y)$). These are equivalence relations on W_n . Note that \sim_{LR}^r is the equivalence relation generated by \sim_L^r and \sim_R^r .

We denote by \approx_L^{r+1} (resp. \approx_R^{r+1} , resp. \approx_{LR}^{r+1}) the equivalence relation generated by \sim_L^r and \sim_L^{r+1} (resp. \sim_R^r and \sim_R^{r+1} , resp. \sim_{LR}^r and \sim_{LR}^{r+1}). Recall the following conjecture from [3, Conjectures A and B]:

Conjecture. *Assume that $a, b > 0$. Let $r \geq 0$ and $? \in \{L, R, LR\}$.*

- (a) *If $ra < b < (r+1)a$, then the relations $\sim_?^r$ and $\sim_?^r$ coincide.*
- (b) *If $r \geq 1$ and $b = ra$, then the relations $\sim_?^r$ and $\approx_?^r$ coincide.*

The main result of this paper is the following partial result towards the previous conjecture:

Theorem 1.24. *Assume that $a, b > 0$. Let $r \geq 0$, $? \in \{L, R, LR\}$ and $x, y \in W_n$. Then:*

- (a) *If $ra < b < (r+1)a$ and $x \sim_?^r y$, then $x \sim_? y$.*
- (b) *If $r \geq 1$, $b = ra$ and $x \approx_?^r y$, then $x \sim_? y$.*

The other sections of this paper are devoted to the proof of this theorem.

COMMENTS - If one assumes Lusztig's Conjectures **P1**, **P2**, ..., **P15** in [10, Chapter 14], then Theorem 1.24 implies that the statement (a) of the Conjecture is true. Indeed, Lusztig's Conjectures imply in this case that the left cell representations are irreducible, and one can conclude by a counting argument. It might be probable that a similar argument applies for the statement (b), using results of Pietraho [11]: however, we are not able to do it.

In the case where $b > (n-1)a$, Theorem 1.24 was proved in [4, Theorem 7.7] (in fact, the conjecture was also proved) by using a counting argument. The proof here does not make use of the counting argument. \square

2. KAZHDAN-LUSZTIG POLYNOMIALS, STRUCTURE CONSTANTS

Hypothesis and notation. *From now on, and until the end of this paper, we assume that a, b are positive. Recall that $Q = e^b$ and $q = e^a$, so that $\mathbb{Z}[Q, Q^{-1}, q, q^{-1}] \subseteq A$. If $p \in A_{\geq 0}$, we denote by $\tau_A(p)$ the coefficient of $1 = e^0$ in the expansion of p in the basis $(e^\gamma)_{\gamma \in \Gamma}$.*

2.A. Recollection of general facts. If x and y are elements of W_n , we set

$$C_x C_y = \sum_{z \in W_n} h_{x,y,z} C_z,$$

where the $h_{x,y,z}$ belong to A and satisfy

$$\overline{h_{x,y,z}} = h_{x,y,z}.$$

We also set

$$C_y = \sum_{x \in W_n} p_{x,y}^* T_x \quad \text{and} \quad p_{x,y} = e^{\varphi(y) - \varphi(x)} p_{x,y}^*.$$

Recall [10, Proposition 5.4] that

$$(2.1) \quad \begin{cases} p_{y,y}^* = p_{y,y} = 1, \\ p_{x,y}^* \in A_{<0} & \text{if } x \neq y, \\ p_{x,y}^* = p_{x,y} = 0 & \text{if } x \not\leq y, \\ p_{x,y} \in A_{\geq 0}, \\ \tau_A(p_{x,y}) = 1 & \text{if } x \leq y. \end{cases}$$

Now, if $s \in S_n$, Lusztig [10, Proposition 6.3] has defined inductively a family of polynomials $(M_{x,y}^s)_{sx < x < y < sy}$ by the following properties:

$$(2.2a) \quad \overline{M_{x,y}^s} = M_{x,y}^s,$$

$$(2.2b) \quad M_{x,y}^s + \sum_{\substack{x < z < y \\ sz < z}} p_{x,z}^* M_{z,y}^s - e^{\varphi(s)} p_{x,y}^* \in A_{<0}.$$

With this notation, we have [10, Theorem 6.6]:

Theorem 2.3 (Kazhdan-Lusztig, Lusztig). *Let $s \in S_n$ and let $y \in W_n$. Then:*

$$(a) \quad C_s C_y = \begin{cases} C_{sy} + \sum_{sx < x < y} M_{x,y}^s C_x & \text{if } sy > y, \\ (e^{\varphi(s)} + e^{-\varphi(s)}) C_y & \text{if } sy < y. \end{cases}$$

(b) If $sy < y$, and if $x \leq y$, then

$$p_{x,y} = \begin{cases} q^2 p_{x,sy} + p_{sx,sy} - \sum_{\substack{x \leq z < sy \\ sz < z}} e^{\varphi(y) - \varphi(z)} p_{x,z} M_{z,sy}^s & \text{if } sx < x, \\ p_{sx,y} & \text{if } sx > x, \end{cases}$$

and

$$p_{x,y}^* = \begin{cases} qp_{x,sy}^* + p_{sx,sy}^* - \sum_{\substack{x \leq z < sy \\ sz < z}} p_{x,z}^* M_{z,sy}^s & \text{if } sx < x, \\ e^{-\varphi(s)} p_{sx,y}^* & \text{if } sx > x. \end{cases}$$

Corollary 2.4. *If $s, s' \in \{s_1, \dots, s_{n-1}\}$ and $x, y \in W_n$ are such that $sx < x < s'x = y < sy$, then $x \sim_L y$.*

Proof. See [9, Proposition 5 (b)]. □

2.B. Special features in type B. The previous results of this section hold for any Coxeter group (finite or not). In this subsection, we shall investigate what is implied by the structure of W_n . The particular ingredient we shall need is the following lemma [4, §4]:

Lemma 2.5. *$\{a_l \mid 0 \leq l \leq n\}$ is the set of elements $w \in W_n$ which have minimal length in $\mathfrak{S}_n w \mathfrak{S}_n$. If $x < a_l$ for some $l \in \{1, 2, \dots, n\}$ and some $x \in W_n$, then $\ell_t(x) < \ell_t(a_l) = l$.*

It has the following consequences (here, if $p \in \mathbb{Z}[q]$, we denote by $\deg_q p$ its degree in the variable q):

Corollary 2.6. *Let x and y be two elements of W_n such that $x \leq y$ and $\ell_t(x) = \ell_t(y)$. Then:*

- (a) $p_{x,y} \in \mathbb{Z}[q]$ and, if $x \neq y$, then $\deg_q p_{x,y} < \ell(y) - \ell(x)$.
- (b) If $1 \leq i \leq n-1$ is such that $s_i x < x < y < s_i y$, then $M_{x,y}^{s_i} \in \mathbb{Z}$: it is equal to $\tau_A(qp_{x,y}^*)$ (note also that $qp_{x,y}^* \in \mathbb{Z}[q^{-1}]$).

Proof. We shall prove (a) and (b) together by induction on the pair $(\ell(y), \ell(y) - \ell(x))$ (with lexicographic order). The result is obvious if $\ell(y) = \ell(x)$ or if $\ell(y) \leq 1$. So assume now that $\ell(y) > 1$, that $\ell(y) - \ell(x) > 0$ and that (a) and (b) hold for all pairs (x', y') such that $(\ell(y'), \ell(y') - \ell(x')) < (\ell(y), \ell(y) - \ell(x))$. First, note that

$$e^{\varphi(y) - \varphi(x)} = q^{\ell(y) - \ell(x)},$$

because $\varphi(y) - \varphi(x) = (\ell_t(y) - \ell_t(x))b + (\ell_s(y) - \ell_s(x))a = (\ell_s(y) - \ell_s(x))a = (\ell(y) - \ell(x))a$.

Let us first prove (a). So we have $x < y$ and $\ell_t(x) = \ell_t(y)$. By Lemma 2.5, this implies that there exists $i \in \{1, 2, \dots, n-1\}$ such that $s_i y < y$ or $ys_i < y$. In the second case, one can exchange y and y^{-1} (and x and x^{-1}) by using [10, §5.6], so that we may assume that $s_i y < y$. Then, Theorem 2.3 (b) can be rewritten as follows:

$$p_{x,y} = \begin{cases} (q^2 p_{x,s_i y} - q^{\ell(y) - \ell(x)} M_{z,s_i y}^{s_i}) + p_{s_i x, s_i y} - \sum_{\substack{x < z < s_i y \\ s_i z < z}} q^{\ell(y) - \ell(z)} p_{x,z} M_{z,s_i y}^{s_i} & \text{if } s_i x < x, \\ p_{s_i x, y} & \text{if } s_i x > x. \end{cases}$$

If $s_i x > x$, then the result follows from the induction hypothesis. If $s_i x < x$, then

$$q^2 p_{x,s_i y} - q^{\ell(y) - \ell(x)} M_{x,s_i y}^{s_i} = q^{\ell(y) - \ell(x)} (qp_{x,s_i y}^* - M_{x,s_i y}^{s_i})$$

belong to $\mathbb{Z}[q]$ and has degree $< \ell(y) - \ell(x)$ by the induction hypothesis. The other terms in the above formula also belong to $\mathbb{Z}[q]$ and also have degree $< \ell(y) - \ell(x)$ by the induction hypothesis. So we get (a).

Let us now prove (b). So we assume that $s_i x < x < y < s_i y$. Then, using the induction hypothesis and 2.1, the condition 2.2 (b) can be rewritten

$$M_{x,y}^{s_i} - qp_{x,y}^* \in A_{<0}.$$

Now, the result follows easily from (a). \square

Now, if $tx < x < y < ty$ are such that $\ell_t(x) = \ell_t(y)$, let us define an element $\mu_{x,y} \in A$ by induction on $\ell(y) - \ell(x)$ by the following formula:

$$\mu_{x,y} = p_{x,y} - \sum_{\substack{x < z < y \\ tz < z}} p_{x,z} \mu_{z,y}.$$

It follows easily from Corollary 2.6 (and an induction argument on $\ell(y) - \ell(x)$) that

$$(2.7) \quad \mu_{x,y} \in \mathbb{Z}[q] \quad \text{and} \quad \deg_q \mu_{x,y} < \ell(y) - \ell(x).$$

Moreover:

Corollary 2.8. *Assume that $tx < x < y < ty$ and that $\ell_t(x) = \ell_t(y)$. Then:*

$$(a) \text{ If } b > (\ell(y) - \ell(x))a, \text{ then } M_{x,y}^t = Qq^{\ell(x) - \ell(y)} \mu_{x,y} + Q^{-1}q^{\ell(y) - \ell(x)} \overline{\mu_{x,y}}.$$

(b) If $b = (\ell(y) - \ell(x))a$, then $M_{x,y}^t = \mu_{x,y} + \overline{\mu_{x,y}} - \tau_A(\mu_{x,y})$.

Proof. Let us assume that $b \geq (\ell(y) - \ell(x))a$. We shall prove the result by induction on $\ell(y) - \ell(x)$. By the induction hypothesis, the condition 2.2 (b) can be written

$$M_{x,y}^t - Qq^{\ell(x)-\ell(y)}p_{x,y} + \sum_{\substack{x < z < y \\ tz < z}} p_{x,z}^* \left(Qq^{\ell(z)-\ell(y)}\mu_{z,y} + Q^{-1}q^{\ell(y)-\ell(z)}\overline{\mu_{z,y}} \right) \in A_{<0}.$$

But, if $x < z < y$ and $tz < z$, then

$$p_{x,z}^* Q^{-1}q^{\ell(y)-\ell(z)}\overline{\mu_{z,y}} \in A_{<0}$$

because $p_{x,z}^* \in A_{<0}$, $\overline{\mu_{z,y}} \in A_{\leq 0}$ and $Q^{-1}q^{\ell(y)-\ell(z)} = e^{-b+(\ell(y)-\ell(z))a} \in A_{<0}$ (since $\ell(y) - \ell(z) < \ell(y) - \ell(x)$). Therefore,

$$M_{x,y}^t - Qq^{\ell(x)-\ell(y)}p_{x,y} + \sum_{\substack{x < z < y \\ tz < z}} Qq^{\ell(x)-\ell(y)}p_{x,z}\mu_{z,y} \in A_{<0}.$$

In other words,

$$M_{x,y}^t - Qq^{\ell(x)-\ell(y)}\mu_{x,y} \in A_{<0}.$$

Let $\mu = Qq^{\ell(x)-\ell(y)}\mu_{x,y}$. Two cases may occur:

- If $b > (\ell(y) - \ell(x))a$, then $\mu \in A_{>0}$ and so the condition 2.2 (a) forces $M_{x,y}^t = \mu + \overline{\mu}$, as required.
- If $b = (\ell(y) - \ell(x))a$, then $\mu = \mu_{x,y} \in A_{\geq 0}$ and now the condition 2.2 (a) forces $M_{x,y}^t = \mu + \overline{\mu} - \tau_A(\mu)$, as required.

The proof of the Corollary is complete. \square

We conclude this subsection with two results involving the decomposition of Lemma 1.15.

Lemma 2.9. *Let x and y be two elements of W_n and let $s \in S_n$ be such that $sx < x < y < sy$, $\ell_t(x) = \ell_t(y)$ and $\beta_x = \beta_y = \beta$. Then $M_{x,y}^s = M_{x\beta,y\beta}^s$ (note that $\beta_x\beta = \beta_y\beta = 1$).*

Proof. See [4, Proposition 7.2]. Strictly speaking, in [4], the authors are generally working with a special choice of a function φ (“asymptotic case”): however, the reader can check that the proof of this particular result, namely [4, Proposition 7.2], remains valid for all choices of parameters. \square

Proposition 2.10. *Let $l \in [0, n]$, let σ and $\sigma' \in \mathfrak{S}_{l, n-l}$ be such that $\sigma \sim_L \sigma'$ and let $\beta \in Y_{l, n-l}$. Then*

$$a_l \sigma \beta^{-1} \sim_L a_l \sigma' \beta^{-1}.$$

Proof. By the description of Kazhdan-Lusztig left cells in the symmetric group [7, Theorem 1.4 and §4], we may assume that there exist two elements s and $s' \in \{s_1, \dots, s_{l-1}, s_{l+1}, \dots, s_{n-1}\}$ such that $\sigma' = s'\sigma$ and $s\sigma < \sigma < \sigma' < s\sigma'$. Let $u = a_l s a_l$ and $u' = a_l s' a_l$. Then u and u' belong to $\{s_1, \dots, s_{l-1}, s_{l+1}, \dots, s_{n-1}\}$ by 1.8, and

$$u a_l \sigma \beta^{-1} < a_l \sigma \beta^{-1} < u' a_l \sigma \beta^{-1} = a_l \sigma' \beta^{-1} < u a_l \sigma' \beta^{-1}.$$

So (*) follows from Corollary 2.4. \square

2.C. *-operation. We shall recall the definition of the *-operation (see [7, §4]) and prove some properties which are particular to the type B . Let us introduce some notation. If $1 \leq i \leq n-2$ and $x \in W_n$, we set

$$\mathcal{R}_i(x) = \{s \in \{s_i, s_{i+1}\} \mid \ell(xs) < \ell(x)\}.$$

We denote by $\mathcal{D}_i(W_n)$ the set of $x \in W_n$ such that $|\mathcal{R}_i(x)| = 1$. If $x \in \mathcal{D}_i(W_n)$, then it is readily seen that the set $\{x s_i, x s_{i+1}\} \cap \mathcal{D}_i(W_n)$ is a singleton. We shall denote by $\gamma_i(x)$ the unique element of this set (it is denoted by x^* in [7, §4], but we want to emphasize that it depends on i). Note that

$$\gamma_i \circ \gamma_i = \text{Id}_{\mathcal{D}_i(W_n)}.$$

We recall Kazhdan-Lusztig result [7, Corollary 4.3]: if x and $y \in \mathcal{D}_i(W_n)$, then

$$(2.11) \quad x \sim_L y \iff \gamma_i(x) \sim_L \gamma_i(y).$$

The fact that t is not conjugate to any of the s_k 's implies the following easy fact:

Proposition 2.12. *Let $x \in W_n$ and let $1 \leq k \leq n-1$. Then $x s_k > x$ if and only if $t x s_k > t x$.*

Proof. Indeed, by Lemma 1.1, we have $x s_k > x$ if and only if $x(k) < x(k+1)$. But, for any $j \in I_n^+$, there is no element $j' \in I_n$ such that $t(j) < j' < j$. So $x(k) < x(k+1)$ if and only if $t x(k) < t x(k+1)$ that is, if and only if $t x s_k > t x$ (again by Lemma 1.1). \square

The proposition 2.12 implies immediately the following result:

Corollary 2.13. *Let $x \in W_n$ and let $1 \leq i \leq n-2$. Then $x \in \mathcal{D}_i(W_n)$ if and only if $t x \in \mathcal{D}_i(W_n)$. If this is the case, then $\gamma_i(t x) = t \gamma_i(x)$.*

2.D. Two relations \xleftarrow{L} . The crucial steps towards the proof of Theorem 1.24 are the following two propositions, whose proofs will be given in sections 3 and 5 respectively.

Proposition 2.14. *Let $l \in \{1, \dots, n-1\}$ and assume that $b \geq (n-1)a$. Then $M_{r_1 \cdots r_l \sigma_{[l+1, n]}, r_2 \cdots r_l r_n \sigma_{[l+1, n]}}^t \neq 0$.*

Proposition 2.15. *Let $l \in \{1, \dots, n-1\}$ and assume that $(n-2)a < b \leq (n-1)a$. Then $a_{l-1} \sigma_{[l, n]} \xleftarrow{L} a_l \sigma_{[l, n]}$.*

3. PROOF OF PROPOSITION 2.14

Notation. *If $u, v \in W_n$ are such that $u \leq v$, we denote by $[u; v]$ the Bruhat interval between u and v . In this section, and only in this section, we assume that $l \geq 1$ and $b \geq (n-1)a$ and we set $x = r_1 \cdots r_l \sigma_{[l+1, n]}$ and $y = r_2 \cdots r_l r_n \sigma_{[l+1, n]}$.*

3.A. Easy reduction. Note that

$$tx < x < y < ty,$$

so it makes sense to compute $M_{x, y}^t$. Moreover, $\ell(y) - \ell(x) = n-1$ so, by Corollary 2.8, we only need to prove that $\mu_{x, y} \neq 0$ (even if $b = (n-1)a$). For this, we only need to show that

$$(?) \quad \tau_A(\mu_{x, y}) \neq 0.$$

3.B. The Bruhat interval $[x; y]$. First, note that

$$x = a_l \sigma_{[l+1, n]} = \sigma_{[l+1, n]} a_l$$

and $y = s_1 \cdots s_{l-1} s_{n-1} \cdots s_l x = s_1 \cdots s_{l-1} s_{n-1} \cdots s_l \sigma_{[l+1, n]} a_l = c_{[1, l-1]} \sigma_{[l, n]} a_l$.

Since a_l has minimal length in $\mathfrak{S}_n a_l$, the map

$$\begin{array}{ccc} [\sigma_{[l+1, n]}; c_{[1, l-1]} \sigma_{[l, n]}] & \longrightarrow & [x; y] \\ z & \longmapsto & z a_l \end{array}$$

is an increasing bijection [10, Lemma 9.10 (f)]. Since the support of $c_{[1, l-1]}$ is disjoint from the support of $\sigma_{[l, n]}$, the map

$$\begin{array}{ccc} [1; c_{[1, l]}] \times [\sigma_{[l+1, n]}; \sigma_{[l, n]}] & \longrightarrow & [\sigma_{[l+1, n]}; c_{[1, l-1]} \sigma_{[l, n]}] \\ (z, z') & \longmapsto & z z' \end{array}$$

is an increasing bijection (for the product order). Now, $\sigma_{[l,n]}$ is the longest element of $\mathfrak{S}_{[l,n]}$ and $\sigma_{[l+1,n]}\sigma_{[l,n]} = c_{[l,n-1]}$. Therefore, the map

$$\begin{array}{ccc} [1; c_{[l,n-1]}] & \longrightarrow & [\sigma_{[l+1,n]}; \sigma_{[l,n]}] \\ z & \longmapsto & z\sigma_{[l,n]} \end{array}$$

is a decreasing bijection. So, if we denote by $\mathcal{P}(E)$ the set of subsets of a set E , then the maps

$$\begin{array}{ccc} \mathcal{P}([1, l-1]) & \longrightarrow & [1; c_{[1, l-1]}] \\ I & \longmapsto & c_I \end{array}$$

and

$$\begin{array}{ccc} \mathcal{P}([l, n-1]) & \longrightarrow & [\sigma_{[l+1,n]}; \sigma_{[l,n]}] \\ J & \longmapsto & c_{\bar{J}}\sigma_{[l,n]} \end{array}$$

are increasing bijections (here, \bar{J} denotes the complement of J). On the other hand, the map

$$\begin{array}{ccc} \mathcal{P}([1, l-1]) \times \mathcal{P}([l, n-1]) & \longrightarrow & \mathcal{P}([1, n-1]) \\ (I, J) & \longmapsto & I \cup J \end{array}$$

is an increasing bijection. Finally, by composing all these bijections, we get an isomorphism of ordered sets

$$\alpha : \begin{array}{ccc} \mathcal{P}([1, n-1]) & \longrightarrow & [x; y] \\ I & \longmapsto & c_{I \cap [1, l-1]} c_{I \cap [l, n-1]} \sigma_{[l,n]} a_I. \end{array}$$

3.C. The elements $z \in [x; y]$ such that $tz < z$. If $I \subseteq [1, n-1]$ is such that $t\alpha(I) < \alpha(I)$, we set $\tilde{\mu}_I = \tau_A(\mu_{\alpha(I), y})$. So we can rephrase (?) as follows:

$$(??) \quad \tilde{\mu}_\emptyset \neq 0.$$

But, by the induction formula that defines the μ -polynomials and by 2.1, we have, for all $I \subseteq [1, n-1]$ such that $t\alpha(I) < \alpha(I)$,

$$(3.1) \quad \tilde{\mu}_I = 1 - \sum_{\substack{I \subsetneq J \subseteq [1, n-1] \\ t\alpha(J) < \alpha(J)}} \tilde{\mu}_J.$$

Let

$$\mathcal{E} = \{I \in \mathcal{P}([1, n-1]) \mid t\alpha(I) < \alpha(I)\}.$$

The set \mathcal{E} is easy to describe:

Lemma 3.2. *Let $I \subseteq [1, n-1]$. Then $t\alpha(I) > \alpha(I)$ if and only if $[1, l-1] \not\subseteq I$.*

Proof of Lemma 3.2. By Lemma 1.1, we just need to show that

$$(\#) \quad \alpha(I)^{-1}(1) > 0 \text{ if and only if } [1, l-1] \not\subseteq I.$$

For simplification, we set $A = [1, l-1] \cap I$ and $B = \overline{I \cap [l, n-1]}$. So $\alpha(I) = c_A c_B \sigma_{[l, n]} a_l$.

First, assume that $[1, l-1] \not\subseteq I$. Then $0 < c_A^{-1}(1) < n$, so $\sigma_{[l, n]}^{-1} c_B^{-1} c_A^{-1}(1) = c_A^{-1}(1)$ and $\alpha(I)^{-1}(1) = a_l^{-1}(c_A^{-1}(1)) < 0$ by 1.6. This shows $(\#)$ in this case.

Now, let us assume that $[1, l-1] = I$. Then $c_A = s_1 \cdots s_{l-1}$ and $c_B = s_l \cdots s_{n-1}$ and so $c_A^{-1}(1) = l$ and $c_B^{-1}(l) = n$. In particular, $\alpha(I)^{-1}(1) = a_l^{-1} \sigma_{[l, n]}(n) = a_l^{-1}(l) = -1 < 0$ by 1.6. This shows $(\#)$ again in this case.

Now, let us assume that $[1, l-1] \subsetneq I$. Then $c_A^{-1}(1) = l$ and $c_B^{-1}(l) < n$ and so $\sigma_{[l, n]}^{-1} c_B^{-1} c_A^{-1}(1) > l$. So $\alpha(I)^{-1}(1) > 0$ by 1.6. The proof of $(\#)$ is complete. \square

3.D. Computation of the $\tilde{\mu}_I$. We shall now compute the family $(\tilde{\mu}_I)_{I \in \mathcal{E}}$ by descending induction on $|I|$, by using the formula 3.1. For this, the following well-known lemma will be useful.

Lemma 3.3. *If S is a finite set and $I \subsetneq S$, then $\sum_{I \subseteq J \subseteq S} (-1)^{|J|} = 0$.*

To obtain the value of $\tilde{\mu}_\emptyset$, the proof goes in three steps.

$$(3.4) \quad \text{If } [l, n-1] \subseteq I \subsetneq [1, n-1], \text{ then } \tilde{\mu}_I = (-1)^{n-|I|}.$$

Proof of 3.4. First, note that $I \in \mathcal{E}$ by Lemma 3.2. We argue by descending induction on $|I|$. If $|I| = n-2$, then $\tilde{\mu}_I = 1$, as desired. Now, let us assume that $[l, n-1] \subseteq I \subsetneq [1, n-1]$ and that $\tilde{\mu}_J = (-1)^{n-|J|}$ for all $I \subsetneq J \subsetneq [1, n-1]$. Then, by 3.1, we have

$$\tilde{\mu}_I = 1 - \sum_{I \subsetneq J \subsetneq [1, n-1]} (-1)^{n-|J|}.$$

Therefore,

$$\tilde{\mu}_I = 1 + (-1)^{n-|I|} + (-1)^{n-(n-1)} - \sum_{I \subseteq J \subseteq [1, n-1]} (-1)^{n-|J|} = (-1)^{n-|I|},$$

the last equality following from Lemma 3.3. \square

$$(3.5) \quad \text{If } I \in \mathcal{E} \text{ is such that } [l, n-1] \not\subseteq I \text{ and } I \not\subseteq [1, l-1], \text{ then } \tilde{\mu}_I = 0.$$

Proof of 3.5. We shall again argue by descending induction on $|I|$. Let $I' = I \cup [l, n-1]$. Then, by 3.1, we have

$$\tilde{\mu}_I = 1 - \sum_{\substack{J \in \mathcal{E} \\ I \subsetneq J \text{ and } I' \subseteq J}} \tilde{\mu}_J - \sum_{\substack{J \in \mathcal{E} \\ I \subsetneq J \text{ and } I' \not\subseteq J}} \tilde{\mu}_J.$$

But, if $J \in \mathcal{E}$ is such that $I \subsetneq J$ and $I' \not\subseteq J$, (or, equivalently, $[l, n-1] \not\subseteq J$), then $\tilde{\mu}_J = 0$ by the induction hypothesis. On the other hand, if $J \in \mathcal{E}$ is such that $I \subsetneq J$ and $I' \subseteq J$, then $\tilde{\mu}_J = (-1)^{n-|J|}$ by 3.4. Therefore,

$$\tilde{\mu}_I = 1 - \sum_{\substack{J \in \mathcal{E} \\ I \subsetneq J \text{ and } I' \subseteq J}} (-1)^{n-|J|} = 1 - \sum_{I' \subseteq J \subsetneq [1, n-1]} (-1)^{n-|J|} = - \sum_{I' \subseteq J \subseteq [1, n-1]} (-1)^{n-|J|} = 0$$

by Lemma 3.3. \square

(3.6) *If $I \subseteq [1, l-1]$, then $\tilde{\mu}_I = (-1)^{l-1-|I|}$.*

Proof of 3.6. Note that $I \in \mathcal{E}$. We shall argue by descending induction on $|I|$. First, for all J such that $I \subsetneq J \subseteq [1, n-1]$, we have $t\alpha(I) > \alpha(I)$. Therefore, $\tilde{\mu}_{[1, l-1]} = 1$, as desired.

Now, let $I \subsetneq [1, l-1]$ and assume that, for all $I \subsetneq J \subseteq [1, l-1]$, we have $\tilde{\mu}_J = (-1)^{l-1-|J|}$. Then

$$\tilde{\mu}_I = 1 - \sum_{\substack{J \in \mathcal{E} \\ I \subsetneq J}} \tilde{\mu}_J.$$

Now, if $J \in \mathcal{E}$ is such that $I \subsetneq J$, then three cases may occur:

- If $J \subseteq [1, l-1]$, then $\tilde{\mu}_J = (-1)^{l-1-|J|}$ by the induction hypothesis.
- If $J \not\subseteq [1, l-1]$ and $[l, n-1] \not\subseteq I$, then $\tilde{\mu}_J = 0$ by 3.5.
- If $[l, n-1] \subseteq J$, then $\tilde{\mu}_J = (-1)^{n-|J|}$.

Therefore, if we set $I' = I \cap [l, n-1]$, then we get

$$\tilde{\mu}_I = 1 - \sum_{I' \subseteq J \subsetneq [l, n-1]} (-1)^{n-|J|} - \sum_{I \subsetneq J \subsetneq [1, l-1]} (-1)^{l-1-|J|}.$$

But

$$1 - \sum_{I' \subseteq J \subsetneq [l, n-1]} (-1)^{n-|J|} = - \sum_{I' \subseteq J \subseteq [l, n-1]} (-1)^{n-|J|} = 0$$

$$\text{and } - \sum_{I \subsetneq J \subsetneq [1, l-1]} (-1)^{l-1-|J|} = (-1)^{l-1-|I|} - \sum_{I \subseteq J \subsetneq [1, l-1]} (-1)^{l-1-|J|} = (-1)^{l-1-|I|}$$

by Lemma 3.3. The proof is now complete. \square

As a special case of 3.6, we get that

$$\tilde{\mu}_\emptyset = (-1)^{l-1}.$$

This shows (?). The proof of the Proposition 2.14 is complete.

4. CONSEQUENCE OF PROPOSITION 2.14

The aim of this section is to prove the following

Proposition 4.1. *Let $l \in \{0, 1, \dots, n\}$, let $\alpha, \beta \in Y_{l, n-l}$ and let σ and $\sigma' \in \mathfrak{S}_{l, n-l}$ be such that $\sigma \sim_L \sigma'$. Assume that $b \geq (n-1)a$. Then*

$$\alpha a_l \sigma \beta^{-1} \sim_L a_l \sigma' \beta^{-1}.$$

REMARKS - (1) The condition $\sigma \sim_L \sigma'$ does not depend on the choice of a and b in Γ . Indeed, by [5, Theorem 1], $\sigma \sim_L \sigma'$ in W_n if and only if $\sigma \sim_L \sigma'$ in $\mathfrak{S}_{l, n-l}$. But this last condition depends neither on the choice of b (since $t \notin \mathfrak{S}_{l, n-l}$) nor on the choice of a (provided that it is in $\Gamma_{>0}$).

(2) If $b > (n-1)a$, then the above proposition is proved in [4, Theorem 7.7] (see also [1, Corollary 5.2] for the exact bound) by a counting argument. The proof below will not use this counting argument but uses instead the proposition 2.14: it allows to extend the scope of validity to the case where $b = (n-1)a$ (this is compatible with [2, Conjecture A (b)]). \square

Proof. First, recall that $a_l \sigma \beta^{-1} \sim_L a_l \sigma' \beta^{-1}$ by Proposition 2.10. This shows that we may (and we will) assume that $\sigma = \sigma'$. We want to show that $\alpha a_l \sigma \beta^{-1} \sim_L a_l \sigma \beta^{-1}$. We shall use induction on n . So let (P_n) denote the following statement:

(P_n) For all $l \in [0, n]$, for all sequences $1 \leq i_1 < \dots < i_l \leq n$, for all $\sigma \in \mathfrak{S}_{l, n-l}$ and for all $\beta \in Y_{l, n-l}$, we have $r_{i_1} r_{i_2} \dots r_{i_l} \sigma \beta^{-1} \sim_L r_1 r_2 \dots r_l \sigma \beta^{-1}$.

The property (P_1) is vacuously true and the property (P_2) can be easily checked by a straightforward computation. So we assume that $n \geq 3$ and (P_m) holds for all $m < n$. Now, let $l \in [0, n]$, let $1 \leq i_1 < \dots < i_l \leq n$ be a sequence of elements of $[1, n]$, let $\sigma \in \mathfrak{S}_{l, n-l}$ and let $\beta \in Y_{l, n-l}$. As a consequence of this induction hypothesis, we get:

Lemma 4.2. *If $k \in [1, l]$ is such that $i_k < n$, then $r_{i_1} r_{i_2} \dots r_{i_l} \sigma \beta^{-1} \sim_L r_1 \dots r_k r_{i_{k+1}} \dots r_{i_l} \sigma \beta^{-1}$.*

Proof. Let $w = r_{i_1} r_{i_2} \cdots r_{i_l} \sigma \beta^{-1}$ and $w' = r_1 \cdots r_k r_{i_{k+1}} \cdots r_{i_l} \sigma \beta^{-1}$. Let us write $w = vx^{-1}$ and $w' = v'x'^{-1}$ with $v, v' \in W_{i_k}$ and $x, x' \in X_n^{(i_k)}$. First, note that

$$ww'^{-1} = (r_{i_1} \cdots r_{i_k}) \cdot (r_1 \cdots r_k)^{-1} \in W_{i_k}.$$

Therefore, $x = x'$ and

$$vv'^{-1} = (r_{i_1} \cdots r_{i_k}) \cdot (r_1 \cdots r_k)^{-1} \in W_{i_k}.$$

Moreover, by Lemma 1.19, we have $0 < x(1) < \cdots < x(i_k)$. So, if $i \in [1, i_k]$, then $v^{-1}(i) < 0$ (resp. $v'^{-1}(i) < 0$) if and only if $i \in \{i_1, \dots, i_k\}$ (resp. $\{1, \dots, k\}$). So, by Lemma 1.15, we have

$$v = r_{i_1} \cdots r_{i_k} \tau \gamma^{-1} \quad \text{and} \quad v' = r_1 \cdots r_k \tau \gamma^{-1},$$

where $\tau \in \mathfrak{S}_{k, i_k - k}$ and $\gamma \in Y_{k, i_k - k}$. But, since $i_k < n$, it follows from the induction hypothesis that $v \sim_L v'$. Here, note that $v \sim_L v'$ in W_{i_k} if and only if $v \sim_L v'$ in W_n (see [5, Theorem 1]). So, by [10, Proposition 9.11], we get that $w \sim_L w'$. \square

Corollary 4.3.

- (a) If $i_l < n$, then $r_{i_1} r_{i_2} \cdots r_{i_l} \sigma \beta^{-1} \sim_L r_1 r_2 \cdots r_l \sigma \beta^{-1}$.
- (b) If $i_l = n$, then $r_{i_1} r_{i_2} \cdots r_{i_l} \sigma \beta^{-1} \sim_L r_1 r_2 \cdots r_{l-1} r_n \sigma \beta^{-1}$.

By Corollary 4.3, we only need to show that

$$(?) \quad r_1 r_2 \cdots r_{l-1} r_n \sigma \beta^{-1} \sim_L r_1 r_2 \cdots r_l \sigma \beta^{-1}.$$

Now, let us write $\sigma = (\lambda, \mu)$, where $\lambda \in \mathfrak{S}_{[1, l]}$ and $\mu \in \mathfrak{S}_{[l+1, n]}$. Three cases may occur:

- *Case 1:* If $\lambda = 1$ and $\mu = \sigma_{[l+1, n]}$, then $\sigma = \sigma_{[l+1, n]}$. Since $r_1 r_2 \cdots r_{l-1} r_k \sigma \beta^{-1} = s_k s_{k-1} \cdots s_{l+1} \sigma \beta^{-1}$ for all $k > l$, we have

$$\begin{aligned} r_1 \cdots r_{l-1} r_n \sigma \beta^{-1} &\leq_L r_1 \cdots r_{l-1} r_{n-1} \sigma \beta^{-1} \leq_L \cdots \\ &\leq_L r_1 \cdots r_{l-1} r_{l+1} \sigma \beta^{-1} \leq_L r_1 \cdots r_{l-1} r_l \sigma \beta^{-1}. \end{aligned}$$

On the other hand, by Proposition 2.14 and Lemma 2.9, we get $r_1 \cdots r_{l-1} r_l \sigma \beta^{-1} \leq_L r_1 \cdots r_{l-1} r_n \sigma \beta^{-1}$. This shows (?) in this particular case.

- *Case 2:* If $\mu \neq \sigma_{[l+1, n]}$, then $n \geq l+2$ and there exists $k \in [l+1, n-1]$ such that $s_k \sigma > \sigma$. Let i be maximal such that $s_i \sigma > \sigma$. We shall prove (?) by descending induction on i . For simplification, let $x = r_1 \cdots r_{l-1} r_n \sigma \beta^{-1}$.

First, if $i = n-1$, then, by 1.4, we have (since $n-2 > l-1$)

$$\begin{aligned} s_{n-2} x &= r_1 \cdots r_{l-1} s_{n-2} r_n \sigma \beta^{-1} = r_1 \cdots r_{l-1} r_n s_{n-1} \sigma \beta^{-1} > s_{n-2} x, \\ s_{n-1} x &= r_1 \cdots r_{l-1} s_{n-1} r_n \sigma \beta^{-1} r_1 \cdots r_{l-1} r_{n-1} \sigma \beta^{-1} < x \end{aligned}$$

and $s_{n-2} s_{n-1} x = r_1 \cdots r_{l-1} s_{n-2} r_{n-1} \sigma \beta^{-1} = r_1 \cdots r_{l-1} r_{n-2} \sigma \beta^{-1} < s_{n-1} x$.

So $x \sim_L s_{n-1} x$ by Corollary 2.4. On the other hand, by Corollary 4.3, we have $s_{n-1} x \sim_L r_1 \cdots r_l \sigma \beta^{-1}$, so we get (?) in this case.

Now, assume that $l + 1 \leq i < n - 1$. Then $s_{i+1}\sigma < \sigma$ (by the maximality of i). Two cases may occur:

- *Subcase 1:* If $s_i s_{i+1}\sigma < s_{i+1}\sigma$, then we set $\tau = s_{i+1}\sigma < \sigma$ and $y = r_1 \cdots r_{l-1} r_n \tau \beta^{-1}$. Then $y = s_i x < x$ by 1.4. Moreover, still by 1.4, we have

$$s_{i-1}x = r_1 \cdots r_{l-1} r_n s_i \sigma \beta^{-1} > x$$

and

$$s_{i-1} s_i x = r_1 \cdots r_{l-1} r_n s_i s_{i+1} \sigma \beta^{-1} < s_i x.$$

So $x \sim_L y$ by Corollary 2.4. But, by the induction hypothesis (and since $s_{i+1}\tau > \tau$), we have $y \sim_L a_l \tau \beta^{-1}$. But $\sigma \sim_L \tau$ (again by Corollary 2.4 and since $s_i \tau < \tau < \sigma = s_{i+1} \tau < s_i \sigma$), so $a_l \sigma \beta^{-1} \sim_L a_l \tau \beta^{-1}$ by (*). This shows (?).

- *Subcase 2:* If $s_i s_{i+1}\sigma > s_{i+1}\sigma$, then $s_{i+1} s_i \sigma > s_i \sigma$ (by an easy application of Lemma 1.1) so, if we set $\tau = s_i \sigma$ and $y = r_1 \cdots r_{l-1} r_n \tau \beta^{-1}$, we have, by the induction hypothesis, $y \sim_L a_l \tau \beta^{-1}$. Moreover, $s_{i+1} \tau > \tau = s_i \sigma > \sigma > s_{i+1} \sigma$ and, by the same argument as in the subcase 1, we have $s_i y > y = s_{i-1} x > x > s_i x$. So $x \sim_L y$, $\sigma \sim_L \tau$. So it follows from (*) and $x \sim_L a_l \sigma \beta^{-1}$, as required.

- *Case 3:* If $\lambda \neq 1$, then we set $x = r_1 \cdots r_{l-1} r_n \sigma \beta^{-1}$ and $y = r_1 \cdots r_l \sigma \beta^{-1}$. We want to show that $x \sim_L y$. For this, let $x' = w_n x$, $y' = w_n y$, $\sigma' = \sigma_n \sigma \sigma_n^{-1} \sigma_{n-l}$ and $\beta' = \beta \sigma_n \sigma_{n-l}$. Then, by Proposition 1.18,

$$x' = r_l r_{l+1} \cdots r_{n-1} \sigma' \beta'^{-1} \quad \text{and} \quad y' = r_l r_{l+1} \cdots r_{n-2} r_n \sigma' \beta'^{-1}.$$

But, by Corollary 4.3, we have

$$x' \sim_L r_1 \cdots r_{n-l} \sigma' \beta'^{-1} \quad \text{and} \quad y' \sim_L r_1 \cdots r_{n-l-1} r_n \sigma' \beta'^{-1}.$$

Now, if we write $\sigma' = (\lambda', \mu')$, with $\lambda' \in \mathfrak{S}_{[1, n-l]}$ and $\mu' \in \mathfrak{S}_{[n-l+1, n]}$, we have $\mu' \neq \sigma_{[n-l+1, n]}$ (because $\lambda \neq 1$). So, by Case 2, we have

$$r_1 \cdots r_{n-l-1} r_n \sigma' \beta'^{-1} \sim_L r_1 \cdots r_{n-l} \sigma' \beta'^{-1}.$$

Therefore, $x' = w_n x \sim_L y' = w_n y$, and so $x \sim_L y$ by [10, Corollary 11.7]. \square

Corollary 4.4. *Let $l \in \{1, \dots, n\}$, let $1 \leq i_1 < \cdots < i_l \leq n$, let $\sigma \in \mathfrak{S}_n$, let $\beta \in Y_{l, n-l}$ and let $k \in [1, l]$ be such that $b \geq (i_k - 1)a$. Then*

$$r_{i_1} \cdots r_{i_l} \sigma \sim_L r_1 \cdots r_k r_{i_{k+1}} \cdots r_{i_l} \sigma.$$

Proof. The proof proceeds essentially as in Lemma 4.2. Let $w = r_{i_1} \cdots r_{i_l} \sigma$, let $w' = r_1 \cdots r_k r_{i_{k+1}} \cdots r_{i_l} \sigma$ and let us write $w = vx^{-1}$ and $w' = v'x'^{-1}$ with $v, v' \in W_{i_k}$ and $x, x' \in X_n^{(i_k)}$.

Since $w'w^{-1} = (r_1 \cdots r_k)^{-1}(r_{i_1} \cdots r_{i_k}) \in W_{i_k}$, we get that $x = x'$. The same argument as in Lemma 4.2 shows that $v = r_{i_1} \cdots r_{i_k} \tau$ and $v' = r_1 \cdots r_k \tau'$ for some $\tau, \tau' \in \mathfrak{S}_{i_k}$. But $v'v^{-1} = w'w^{-1} = (r_1 \cdots r_k)^{-1}(r_{i_1} \cdots r_{i_k})$, so $\tau = \tau'$. Now, by Proposition 4.1, $v \sim_L v'$. So $w \sim_L w'$ by [10, Proposition 9.11]. \square

5. PROOF OF PROPOSITION 2.15

Notation. *In this section, and only in this section, we assume that $1 \leq l \leq n-1$ and that $(n-2)a < b \leq (n-1)a$.*

We define a sequence $(C_j)_{l-1 \leq j \leq n-1}$ by induction as follows:

$$\begin{cases} C_{l-1} = 1, \\ C_l = C_{s_l}, \\ C_{j+1} = C_{s_{j+1}}C_j - C_{j-1}, \quad \text{if } l \leq j \leq n-2. \end{cases}$$

Let μ denote the coefficient of $C_{a_{l-1}\sigma_{[l,n]}}$ in the expansion of $C_{n-1}C_{a_l\sigma_{[l,n]}}$ in the Kazhdan-Lusztig basis. To prove Proposition 2.15, it is sufficient to show the following statement:

$$(5.1) \quad \mu = \begin{cases} 1 & \text{if } b = (n-1)a, \\ Q^{-1}q^{n-1} + Qq^{1-n} & \text{if } (n-2)a < b < (n-1)a. \end{cases}$$

Proof of 5.1. If $r \in \mathbb{Z}$, we set

$$\mathcal{H}_n[r] = \bigoplus_{\ell_t(w) \leq r} AT_w = \bigoplus_{\ell_t(w) \leq r} AC_w.$$

We shall show that

$$(5.2) \quad C_{n-1}C_{a_l\sigma_{[l,n]}} \equiv T_{s_{n-1} \cdots s_{l+1} s_l a_l \sigma_{[l,n]}} + Q^{-1}q^{n-1}T_{a_{l-1}\sigma_{[l,n]}} \pmod{\left(\mathcal{H}_n[l-2] + \mathcal{H}_n^{<0}\right)}.$$

The statement 5.2 will be proved at the end of this section. Let us conclude the proof of 5.1, assuming that 5.2 holds.

Let

$$\tilde{\mu} = \begin{cases} 1 & \text{if } b = (n-1)a, \\ Q^{-1}q^{n-1} + Qq^{1-n} & \text{if } (n-2)a < b < (n-1)a. \end{cases}$$

We want to show that $\mu = \tilde{\mu}$. But, by 5.2, we have

$$C_{n-1}C_{a_l\sigma_{[l,n]}} - C_{s_{n-1} \cdots s_{l+1} s_l a_l \sigma_{[l,n]}} - \tilde{\mu}C_{a_{l-1}\sigma_{[l,n]}} \in \mathcal{H}_n[l-2] + \mathcal{H}_n^{<0} + \bigoplus_{w < a_{l-1}\sigma_{[l+1,n]}} AT_w.$$

Since

$$\bigoplus_{w < a_{l-1}\sigma_{[l+1,n]}} AT_w = \bigoplus_{w < a_{l-1}\sigma_{[l+1,n]}} AC_w,$$

there exists a family $(\nu_w)_{\ell_t(w) \leq l-2 \text{ or } w < a_{l-1}\sigma_{[l+1,n]}}$ of elements of $A_{\geq 0}$ such that

$$C_{n-1}C_{a_l\sigma_{[l,n]}} - C_{s_{n-1}\cdots s_{l+1}s_l a_l\sigma_{[l,n]}} - \tilde{\mu}C_{a_{l-1}\sigma_{[l,n]}} - \sum_{\substack{\ell_t(w) \leq l-2 \\ \text{or } a_{l-1}\sigma_{[l+1,n]}}} \nu_w C_w \in \mathcal{H}_n^{<0}.$$

Let $\nu'_w = \nu_w + \bar{\nu}_w - \tau_A(\nu_w)$. Then

$$C_{n-1}C_{a_l\sigma_{[l,n]}} - C_{s_{n-1}\cdots s_{l+1}s_l a_l\sigma_{[l,n]}} - \tilde{\mu}C_{a_{l-1}\sigma_{[l,n]}} - \sum_{\substack{\ell_t(w) \leq l-2 \\ \text{or } a_{l-1}\sigma_{[l+1,n]}}} \nu'_w C_w \in \mathcal{H}_n^{<0}$$

and $\bar{\nu}'_w = \nu_w$. So, if we set

$$C = C_{n-1}C_{a_l\sigma_{[l,n]}} - C_{s_{n-1}\cdots s_{l+1}s_l a_l\sigma_{[l,n]}} - \tilde{\mu}C_{a_{l-1}\sigma_{[l,n]}} - \sum_{\substack{\ell_t(w) \leq l-2 \\ \text{or } a_{l-1}\sigma_{[l+1,n]}}} \nu'_w C_w,$$

then

$$\bar{C} = C \quad \text{and} \quad C \in \mathcal{H}_n^{<0}.$$

So $C = 0$ by [10, Theorem 5.2], and so $\mu = \tilde{\mu}$, as expected. \square

So it remains to prove the statement 5.2:

Proof of 5.2. First of all, we have $C_{a_l\sigma_{[l,n]}} = C_{a_l}C_{\sigma_{[l,n]}}$, since the supports of a_l and $\sigma_{[l,n]}$ (in S_n) are disjoint. Moreover, since $l \leq n-1$ (i.e. $a_l \in W_{n-1}$) and $b > (n-2)a$, it follows from [1, Propositions 2.5 and 5.1] that

$$(5.3) \quad C_{a_l} = (T_{t_1} + Q^{-1})(T_{t_2} + Q^{-1}) \cdots (T_{t_l} + Q^{-1})T_{\sigma_l}^{-1}.$$

Let $\mathcal{H}(\mathfrak{S}_n)$ denote the sub- A -algebra of \mathcal{H}_n generated by $T_{s_1}, \dots, T_{s_{n-1}}$. It is the Hecke algebra of \mathfrak{S}_n (with parameter a). Then $\mathcal{H}_n[l-2]$ is a sub- A -module of \mathcal{H}_n . Therefore, it follows from 5.3 that

$$C_{a_l} \equiv (T_{w_l} + Q^{-1} \sum_{1 \leq i \leq l} T_{t_1 \cdots t_{i-1} t_{i+1} \cdots t_l}) T_{\sigma_l}^{-1} \pmod{\mathcal{H}_n[l-2]}.$$

But, if $1 \leq i \leq l$, then

$$t_1 \cdots t_{i-1} t_{i+1} \cdots t_l = s_i s_{i+1} \cdots s_{l-1} a_{l-1} \sigma_{l-1} s_{l-1} \cdots s_{i+1} s_i,$$

and $\sigma_l = s_{l+1-i} \cdots s_{l-2} s_{l-1} \sigma_{l-1} s_{l-1} \cdots s_{i+1} s_i$. Moreover,

$$\ell(\sigma_l) = \ell(s_{l+1-i} \cdots s_{l-2} s_{l-1}) + \ell(\sigma_{l-1} s_{l-1} \cdots s_{i+1} s_i).$$

Therefore,

$$C_{a_l} \equiv T_{a_l} + Q^{-1} \sum_{1 \leq i \leq l} T_{s_i s_{i+1} \cdots s_{l-1}} T_{a_{l-1}} (T_{s_{l+1-i} \cdots s_{l-2} s_{l-1}})^{-1} \pmod{\mathcal{H}_n[l-2]}.$$

Finally, we get

$$C_{a_l\sigma_{[l,n]}} \equiv T_{a_l}C_{\sigma_{[l,n]}} + Q^{-1} \sum_{1 \leq i \leq l} T_{c_{[i,l-1]}} T_{a_{l-1}} (T_{c_{[l+1-i,l-1]}})^{-1} C_{\sigma_{[l,n]}} \pmod{\mathcal{H}_n[l-2]}.$$

Now, if $l-1 \leq j \leq n-1$, then

$$(5.4) \quad \begin{aligned} C_j C_{a_l\sigma_{[l,n]}} &\equiv \sum_{i=l-1}^j q^{i-j} T_{d_{[l,i]}} T_{a_l} C_{\sigma_{[l,n]}} \\ &+ Q^{-1} C_j \sum_{1 \leq i \leq l} T_{c_{[i,l-1]}} T_{a_{l-1}} (T_{c_{[l+1-i,l-1]}})^{-1} C_{\sigma_{[l,n]}} \pmod{\mathcal{H}_n[l-2]}. \end{aligned}$$

Proof of 5.4. We shall argue by induction on j . The cases where $j = l-1$ or $j = l$ are obvious. So assume that $j \in [l, n-2]$ and that 5.4 holds for j . By the induction hypothesis, we get

$$\begin{aligned} C_{j+1} C_{a_l\sigma_{[l,n]}} &\equiv C_{s_{j+1}} \sum_{i=l-1}^j q^{i-j} T_{d_{[l,i]}} T_{a_l} C_{\sigma_{[l,n]}} - \sum_{i=l-1}^{j-1} q^{i-j+1} T_{d_{[l,i]}} T_{a_l} C_{\sigma_{[l,n]}} \\ &+ Q^{-1} C_{j+1} \sum_{1 \leq i \leq l} T_{c_{[i,l-1]}} T_{a_{l-1}} (T_{c_{[l+1-i,l-1]}})^{-1} C_{\sigma_{[l,n]}} \pmod{\mathcal{H}_n[l-2]}. \end{aligned}$$

Now,

$$C_{s_{j+1}} T_{d_{[l,j]}} T_{a_l} C_{\sigma_{[l,n]}} = T_{d_{[l,j+1]}} T_{a_l} C_{\sigma_{[l,n]}} + q^{-1} T_{d_{[l,j]}} T_{a_l} C_{\sigma_{[l,n]}}$$

and, if $l-1 \leq i < j$, then

$$C_{s_{j+1}} T_{d_{[l,i]}} T_{a_l} C_{\sigma_{[l,n]}} = T_{d_{[l,i]}} T_{a_l} C_{s_{j+1}} C_{\sigma_{[l,n]}} = (q + q^{-1}) T_{d_{[l,i]}} T_{a_l} C_{\sigma_{[l,n]}}.$$

Now 5.4 follows from a straightforward computation. \square

Since $d_{[l,i]} \in Y_{l,n-l}$, we have

$$T_{d_{[l,i]}} T_{a_l} C_{\sigma_{[l,n]}} = T_{d_{[l,i]a_l}} C_{\sigma_{[l,n]}} \equiv T_{d_{[l,i]a_l\sigma_{[l,n]}}} \pmod{\mathcal{H}_n^{<0}},$$

so, by 5.4, we get

$$\begin{aligned} C_{n-1} C_{a_l\sigma_{[l,n]}} &\equiv T_{d_{[l,n-1]a_l\sigma_{[l,n]}}} \\ &+ Q^{-1} C_{n-1} \sum_{1 \leq i \leq l} T_{c_{[i,l-1]}} T_{a_{l-1}} (T_{c_{[l+1-i,l-1]}})^{-1} C_{\sigma_{[l,n]}} \pmod{(\mathcal{H}_n[l-2] + \mathcal{H}_n^{<0})}. \end{aligned}$$

For $1 \leq i \leq l$, let $\mathcal{X}_i = Q^{-1} C_{n-1} T_{c_{[i,l-1]}} T_{a_{l-1}} (T_{c_{[l+1-i,l-1]}})^{-1} C_{\sigma_{[l,n]}}$. There exists a family $(f_I)_{I \subseteq [l,n-1]}$ of elements of \mathbb{Z} such that $C_{n-1} = \sum_{I \subseteq [l,n-1]} f_I C_{d_I}$. Moreover, $f_{[l,n-1]} = 1$. Also,

$$(T_{c_{[l+1-i,l-1]}})^{-1} = \sum_{J \subseteq [l+1-i,l-1]} (q - q^{-1})^{i-1-|J|} T_{c_J}.$$

Therefore,

$$\mathcal{X}_i = \sum_{\substack{I \subseteq [l, n-1] \\ J \subseteq [l+1-i, l-1]}} f_I Q^{-1}(q - q^{-1})^{i-1-|J|} C_{d_I} T_{c_{[i, l-1]}} T_{a_{l-1}} T_{c_J} C_{\sigma_{[l, n]}}.$$

Let $\Delta_{i, I, J} = f_I Q^{-1}(q - q^{-1})^{i-1-|J|} C_{d_I} T_{c_{[i, l-1]}} T_{a_{l-1}} T_{c_J} C_{\sigma_{[l, n]}}$. If we express $\Delta_{i, I, J}$ in the standard basis $(T_w)_{w \in W_n}$, then the degree of the coefficients are bounded by $-b + (i - 1 - |J| + |I|)a$. Since $b > (n - 2)a$, this degree is in $\Gamma_{<0}$, except if $i = l$, $J = \emptyset$ and $I = [l, n - 1]$. Therefore,

$$C_{n-1} C_{a_l \sigma_{[l, n]}} \equiv T_{d_{[l, n-1]} a_l \sigma_{[l, n]}} + \Delta_{l, [l, n-1], \emptyset} \pmod{(\mathcal{H}_n[l - 2] + \mathcal{H}_n^{<0})}.$$

But

$$\begin{aligned} \Delta_{l, [l, n-1], \emptyset} &= Q^{-1}(q - q^{-1})^{l-1} C_{s_{n-1} \cdots s_l} T_{a_{l-1}} C_{\sigma_{[l, n]}} \\ &= Q^{-1}(q - q^{-1})^{l-1} T_{a_{l-1}} C_{s_{n-1} \cdots s_l} C_{\sigma_{[l, n]}} \\ &= Q^{-1}(q - q^{-1})^{l-1} (q + q^{-1})^{n-l} T_{a_{l-1}} C_{\sigma_{[l, n]}}, \end{aligned}$$

the last equality following from Theorem 2.3 (a). So $\Delta_{l, [l, n-1], \emptyset} \equiv Q^{-1} q^{n-1} T_{a_{l-1}} C_{\sigma_{[l, n]}} \pmod{\mathcal{H}_n^{<0}}$. The proof of 5.2 is complete. \square

6. CONSEQUENCES OF PROPOSITION 2.15

The aim of this section is to prove the following proposition:

Proposition 6.1. *Let $l \in \{1, \dots, n\}$ and assume that $b \leq (n - 1)a$. Then*

$$s_1 s_2 \cdots s_{n-1} a_{l-1} \sigma_{[l, n-1]} \sim_L t s_1 s_2 \cdots s_{n-1} a_{l-1} \sigma_{[l, n-1]}.$$

Proof. Let $u_{l, n} = t s_1 s_2 \cdots s_{n-1} a_{l-1} \sigma_{[l, n-1]} = t s_1 \cdots s_{l-1} a_{l-1} s_l \cdots s_{n-1} \sigma_{[l, n-1]} = a_l \sigma_{[l, n]}$. We need to show that $t u_{l, n} \sim_L u_{l, n}$ (note that $t u_{l, n} \leq u_{l, n}$). We shall argue by induction on n , the cases where $n = 1$ or 2 being obvious. So assume that $n \geq 3$ and that $t u_{l, n-1} \sim_L u_{l, n-1}$ if $b \leq (n - 2)a$.

First, assume that $b \leq (n - 2)a$. Then

$$\begin{cases} u_{l, n} = u_{l, n-1} s_{n-1} \cdots s_{l+1} s_l & \text{if } l \leq n - 1, \\ u_{l, n} = a_n = u_{l-1, n-1} s_{n-1} \cdots s_2 s_1 t & \text{if } l = n. \end{cases}$$

By the induction hypothesis, we have $t u_{k, n-1} \sim_L u_{k, n-1}$ so, since $s_{n-1} \cdots s_{l+1} s_l$ and $s_{n-1} \cdots s_2 s_1 t$ belong to X_n^{-1} , it follows from [10, Proposition 9.11] that $t u_{l, n} \sim_L u_{l, n}$.

This means that we may, and we will, assume that $(n - 2)a < b \leq (n - 1)a$. But, by Proposition 2.15, we have $a_{l-1} \sigma_{[l, n]} \leq_L a_l \sigma_{[l, n]} = u_{l, n}$. On the other hand,

$$t u_{l, n} = c_{[1, l-1]} a_{l-1} \sigma_{[l, n]} \leq_L c_{[2, l-1]} a_{l-1} \sigma_{[l, n]} \leq_L \cdots \leq_L s_{l-1} a_{l-1} \sigma_{[l, n]} \leq_L a_{l-1} \sigma_{[l, n]}.$$

So $t u_{l, n} \sim_L u_{l, n}$, as desired. \square

REMARK 6.2 - Note that the converse of Proposition 6.1 also holds. Indeed, if $b > (n-1)a$ and if $x \sim_L y$ for some x and y in W_n , then $\ell_t(x) = \ell_t(y)$ (see [4, Theorem 7.7] and [1, Corollary 5.2]). \square

Corollary 6.3. *Let $l \in \{1, 2, \dots, n\}$ and let $\beta \in Y_{l-1, n-l}$. Then*

$$s_1 s_2 \cdots s_{n-1} a_{l-1} \sigma_{[l, n-1]} \beta^{-1} \sim_L t s_1 s_2 \cdots s_{n-1} a_{l-1} \sigma_{[l, n-1]} \beta^{-1}.$$

Proof. Let $w = s_1 s_2 \cdots s_{n-1} a_l \sigma_{[l+1, n-1]} \beta^{-1}$. We want to show that $w \sim_L tw$. We shall argue by induction on $\ell(\beta)$. If $\ell(\beta) = 0$ (i.e. $\beta = 1$), this is just the proposition 6.1. Sp we assume now that $\ell(\beta) \geq 1$. We shall use the $*$ -operation (see §2.C). For this, we need to study the action of the γ_i 's on w , when possible.

We have $\sigma_{[l, n-1]} a_{l-1} = a_{l-1} \sigma_{[l, n-1]}$, so

$$w = s_1 s_2 \cdots s_{n-1} \sigma_{[l, n-1]} (a_{l-1} \beta)^{-1} = s_1 s_2 \cdots s_{l-1} \sigma_{[l, n]} (a_{l-1} \beta)^{-1}.$$

Let $1 \leq j_1 < \cdots < j_{l-1} \leq n-1$ be the unique sequence such that $a_l \beta = r_{j_1} r_{j_2} \cdots r_{j_{l-1}}$. Since $\ell(\beta) > 0$, we have $(j_1, j_2, \dots, j_{l-1}) \neq (1, 2, \dots, l-1)$, so there exists $k \in [1, l-1]$ such that $j_k - j_{k-1} \geq 2$ (where $j_0 = 0$ by convention). Note that $j_k < n$ so $j_k + 1 \in [2, n]$. We have, by 1.16

$$\begin{aligned} w(j_k) s_1 \cdots s_{l-1} \sigma_{[l, n]} (r_{j_1} \cdots r_{j_{l-1}})^{-1} (j_k) &= s_1 \cdots s_{l-1} \sigma_{[l, n]} (k-l) \\ &= -s_1 \cdots s_{l-1} (l-k) \\ &= -(l+1-k) < 0 \end{aligned}$$

and

$$\begin{aligned} w(j_k - 1) s_1 \cdots s_{l-1} \sigma_{[l, n]} (r_{j_1} \cdots r_{j_{l-1}})^{-1} (j_k - 1) &= s_1 \cdots s_{l-1} \sigma_{[l, n]} (l+q) \\ &= s_1 \cdots s_{l-1} (n+1-q) \\ &= n+1-q > 0 \end{aligned}$$

for some $q \in [1, n+1-l]$. Moreover, a similar computation shows that (with the convention that $j_l = n+1$)

$$w(j_k) = \begin{cases} -(l-k) & \text{if } j_{k+1} = j_k + 1, \\ n-q & \text{if } j_{k+1} \geq j_k + 2. \end{cases}$$

In any case, we have

$$w(j_k) < w(j_k + 1) < w(j_k - 1).$$

This shows that

$$w s_{j_k-1} s_{j_k} < w s_{j_k-1} < w < w s_{j_k},$$

So $w \in \mathcal{D}_{j_k-1}(W_n)$ and $\gamma_{j_k-1}(w) = w s_{j_k-1} < w$. Now, let $\beta' = s_{j_k} \beta$. An easy computation as above shows that $\beta' < \beta$, so that $\beta' \in Y_{l, n-1-l}$ by Deodhar's Lemma

(see [6, Lemma 2.1.2]). So $\gamma_{j_k}(w) = s_1 \cdots s_{n-1} a_{l-1} \sigma_{[l, n-1]} \beta'^{-1}$ where $\beta' \in Y_{l, n-1-l}$ is such that $\ell(\beta') = \ell(\beta) - 1$. But, by Corollary 2.13, we have $t\gamma_i(w) = \gamma_i(tw)$. So, by 2.11 and by the induction hypothesis, we get that $w \sim_L tw$, as desired. \square

7. PROOF OF THEOREM 1.24

7.A. Knuth relations. By recent results of Taskin [13, Theorems 1.2 and 1.3], the equivalence relations \sim_R^r and \simeq_R^r can be described using generalisations of Knuth relations (for the relation \simeq_R^r , a similar result has been obtained independently by Pietraho [12, Theorems 3.8 and 3.9] using other kinds of Knuth relations). We shall recall here Taskin's construction. For this, we shall need the following notation: if $0 \leq r \leq n-2$, we denote by $\mathcal{E}_n^{(r)}$ the set of elements $w \in W_n$ such that $|w(1)| > |w(i)|$ for $i \in \{2, 3, \dots, r+2\}$ and such that the sequence $(w(2), w(3), \dots, w(r+2))$ is a shuffle of a positive decreasing sequence and a negative increasing sequence. If $r \geq n-1$, we set $\mathcal{E}_n^{(r)} = \emptyset$. Following [13, Definition 1.1], we introduce three relations which will be used to generate the relations \sim_R^r and \simeq_R^r .

Let $w, w' \in W_n$ and let $r \geq 0$:

- We write $w \smile_1 w'$ if there exists $i \geq 2$ (respectively $i \leq n-2$) such that $w(i) < w(i-1) < w(i+1)$ (respectively $w(i) < w(i+2) < w(i+1)$) and $w' = ws_i$.
- We write $w \smile_2^r w'$ if there exists $i \leq \min(r, n-1)$ such that $w(i)w(i+1) < 0$ and $w' = ws_i$. The relation \smile_2^0 never occurs.
- We write $w \smile_3^r w'$ if $w \in \mathcal{E}_n^{(r)}$ and $w' = wt$. If $r \geq n-1$, the relation \smile_3^r never occurs.

REMARK - If $w \smile_2^r w'$, then $w \smile_2^{r+1} w'$. If $w \smile_3^r w'$, then $w \smile_3^{r-1} w'$ (indeed, $\mathcal{E}_n^{(r)} \subseteq \mathcal{E}_n^{(r-1)}$). \square

Taskin's Theorem. *With the above notation, we have:*

- (a) *The relation \sim_R^r is the equivalence relation generated by the relations \smile_1 , \smile_2^r and \smile_3^r .*
- (b) *The relation \simeq_R^r is the equivalence relation generated by the relations \smile_1 , \smile_2^r and \smile_3^{r-1} .*

7.B. Proof of Theorem 1.24. Recall that the relation \sim_{LR}^r (respectively \simeq_{LR}^r) is the equivalence relation generated by \sim_L^r and \sim_R^r (respectively \simeq_L^r and \simeq_R^r). Recall also that $x \sim_L^r y$ (respectively $x \simeq_L^r y$, respectively $x \sim_L y$) if and only if $x^{-1} \sim_R^r y^{-1}$ (respectively $x^{-1} \simeq_R^r y^{-1}$, respectively $x^{-1} \sim_R y^{-1}$). So it is sufficient to show that Theorem 1.24 holds whenever $? = R$. It is then easy to see that Theorem 1.24 will

follow from Taskin's Theorem and from the following three lemmas (which will be proved in subsections 7.C, 7.D and 7.E).

Lemma 7.1. *Let $w, w' \in W_n$ be such that $w \smile_1 w'$. Then $w \sim_R w'$.*

Lemma 7.2. *Let $w, w' \in W_n$ and let $r \geq 0$ be such that $b \geq ra$ and $w \smile_2^r w'$. Then $w \sim_R w'$.*

Lemma 7.3. *Let $w \in W_n$ and let $r \geq 0$ be such that $b \leq (r+1)a$ and $w \smile_3^r w'$. Then $w \sim_R w'$.*

7.C. Proof of Lemma 7.1. Let $w, w' \in W_n$ be such that $w \smile_1 w'$. Let $i \in I_{n-1}^+$ be such that $w' = ws_i$. Then $i \geq 2$ and $w(i) < w(i-1) < w(i+1)$, or $i \leq n-2$ and $w(i) < w(i+2) < w(i+1)$. In the first case, we have $ws_i s_{i-1} > ws_i > w > ws_{i-1}$ while, in the second case, we have $ws_i s_{i+1} > ws_i > w > ws_{i+1}$. So $w' = ws_i \sim_R w$ by 1.23 and Corollary 2.4. The proof of Lemma 7.1 is complete.

7.D. Proof of Lemma 7.2. Let $w, w' \in W_n$ and let $r \geq 0$ be such that $b \geq ra$ and $w \smile_2^r w'$. Let $i \in I_{n-1}^+$ be the element such that $w' = ws_i$. Then $i \leq r$ and $w(i)w(i+1) < 0$. By exchanging w and w' if necessary, we may assume that $w(i) < 0$ and $w(i+1) > 0$.

Let us write $w = xv$, with $x \in X_n^{(i+1)}$ and $v \in W_{i+1}$. Then $vs_i \in W_{i+1}$ and $ws_i = xvs_i$. Therefore, by [10, Proposition 9.11], we only need to show that $vs_i \sim_L v$. But $0 < x(1) < \dots < x(r+1)$ (see Lemma 1.19), and $v(j) \in I_{i+1}$ for all $j \in I_{i+1}$. So $v(i) > 0$ and $v'(i+1) < 0$. In particular, $v \smile_2^r vs_i$ (and even $v \smile_2^i v'$). This means that we may (and we will) assume that $i = n-1$. So we have

$$b \geq (n-1)a, \quad w(n-1) < 0 \quad \text{and} \quad w(n) > 0,$$

and we want to show that $w \sim_R ws_{n-1}$ or, in other words, that

$$(?) \quad w^{-1} \sim_L s_{n-1} w^{-1}.$$

Let $\alpha = \alpha_{w^{-1}}$, $\sigma = \sigma_{w^{-1}}$ and $\beta = \beta_{w^{-1}}$. Then

$$w^{-1} = \alpha a_l \sigma a_l \beta^{-1}.$$

By Lemma 1.15, there exists a unique sequence $1 \leq i_1 < \dots < i_l \leq n$ such that $\alpha a_l = r_{i_1} \dots r_{i_l}$ so

$$w^{-1} = r_{i_1} \dots r_{i_l} \sigma \beta^{-1}.$$

But, again by Lemma 1.15, we have $w^{-1}(i) < 0$ if and only if $i \in \{i_1, \dots, i_l\}$. So

$$i_l = n-1.$$

So

$$w^{-1} = r_{i_1} \cdots r_{i_{n-1}} r_{n-1} \sigma \beta^{-1}.$$

and

$$s_{n-1} w^{-1} = r_{i_1} \cdots r_{i_{n-1}} r_n \sigma \beta^{-1}.$$

So the result follows from Proposition 4.1.

7.E. Proof of Lemma 7.3. Let $w \in W_n$ and let $r \geq 0$ be such that $b \leq (r+1)a$ and $w^{-1} \smile_3^r w'^{-1}$. We want to show that $w \sim_L w' = tw$. The proof goes through several steps.

First step: easy reductions. First, note that $r \leq n-2$. Let us write $w = vx^{-1}$, with $v \in W_{r+2}$ and $x \in X_n^{(r+2)}$. Then $0 < x(1) < \cdots < x(r+2)$ by Lemma 1.19, so $v^{-1} \in \mathcal{E}_{r+2}^{(r)}$. Then $tw = (tv)x^{-1}$ with $tv \in W_{r+2}$ so, by [10, Proposition 9.11], it is sufficient to show that $tv \sim_L v$. This shows that we may (and we will) assume that $r = n-2$.

By [10, Corollary 11.7], this is equivalent to show that $tw_n w \sim_L w_n w$. Since $w_n w \in \mathcal{E}_n^{(n-2)}$ we may, by replacing w by tw , $w_n w$ or $tw_n w$, assume that $w^{-1}(1) > 0$ and $w^{-1}(n) > 0$. Since moreover $|w^{-1}(1)| > |w^{-1}(i)|$ for all $i \in \{2, 3, \dots, n = r+2\}$, we have $w^{-1}(1) = n$.

As a conclusion, we are now working under the following hypothesis:

Hypothesis. *From now on, and until the end of this subsection, we assume that*

- (1) $w^{-1}(1) = n$ and $w^{-1}(n) > 0$, and
- (2) $w^{-1} \in \mathcal{E}_n^{(n-2)}$.

And recall that we want to show that

$$(?) \quad tw \sim_L w.$$

Second step: decomposition of w . Let $v = s_{n-1} \cdots s_2 s_1 w$. Then $v^{-1}(n) = w^{-1}(1) = n$ by (3), so $v \in W_{n-1}$. Therefore,

$$(7.4) \quad w = s_1 s_2 \cdots s_{n-1} v, \quad s_1 s_2 \cdots s_{n-1} \in X_n \quad \text{and} \quad v \in W_{n-1}.$$

Note that

$$(7.5) \quad v^{-1}(k) = w^{-1}(k+1)$$

for all $k \in [1, n-1]$, so that

$$(7.6) \quad v \in \mathcal{E}_{n-1}^{(n-3)}$$

and, by (2),

$$(7.7) \quad v^{-1}(n-1) > 0.$$

Let us write $v = r_{i_1} \cdots r_{i_l} \sigma \beta^{-1}$, with $l = \ell_t(v) = \ell_t(w)$, $1 \leq i_1 < \cdots < i_l \leq n-1$, $\sigma \in \mathfrak{S}_{l, n-1-l}$ and $\beta \in Y_{l, n-1-l}$. By 7.7 and Lemma 1.15, we have

$$(7.8) \quad i_l \leq n-2.$$

Finally, note that

$$(7.9) \quad \sigma = \sigma_{[l+1, n-1]}.$$

Proof of 7.9. By 7.6, we have $|v^{-1}(i_1)| > |v^{-1}(i_2)| > \cdots > |v^{-1}(i_l)|$. Therefore, it follows from 1.16 that $\beta(\sigma^{-1}(l)) > \beta(\sigma^{-1}(l-1)) > \cdots > \beta(\sigma^{-1}(1))$. Since σ stabilizes the interval $[1, l]$ and since β is increasing on $[1, l]$ (because it lies in $Y_{l, n-l}$), this forces $\sigma(k) = k$ for all $k \in [1, l]$.

Similarly, if $1 \leq j_1 < \cdots < j_{n-l} \leq n$ denotes the unique sequence such that $[1, n] = \{i_1, \dots, i_l\} \cup \{j_1, \dots, j_{n-l}\}$, then $|v^{-1}(j_1)| > |v^{-1}(j_2)| > \cdots > |v^{-1}(j_{n-l})|$ by 7.6. So it follows from 1.16 that $\beta(\sigma^{-1}(l+1)) > \beta(\sigma^{-1}(l+2)) > \cdots > \beta(\sigma^{-1}(n))$ and, since σ stabilizes the interval $[l+1, n]$ and β is increasing on the same interval, this forces $\sigma(l+k) = n+1-k$ for $k \in [1, n-l]$. \square

Third step: conclusion. We first need the following elementary result:

$$(7.10) \quad s_1 s_2 \cdots s_{n-1} r_{i_1} \cdots r_{i_l} = r_{i_1+1} \cdots r_{i_l+1} s_{l+1} s_{l+2} \cdots s_{n-1}.$$

Proof of 7.10. This follows easily from 1.3 or from 1.4. \square

Now, let $\tau = s_{l+1} s_{l+2} \cdots s_{n-1} \sigma_{[l+1, n]} \beta^{-1} = \sigma_{[l+1, n]} \beta^{-1} \in \mathfrak{S}_n$. Then, by 7.10, we have

$$w = r_{i_1+1} r_{i_2+1} \cdots r_{i_l+1} \tau \quad \text{and} \quad tw = r_1 r_{i_1+1} r_{i_2+1} \cdots r_{i_l+1} \tau.$$

By 7.8, we have $b \geq (i_l + 1 - 1)a$, so, by Corollary 4.4, we have

$$w \sim_L r_2 r_3 \cdots r_{l+1} \tau \quad \text{and} \quad tw \sim_L r_1 r_2 \cdots r_{l+1} \tau.$$

So we only need to show that $r_2 r_3 \cdots r_{l+1} \tau \sim_L r_1 r_2 \cdots r_{l+1} \tau = tr_2 r_3 \cdots r_{l+1} \tau$. But $r_2 \cdots r_{l+1} \sigma_{[l+1, n]} \beta^{-1} s_1 \cdots s_{n-1} a_l \sigma_{[l+1, n-1]} \beta^{-1}$, with $\beta \in Y_{l, n-1-l}$. So the result follows from Corollary 6.3.

The proof of Lemma 7.3 is complete, as well as the proof of Theorem 1.24.

REFERENCES

- [1] C. BONNAFÉ, Two-sided cells in type B (asymptotic case), *J. Algebra* **304** (2006), 216–236.
- [2] C. BONNAFÉ, Semicontinuity properties of Kazhdan-Lusztig cells, preprint (2008), available at [arXiv:0808.3522](https://arxiv.org/abs/0808.3522).
- [3] C. BONNAFÉ, M. GECK, L. IANCU & T. LAM, On domino insertion and Kazhdan-Lusztig cells in type B_n , to appear in *Progress in Mathematics (Birkhäuser)*, available at [arXiv:math/0609279](https://arxiv.org/abs/math/0609279).
- [4] C. BONNAFÉ & L. IANCU, Left cells in type B_n with unequal parameters, *Represent. Theory* **7** (2003), 587–609.

- [5] M. GECK, *On the induction of Kazhdan-Lusztig cells*, Bull. London Math. Soc. **35** (2003), 608–614.
- [6] M. GECK AND G. PFEIFFER, *Characters of finite Coxeter groups and Iwahori–Hecke algebras*, London Math. Soc. Monographs, New Series **21**, Oxford University Press, 2000.
- [7] D. A. KAZHDAN AND G. LUSZTIG, Representations of Coxeter groups and Hecke algebras, *Invent. Math.* **53** (1979), 165–184.
- [8] T. LAM, Growth diagrams, domino insertion, and sign-imbalance, *J. Comb. Theory Ser. A.* **107** (2004), 87–115.
- [9] G. LUSZTIG, *Left cells in Weyl groups*, Lie Group Representations I (R. L. R. Herb and J. Rosenberg, eds.), Lecture Notes in Math., vol. 1024, Springer-Verlag, 1983, pp. 99–111.
- [10] G. LUSZTIG, *Hecke algebras with unequal parameters*, CRM Monograph Series **18**, American Mathematical Society, Providence, RI (2003), 136 pp.
- [11] T. PIETRAHO, Notes on combinatorial cells and constructible representations in type B , preprint (2007), available on [arXiv:0710.3846](https://arxiv.org/abs/0710.3846).
- [12] T. PIETRAHO, Knuth relations for the hyperoctahedral groups, preprint (2008), available at [arXiv:0803.3335](https://arxiv.org/abs/0803.3335).
- [13] M. TASKIN, Plactic relations for r -domino tableaux, preprint (2008), available at [arXiv:0803.1148](https://arxiv.org/abs/0803.1148).
- [14] M. VANLEEUVEN, The Robinson-Schensted and Schutzenberger algorithms, an elementary approach, The Foata Festschrift, *Electron. J. Combin.* **3** (1996), Research Paper 15.
- [15] M. SHIMOZONO AND D. E. WHITE, Color-to-spin ribbon Schensted algorithms, *Discrete Math.* **246** (2002), 295–316.

LABO. DE MATH. DE BESANÇON (CNRS: UMR 6623), UNIVERSITÉ DE FRANCHE-COMTÉ,
16 ROUTE DE GRAY, 25030 BESANÇON CEDEX, FRANCE
E-mail address: cedric.bonnafe@univ-fcomte.fr