Metal and metalloid concentrations in the giant squid *Architeuthis dux* from Iberian waters

Paco Bustamante, Angel Gonzalez, Francisco Rocha, Pierre Miramand, Angel
Guerra

► To cite this version:

Paco Bustamante, Angel Gonzalez, Francisco Rocha, Pierre Miramand, Angel Guerra. Metal and metalloid concentrations in the giant squid *Architeuthis dux* from Iberian waters. *Marine Environmental Research*, 2008, 66 (2), pp.278-287. 10.1016/j.marenvres.2008.04.003 . hal-00289572

HAL Id: hal-00289572

<https://hal.science/hal-00289572>

Submitted on 22 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Metal and metalloid concentrations in the giant squid *Architeuthis dux* from
Iberian waters**

Bustamante P^{1*}, González AF², Rocha F³, Miramand P¹, Guerra A²

1. Littoral, Environnement et Sociétés (LIENSs), UMR 6250, CNRS-Université de La
Rochelle, 2 rue Olympe de Gouges, F-17042 La Rochelle cedex 01, France

2. Instituto de Investigaciones Marinas (CSIC), ECOBIOMAR, Eduardo Cabello 6, 36208
Vigo, Spain

3. Departamento de Ecología y Biología Animal, Facultad de Ciencias, Campus As Lagoas-
Marcosende, Universidad de Vigo, 36310 Vigo, Spain

* Corresponding author; Phone : +33 546 507 625 ; Fax : +33 546 458 264; E-mail : pbustama@univ-lr.fr

Abstract: This study investigated 14 trace elements (Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, V and Zn) in the tissues of the giant squid *Architeuthis dux* from the Mediterranean and Atlantic Spanish waters. As for other families of cephalopods, the digestive gland and the branchial hearts of *Architeuthis* showed the highest concentrations of Ag, Cd, Co, Cu, Fe, Ni, Se, V and Zn, highlighting their major role in the bioaccumulation and detoxification processes. With the exception of Hg, the muscles showed relatively low trace element concentrations. Nevertheless, this tissue contained the main proportion of the total As, Cr, Hg, Mn, Ni, and Zn body burden because muscles represent the main proportion of the squid mass. These findings suggest that the metal metabolism is overall the same as other cephalopod families from neritic waters. In females, Zn concentrations increased in the digestive gland with the squid's weight likely reflecting physiological changes during sexual maturation. Comparing the trace element concentrations in the tissues of *Architeuthis*, higher Ag, Cu, Hg and Zn concentrations in the squid from the Mediterranean reflected different exposure conditions. In comparison to other meso-pelagic squids from the Bay of Biscay, Cd concentrations recorded in the digestive gland suggest that *Architeuthis* might feed on more contaminated prey or that it displays a longer life span than other cephalopods.

Key words: trace element; bioaccumulation; cephalopod; giant squid, *Architeuthis*

1. Introduction

Cephalopods play a key role in marine ecosystems both as predators and prey. They constitute a class of marine molluscs which are found in a great variety of habitats from coastal waters to very deep-ocean environments (Boyle and Rodhouse, 2005). Independently of the species, the habitat or the life span they display, cephalopods share the ability to accumulate inorganic and organic pollutants such as metals, PCBs or organochlorine pesticides (e.g. Martin and Flegal, 1975; Tanabe et al., 1984; Miramand and Bentley, 1992; Yamada et al., 1997; Bustamante et al., 2000, 2006a; Ueno et al., 2003; Storelli et al., 2006). Consequently, they were reported to constitute a significant vector of contaminants to the species feeding on them, in particular seabirds and marine mammals (e.g. Honda et al., 1983; Muirhead and Furness 1988; Bustamante et al., 1998; Lahaye et al., 2005). However, most studies focused on commercially targeted species which are 1) easy to sample and 2) of high economic and health interest concerning human consumption. Many of these cephalopod species are also consumed by top marine predators, but there is a gap in the information concerning non-targeted species. This lack of data on bioaccumulation of contaminants is particularly obvious for oceanic and deep-sea species, like the giant squid *Architeuthis*.

Overall, the biology, behaviour and life cycle of *Architeuthis* are still poorly known in many aspects even if this squid has received considerable attention over the last decade (see González et al., 2002; Guerra et al., 2004, 2006; Kubodera and Mori 2005). Most of the information available on this squid comes from dead stranded animals and from predator trophic ecology studies. Indeed, squid flesh and beaks are often recorded in the stomach of sperm whales, but also of seabirds and sharks (Roper and Boss 1982; Clarke 1996; Santos et al., 2002; Cherel and Hobson 2005). Giant squids are also increasingly captured by trawling

57 nets because of the development of deep-sea fisheries (Guerra et al., 2006). In deep water
58 conditions, the giant squid would have a particular diet and exposure conditions to trace
59 elements and metals. For example, Hg bioavailability seems to be enhanced in these deep
60 environments because the absence of solar radiation and the low oxygen concentrations in
61 the deep environment favors a high methylation rate by bacteria (Monteiro et al., 1996). Also,
62 Cd is enriched in mesopelagic waters while depleted in the surface ocean because of its
63 regeneration from sinking biological debris from epipelagic zone and its uptake by organisms
64 at the surface (Boyle et al., 1976).

65
66 In this framework, the objectives of this study were to provide baseline data on a wide range
67 of trace elements in the giant squid *Architeuthis dux* from the Spanish waters. To this end, the
68 concentrations and tissue distribution of 12 metals (Ag, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb,
69 V and Zn) and 2 metalloids (As and Se) were determined in the tissues and organs of fished
70 and stranded specimens. The recorded values were then compared to the data from the
71 current literature for other cephalopod species.

72 73 **2. Material and methods**

74 75 **2.1. Sampling and sample preparation**

76 Six specimens of giant squid were collected between 2001 and 2005. Table 1 shows the main
77 data of the specimens as well as the sampling site, date, and mode of capture. Five specimens
78 were caught in the Bay of Biscay (Asturias, North Spain) and one in the western
79 Mediterranean Sea. Two of them were mature males and four were immature or maturing
80 females.

Collected specimens were immediately frozen prior to dissection. In the laboratory, each individual was weighed and measured (mantle length ML, total length TL), and the sex and maturity stage determined. The digestive gland, gills, ink sack, branchial hearts and their appendages, systemic heart and brain were totally removed. In addition, pieces of muscle, skin, digestive and genital tissues (i.e. oviduct gland, ovary and testis), were sampled to determine trace element concentrations. As it was not possible to separate the different tissues and to weigh them, the total concentrations in the whole *Architeuthis* specimens were estimated according to the measured concentrations in the different tissues and to their relative weight in fishery targeted squids.

2.2. Analytical procedure

All samples were freeze-dried for several days then grounded. Aliquots of the samples ranging from 50 to 300 mg were digested using a 3:1 v:v nitric-hydrochloric acid mixture with 65% HNO₃ (Merck, suprapur quality) and 70% HCl (Merck, suprapur quality). Acidic digestion was performed overnight under ambient temperature and then heated in a microwave during 30 min with increasing temperature until 105°C, and 15 min at 105°C (1200 W). After the mineralization process, each sample was diluted to 30 or 50 ml with milli-Q quality water, according to the volume of acid added to the mineralization (3.0 ml or 4.5 ml).

Ag, As, Cd, Co, Cr, Cu, Fe, Mn, Ni Pb, Se, V and Zn were analysed either by Inductively Coupled Plasma-Optical Emission Spectrometry (Varian[®] Vista-Pro) or Inductively Coupled Plasma-Mass Spectrometry (Varian[®] Ultra Mass 700). For Hg, two aliquots ranging from 10 to 50 mg of dried material were directly analysed in an Advanced Mercury Analyser spectrophotometer (Altec[®] AMA 254).

Reference tissues (dogfish liver DOLT-3, NRCC, and lobster hepatopancreas TORT-2, NRCC) were treated and analysed in the same way as the samples. Results were in good agreement with the certified values, and the standard deviations were low, proving good repeatability of the method (Table 2). The results for standard reference materials displayed recoveries of the elements ranging from 88% to 116% (n=10).

The detection limits ($\mu\text{g.g}^{-1}$ dry wt) for ICP-OES were 8.3 (As, Fe, Zn), 3.3 (Ag, Se), 1.67 (Pb, V), 0.83 (Cd, Co, Cr, Cu, Mn, Ni), and were 0.167 (Ni, V), 0.083 (Cd, Co, Cr, Cu, Mn, Pb), 0.033 (Ag) for ICP-MS. Trace element concentrations are given relative to the dry weight ($\mu\text{g.g}^{-1}$ dry wt) while the distribution percentages are calculated for wet weight.

3. Results

3.1. Trace element concentrations in soft tissues

Trace element concentrations in the tissues and organs of *Architeuthis* are reported in Figure 1. Among the sampled tissues, the digestive gland was the major site of concentration for Ag, Cd, Co, Cu, Fe and Se with 1.13-14.0 $\mu\text{g Ag.g}^{-1}$, 27.2-134 $\mu\text{g Cd.g}^{-1}$, 2.0-4.8 $\mu\text{g Co.g}^{-1}$, 64-1218 $\mu\text{g Cu.g}^{-1}$, 52-1862 $\mu\text{g Fe.g}^{-1}$ and 9.7-19.5 $\mu\text{g Se.g}^{-1}$ (Figure 1). The digestive gland also concentrated Ni, V and Zn at concentrations closed to the highest concentrations recorded in the other tissues with 0.11-1.42 $\mu\text{g Ni.g}^{-1}$, 0.60-4.34 $\mu\text{g V.g}^{-1}$, and 34-219 $\mu\text{g Zn.g}^{-1}$. Interestingly, the digestive gland also exhibited the lowest concentrations of As, Cr and Hg with 31-65 $\mu\text{g As.g}^{-1}$, 0.41-1.40 $\mu\text{g Cr.g}^{-1}$, and 0.32-1.56 $\mu\text{g Hg.g}^{-1}$ (Figure 1).

The concentrations of As, Co, Hg, Ni, Se and V were also remarkable in branchial hearts, which play an important excretory role in cephalopods. In this tissue, the concentrations of trace elements were the highest for Ni or very close to the highest for As, Co, Hg, Se and V. Branchial hearts concentrated As from 86 to 111 $\mu\text{g.g}^{-1}$, Co from 3.15 to 3.22 $\mu\text{g.g}^{-1}$, Hg

from 0.71 to 4.37 $\mu\text{g.g}^{-1}$, Ni from 0.74 to 2.60 $\mu\text{g.g}^{-1}$, Se from 8.2 to 9.3 $\mu\text{g.g}^{-1}$ and V from 1.50 to 2.46 $\mu\text{g.g}^{-1}$ (Figure 1).

Muscles exhibited generally among the lowest concentrations for all trace elements except for Hg which exhibited the highest values in this tissue (1.86-3.32 $\mu\text{g Hg.g}^{-1}$ dry weight; Figure 1). Cr and Zn were highly concentrated in the ovary with concentrations ranging from 0.65 to 6.12 $\mu\text{g.g}^{-1}$ and 131 to 149 $\mu\text{g.g}^{-1}$, respectively. The oviduct gland also displayed high concentrations of As and Mn as well as the digestive gland appendages had high concentrations of Cr and Pb (Figure 1).

3.2. Distribution of the trace elements in soft tissues

The proportions of the whole body burden of the trace elements contained in each organ and tissue are shown in Figure 2. With the exception of As, Cr, Hg and Mn which were mainly found in the body muscular parts ($69 \pm 9\%$, $68 \pm 20\%$, $87 \pm 5\%$, and $55 \pm 22\%$, respectively), the digestive gland contained the largest quantities of all trace elements : $98 \pm 1\%$ of Ag, $99 \pm 1\%$ of Cd, $97 \pm 2\%$ of Co, $92 \pm 5\%$ of Cu, $85 \pm 16\%$ of Fe, $53 \pm 28\%$ of Ni, $69 \pm 22\%$ of Pb, $77 \pm 5\%$ of Se, $71 \pm 19\%$ of V, and $53 \pm 17\%$ of Zn (Figure 2).

Although the concentrations of some trace elements were high in the branchial hearts or in the gills, these tissues contained in fact low amounts of the considered elements because of their small masses relative to the whole body weight (Figure 2).

3.3. Influence of the size/weight and origin

The size/weight only influenced Zn concentrations in the digestive gland ($R^2=0.934$, $p=0.020$) and Cr and Ni concentrations in the gills ($R^2=0.968$ $p=0.007$ and $R^2=0.969$ $p=0.007$).

No statistical tests were performed to compare the influence of the origin on the accumulated trace elements because our sampling only included one specimen from the Mediterranean.

However, this giant squid clearly displayed higher concentrations of Hg than any of those from the Bay of Biscay with $1.56 \text{ vs } 0.47 \pm 0.13 \text{ } \mu\text{g.g}^{-1}$ in the digestive gland (see Table 3), $2.97 \text{ vs } 1.38 \pm 0.34 \text{ } \mu\text{g.g}^{-1}$ in the gills, and $3.32 \text{ vs } 2.07 \pm 0.19 \text{ } \mu\text{g.g}^{-1}$ in the mantle muscle (data not shown). Important differences also appeared for Ag, Cu and Zn with $14.0 \text{ vs } 1.90 \pm 0.47 \text{ } \mu\text{g Ag.g}^{-1}$, $1218 \text{ vs } 108 \pm 83 \text{ } \mu\text{g Cu.g}^{-1}$, and $219 \text{ vs } 103 \pm 51 \text{ } \mu\text{g Zn.g}^{-1}$ in the digestive gland (Table 3), and $4.80 \text{ vs } 0.31 \pm 0.11 \text{ } \mu\text{g Ag.g}^{-1}$, $206 \text{ vs } 31 \pm 24 \text{ } \mu\text{g Cu.g}^{-1}$ and $111 \text{ vs } 60 \pm 35 \text{ } \mu\text{g Zn.g}^{-1}$ in the gills (data not shown).

4. Discussion

Previous studies have demonstrated the ability of cephalopods to accumulate high concentrations of trace elements in their tissues but very little data is available in the current literature for non-targeted and/or deep-waters species such as the giant squid *Architeuthis*. Although globally poorly documented, metal and metalloid concentrations in cephalopod tissues have received increasing interest over the last decades, particularly in Europe and Japan, as these molluscs play a major role both as predators and food items in marine ecosystems (see the reviews by Clarke 1996; Croxall and Prince 1996; Klages 1996; Smale 1996; Boyle and Rodhouse, 2005). The central role of the digestive gland in trace element bioaccumulation and detoxification has been highlighted many times, particularly for toxic metals such as Ag and Cd (e.g. Martin and Flegel 1975; Miramand and Bentley 1992; Bustamante et al., 2002, 2004; Ichiashi et al., 2001a; Miramand et al., 2006). Beside the digestive gland, which also plays a major role in the energetic metabolism of cephalopods (e.g. Rosa et al., 2005; Moltschaniwskyj and Johnston 2006), the branchial hearts and their appendages are involved in trace element excretion processes, allowing the depuration and/or the storage of various metals and radionuclides (e.g. Nardi and Steinberg 1974; Miramand

181 and Guary 1980; Guary et al., 1981; Miramand and Bentley 1992; González et al., 1998;
182 Bustamante et al., 2002, 2006b). As for coastal and/or oceanic targeted cephalopods, the
183 digestive gland and the branchial hearts of *Architeuthis* generally contained the highest
184 concentrations of most of the considered elements, i.e. Ag, Cd, Co, Cu, Fe, Ni, Se, V and Zn
185 (Figure 1). This finding strongly suggests that the metabolism of trace elements in
186 *Architeuthis* is very close to, or even the same as other families of cephalopods. This is
187 supported by the fact that trace element concentrations in the digestive gland of *Architeuthis*
188 closely fall within the same range than for other cephalopod species (Table 3), indicating that
189 the potential of *Architeuthis* for their bioaccumulation is relatively similar. Moreover,
190 according to the elevated proportions of the total element body burden, the digestive gland of
191 *Architeuthis* might play a central role in the detoxification and storage of most of the
192 analysed elements, i.e. Ag, Cd, Co, Cu, Fe, Ni, Pb, Se, V and Zn (Figure 2). It would be a
193 great interest to investigate the detoxification strategies in the digestive gland of *Architeuthis*
194 in comparison to that of other families of cephalopods.

195 Besides the digestive gland, muscles contained very high proportions of total body burdens
196 of As, Cr, Hg, Mn, Ni, and Zn (Figure 2). With the exception of Hg, trace element
197 concentrations in *Architeuthis* muscles were relatively low and these high proportions
198 resulted from the elevated muscular mass respective to the whole body weight. For Hg, the
199 concentrations recorded in the muscles were the highest among the different organs and
200 tissues (Figure 1). Previous studies have reported relatively similar Hg concentrations
201 between the different tissues of different squid species from the Northern Atlantic waters
202 (Bustamante et al., 2006a; Pierce et al., 2008). It is therefore noteworthy that muscular Hg
203 concentrations in *Architeuthis* were 2 to 4 times higher than in the digestive gland. In
204 comparison with Ag, Cd, Co, Cu, Fe, Ni, Pb, Se, V and Zn, the role of the digestive gland in
205 the storage of Hg appeared to be relatively limited in *Architeuthis*. This may be due to an

206 excretion function of Hg by the digestive gland, and/or a preferential redistribution of Hg to
207 muscular tissues where it binds to the sulphhydryl groups of proteins (Bloom, 1992;
208 Bustamante et al., 2006a). Such a redistribution might indicate that most of the Hg ingested
209 from the prey would be in the organic form such as fish in which Hg content is virtually
210 100% in the methylated form (Bloom, 1992). Little is known about the diet of *Architeuthis*,
211 it includes other cephalopods (Pérez-Gándaras and Guerra, 1978, 1989), crustaceans (e.g.
212 *Nephros norvergicus*) as well as a large proportion of fish of different families accordingly to
213 the available prey in the area (for instance *Trachurus trachurus*, *Maurolicus muelleri* and
214 *Micromesistius poutassou* in Ireland waters and equivalent species from Namibia and New
215 Zealand) (see Guerra et al., 2006 for a review). Further studies therefore should focus on
216 trace elements in the typical prey of *Architeuthis* to provide insights on this aspect.

217 As in other cephalopod species, trace element concentrations in *Architeuthis* may vary with
218 biological and environmental factors such as age (size/weight), sex, and geographical origin
219 (e.g. Monteiro et al., 1992; Bustamante et al., 1998; Raimundo et al., 2004; Pierce et al.,
220 2008). Our limited sampling did not allow making comparisons for all these factors.
221 However, considering the 5 specimens from the Bay of Biscay, it appears that size/weight
222 poorly influenced trace element concentrations in *Architeuthis* tissues. The increase of Zn
223 concentrations from 34 to 160 $\mu\text{g.g}^{-1}$ dwt in the digestive gland might be related to metal
224 physiological changes related to the sexual maturation as reported for other cephalopod
225 species like *Sepia officinalis* from the English Channel (Miramand et al., 2006). In females,
226 high concentrations of Zn in the ovary of *Architeuthis* ($120 \pm 38 \mu\text{g.g}^{-1}$ dwt) were close to
227 that in the genital tract of *Sepia officinalis* ($123 \pm 3 \mu\text{g.g}^{-1}$ dwt; Miramand and Bentley,
228 1992). Within the ovary, essential elements such as Zn are stored in metal-containing
229 enzymes and metalloproteins (Gerpe et al., 2000) and transferred to the yolk of the eggs
230 (Villanueva and Bustamante, 2006, Lacoue-Labarthe et al., 2008). Cr and Ni in the gills also

displayed a significant increase with size/weight. In the current literature, very little data is available on the variation of Cr and Ni concentrations in cephalopod tissues. For example, in the squid *Sthenoteuthis oualaniensis* Cr concentrations were higher in juveniles than in adults, whereas juveniles displayed lower Ni concentrations than adults (Ichihashi et al., 2001a). Such a difference for *S. oualaniensis* was explained by the evolution of food habits between the juvenile and adult stages, juvenile feeding more on crustaceans while adults primarily preyed on fish. According to the lack of significant variation in the digestive gland, such a switch is not likely to occur in the size range of *Architeuthis* we analysed. Furthermore, even if the diet could represent the main pathway for many elements - as experimentally shown for Am, Cd, Co and Zn (Koyama et al., 2000; Bustamante et al., 2002, 2004, 2006b) - seawater could also be an important uptake pathway, as elements pass through the skin and through the gills. For instance, seawater represents the main pathway for Ag in *Sepia officinalis* (Bustamante et al., 2004). Therefore, Cr and Ni bioaccumulation in gills might also result from a direct uptake from seawater all along the lifespan of *Architeuthis*.

Trace element concentrations in cephalopods could also vary according the location where individuals were captured (Bustamante et al., 1998; Seixas et al., 2005ab; Pierce et al., 2008). In the case of *Architeuthis*, this is clearly exemplified by the much higher Hg concentrations in the tissue of the specimen from the Mediterranean. Higher Hg concentrations in Mediterranean organisms have been highlighted many times and are typically explained by high temperatures and absence of solar radiation in the deep environment. These conditions favor a high methylation rate of the metal, methyl-Hg being highly bioavailable for marine biota, which consistently biomagnifies through the food chain. Moreover, natural sources of Hg in the Mediterranean Sea may contribute to Hg enrichment through the benthic food webs, as it constitutes the richest natural reserve of this element (Bacci, 1989). Higher Ag, Cu and

256 Zn concentrations in the digestive gland and in the gills also indicated different exposure
257 conditions of this specimen compared to those from the Bay of Biscay. In cephalopods, Ag
258 bioaccumulation in the digestive gland might reflect the global contamination of the
259 surrounding waters (e.g. Martin and Flegal, 1975; Miramand et al., 2006), seawater being the
260 main pathway of exposure and Ag having a fast turn-over in the tissues (Bustamante et al.,
261 2004). Interestingly, Ag, Cu, Hg and Zn are metals that bind to metallothionein proteins
262 (MTs), which play a role in the homeostasis of the essential metals (i.e. Cu and Zn) and in
263 the detoxification of non-essential metals (i.e. Ag, Cd and Hg) (Roesijadi, 1992, 1996;
264 Viarengo and Nott, 1993). The role of MTs in metal sequestration in cephalopods is not
265 completely clear (Bustamante et al., 2006c) and this issue clearly deserves further
266 investigation.

267 Even though trace elements are generally considered for their potential toxicity in
268 ecotoxicological studies and biomonitoring surveys, there is increasing interest in their use
269 for providing information on life history and trophic ecology of cephalopods (Jackson et al.,
270 2007). Thus, Cd is of particular interest because it is highly bioaccumulated by cephalopods.
271 Indeed, Cd is efficiently absorbed and strongly retained in the digestive gland (Bustamante et
272 al., 1998, 2002). Even if most of cephalopod species display short life spans i.e. typically less
273 than 2 years, they can accumulate very high Cd concentrations in their digestive gland
274 reaching up 1000 $\mu\text{g g}^{-1}$ wet wt in the squid *Illex argentinus* (Dorneles et al., 2007). Because
275 of Cd incorporation by organisms in epipelagic waters and its regeneration from sinking
276 biological debris in the mesopelagic environment (Boyle et al., 1976), deep-water
277 cephalopods might show relatively high Cd concentrations. In the Bay of Biscay, Cd
278 concentrations reached 9.1 and 33.1 $\mu\text{g.g}^{-1}$ dry wt in the digestive gland of the mesopelagic
279 squids *Histioteuthis reversa* and *Teuthowenia megalops*, respectively (unpublished data) and
280 would be due to the consumption of prey highly contaminated with Cd. Even elevated

compared to neritic squid species (Bustamante et al., 1998), these concentrations are lower than those measured in the digestive gland of *Architeuthis*, suggesting that 1) the giant squid feed on more contaminated prey than Histioteuthidae and Cranchidae, or 2) it displays a much longer life span than other cephalopods. Age estimation and growth rates of giant squids are still open questions. Indeed, isotopic analysis indicated that the age for the giant squid *Architeuthis sanctipauli* from Tasmania was 14 years for specimens ranging from 191 to 240 cm ML (Landman et al., 2004). This completely disagrees with the age estimated from growth increment counts in statoliths of *Architeuthis dux* and *Architeuthis* sp. from the Atlantic and caught off New Zealand. Specimens ranging from 43 to 161 cm ML had between 153 and 435 increments (Jackson et al., 1991; Gauldie et al., 1994; Lordan et al., 1998; González et al., 2002). If the increments were daily deposited, as it occurs in other cephalopods, the age of these animals will not exceed two years. This finding implies that *Architeuthis* would have a very fast growth rate with intense food intakes that in turn would lead to the bioaccumulation of relatively high Cd concentrations in its digestive gland.

Acknowledgements

We thank Luis Laria and other members of CEPESMA who collected the squids, Dr. Alberto Fernández Costa for his technical assistance in the dissections, and the staff of the Centre Commun d'Analyses (University of La Rochelle) for their help in the analytical process. This study was supported financially by LIENSs (University of La Rochelle).

References

Bacci, E. 1989. Mercury in the Mediterranean. Marine Pollution Bulletin 20, 59-63.

306

307 Bloom, N.S. 1992. On the chemical form of mercury in edible fish and marine invertebrate
 308 tissue. Canadian Journal of Fisheries and Aquatic Sciences 49, 1010-1017.

309

310 Boyle, E.A., Sclater, F., Edmond, J.M. 1976. On the marine geochemistry of cadmium. Nature
 311 263, 42-44.

312

313 Boyle, P.R., Rodhouse, P.G. 2005. *Cephalopods: Ecology and Fisheries*. Blackwell Science, Oxford,
 314 United Kingdom.

315

316 Bustamante, P., Caurant, F., Fowler, S.W., Miramand, P. 1998. Cephalopods as a vector for the
 317 transfer of cadmium to top marine predators in the north-east Atlantic Ocean. Science of the
 318 Total Environment 220, 71-80.

319

320 Bustamante, P., Grigioni, S., Boucher-Rodoni, R., Caurant, F., Miramand, P. 2000.
 321 Bioaccumulation of 12 trace elements in the tissues of the nautilus *Nautilus macromphalus*
 322 from New Caledonia. Marine Pollution Bulletin 40, 688-696.

323

324 Bustamante, P., Teyssié, J-L., Fowler, S.W., Cotret, O., Danis, B., Miramand, P., Warnau, M.
 325 2002. Biokinetics of zinc and cadmium accumulation and depuration at different stages in the
 326 life cycle of the cuttlefish *Sepia officinalis*. Marine Ecology Progress Series 231, 167-177.

327

328 Bustamante, P., Teyssié, J-L., Fowler, S.W., Danis, B., Cotret, O., Miramand, P., Warnau, M.
 329 2004. Uptake, transfer and distribution of silver and cobalt in the tissues of the common
 330 cuttlefish *Sepia officinalis* at different stages of its life cycle. Marine Ecology Progress Series
 331 269, 185-195.

332

333 Bustamante, P., Lahaye, V., Durnez, C., Churlaud, C., Caurant, F. 2006a. Total and organic Hg
 334 concentrations in cephalopods from the North East Atlantic waters: influence of geographical
 335 origin and feeding ecology. *Science of the Total Environment*, 368: 585-596.

336

337 Bustamante, P., Teyssié, J-L., Fowler, S.W., Warnau, M. 2006b. Assessment of the exposure
 338 pathway in the uptake and distribution of americium and cesium in cuttlefish (*Sepia officinalis*)
 339 at different stages of its life cycle. *Journal of Experimental Marine Biology and Ecology* 331,
 340 198-207.

341

342 Bustamante, P., Bertrand, M., Boucaud-Camou, E., Miramand, P. 2006c. Subcellular
 343 distribution of Ag, Cd, Co, Cu, Fe, Mn, Pb and Zn in the digestive gland of the common
 344 cuttlefish *Sepia officinalis*. *Journal of Shellfish Research* 3, 987-994.

345

346 Cherel, Y., Hobson, K.A. 2005. Stable isotopes, beaks and predators: a new tool to study the
 347 trophic ecology of cephalopods, including giant and colossal squids. *Proceedings of the Royal*
 348 *Society B* 272, 1601-1607.

349

350 Clarke, M.R. 1996. Cephalopods as prey. III. Cetaceans. *Philosophical Transactions of the*
 351 *Royal Society of London. Series B, Biological Sciences* 351, 1053-1065.

352

353 Croxall, J.P., Prince, P.A. 1996. Cephalopods as prey. I. Seabirds. *Philosophical transactions of*
 354 *the Royal Society of London. Series B, Biological Sciences* 351, 1023-1043.

355

356 Dorneles, P.R., Lailson-Brito, J., dos Santos, R.A., Silva da Costa, P.A., Malm, O., Azevedo,
 357 A.F., Machado Torres, J.P. 2007. Cephalopods and cetaceans as indicators of offshore
 358 bioavailability of cadmium off Central South Brazil Bight. *Environmental Pollution* 148, 352-
 359 359.
 360
 361 Finger, J.M., Smith, J.D. 1987. Molecular association of Cu, Zn, Cd and ^{210}Po in the digestive
 362 gland of the squid *Nototodarus gouldi*. *Marine Biology* 95, 87-91.
 363
 364 Gauldie, R.W., West, I.F., Förch, E.C. 1994. Statocyst, statolith, and age estimation of the
 365 giant squid, *Architeuthis kirki*. *The Veliger* 37(1), 93-109.
 366
 367 Gerpe, M.S., de Moreno, J.E.A., Moreno, V.J., Patat, M.L. 2000. Cadmium, zinc and copper
 368 accumulation in the squid *Illex argentinus* from the Southwest Atlantic Ocean. *Marine Biology*
 369 136, 1039–1044.
 370
 371 González, A.F., Guerra, A., Pascual, S. & Briand, P. 1998. *Vulcanoctopus hydrothermalis* gen.
 372 et sp. nov. (Mollusca, Cephalopoda): a new octopod from a deep sea hydrothermal vent.
 373 *Cahiers de Biologie Marine* 39, 169-184.
 374
 375 González, A.F., Guerra, A., Rocha, F., Gracia, J. 2002. Recent findings of the giant squid
 376 *Architeuthis* in northern Spanish waters. *Journal of the Marine Biological Association of the*
 377 *United Kingdom* 82, 859-861.
 378

379 Guary, J-C., Higgs, J.J.W., Cherry, R.D., Heyraud, M. 1981. High concentrations of
 380 transuranic and natural radioactive elements in the branchial hearts of the cephalopods *Octopus*
 381 *vulgaris*. Marine Ecology Progress Series 4, 123-126.
 382
 383 Guerra, A., González, A.F., Dawe, E.G., Rocha, F. 2004. A review of records of giant squid in
 384 the north-eastern Atlantic, with a note on the two first records of males *Architeuthis* sp. off the
 385 Iberian Peninsula. Journal of the Marine Biological Association of the United Kingdom 84,
 386 427-431.
 387
 388 Guerra, A., González, A.F., Rocha, F., Laria, L., Gracia, J. 2006. Enigmas de la ciencia: el
 389 calamar gigante. Guerra, A. et al., (eds.). Instituto de Investigaciones Marinas (CSIC, Vigo),
 390 313 pp., 88 figs.
 391
 392 Honda, K., Tatsukawa, R., Itano, K., Miyazaki, N., Fujiyama, T. 1983. Heavy metal
 393 concentrations in muscle, liver and kidney tissue of striped dolphin, *Stenella coeruleoalba*, and
 394 their variations with body length, weight, age and sex. Agric Biol Chem 47(6), 1219-1228.
 395
 396 Ichihashi, H., Kohno, H., Kannan, K., Tsumura, A., Yamasaki, S.I. 2001a. Multielemental
 397 analysis of purpleback flying squid using high resolution inductively coupled plasma-mass
 398 spectrometry (HR ICP-MS). Environmental Science and Technology 35, 3103-3108.
 399
 400 Ichihashi, H., Nakamura, Y., Kannan, K., Tsumura, A., Yamasaki, S.I. 2001b. Multi-elemental
 401 concentrations in tissues of Japanese common Squid (*Todarodes pacificus*). Archives of
 402 Environmental Contamination and Toxicology 41, 483-490.
 403

404 Jackson, G.D., Lu, C.C., Dunning, M.C. 1991. Growth rings within the statolith microstructure
 405 of the giant squid *Architeuthis*. The Veliger 34 (4), 331-334.
 406

407 Jackson, G.D., Bustamante, P., Cherel, Y., Fulton, A., Grist, E.P.M., Jackson, C.H., Nichols,
 408 P.D., Pethybridge, H., Phillips, K., Ward, R.D., Xavier, J.C. 2007. Applying new tools to
 409 cephalopod trophic dynamics and ecology: perspectives from the Southern Ocean Cephalopod
 410 Workshop, February 2-3, 2006. Reviews in Fish Biology and Fisheries 17, 79-99.
 411

412 Klages, N.T.W. 1996. Cephalopods as prey. II. Seals. Philosophical transactions of the Royal
 413 Society of London. Series B, Biological Sciences 351, 1045-1052.
 414

415 Koyama, J., Nanamori, N., Segawa, S. 2000. Bioaccumulation of waterborne and dietary
 416 cadmium by oval squid *Sepioteuthis lessoniana*, and its distribution among organs. Marine
 417 Pollution Bulletin 40, 961-967.
 418

419 Kubodera, T., Mori, K. 2005. First-ever observations of a live giant squid in the wild.
 420 Proceedings of the Royal Society B 272, 2583-2586.
 421

422 Lacoue-Labarthe, T., Warnau, M., Oberhänsli, F., Teyssié, J-L., Jeffree, R., Bustamante, P.
 423 2008. First experiments on the maternal transfer of heavy metals in the cuttlefish, *Sepia*
 424 *officinalis*. Marine Pollution Bulletin 57, 826-831.
 425

426 Lahaye, V., Bustamante, P., Spitz, J., Das, K., Meynier, L., Magnin, V., Dabin, W., Caurant, F.
 427 2005. Long-term dietary preferences of common dolphins in the Bay of Biscay using a metallic
 428 tracer. Marine Ecology Progress Series 305, 275-285.

429

430 Landman, N.H., Cochran, J.K, Cerrato, R., Mak, J., Roper, C.F.E., Lu, C.C. 2004. Habitat and
 431 age of the giant squid (*Architeuthis sanctipauli*) inferred from isotopic analyses. Marine
 432 Biology 144, 685-691.

433

434 Lordan, C., Collins, M.A., Raya, C.P. 1998. Observations on morphology, age and diet of three
 435 *Architeuthis* caught off the west coast of Ireland in 1995. Journal of the Marine Biological
 436 Association of the United Kingdom 78, 903-917.

437

438 Martin, J.H., Flegal, A.R. 1975. High copper concentrations in squid livers in association with
 439 elevated levels of silver, cadmium, and zinc. Marine Biology 30, 51-55.

440

441 Miramand, P., Guary, J-C. 1980. High concentrations of some heavy metals in tissues of the
 442 Mediterranean octopus. Bulletin of Environmental Contamination and Toxicology 24, 783-788.

443

444 Miramand, P., Bentley, D. 1992. Concentration and distribution of heavy metals in tissues of
 445 two cephalopods, *Eledone cirrhosa* and *Sepia officinalis*, from the French coast of the English
 446 Channel. Marine Biology 114, 407-414.

447

448 Miramand, P., Bustamante, P., Bentley, D., Koueta, N. 2006. Variation of heavy metal
 449 concentrations (Ag, Cd, Co, Cu, Fe, Pb, V, Zn) during the life cycle of the common cuttlefish
 450 *Sepia officinalis*. Science of the Total Environment 361, 132-143.

451

452 Moltschaniwskyj, N., Johnston, D. 2006. Evidence that lipid can be digested by the dumpling
 453 squid *Euprymna tasmanica*, but is not stored in the digestive gland. *Marine Biology* 149, 565-
 454 572.

455

456 Monteiro, L.R., Porteiro, F.M., Gonçalves, J.M. 1992. Inter- and intra-specific variation of
 457 mercury levels in muscle of cephalopods from the Azores. *Arquipelago* 10, 13–22.

458

459 Monteiro, L.R., Costa, V., Furness, R.W., Santos, R.S. 1996. Mercury concentrations in prey
 460 fish indicate enhanced bioaccumulation in mesopelagic environments. *Marine Ecology*
 461 *Progress Series* 141, 21-25.

462

463 Muirhead, S.J., Furness, R.W. 1988. Heavy metal concentrations in the tissues of seabirds from
 464 Gough Island, South Atlantic Ocean. *Marine Pollution Bulletin* 19, 278-283.

465

466 Nardi, G., Steinberg, H. 1974. Isolation and distribution of adenochrome(s) in *Octopus*
 467 *vulgaris*. *Comparative Biochemistry and Physiology* 48 B, 453-461.

468

469 Pérez-Gándaras, G., Guerra, A. 1978. Nueva cita de *Architeuthis* (Cephalopoda: Teuthoidea):
 470 descripción y alimentación. *Investigación Pesquera* 42 (2), 401-414.

471

472 Pérez-Gándaras, G., Guerra, A. 1989. *Architeuthis* de Sudafrica: nuevas citas y consideraciones
 473 biológicas. *Scientia Marina* 53, 113-116.

474

475 Pierce, G.J., Stowasser, G., Hastie, L.C., Bustamante, P. 2008. Geographic, seasonal and
 476 ontogenetic variation in cadmium and mercury concentrations in squid (Cephalopoda:
 477 Teuthoidea) from UK waters. *Ecotoxicology and Environmental Safety* 70, 422-432
 478
 479 Raimundo, J., Caetano, M., Vale, C. 2004. Geographical variation and partition of metals in
 480 tissues of *Octopus vulgaris* along the Portuguese coast. *Science of the Total Environment* 325,
 481 71-81.
 482
 483 Roesijadi, G. 1992. Metallothionein in metal regulation and toxicity in aquatic animals.
 484 *Aquatic Toxicology* 22, 81-114.
 485
 486 Roesijadi, G. 1996. Metallothionein and its role in toxic metal regulation. *Comparative*
 487 *Biochemistry and Physiology* 113C, 117-123.
 488
 489 Roper, C.F.E., Boss, K.J. 1982. The giant squid. *Scientific American* 246, 96-105.
 490
 491 Rosa, R., Pereira, J.M.F., Nunes, M.L. 2005. Biochemical composition of cephalopods with
 492 different life strategies, with special reference to a giant squid, *Architeuthis* sp. *Marine Biology*
 493 146, 739-751.
 494
 495 Santos, M.B., Pierce, G.J., García Hartmann, M., Smeenk, C., Addink, M.J., Kuiken, T., Reid,
 496 R.J., Patterson, I.A.P., Lordan, C., Rogan, E., Mente, E. 2002. Additional notes on stomach
 497 contents of sperm whales *Physeter macrocephalus* stranded in the north-east Atlantic. *Journal*
 498 *of the Marine Biological Association of the United Kingdom* 82(3), 501-507.
 499

500 Seixas, S., Pierce, G.J. 2005a. Vanadium, rubidium and potassium in *Octopus vulgaris*
 501 (Mollusca: Cephalopoda). *Scientia Marina* 69(2), 215-222.
 502

503 Seixas, S., Pierce, G.J. 2005b. Bioaccumulation of lead, calcium and strontium and their
 504 relationships in the octopus *Octopus vulgaris*. *Water, Air, and Soil Pollution* 163, 137-152.
 505

506 Seixas, S., Bustamante, P., Pierce, G.J. 2005a. Accumulation of mercury in the tissues of the
 507 common octopus *Octopus vulgaris* (L.) in two localities on the Portuguese coast. *Science of*
 508 *the Total Environment* 340, 113-122.
 509

510 Seixas, S., Bustamante, P., Pierce, G.J. 2005b. Interannual patterns of variation in
 511 concentrations of trace elements in arms of *Octopus vulgaris*. *Chemosphere* 59, 1113-1124.
 512

513 Smale, M.J. 1996. Cephalopods as prey. IV. Fishes. *Philosophical transactions of the Royal*
 514 *Society of London Series B, Biological Sciences* 351, 1067-1081.
 515

516 Smith, J.D., Plues, L., Heyraud, M., Cherry, R.D. 1984. Concentrations of the elements Ag, Al,
 517 Ca, Cd, Cu, Fe, Mg, Pb and Zn, and the radionuclides ²¹⁰Pb and ²¹⁰Po in the digestive gland of
 518 the squid *Nototodarus gouldi*. *Marine Environmental Research* 13, 55-68.
 519

520 Storelli, M.M., Barone, G., D'Addabbo R., Marcotrigano, G.O. 2006. Concentrations and
 521 composition of organochlorine contaminants in different species of cephalopod molluscs from
 522 the Italian waters (Adriatic Sea). *Chemosphere* 64, 129-134.
 523

524 Tanabe, S., Tanaka, H., Tatsukawa, R. 1984. Polychlorobiphenyls, DDT, and
 525 hexachlorocyclohexane isomers in the western North Pacific ecosystem. Archives of
 526 Environmental Contamination and Toxicology 13, 731-738.

527

528 Ueda, T., Nakahara, M., Ishii, T., Suzuki, Y., Suzuki, H. 1979. Amounts of trace elements in
 529 marine cephalopods. Journal of Radioactivity Research 20, 338-342.

530

531 Ueno, D., Inoue, S., Ikeda, K., Tanaka, H., Yamada, H., Tanabe, S. 2003. Specific
 532 accumulation of polychlorinated biphenyls and organochlorine pesticides in Japanese common
 533 squid as a bioindicator. Environmental Pollution 125, 227-235.

534

535 Viarengo, A., Nott, J.A. 1993. Mini-review. Mechanisms of heavy metal cation homeostasis in
 536 marine invertebrates. Comparative Biochemistry and Physiology 104C, 355-372.

537

538 Villanueva, R., Bustamante, P. 2006. Composition in essential and non-essential elements of
 539 early stages of cephalopods and dietary effects on the elemental profiles of *Octopus vulgaris*
 540 paralarvae. Aquaculture 261, 225-240.

541

542 Yamada, H., Takayanagi, K., Tateishi, M., Tagata, H., Ikeda, K. 1997. Organotin compounds
 543 and polychlorinated biphenyls of livers in squid collected from coastal waters and Open Ocean.
 544 Environmental Pollution 96(2), 217-226.

Table 1. Sampling information and squid biological characteristics. T: trawling; S: Stranded; F: Floating at the surface; ML: Dorsal mantle length; TL: Total length.

N°	Sampling site	Sampling date	Mode of capture	Weight (kg)	ML (cm)	TL (cm)	Sex / Maturation
1	Off Luarca (Asturias, North Spain)	12 September 2001	T	90	127	710	F / Immature
2	Ribadesella (Asturias, North Spain)	23 October 2001	S	104	150	800	F / Maturing
3	Colunga (Asturias, North Spain)	15 September 2003	S	80	152	1200	F / Immature
4	Off Gijón (Asturias, North Spain)	16 September 2003	F	66	122	620	M / Mature
5	Off Gandía (Valencia, Western Mediterranean)	19 July 2005	T	50	107	600	M / Mature
6	Off Gijón (Asturias, North Spain)	22 July 2005	T	139	146	820	F/ Maturing

Table 2. Comparison of certified trace elements concentrations ($\mu\text{g g}^{-1}$ dry weight) in reference materials (n=10) with the values determined in the present study (nc: not certified value, nd: not determined). ICP-MS - Inductively Coupled Plasma Mass Spectrometry; ICP-OES - Inductively Coupled Plasma Optical Emission Spectrometry; AMA – Advanced Mercury Analyser.

Element	Method	TORT-2				DOLT-3			
		Measured		Certified	Recovery	Measured		Certified	Recovery
		Mean	± SD	Mean ± SD		Mean ± SD	Mean ± SD		
Ag	ICP-MS	6.21	± 1.69	nc	-	1.21	± 0.10	1.20 ± 0.07	101
As	ICP-OES	21.8	± 2.4	21.6 ± 1.8	101	9.9	± 0.3	10.2 ± 0.5	97
Cd	ICP-MS	26.4	± 2.2	26.7 ± 0.6	99	19.3	± 0.7	19.4 ± 0.6	99
Co	ICP-MS	0.45	± 0.09	0.51 ± 0.09	88	0.29	± 0.05	nc	-
Cr	ICP-OES	0.69	± 0.18	0.77 ± 0.15	90	4.02	± 0.93	nc	-
Cu	ICP-OES	95	± 15	106 ± 10	90	31.9	± 0.7	31.2 ± 1.0	99
Fe	ICP-OES	100	± 10	105 ± 13	95	1349	± 76	1484 ± 57	91
Hg	AMA	0.27	± 0.01	0.27 ± 0.06	100	3.36	± 0.08	3.37 ± 0.14	100
Mn	ICP-OES	13.5	± 2.0	13.6 ± 1.2	99	9.73	± 0.14	nc	-
Ni	ICP-OES	2.44	± 0.56	2.50 ± 0.19	98	2.46	± 0.45	2.72 ± 0.35	90
Pb	ICP-MS	0.32	± 0.17	0.35 ± 0.13	91	0.294	± 0.056	0.319 ± 0.045	92
Se	ICP-MS	6.48	± 0.48	5.63 ± 0.67	115	7.56	± 0.65	7.06 ± 0.48	107
V	ICP-MS	1.55	± 0.24	1.64 ± 0.19	95	nd		nc	-
Zn	ICP-OES	188	± 20	180 ± 6	104	97.3	± 1.4	86.6 ± 2.4	116

Table 3. Reported metal concentrations (Mean $\pm$ SD, $\mu\text{g g}^{-1}$ dry weight) in the digestive gland of different cephalopod species.

Species	Ag	As	Cd	Co	Cr	Cu	Fe	Hg	Mn	Ni	Pb	V	Zn	Reference
Architeuthidae														
<i>Architeuthis dux</i>	1.90 $\pm$ 0.47	48 $\pm$ 14	60.8 $\pm$ 46.2	3.27 $\pm$ 1.76	0.93 $\pm$ 0.41	108 $\pm$ 83	497 $\pm$ 779	0.47 $\pm$ 0.13	2.72 $\pm$ 2.12	0.62 $\pm$ 0.54	0.41 $\pm$ 0.33	2.24 $\pm$ 1.91	103 $\pm$ 51	a
<i>A. dux</i>	14	44	90.7	4.8	0.49	1218	158	1.56	2.26	1.37	0.85	1.72	219	b
Nautilidae														
<i>Nautilus macromphalus</i>	4.43 $\pm$ 1.95	185 $\pm$ 64	45 $\pm$ 13	5.9 $\pm$ 3.6	4.2 $\pm$ 0.8	106 $\pm$ 46	554 $\pm$ 238	-	8.9 $\pm$ 2.0	11.9 $\pm$ 7.8	-	8.0 $\pm$ 2.3	672 $\pm$ 208	c
Sepiidae														
<i>Sepia officinalis</i>	6.15 $\pm$ 1.75	-	12.7 $\pm$ 0.4	3.3 $\pm$ 0.6	1.1 $\pm$ 0.1	315 $\pm$ 3	244 $\pm$ 28	-	3.3 $\pm$ 0.1	1.3 $\pm$ 0.4	1.14 $\pm$ 0.06	5.0 $\pm$ 1.3	571 $\pm$ 47	d
<i>S. officinalis</i>	13 $\pm$ 2	-	25 $\pm$ 5	10 $\pm$ 2	-	600 $\pm$ 10	390 $\pm$ 10	-	-	-	2.2 $\pm$ 0.5	3.3 $\pm$ 0.1	1400 $\pm$ 500	e
Loliginidae														
<i>Loligo opalescens</i>	25.1 $\pm$ 12.6	-	85.0 $\pm$ 51.6	-	-	5350 $\pm$ 3210	111 $\pm$ 73	-	-	-	-	-	247 $\pm$ 131	f
<i>L. opalescens</i>	45.9 $\pm$ 19.0	-	122 $\pm$ 58	-	-	8370 $\pm$ 3130	87 $\pm$ 49	-	-	-	-	-	449 $\pm$ 201	f
Ommastrephidae														
<i>Nototodarus gouldi</i>	-	-	33 $\pm$ 30	-	-	363 $\pm$ 238	-	-	-	-	-	-	830 $\pm$ 355	g
<i>N. gouldi</i>	3.3 $\pm$ 1.4	-	50 $\pm$ 25	-	-	246 $\pm$ 298	745 $\pm$ 440	-	4.2 $\pm$ 1.1	-	-	-	696 $\pm$ 295	h
<i>Ommastrephes bartrami</i>	12.1 $\pm$ 8.6	-	287 $\pm$ 202	-	-	195 $\pm$ 212	399 $\pm$ 204	-	-	-	-	-	163 $\pm$ 55	f
<i>Stenoteuthis oualaniensis</i>	24.1 $\pm$ 10.9	-	782 $\pm$ 255	-	-	1720 $\pm$ 151	319 $\pm$ 67	-	-	-	-	-	513 $\pm$ 288	f
<i>S. oualaniensis</i> *	14.0	22.4	199	3.28	0.163	558	293	0.125	1.36	1.91	1.10	1.85	128	i
<i>Todarodes pacificus</i> *	3.5	7.5	60	0.78	0.375	27.5	325	0.133	3.5	7.0	0.60	13.8	195	j
Octopodidae														
<i>Eledone cirrhosa</i>	3.20 $\pm$ 1.74	-	24.0 $\pm$ 1.8	2.06 $\pm$ 0.08	0.8 $\pm$ 0.1	456 $\pm$ 11	287 $\pm$ 13	-	4.2 $\pm$ 1.6	2.5 $\pm$ 0.1	1.17 $\pm$ 0.09	3.3 $\pm$ 0.5	646 $\pm$ 86	d
<i>Octopus vulgaris</i>	-	-	-	8.8	-	275	275	-	2.7	-	-	-	1300	k
<i>O. vulgaris</i>	-	-	50 $\pm$ 10	-	-	2500 $\pm$ 700	700 $\pm$ 130	-	7.0 $\pm$ 0.5	-	-	4.5 $\pm$ 1.0	1450 $\pm$ 400	l
<i>O. vulgaris</i>								0.58 $\pm$ 0.08			4.9 $\pm$ 1.9	7.2 $\pm$ 6.9		m

a: Present study (Bay of Biscay); b: Present study (Mediterranean); c: Bustamante et al., (2000); d: Miramand and Bentley (1992); e: Miramand et al., (2006); f: Martin and Flegal (1975); g: Finger and Smith (1987); h: Smith et al., (1984); i: Ichihashi et al., (2001a); j: Ichihashi et al., (2001b); k: Ueda et al., (1979); l: Miramand and Guary (1980); m: Seixas and Pierce (2005ab) and Seixas et al., (2005a)

* converted to dwt using a factor of 2.5

in italics: median

Figure 1.

Figure 2.