



The impact of mid-latitude intrusions into the polar vortex on ozone loss estimates

J.-U. Grooss, R. Müller

► To cite this version:

J.-U. Grooss, R. Müller. The impact of mid-latitude intrusions into the polar vortex on ozone loss estimates. *Atmospheric Chemistry and Physics Discussions*, 2002, 2 (6), pp.2489-2506. hal-00301024

HAL Id: hal-00301024

<https://hal.science/hal-00301024>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Impact of vortex
intrusions on ozone
loss estimates**

J.-U. Grooß and R. Müller

The impact of mid-latitude intrusions into the polar vortex on ozone loss estimates

J.-U. Grooß and R. Müller

Institut für Chemie und Dynamik der Geosphäre I: Stratosphäre (ICG-I), Forschungszentrum Jülich, Jülich, Germany

Received: 5 November 2002 – Accepted: 27 November 2002 – Published: 16 December 2002

Correspondence to: J.-U. Grooß (j.-u.grooss@fz-juelich.de)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

Abstract

Current stratospheric chemical model simulations underestimate substantially the large ozone loss rates that are derived for the Arctic from ozone sondes for January of some years. Until now, no explanation for this discrepancy has been found. Here, we examine the influence of intrusions of mid-latitude air into the polar vortex on these ozone loss estimates. This study focuses on the winter 1991/92. It is based on simulations performed with the Chemical Lagrangian Model of the Stratosphere (CLaMS). The simulations for January 1992 show that the intrusions induce a reduction of vortex average ozone mixing ratio corresponding to a systematic offset of the ozone loss rate of about 12 ppb per day. Further, the results of the Match method are influenced by the intrusions, since the intruded air masses are deformed and reach dimensions below the Match radius. From our calculations we deduce a systematic offset of the Match ozone loss rate by about 10 ppb/day, which may explain about 28% of the published discrepancy between Match and box model simulations for the winter 1991/92.

1. Introduction

It is difficult to determine the chemical ozone loss in the Arctic stratosphere because chemical ozone loss is masked by dynamic variability caused by reversible vertical and horizontal transport and mixing of air masses. Different methods are known to derive the chemical ozone loss from observations such as a comprehensive set of ozone sonde measurements (Harris et al., 2002). Here we discuss the so-called vortex average method (Knudsen et al., 1998; EU, 2001) and the Match approach (von der Gathen et al., 1995; Rex et al., 1998). Both methods use ozone sonde data together with meteorological analyses and estimates of the diabatic descent to derive chemical ozone loss estimates for the polar vortex. Briefly, the vortex average approach calculates the average of all vortex ozone data on an isentropic surface as a function of time. Using a correction for diabatic descent, the chemical ozone loss is derived. The Match method

ACPD

2, 2489–2506, 2002

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

uses pairs of observations in the same air mass at different times, so called matches. These matches are determined by trajectory calculations on the basis of meteorological analyses and estimates of the diabatic descent. A statistical analysis is conducted to deduce the rate of ozone loss in the vortex from such a dataset. Only matches for which the distance between the positions of the air parcel trajectory and the second observation is below the so-called match radius (typically 500 km) contribute to the analysis.

During January in some years, large discrepancies were found between Match and the corresponding simulated ozone depletion rates (Becker et al., 1998, 2000; Kilbane-Dawe et al., 2001; Rex et al., 2002): e.g. in 1992 the Match results for 24 January ± 7 days yield ozone loss rates of 10.0 ± 1.3 ppb/sunlight hour (54 ± 7 ppb/day) whereas the corresponding box model simulations yield only 3.3 ± 0.2 ppb/sunlight hour (18 ± 1 ppb/day). Becker et al. (1998) further demonstrated that this discrepancy cannot be explained by known model uncertainties. Rex et al. (2002) show that this is the largest discrepancy of all the Match campaigns between 1992 and 2000: to explain the ozone loss rates in January 1992 with standard photochemistry the amount of active chlorine must exceed the maximum available chlorine Cl_y by a factor of 2.5.

The air masses within the polar vortex in early winter are often viewed as well isolated from mid-latitudes. However, this assumption is not true in all winters at all times. Plumb et al. (1994) show an example of a large-scale intrusion of mid-latitude air masses into the polar vortex in January 1992. Here, we use simulations of the Chemical Lagrangian Model of the Stratosphere (CLaMS) (McKenna et al., 2002a,b) to investigate the impact of such intrusions in January 1992 on ozone loss estimates because for this time both large scale intrusions took place and the largest discrepancy between Match and model was found.

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

2. Model simulations

The Chemical Lagrangian Model of the Stratosphere (CLaMS) is described in detail by McKenna et al. (2002a,b). The simulation was performed for the 475 K potential temperature level for the time period between 12 January and 4 February 1992. The horizontal resolution is 90 km between 30° and 90° N and 450 km south of 30° N corresponding to approximately 18 000 air parcels. Mixing between air parcels is induced when a critical Lyapunov coefficient λ exceeds the value of 1.5 using a mixing time step of 12 h (see McKenna et al., 2002a; Konopka et al., 2002, for details). Wind and temperature information was taken from ECMWF data with high spatial and temporal resolution ($1.125^\circ \times 1.125^\circ$, 6 h).

For the problem under investigation, a two-dimensional isentropic simulation is justified, because the simulation period is rather short (23 days). To estimate the amount of diabatic descent that the air parcels would have experienced, 16 600 3-D trajectories were calculated (without mixing) for the simulation period. The diabatic descent was calculated by a radiation scheme (Morcrette, 1991; Zhong and Haigh, 1995) using a 10-year climatology of HALOE ozone data as input (Grooß et al., manuscript in preparation). For this estimate, temperature and wind data were taken from the UKMO analyses that cover the complete stratosphere up to 0.3 hPa. The derived amount of descent for air parcels in the vortex is $21.5 \pm 2.4 \text{ K}(1\sigma)$ in 23 days which is less than 1 km. In the two-dimensional isentropic simulation we do not simulate the exact positions of the air parcels correctly because of the neglect of diabatic descent. Thus, an exact sampling of the match trajectories within the model domain as by Kilbane-Dawe et al. (2001) can not be performed here. However, the simulation still demonstrates the effect of the vortex intrusions very clearly.

To derive an initial ozone field at 475 K for the CLaMS simulation, we used version 5 data from the Microwave Limb Sounder (MLS) on board the Upper Atmosphere Research Satellite (UARS) (Barath et al., 1993) between 8 and 13 January. Using backward and forward trajectories, the locations of the observations were mapped to

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

the assimilation time (12 January 1992, 12:00 UT). These data were then combined on a regular grid with 1° latitude $\times$ 3° longitude spacing. Each grid point summarizes all data points within a distance less than 220 km using a cosine-square weighting with 110 km distance half-width. Figure 1 shows the derived ozone mixing ratio. The other species are initialized using the Mainz photochemical 2-D model (Gidel et al., 1983; Grooß, 1996) mapped onto equivalent latitude.

Figure 2 shows an example of the simulated ozone mixing ratio for 22 and 27 January on a stereographic projection. For illustration, Fig. 2 also shows the location of two Match trajectories with over 0.5 ppm diagnosed ozone loss and over 50 sunlight hours (violet symbols). On 22 January, a largely extended area of air masses outside the vortex between Scandinavia and Greenland with ozone mixing ratios below about 2.5 ppmv (blue colors) is visible. These air masses are drawn into the vortex as also simulated by Plumb et al. (1994). They can be seen as green colors over Newfoundland and eastward of Greenland on 27 January. The simulations show that with time the intruded air masses are deformed and elongated. In this context it is important that the horizontal scale of intruded air masses shrinks within one week below about 500 km which is the maximum allowed Match radius. Figure 2 also shows the individual Match radii of selected Match trajectories (≈ 300 km) as violet circles. On 27 January, a second tongue of air around 75° N, 180° E starts to be drawn into the vortex.

Figure 3 shows an enlargement of Fig. 2 over Scandinavia on 27 January around one Match trajectory location. This is within 24 h before it matches the second ozone sonde. The ozone variance within the match radius – 293 km for this trajectory – is evident. The impact of the variability of ozone mixing ratios within the Match radius on the deduced ozone loss rates is discussed below.

To be comparable with the Match analyses, we calculate the vortex average ozone mixing ratio from the model results using exactly the definition of the vortex edge that is used in the Match analysis Rex et al. (1998), i.e. the 36 PVU line from ECMWF data on the 475 K level. Figure 4a shows the ozone mixing ratio averaged over the vortex area defined in this way for the time of this simulation as a blue line. The red line shows the

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

vortex average of a passive tracer that was initialized identically to the ozone mixing ratio. As the tracer is not influenced by chemistry, its vortex average should be constant with time. However, from this analysis it is evident that the major fraction of the change in the vortex average ozone mixing ratio is due to the dynamically caused intrusions into the vortex but not due to chemical ozone depletion. The change of the dynamical ozone tracer in this simulation is a measure for the bias in derived chemical ozone depletion. To derive the chemical ozone loss, one should correct for this dynamical effect. In the following, only the simulated passive ozone tracer will be discussed.

In addition to the regular CLaMS simulation described above we performed a “virtual Match” calculation in order to quantify the expected impact of the vortex intrusions on the Match results. Figure 5 illustrates this method: At each CLaMS air parcel within the polar vortex on one day a trajectory (green line) was started to predict the location of the air parcel on a later day (large green circle). For each of the calculated locations, 8 new points were determined with the distance of a given match radius equally distributed in all directions (red circles). If a new point is still within the vortex, it was considered a Match with the first air parcel. The corresponding passive ozone tracer mixing ratios were then determined from the CLaMS simulation using the data from the closest CLaMS point (blue circles) to each new point. The average vortex ozone depletion rate was then determined from these Matches exactly as in the Match analyses.

3. Discussion

From Figs. 2 and 3 it is evident that the ozone mixing ratios in the polar vortex are not necessarily distributed uniformly on a potential temperature surface. Ozone variability within the polar vortex exists because of various reasons. In spring 1997, for example, a particularly pronounced inhomogeneity of ozone depletion within the polar vortex was both observed and simulated (Müller et al., 1997; Schulz et al., 2000; McKenna et al., 2002b; Tilmes et al., 2002). Also inhomogeneous diabatic descent rates and

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

intrusions at earlier times that have not yet been completely mixed with vortex air are possible reasons for variability of ozone inside the vortex on an isentropic surface.

5 Principally ozone variability should not significantly influence estimates of vortex averaged ozone loss in a confined vortex except that it increases the statistical error of the result. However, intrusions of a significant amount of air from mid-latitudes into the vortex cause a systematic offset of vortex-averaged ozone mixing ratios. Since the vortex-average method depends on the assumption that the vortex air masses are well isolated from mid-latitudes, it should lead to a bias in the derived ozone loss rate during a time period with vortex intrusions. In mid-January 1992, the ozone mixing ratios
10 below potential temperature levels of ≈ 500 K are lower outside the vortex than inside and vice versa above. Thus, the intrusions into the polar vortex would lead to an overestimate of chemical ozone depletion detected by this method at the 475 K level that is discussed here. Some studies employing the vortex-average method try to correct for cross vortex edge transport by trajectory simulations, e.g. Knudsen et al. (1998) find
15 no significant effect in the year 1997. However, that is clearly not true for other winters, especially for the winter 1991/92 discussed here.

From the time derivative of the vortex average passive ozone tracer mixing ratio (Fig. 4, bottom panel) the bias of derived ozone depletion rates due to vortex intrusions can be estimated. It shows the simulated vortex-averaged depletion rate for ozone
20 (red line) and the passive ozone tracer (blue line) for ± 2.5 days, i.e. for intervals of 5 day length. (We chose this length because it is comparable to the length of the match trajectories, since e.g. the trajectories that contribute to the data point for 24 January have an average length of 4.5 days.) The red line corresponds to the derived ozone depletion rate due to the intrusions only, while the blue line also includes the simulated
25 chemical ozone loss. Around 24 January, the derived dynamical bias is about 11 to 14 ppb/day.

The bias derived from vortex-averaged ozone tracer loss rate depends also on the definition of the vortex edge. If the vortex edge is moved poleward by using a greater PV value, the derived ozone depletion rate is lower and vice versa (e.g. ≈ 8 ppb/day for

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

$PV^{\text{edge}} = 40$ PVU and ≈ 24 ppb/day for $PV^{\text{edge}} = 30$ PVU on 24 January). However, the general shape of Fig. 4b does not change significantly.

The Match method is a more sophisticated approach than the vortex average method. It aims at avoiding such errors as discussed above since individual air parcels are examined that have been probed twice by an ozone sonde inside the vortex. It is therefore unlikely that earlier vortex observations match with air masses that have crossed the vortex edge during the time of the Match trajectory. However, there are displacements between the probed air mass and the location of the matching air mass given by the match radius and furthermore by uncertainties induced by the trajectory calculation. The Match trajectory errors due to uncertainties in the wind data can not be examined by this simulation since both the simulation and the Match trajectory calculation are based on the same meteorological analyses (ECMWF).

Figures 2 and 3 show that in 1992 the large-scale intrusions caused filaments with a width below the Match radius (Match resolution) within about one week. Therefore, the systematic offset described above should also influence the Match results in a similar manner. The results of the Match method can only account for vortex intrusions as long as the horizontal scale of the intrusions is much greater than the Match radius. The actual bias of the ozone depletion rate derived by the Match method depends further on the sampling of these filaments by the Match trajectories. If the matches are distributed equally over the vortex domain, the estimate of a systematic offset of vortex average ozone loss due to intrusions such as those discussed in this study should appear as a systematic error of the Match results as well.

In order to quantify the expected impact of the vortex intrusions on the Match results we performed an additional “virtual Match” calculation as described above. The trajectory length was chosen to be 4 days from 22 January to 26 January. The chosen Match radius was 300 km. For this case 3717 trajectories were started within the vortex of which 2978 stayed within the vortex over the 4 day period resulting in 21 768 virtual match events. Here, the dynamical bias was also determined from the passive ozone tracer from which the ozone depletion rate for the passive ozone tracer was derived

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

exactly as in the Match analysis. The resulting dynamical bias was 10.0 ± 0.4 ppb/day (1.94 ± 0.07 ppb per sunlight hour) which is 28% of the discrepancy found by Becker et al. (1998). This result does not seem to depend strongly on the chosen Match radius, e.g. for a Match radius of 450 km, the ozone depletion rate derived from the passive ozone tracer is 11.7 ± 0.4 ppb/day.

In the Match analysis for 24 January 1992, there are only 19 Match events. With this simulation we also tested what statistical error one should expect from this number of events. This was done by randomly choosing a subset of 20 Matches out of the 21 768 virtual Matches which was repeated 200 000 times. From each subset of 20 Matches the ozone depletion rate is evaluated exactly as in the regular Match analysis. The standard deviation of the so derived results is 2.9 ppb per sunlight hour which is a factor 2.2 greater than the derived Match error bar for 24 January. Figure 6 shows a histogram of the derived ozone loss rate bias for the subsets of 20 Matches. For 7.8% of the Match subsets, the deduced ozone depletion rate is equal or even larger than the published discrepancy (6.7 ppb per sunlight hour). However, it seems unlikely that an unfavorable sampling of the Match events causes the complete January ozone loss discrepancy because it was found similarly for other years as well (Becker et al., 2000; Kilbane-Dawe et al., 2001; Rex et al., 2002).

4. Conclusions

We have shown that intrusions of mid-latitude air into the polar vortex introduce a systematic offset of vortex average ozone mixing ratios. Estimates of ozone loss rates are influenced by this effect. If the intrusions reach horizontal dimensions below the Match resolution (Match radius) this systematic offset influences also the Match analyses. This effect does not solve the published discrepancy between Match deduced ozone loss rates and those derived from simulations but it may explain about 28% of the discrepancy that was found for January 1992 (Becker et al., 1998). Therefore the estimates of chemical ozone depletion derived from ozone sonde data need to be cor-

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

rected for this effect. Further it was shown that the error of the Match results derived from only 20 match events in an unhomogeneous ozone distribution caused by the intrusions in January 1992 must be a factor 2.2 greater. Although the intrusions of mid-latitude air into the polar vortex do not explain the discrepancy completely it is a possible reason why for January 1992 the discrepancy is the greatest so far (Rex et al., 2002).

Acknowledgements. The authors thank Peter von der Gathen, Markus Rex and other participants of a workshop on stratospheric ozone loss in Potsdam in March 2002 for fruitful discussions. We acknowledge the MLS team led by Joe Waters. We also thank the European Centre for Medium-Range Weather Forecasts and the United Kingdom Meteorological Office for providing meteorological analyses.

References

- Barath, F. T., Chavez, M. C., Cofield, R. E., Flower, D. A., Frerking, M. A., Gram, M. B., Harris, W. M., Holden, J. R., Jarnot, R. F., Kloezezan, W. G., Klose, G. J., Lau, G. K., Loo, M. S., Maddison, B. J., Mattauch, R. J., McKinney, R. P., Peckham, G. E., Pickett, H. M., Siebes, G., Soltis, F. S., Suttie, R. A., Tarsala, J. A., Waters, J. W., and Wilson, W. J.: The Upper Atmosphere Research Satellite microwave limb sounder instrument, J. Geophys. Res., 98, 10 751–10 762, 1993. [2492](#)
- Becker, G., Müller, R., McKenna, D. S., Rex, M., and Carslaw, K. S.: Ozone loss rates in the Arctic stratosphere in the winter 1991/92: Model calculations compared with Match results, Geophys. Res. Lett., 25, 4325–4328, 1998. [2491](#), [2497](#), [2506](#)
- Becker, G., Müller, R., McKenna, D. S., Rex, M., Carslaw, K. S., and Oelhaf, H.: Ozone loss rates in the Arctic stratosphere in the winter 1994/1995: Model simulations underestimate results of the Match analysis, J. Geophys. Res., 105, 15 175–15 184, 2000. [2491](#), [2497](#)
- EU: European research in the stratosphere 1996–2000, chap. 3.5, pp. 112–117, EUR 19867, Brussels, 2001. [2490](#)
- Gidel, L. T., Crutzen, P. J., and Fishman, J.: A two-dimensional photochemical model of the atmosphere; 1: Chlorocarbon emissions and their effect on stratospheric ozone, J. Geophys. Res., 88, 6622–6640, 1983. [2493](#)

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- Grooß, J.-U.: Modelling of Stratospheric Chemistry based on HALOE/UARS Satellite Data, PhD thesis, University of Mainz, 1996. [2493](#)
- Harris, N., Rex, M., Knudsen, B., Manney, G., Müller, R., and von der Gathen, P.: Comparison of empirically derived ozone loss rates in the Arctic vortex, *J. Geophys. Res.*, 107, 10.1029/2001JD000482, 2002. [2490](#)
- Kilbane-Dawe, I., Harris, N., Pyle, J., Rex, M., Lee, A., and Chipperfield, M.: A comparison of Match and 3D model ozone loss rates in the Arctic polar vortex during the winters of 1994/95 and 1995/96, *J. Atmos. Chem.*, 39, 123–138, 2001. [2491](#), [2492](#), [2497](#)
- Knudsen, B. M., Larsen, N., Mikkelsen, I. S., Morcrette, J.-J., Braathen, G. O., Kyrö, E., Fast, H., Gernandt, H., Kanzawa, H., Nakane, H., Dorokhov, V., Yushkov, V., Hansen, G., Gil, M., and Shearman, R. J.: Ozone depletion in and below the Arctic vortex for 1997, *Geophys. Res. Lett.*, 25, 627–630, 1998. [2490](#), [2495](#)
- Konopka, P., Grooß, J. U., Günther, G., McKenna, D. S., Müller, R., Elkins, J. W., Fahey, D., Popp, P., and Stimpfle, R. M.: Weak influence of mixing on the chlorine deactivation during SOLVE/THESEO2000: Lagrangian modeling (CLaMS) versus ER-2 in situ observations, *J. Geophys. Res.*, accepted, 2002. [2492](#)
- McKenna, D. S., Konopka, P., Grooß, J.-U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): Part I Formulation of advection and mixing, *J. Geophys. Res.*, 107, 10.1029/2000JD000114, 2002a. [2491](#), [2492](#)
- McKenna, D. S., Grooß, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): Part II Formulation of chemistry-scheme and initialisation, *J. Geophys. Res.*, 107, 10.1029/2000JD000113, 2002b. [2491](#), [2492](#), [2494](#)
- Morcrette, J.-J.: Radiation and cloud radiative properties in the European Centre for Medium-Range Weather Forecasts forecasting system, *J. Geophys. Res.*, 96, 9121–9132, 1991. [2492](#)
- Müller, R., Grooß, J.-U., McKenna, D., Crutzen, P. J., Brühl, C., Russell, J. M., and Tuck, A. F.: HALOE observations of the vertical structure of chemical ozone depletion in the Arctic vortex during winter and early spring 1996–1997, *Geophys. Res. Lett.*, 24, 2717–2720, 1997. [2494](#)
- Plumb, R. A., Waugh, D. W., Atkinson, R. J., Newman, P. A., Lait, L. R., Schoeberl, M. R., Browell, E. V., Simmons, A. J., and Loewenstein, M.: Intrusions into the lower stratospheric Arctic vortex during the winter of 1991–1992, *J. Geophys. Res.*, 99, 1089–1105, 1994. [2491](#),

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- Rex, M., von der Gathen, P., Harris, N. R. P., Lucic, D., Knudsen, B. M., Braathen, G. O., Reid, S. J., De Backer, H., Claude, H., Fabian, R., Fast, H., Gil, M., Kyrö, E., Mikkelsen, I., Rummukainen, M., Smit, H. G., Stähelin, J., Varotsos, C., and Zaitcev, I.: In situ measurements of stratospheric ozone depletion rates in the Arctic winter 1991/92: A Lagrangian approach, *J. Geophys. Res.*, 103, 5843–5853, 1998. [2490](#), [2493](#)
- Rex, M., Salawitch, R. J., Santee, M. L., Waters, J. W., Hoppel, K., and Bevilacqua, R.: On the unexplained stratospheric ozone losses during cold Arctic Januaries, *Geophys. Res. Lett.*, in press, 2002. [2491](#), [2497](#), [2498](#)
- Schulz, A., Rex, M., Steger, J., Harris, N. R. P., Braathen, G. O., Reimer, E., Alfier, R., Beck, A., Alpers, M., Cisneros, J., Claude, H., De Backer, H., Dier, H., Dorokhov, V., Fast, H., Godin, S., Hansen, G., Kanzawa, H., Kois, B., Kondo, Y., Kosmidis, E., Kyrö, E., Litynska, Z., Molyneux, M. J., Murphy, G., Nakane, H., Parrondo, C., Ravagnani, F., Varotsos, C., Vialle, C., Viatte, P., Yushkov, V., Zerefos, C., and von der Gathen, P.: Match observations in the Arctic winter 1996/97: High stratospheric ozone loss rates correlate with low temperatures deep inside the polar vortex, *Geophys. Res. Lett.*, 27, 205–208, 2000. [2494](#)
- Tilmes, S., Müller, R., Grooß, J.-U., McKenna, D., Russell, J., and Sasano, Y.: Calculation of chemical ozone loss in the Arctic winter 1996–1997 using ozone-tracer correlations: comparison of ILAS and HALOE results, *J. Geophys. Res.*, in press, 2002. [2494](#)
- von der Gathen, P., Rex, M., Harris, N. R. P., Lucic, D., Knudsen, B. M., Braathen, G. O., De Backer, H., Fabian, R., Fast, H., Gil, M., Kyrö, E., Mikkelsen, I. S., Rummukainen, M., Stähelin, J., and Varotsos, C.: Observational evidence for chemical ozone depletion over the Arctic in winter 1991–92, *Nature*, 375, 131–134, 1995. [2490](#)
- Zhong, W. and Haigh, J. D.: Improved broadband emissivity parameterization for water vapor cooling rate calculations, *J. Atmos. Sci.*, 52, 124–138, 1995. [2492](#)

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller




Fig. 1. Initialization for ozone on the 475 K level derived from MLS data. White areas indicate mixing ratios below 2.0 ppm. Also plotted is the PV contour of 36 PVU that is employed here as the definition of the vortex edge.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller




Fig. 2. Simulated ozone mixing ratio for 22 January (top) and 27 January (bottom). The violet stars and circles show the location and Match radius of two individual Match trajectories. The black contour corresponds to the 36 PVU line which is used here as the definition of the vortex edge. White areas indicate ozone mixing ratios below 2 ppm.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller




Fig. 3. Enlargement of Fig. 2, bottom panel in the vicinity one Match trajectory location.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller




Fig. 4. Vortex average of the simulations. The top panel shows the vortex average ozone mixing ratio (blue line) and the average passively advected tracer that was initialized identically as ozone (red line). The bottom panel shows the time derivative of the top panel, i.e. the ozone (or tracer) change per day for ± 2.5 day intervals.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller




Fig. 5. Sketch of the “virtual Match” calculation. The green line corresponds to one example trajectory ending at the large green circle. The Match radius here is 300 km indicated by the red circle. 8 new locations are marked by the red circles for which only those inside the vortex (filled circles) are considered. The vortex edge is plotted as a thick black line. Grey circles correspond to the individual CLaMS air parcel locations and those chosen for the Match are marked blue.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Impact of vortex intrusions on ozone loss estimates

J.-U. Grooß and R. Müller

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002




Fig. 6. Histogram of the derived ozone loss rate bias for 200000 subsets of 20 out of 21 768 virtual matches. The solid line shows the ozone loss rate bias derived from the passive ozone tracer for all virtual matches. The right dotted line shows the discrepancy between simulation and Match after Becker et al. (1998). 7.8% of the subsets have equal or larger values than this discrepancy.