Height of convective layer in planetary atmospheres with condensable and non-condensable greenhouse substances

A. M. Makarieva, V. G. Gorshkov, T. Pujol

► To cite this version:

A. M. Makarieva, V. G. Gorshkov, T. Pujol. Height of convective layer in planetary atmospheres with condensable and non-condensable greenhouse substances. *Atmospheric Chemistry and Physics Discussions*, 2003, 3 (6), pp.6701-6720. hal-00301398

HAL Id: hal-00301398

<https://hal.science/hal-00301398>

Submitted on 16 Dec 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Height of convection

A. M. Makarieva et al.

Height of convective layer in planetary atmospheres with condensable and non-condensable greenhouse substances

A. M. Makarieva¹, V. G. Gorshkov¹, and T. Pujol²

¹Theoretical Physics Division, Petersburg Nuclear Physics Institute, Gatchina, St.-Petersburg, Russia

²Department of Physics, University of Girona, Girona, Catalonia, Spain

Received: 25 October 2003 – Accepted: 8 December 2003 – Published: 16 December 2003

Correspondence to: A. M. Makarieva (elba@infopro.spb.su)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

Abstract

Convection reduces greenhouse effect by transporting a certain amount of non-radiative dynamic energy to the upper atmosphere, where this energy dissipates and radiates into space without interaction with greenhouse substances in the lower atmosphere. In this paper we show that the height of the convective layer z_c is finite and independent of atmospheric optical thickness τ_s at large values of the latter. We derive an analytical formula for z_c at large values of τ_s for condensable and non-condensable greenhouse substances. The formula obtained yields reasonable quantitative estimates of the observed height of convective layer on Venus and at low latitudes on Earth, where atmospheric thickness of water vapor is maximum.

The dissipative power of dynamic convective processes is limited by the incoming flux of solar radiation. Height of convective layer being finite, values of optical depth at the top of the convective layer and at the mean height of convective energy dissipation increase proportionally to the atmospheric optical thickness, while the contribution of convective energy fluxes to formation of the outgoing flux of thermal radiation proportionally diminishes. As far as optical thickness of condensable greenhouse substances grows exponentially with increasing surface temperature, the obtained results lead to the conclusion that the outgoing thermal radiation into space in the presence of convection tends exponentially to zero with increasing surface temperature, instead of reaching a finite plateau as suggested by earlier radiative-convective studies.

1. Introduction

In radiative equilibrium, when energy is transported in the atmosphere by means of radiative processes only, the ratio of the outgoing flux of thermal radiation into space to the upward flux of thermal radiation at the Earth's surface is, in accordance with Eddington's approximation, inversely proportional to the amount of greenhouse substances in the atmosphere. The atmospheric content of the major greenhouse substance on Earth

Height of convection

A. M. Makarieva et al.

Title Page

AbstractIntroduction

ConclusionsReferences

TablesFigures

◀▶

◀▶

BackClose

Full Screen / Esc

Print Version

Interactive Discussion

– water vapor – in the presence of liquid hydrosphere rises exponentially with increasing surface temperature. With an account made for the relevant increase in cloudiness absorbing thermal radiation over entire thermal spectrum, this corresponds to an exponential decline of the outgoing flux of thermal radiation into space with increasing surface temperature.

Non-radiative convective processes could transport a certain amount of dynamic energy to the upper radiative layer of the atmosphere. There this energy would dissipate and radiate directly into space, avoiding interaction with greenhouse substances in any atmospheric layers lower than the upper radiative one. If this convective contribution to the flux of thermal radiation into space were independent of the amount of greenhouse substances in the atmosphere, the thermal flux into space would not diminish exponentially with surface temperature, as in the case of radiative equilibrium, but ultimately reached a non-zero plateau. Such a scenario was suggested by several radiative-convective studies (Komabayasi, 1967; Ingersoll, 1969; Kasting, 1988; Goody and Yung, 1989; Nakajima et al., 1992; Rennó, 1997; Pujol and North, 2002). Such a scenario would have profoundly different implications for the thermal evolution of the planets than the exponentially diminishing thermal flux into space. It is important therefore to verify carefully the physical assumptions underlying the two scenarios.

A constant convective contribution to thermal flux into space implies that at any value of atmospheric optical thickness convection propagates up to a fixed optical depth, e.g. to the upper radiative layer of the atmosphere, which corresponds to optical depth of about unity. In this paper we demonstrate that convection propagates up to a fixed atmospheric height rather than to a fixed optical depth. The optical depth corresponding to the top of the convective layer grows proportionally to atmospheric optical thickness. Given that the dissipative power of the dynamic convective processes is limited by the incoming solar flux, this means that the convective contribution to thermal flux into space diminishes inversely proportionally to the growing atmospheric optical thickness and does not reach a finite plateau at high surface temperatures. This result allows to conclude that at large values of atmospheric optical thickness the condition of ra-

Height of convectionA. M. Makarieva et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

diative equilibrium described by Eddington's approximation is fulfilled not only in the stratosphere, but spreads to the entire atmosphere.

5 In Sect. 2 we consider how height of convective layer changes with increasing optical thickness of an atmosphere containing non-condensable greenhouse gases only (like CO₂). We show that the height of convective layer is finite and independent of atmospheric optical thickness at large values of the latter. We demonstrate that the developed theoretical approach yields the exact numerical value for the height of convective layer on Venus.

10 In Sect. 3.1 we extend our consideration to condensable greenhouse substances and show that the results obtained are valid for this case as well. We estimate the maximum height of convective layer in an atmosphere saturated with water vapor. The obtained estimate displays a good agreement with the observed height of convection at low latitudes on Earth, where atmospheric content of water vapor is maximum.

15 In Sect. 3.2 we show that the presence of convection in the terrestrial atmosphere does not decelerate the exponential decline of the outgoing flux of thermal radiation into space with growing surface temperature.

2. Convection in an atmosphere with non-condensable greenhouse substances

2.1. Brightness temperature

20 In the state of radiative equilibrium the dependence between the upward fluxes of thermal radiation into space, F_e , and at optical depth τ in the atmosphere, $F^+(\tau)$, in a spectral interval of resonance absorption of thermal radiation by atmospheric greenhouse substances is accurately described by Eddington's approximation (Michalas and Michalas, 1984):

$$F^+(\tau) = F_e(1 + k\tau), \quad k = \frac{3}{4}; \quad F_s^+ = F_e(1 + k\tau_s). \quad (2.1)$$

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

Here and below low index s refer to values at the Earth's surface. At the Earth's surface the upward flux of thermal radiation is related to surface temperature T_s by Stephen-Boltzmann formula for the black body radiation. For any optical depth τ (and corresponding height z) the so-called brightness temperature T_b in a given spectral interval

5 $\Delta\lambda \equiv \lambda_2 - \lambda_1$ is defined as follows:

$$F^+(\tau) = \int_{\lambda_1}^{\lambda_2} I_P(\lambda, T_b) d\lambda \equiv \delta_{\Delta\lambda} \sigma T_b^4, \quad F_s^+ = \delta_{\Delta\lambda} \sigma T_s^4. \quad (2.2)$$

Here T_s is the surface temperature, σ is the Stephen-Boltzmann constant, $I_P(\lambda, T_b)$ is Planck's function, $\delta_{\Delta\lambda}$ is the relative share of the observed flux $F^+(\tau)$ of thermal radiation within spectral interval $\Delta\lambda$ in the total flux of black body radiation of temperature T_b . We assume $\delta_{\Delta\lambda}$ to be independent of T_b and height z . This assumption is valid
10 for a grey greenhouse substance (e.g. clouds or gas above the critical point) for which $\delta_{\Delta\lambda} \sim 1$. It is also valid when brightness temperature in the interval $\Delta\lambda$ changes relatively little within the convective layer.

Using Eqs. (2.1) and (2.2) we obtain the following expression of brightness temperature $T_b \equiv T_b(z)$ in terms of optical depth $\tau \equiv \tau(z)$:
15

$$T_b = T_s \left(\frac{1 + k\tau}{1 + k\tau_s} \right)^{1/4}. \quad (2.3)$$

Brightness temperature is maximum at the Earth's surface and decreases monotonically with growing height z due to the monotonical decrease of τ with height z . The lapse rate of brightness temperature $G_b \equiv -dT_b/dz$ also monotonically decreases
20 with height.

We note that at any height $z > 0$ the local air temperature T is higher than the brightness temperature T_b , $T(z) > T_b(z)$. This follows from the direction of energy conversion: the energy of non-radiative fluxes which, upon its dissipation, is characterized by air temperature T , is converted to the energy of the excited levels of molecules of

greenhouse substances with subsequent emission of thermal radiation characterized by brightness temperature T_b . The relation $T > T_b$ for $z > 0$ and the monotonical smooth decline of both T and T_b with height mean that the lapse rate of air temperature $G \equiv -dT/dz$ is always lower than the lapse rate of brightness temperature

5 $G_b \equiv -dT_b/dz$.

2.2. Temperature lapse rate and the onset of convection

The gaseous atmosphere in the gravitational field of Earth conforms to the equation of hydrostatic equilibrium, $-dp/dz = MNg$, and the equation of state for atmospheric air, $p = NRT$, where p , N and T are pressure, molar concentration and temperature of air

10 at height z , M is air molar mass assumed to be independent of height z in accordance with observations, $R = 8.3 \text{ J mole}^{-1} \text{ K}^{-1}$ is gas constant, g is acceleration of gravity. The two equations can be combined in the following well-known form:

$$\frac{dp}{dz} = -\frac{p}{h}; \quad h \equiv \frac{RT}{Mg}. \quad (2.4)$$

Relation (2.4) is valid for all gases with constant mass fraction, in particular, for non-condensable greenhouse substances like CO_2 . Due to the observed constant molar mass of air, the ratio p/p_s in Eq. (2.4) is, as is well-known, equal to the corresponding ratio of partial pressures of such gases and to the ratio τ/τ_s , i.e. $p/p_s = \tau/\tau_s$. Using this equality together with Eqs. (2.3) and (2.4) we obtain:

15

$$G_b = -\frac{dT_b}{dz} = G_{\max} \alpha_b, \quad G_{\max} \equiv \frac{T_s}{4h_s} = \frac{Mg}{4R},$$

$$\alpha_b = \frac{k\tau}{(1+k\tau)^{3/4}(1+k\tau_s)^{1/4}} \cdot \frac{T_s}{T}. \quad (2.5)$$

Here h_s is the scale height of distribution of non-condensable gases in the atmosphere. The fundamental value of $G_{\max}$ is a convenient scale for representing any temperature lapse rates G_i in the form of dimensionless variables $\alpha_i \equiv G_i/G_{\max}$.

20

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

Convective processes in the atmosphere set in when the lapse rate of air temperature exceeds some critical value Γ (Chandrasekhar, 1957; Ramanathan and Coakley, 1978; Weaver and Ramanathan, 1995). In the terrestrial atmosphere $G_{\max} = 8.56 \text{ K km}^{-1}$, while convection exists at the observed lapse rate $\Gamma = 6.5 \text{ K km}^{-1}$. This value can be viewed as the critical lapse rate for the advent of convection in the terrestrial atmosphere (Ramanathan and Coakley, 1978). The condition $\Gamma < G_{\max}$ implied by Eq. (2.5) is therefore fulfilled in the terrestrial atmosphere.

With increasing amount of greenhouse substances in the atmosphere the lapse rate of brightness temperature starts to grow, first of all at the surface, as far as $G_{bs} > G_b(z)$ for any $z > 0$. At some value of atmospheric optical thickness $\tau_s = \tau_{s \min}$ the lapse rate of brightness temperature at the surface may reach the critical value, $G_{bs} = \Gamma$, so that it becomes possible for convection to originate. It happens if the lapse rate of air temperature follows G_b closely enough. With increasing $\tau_s > \tau_{s \min}$ the convective layer spreads upwards till yet larger and larger heights.

Once convection sets in, it works to keep the lapse rate of air temperature G close to the critical value of Γ (Ramanathan and Coakley, 1978). Thus, within the convective layer $z \leq z_c$ the lapse rate of air temperature is approximately constant, $G \approx \Gamma$, and air temperature obeys the observed relation:

$$T = T_s - \Gamma z = T_s \left(1 - \frac{\alpha z}{4 h_s} \right), \quad \alpha \equiv \frac{\Gamma}{G_{\max}}. \quad (2.6)$$

For Earth $\Gamma = 6.5 \text{ K km}^{-1}$ and $\alpha = 0.77$. At $z > z_c$ there is no convection, and Eq. (2.6), as well as any of its derivations, may not be used for description of air temperature.

Using Eqs. (2.4) and (2.6) we obtain the following well-known relation for $z \leq z_c$:

$$\frac{\tau}{\tau_s} = \frac{\rho}{\rho_s} = \left(1 - \frac{\alpha z}{4 h_s} \right)^{4/\alpha}. \quad (2.7)$$

Convection exists in that part of the atmosphere where air temperature lapse rate G exceeds the critical value Γ , $G > \Gamma$. As far as $G < G_b$, the maximum height z_c of

convective layer at a given value of optical thickness τ_s and surface temperature T_s can be obtained from the equation $G_b(z_c) = \Gamma$ or, using Eqs. (2.5) and (2.6), from the equation

$$\alpha_b = \alpha. \quad (2.8)$$

- 5 Variables τ and T in Eq. (2.8) are related to height z by Eqs. (2.7) and (2.6), respectively. Equation (2.8) determines the maximum height of the convective layer $z = z_c$ at different values of atmospheric optical thickness τ_s , Fig. 1.

The threshold atmospheric optical thickness $\tau_{s \min}$ is obtained from Eq. (2.8) by putting $z = 0$, which means $\tau = \tau_s$ and $T = T_s$:

$$10 \quad \tau_{s \min} = \frac{1}{k} \frac{\alpha}{1 - \alpha}. \quad (2.9)$$

- For Earth we obtain $\tau_{s \min} = 4.5$. The major non-condensable greenhouse substance in the terrestrial atmosphere is CO_2 . Atmospheric optical thickness of CO_2 in the $15 \mu\text{m}$ spectral interval is estimated as $k\tau_{s \text{CO}_2} \approx 1.9$ (Gorshkov and Makarieva, 2002), which corresponds to $\tau_{s \text{CO}_2} = 2.5$ at $k = 3/4$. Thus, the atmospheric CO_2 concentration on Earth is insufficient for switching convection on. As we show in the next section, convection on Earth is switched on by atmospheric water vapor. On Venus, on the contrary, atmospheric CO_2 concentration appears to be by far sufficient for convection to originate, see below.

2.3. Height of convective layer at large values of τ_s

- 20 As one can see from Fig. 1, the height of convective layer z_c reaches a finite limit at large values of atmospheric optical thickness. For $\tau_s \rightarrow \infty$ ($\tau_s \gg 1$) one can use Eqs. (2.5)–(2.7) to re-write Eq. (2.8) as

$$\left(\frac{p}{p_s}\right)^{1/4} \frac{T_s}{T} = \alpha. \quad (2.10)$$

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Putting Eqs. (2.6) and (2.7) into Eq. (2.10) we obtain an exact finite value for the height of the convective layer $z_{c \max}$ at $\tau_s \rightarrow \infty$:

$$\frac{z_{c \max}}{h_s} = 4 \frac{1}{\alpha} \left(1 - \alpha^{\alpha/(1-\alpha)} \right) = 3.03. \quad (2.11)$$

For modern Earth we have $T_s = 288$ K, $h_s = 8.4$ km and $z_{c \max} = 25$ km. Eq. (2.11) describes the maximum height of the convective layer that could form in the terrestrial atmosphere with non-condensable greenhouse substances as the only absorbers of thermal radiation.

Minimum values of air temperature $T_c \equiv T(z_c)$ and brightness temperature $T_{bc} \equiv T_b(z_c)$ at the top of the convective layer, $z_c = z_{c \max}$, at large values of atmospheric optical thickness, $\tau_s \gg 1$, are obtained by substituting Eq. (2.11) into Eq. (2.6), and Eq. (2.11) into Eqs. (2.7) and (2.3), respectively:

$$\begin{aligned} \frac{T_c}{T_s} &= \alpha^{\alpha/(1-\alpha)}, \quad \frac{T_{bc}}{T_s} = \alpha^{1/(1-\alpha)}; \\ \frac{T_c}{T_s} &= 0.42, \quad \frac{T_{bc}}{T_s} = 0.32, \quad \alpha = 0.77. \end{aligned} \quad (2.12)$$

The numeric values in the second line of Eq. (2.12) correspond to modern Earth.

As is clear from Eq. (2.12), brightness temperature T_{bc} at the top of the convective layer grows proportionally to surface temperature T_s . Upward flux F_c^+ of thermal radiation at the top of the convective layer increases therefore proportionally to T_s^4 . The dissipative power of non-radiative convective fluxes of energy and, consequently, collisional excitation of the absorption bands of greenhouse substances, are limited by the incoming flux of solar radiation F_{in} , which does not change with increasing τ_s or T_s . (We note that in the non-stationary case the absorbed flux of solar radiation F_{in} is not equal to the outgoing flux of thermal radiation F_e , $F_{in} \neq F_e$, and also that $F_s^+/F_{in} \neq 1 + k\tau_s$.) When the upward flux of thermal radiation at the surface is much larger than the absorbed flux of solar radiation, $F_s^+ = \sigma T_s^4 \gg F_{in}$, convective processes

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

are unable to significantly perturb the upward fluxes of thermal radiation and brightness temperature within the convective layer as compared to the case of radiative equilibrium. With growing τ_s and T_s the radiative equilibrium spreads from the stratosphere to the entire atmosphere, so that Eddington's approximation becomes valid for the entire atmosphere as well. Therefore, Eqs. (2.11) and (2.12) obtained with use of Eddington's approximation yield at $F_s^+ \gg F_{in}$ and $\tau_s \gg 1$ the exact values of z_c , T_{bc} and T_c .

On Venus the ratio of the upward flux of thermal radiation at the surface F_s^+ to the outgoing flux of thermal radiation into space $F_e = F_{in}$ equals $F_s^+/F_e = 98$ (Mitchell, 1989). In accordance to Eddington's approximation (2.1) this corresponds to $\tau_s = \frac{4}{3}(F_s^+/F_e) = 131 \gg 1$. Thus, Eqs. (2.11) and (2.12) can be applied to Venus with a high accuracy. Taking for Venus $h_s = 15.6$ km, $G_{max} = 11.4$ K km⁻¹ ($T_s = 730$ K, $M_{CO_2} = 44$ g mole⁻¹, $g = 8.6$ m s⁻² (Allen, 1973; Kasting, 1988; Mitchell, 1989)), the observed air temperature lapse rate $\Gamma \approx 8$ K km⁻¹ (Goody and Yung, 1989), we obtain for Venus $\alpha \equiv \Gamma/G_{max} = 0.70$. Putting these values into Eq. (2.11), we obtain the following values for the height $z_{c\ max}$ of the convective layer on Venus, $z_{c\ max} = 50.4$ km. This value agrees very well with the observed height of cloudiness on Venus (~50 km), where, according to observations (Goody and Yung, 1989), the air temperature lapse rate starts to diminish becoming less than the critical convective value. From (2.12) we calculate air temperature at $z_{c\ max}$ for Venus to be $T_c = 318$ K (44°C).

We note that the height of convection z_c is generally less than the height of the tropopause z_{tp} , $z_c < z_{tp}$. The latter is usually defined as the point where the lapse rate of air temperature approaches zero. As suggested by Fig. 1, the lapse rate of brightness temperature and, consequently, lapse rate of air temperature may remain substantial for $z > z_c$, being nevertheless smaller than the critical value Γ responsible for the presence of convection.

3. Convection in an atmosphere with condensable greenhouse substances

3.1. Height of convective layer

The dependence of saturated water vapor pressure p_L on temperature is described by Clausius-Clapeyron equation (everywhere below the low index L denotes variables corresponding to water vapor):

$$p_L = p_{sL} \exp \left(\frac{T_L}{T_s} - \frac{T_L}{T} \right), \quad T_L \equiv \frac{Q}{R} \approx 5300 \text{ K}, \quad (3.1)$$

where $Q \approx 44 \text{ KJ mole}^{-1}$ is the latent heat of evaporation of water vapor, p_{sL} is partial pressure of water vapor at the surface.

In an isothermal atmosphere water vapor can be saturated only immediately above the water surface, $z = 0$. In the gravitational field the concentration of water vapor drops exponentially with height, so that at $z > 0$ water vapor becomes unsaturated.

If we demand that water vapor in the atmosphere is saturated and is in hydrostatic equilibrium at all heights z , the corresponding lapse rate Γ_L of air temperature is obtained from the condition that both Eqs. (2.4) and (3.1) hold:

$$\Gamma_L \equiv -\frac{dT}{dz} = \frac{T_s}{H_L} \exp \left(-\frac{z}{H_L} \right) \approx \frac{T_s}{H_L}, \quad (3.2)$$
$$H_L \equiv \frac{RT_L}{gM} = 154 \text{ km}.$$

Due to the large value of H_L Eq. (3.1), for a relatively wide interval of surface temperature changes Γ_L is practically constant (independent of z) in the entire troposphere.

If the lapse rate of air temperature G were smaller than Γ_L , atmospheric water vapor would be unsaturated at all heights z .

In the modern atmosphere of Earth we have

$$G \approx \Gamma = \beta_s \Gamma_L, \quad \beta_s = \frac{\alpha H_L}{4h_s} = \alpha \frac{T_L}{4T_s} = 3.5. \quad (3.3)$$

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

As far as the observed lapse rate $G \approx \Gamma$ of air temperature is steeper than Γ_L , $\Gamma > \Gamma_L$, the distribution of water vapor in the atmosphere differs from the hydrostatic equilibrium distribution (2.4). Calculating dp_L/dz with use of Eqs. (3.1) and (2.4) we find for the convective layer $z < z_{cL}$:

$$\frac{dp_L}{dz} = -\frac{p_L}{h_L}, \quad h_L = \frac{h}{\beta}, \quad \beta \equiv \alpha \frac{T_L}{4T} \approx \beta_s = 3.5. \quad (3.4)$$

As is clear from Eq. (3.4), the modern distribution of water vapor in the atmosphere is compressed by $\beta \approx 3.5$ times as compared to distribution of air (Weaver and Ramanathan, 1995). The scale height of water vapor distribution in the modern atmosphere equals $h_{sL} \equiv h_s/\beta_s \approx 2.4$ km, which agrees well with observations (Goody and Yung, 1989; Weaver and Ramanathan, 1995). Compression of water vapor in the modern atmosphere significantly violates hydrostatic equilibrium and brings about intensive dynamic fluxes of hydrosphere evaporation, upward transport of evaporated water and compensating downward fluxes of precipitation.

Assuming approximate constancy of β in Eq. (3.4) we obtain an approximate relation similar to Eq. (2.7):

$$\frac{\tau_L}{\tau_{sL}} \approx \frac{p_L}{p_{sL}} \approx \left(1 - \frac{\alpha_L}{4} \frac{z}{h_{sL}}\right)^{4/\alpha_L} = \left(\frac{p}{p_s}\right)^{\beta_s}, \quad z < z_c, \quad (3.5)$$

$$h_{sL} \equiv \frac{h_s}{\beta_s}, \quad \alpha_L \equiv \frac{\alpha}{\beta_s}.$$

Due to the large value of β_s and the rapid drop of pressure with height, relation Eq. (3.5) differs from Eq. (3.1) for the area of interest $z \leq h_{sL}$ by magnitudes of the order of $\alpha_L/4 = T_s/T_L \sim 0.05$, i.e. by less than 5%.

Relation (3.5) for condensable greenhouse substances appear to be identical to the corresponding relations (2.4) for non-condensable greenhouse substances if one substitutes α and h_s by α_L and h_{sL} , respectively, the latter being β_s times lower than the for-

mer. Similarly, the maximum value of brightness temperature lapse rate $G_{\max} = T_s/4h_s$ (2.5) becomes for water vapor β_s times larger:

$$G_{L \max} = T_s/4h_{sL} = G_{\max}\beta_s, \quad (3.6)$$

i.e. equal to 30 K km^{-1} for Earth.

5 The expression for maximum height of the convective layer determined by water vapor, $z_{cL \max}$, is obtained from Eq. (2.8) by a similar substitution $\alpha \rightarrow \alpha_L$, $\tau \rightarrow \tau_L$ and $\tau_s \rightarrow \tau_{sL}$, where $\alpha_L = \alpha/\beta_s = 0.22$, τ_L is determined by Eq. (3.5) and temperature T is determined by Eq. (2.6). The onset of convection in the terrestrial atmosphere due to the greenhouse effect of water vapor occurs at $\tau_{sL \min} = 0.38$, Fig. 2, which is more
10 than ten times smaller than $\tau_{s \min} = 4.5$ for non-condensable greenhouse substances like CO_2 , Fig. 1. The atmospheric optical thickness of water vapor in the terrestrial atmosphere is estimated as $k\tau_{sL} \approx 0.53$ (Gorshkov and Makarieva, 2002), which exceeds the value of $\tau_{sL \min}$ by almost twofold. This makes it possible to conclude that convection in the terrestrial atmosphere is switched on and maintained by water va-
15 por. The idea that water vapor is more efficient with respect to switching on convection than non-condensable absorbers like CO_2 due to the above discussed compression of water vapor in the modern atmosphere, was put forward by Weaver and Ramanathan (1995).

Note that in the intervals of $\tau_{sL} < \tau_{sL \min}$ corresponding to $\Gamma > G > \Gamma_L$ water va-
20 por is already not in hydrostatic equilibrium (i.e. there are fluxes of evaporation and precipitation), while atmospheric convection is still absent.

At large values of optical thickness $\tau_s \gg 1$ Eqs. (2.11) and (2.12) retain their form for water vapor if one substitutes $h_s \rightarrow h_{sL} = h_s/\beta_s$ and $\alpha \rightarrow \alpha_L = \alpha/\beta_s$:

$$\begin{aligned} z_{cL \max} &= 4h_{sL} \frac{1}{\alpha_L} \left(1 - \alpha_L^{\alpha_L/(1-\alpha_L)}\right) \approx \\ &\approx 4 \frac{h_s}{\beta_s} \ln \frac{\beta_s}{\alpha} = 6.3 \frac{h_s}{\beta_s} = 15 \text{ km}. \end{aligned} \quad (3.7)$$

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

The values of air temperature T_{cL} and brightness temperature T_{bcL} at the top of convective layer at $z = z_{cL \max}$ are obtained from Eqs. (2.12) by substituting α for α_L . At $T_s = 288$ K and $\alpha_L = 0.22$ we have $T_{cL} = 188$ K = -85°C .

On Earth the atmospheric content of water vapor is strongly dependent on surface temperature rising approximately twofold per each ten degrees of surface temperature increase, see Eq. (3.1). The maximum values of atmospheric optical thickness of water vapor are therefore to be observed in the equatorial regions (Raval and Ramanathan, 1989). In accordance with Eq. (3.7), in these regions one also observes maximum height of the convective layer of the order of 15 km and minimum air temperatures of the order of -80°C . In the polar regions where surface temperature and, consequently, atmospheric water vapor content, are much lower, one should expect to observe significantly lower convective layer and higher temperatures at its top, which is indeed the pattern observed.

3.2. Outgoing flux of thermal radiation into space

Clausius-Clapeyron equation (3.1) can be viewed as determining partial pressure of saturated water vapor at an arbitrary height z at an arbitrary moment of time t . It is easy to see that Eq. (3.1) conforms to the following invariance rule:

$$p_{sL} \exp\left(\frac{T_L}{T_s}\right) = p_{sL0} \exp\left(\frac{T_L}{T_{s0}}\right), \quad (3.8)$$

where T_{s0} and p_{sL0} are the values of surface temperature and partial pressure at the initial moment of time $t = 0$, which can be taken corresponding to the state of thermal equilibrium $F_e = F_{in}$. It is easy to see from Eq. (3.8) that optical thickness of atmospheric water vapor τ_{sL} , as well as partial pressure of water vapor at the Earth's surface p_{sL} , see Eq. (3.5), increase exponentially with growing increment of surface

temperature $\theta \equiv T_s - T_{s0}$ ($T_{s0} = 288$ K):

$$p_{sL} = p_{sL0} \exp\left(\frac{\theta}{\theta_s}\right), \quad \theta_s \equiv \frac{T_{s0} T_s}{T_L}, \quad \theta_{s0} \equiv \frac{T_{s0}^2}{T_L} = 18 \text{ K}. \quad (3.9)$$

This means that at $\tau_{sL} \gg 1$ the ratio of the outgoing flux of thermal radiation F_{Le} to the upward flux of thermal radiation at the surface F_{sL}^+ within the absorption interval of water vapor drops exponentially with growing θ :

$$\frac{F_{eL}}{F_{sL}^+} \propto \frac{1}{\tau_{sL}} = \frac{1}{\tau_{sL0}} \exp\left(-\frac{\theta}{\theta_s}\right). \quad (3.10)$$

In a number of radiative-convective studies where semigrey atmospheres with condensable greenhouse substances were considered, the outgoing flux of thermal radiation into space at high surface temperatures was suggested to remain practically constant (Komabayasi, 1967; Ingersoll, 1969; Kasting, 1988; Goody and Yung, 1989; Nakajima et al., 1992; Pujol and North, 2002). The major assumption responsible for this result is that the entire atmosphere, including the upper radiative layer, is in hydrostatic equilibrium and saturated with respect to water vapor.

The constancy of F_{Le} follows immediately from this assumption. Indeed, in the case of hydrostatic equilibrium the condition $\tau_L = 1$ (upper radiative layer, where the outgoing flux is formed) unambiguously determines partial pressure of water vapor at this height, which appears to be independent of surface temperature and atmospheric optical thickness, see Eq. (2.4). On the other hand, as far as partial pressure of saturated water vapor is unambiguously related to ambient air temperature in accordance with Clausius-Clapeyron equation (3.1), the temperature T_1 at the upper radiative layer appears independent of surface temperature either. In the case of constant T_1 (and assuming local thermodynamic equilibrium), the outgoing thermal radiation into space does not change with growing surface temperature and atmospheric optical thickness.

However, height of convective layer being finite, optical depth at the top of the convective layer τ_c increases with growing τ_s . At some value of the latter the upper radiating

layer of the atmosphere $\tau \sim 1$ goes outside the convective layer. There the lapse rate of air temperature at $\tau \sim 1$ drops radically and atmospheric water vapor becomes unsaturated. The dependence between its partial pressure and ambient air temperature, prescribed by Clausius-Clapeyron equation in the case of saturation, vanishes.

5 The fact that the height of convective layer at low latitudes in the modern Earth's climate is close to its maximum value (3.7) testifies that the upper radiative layer corresponding to certain absorption lines of the water vapor absorption interval is located above the convective layer. This suggests that the surface temperatures at which the exponential drop of outgoing radiation is to be observed are not drastically different
10 from those at modern Earth.

4. Conclusions

The main result of this paper is that, in contrast to the results obtained in several previous studies (Komabayasi, 1967; Ingersoll, 1969; Kasting, 1988; Goody and Yung, 1989; Nakajima et al., 1992; Rennó, 1997; Pujol and North, 2002), on a planet with
15 an infinite liquid hydrosphere the outgoing flux of thermal radiation into space does not reach a finite plateau with growing surface temperature but diminishes exponentially.

To arrive to this result, we showed that the height of convective layer in an atmosphere with condensable and non-condensable absorbers is finite (in terms of atmospheric scale height h_s) and independent of optical thickness of the atmosphere τ_s at large values of the latter. As far as the dissipative power of the dynamic convective processes is limited by the absorbed solar radiation, the obtained result means that the contribution of convective processes to formation of the outgoing flux of thermal radiation into space tends to zero with increasing atmospheric optical thickness inversely proportionally to the latter. The radiative equilibrium heat transfer described
20 by Eddington approximation becomes the dominant form of heat transfer in the entire atmosphere including the troposphere.

25 We obtained the exact numerical value of the height of convective layer at the con-

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

dition of large values of optical thickness τ_s , which applies to Venus and the equatorial region of Earth (for some part of thermal spectrum). We also showed that at large values of atmospheric optical thickness water vapor at the upper radiative layer will be unsaturated.

5 The assumption of saturation of water vapor in the entire atmosphere, including upper radiative layer, which was used in several radiative-convective studies (Komabayasi, 1967; Ingersoll, 1969; Kasting, 1988; Goody and Yung, 1989; Nakajima et al., 1992; Pujol and North, 2002), may not be used therefore for calculation of the outgoing flux of thermal radiation into space.

10 In the presence of an infinite liquid hydrosphere the outgoing thermal radiation into space should decrease exponentially with growing surface temperature. This is a consequence of an exponential growth with increasing surface temperature of the amount of water in the atmosphere, including cloudiness absorbing thermal radiation over entire thermal spectrum. We conclude that the account of convective processes does not
15 alter this basic physical mechanism of climate change on a planet with liquid hydrosphere.

References

- Allen, C. W.: Astrophysical Quantities, Athlone Press, London, 1973.
Chandrasekhar, S.: An Introduction in the Study of Stellar Structure, Dover, New York, 1957.
20 Goody, R. M. and Yung, Y. L.: Atmospheric radiation, theoretical basis, 2nd edn., Oxford Univ. Press, New York, 1989.
Gorshkov, V. G. and Makarieva, A. M.: Greenhouse effect dependence on atmospheric concentrations of greenhouse substances and the nature of climate stability on Earth, Atmos. Chem. Phys. Discuss., 2, 289–337, 2002.
25 Ingersoll, A. P.: The runaway greenhouse: A history of water on Venus, J. Atmos. Sci., 26, 1191–1198, 1969.
Kasting, J. F.: Runaway and moist greenhouse atmospheres and the evolution of Earth and Venus, Icarus, 74, 472–494, 1988.

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Komabayasi, M.: Discrete equilibrium temperatures of a hypothetical planet with the atmosphere and the hydrosphere of one component-two phase system under constant solar radiation, J. Meteorol. Soc. Japan 45, 137–138, 1967.

Michalas, D. and Michalas, B. W.: Foundations of radiation hydrodynamics, Oxford Univ. Press, New York, 1984.

Mitchell, J.: The “greenhouse” effect and climate change, Rev. Geophys., 27, 115–139, 1989.

Nakajima, S., Hayashi, Y. -Y. and Abe, Y.: A study on the “runaway greenhouse effect” with a one-dimensional radiative-convective equilibrium model, J. Atmos. Sci., 49, 2256–2266, 1992.

Pujol, T. and North, G. R.: Runaway greenhouse effect in a semigray radiative-convective model, J. Atm. Sci., 59, 2801–2810, 2002.

Ramanathan, V. and Coakley J. A.: Climate modeling through radiative-convective models, Rev. Geophys. Space Phys., 16, 465–489, 1978.

Raval, A. and Ramanathan, V.: Observational determination of the greenhouse effect, Nature, 342, 758–761, 1989.

Rennó, N. O.: Multiple equilibria in radiative-convective atmospheres, Tellus 49A, 423–438, 1997.

Weaver, C. P. and Ramanathan, V.: Deductions from a simple climate model: Factors governing surface temperature and atmospheric thermal structure, J. Geophys. Res., 100D, 11585–11591, 1995.

Height of convection

A. M. Makarieva et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Height of convection

A. M. Makarieva et al.

Fig. 1. Lapse rate of brightness temperature G_b versus height z for different values of atmospheric optical thickness τ_s of non-condensable greenhouse substances, see Eqs. (2.5) and (2.6). The corresponding values of τ_s are shown below each curve; $h_s \equiv RT_s/Mg$. The upper dashed line denotes maximum lapse rate $G_{\max} = Mg/4R$, the lower dashed line denotes the observed modern lapse rate of air temperature within the convective layer on Earth, $\Gamma = 0.77 G_{\max} = 6.5 \text{ K km}^{-1}$. The height z_c of the convective layer for different values of τ_s is determined by the points of intersection (empty circles) of the curves $G_b(z)$ with the straight line $\Gamma = 0.77 G_{\max}$, see Eq. (2.8). The value of $z_{c\max} = 3.03 h_s$ (shown by the vertical dash-dotted line), see Eq. (2.11), corresponds to the maximum height of the convective layer observed at $\tau \gg 1$. As far as Eq. (2.6) is valid at $z \leq z_c$, parts of curves $G_b(z)$ at $z > z_c$ are denoted by dots. The onset of convection in the atmosphere occurs at $\tau_s = \tau_{s\min} = 4.5$, for which $z_c = 0$. This figure is invariant with respect to changes of surface temperature T_s .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003

Height of convection

A. M. Makarieva et al.

Fig. 2. Lapse rate of brightness temperature G_{bL} versus height z for different values of atmospheric optical thickness τ_{sL} of water vapor, see Eq. (3.7). The corresponding values of τ_{sL} are shown below each curve; scale height $h_{sL} = 2.4$ km. The upper dashed line denotes maximum lapse rate $G_{L\max} = G_{\max}\beta = 30 \text{ K km}^{-1}$, $\beta = 3.5$; the lower dashed line denotes the observed modern lapse rate of air temperature within the convective layer on Earth, $\Gamma = \alpha_L G_{L\max} = 6.5 \text{ K km}^{-1}$, $\alpha_L = 0.22$. Height z_{cL} of the convective layer for different values of τ_{sL} is determined by the points of intersection of the curves $G_{bL}(z)$ with the straight line $\Gamma = \alpha_L G_{L\max}$ (denoted as empty circles). The value of $z_{cL\max} = 6.3h_{sL}$ (shown by the vertical dash-dotted line), see Eq. (3.7), corresponds to the maximum height of the convective layer observed at $\tau_{sL} \gg 1$ and $T_s = 288$ K. As far as Eq. (2.6) is valid within the convective layer, parts of curves $G_{bL}(z)$ at $z > z_{cL}$ are denoted by dots. The onset of convection in the atmosphere occurs at $\tau_{sL} = \tau_{sL\min} = 0.38$, for which $z_{cL} = 0$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2003