



# Lightning-produced NO<sub>2</sub> observed by two ground-based UV-visible spectrometers at Vanscoy, Saskatchewan in August 2004

A. Fraser, Florence Goutail, C. A. McLinden, S. M. L. Melo, K. Strong

## ► To cite this version:

A. Fraser, Florence Goutail, C. A. McLinden, S. M. L. Melo, K. Strong. Lightning-produced NO<sub>2</sub> observed by two ground-based UV-visible spectrometers at Vanscoy, Saskatchewan in August 2004. *Atmospheric Chemistry and Physics Discussions*, 2006, 6 (5), pp.10063-10086. hal-00328014

**HAL Id: hal-00328014**

**<https://hal.science/hal-00328014>**

Submitted on 18 Jun 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.

# Lightning-produced NO<sub>2</sub> observed by two ground-based UV-visible spectrometers at Vanscoy, Saskatchewan in August 2004

**A. Fraser<sup>1</sup>, F. Goutail<sup>2</sup>, C. A. McLinden<sup>3</sup>, S. M. L. Melo<sup>4</sup>, and K. Strong<sup>1</sup>**

<sup>1</sup>Department of Physics, University of Toronto, Toronto, Ontario, Canada

<sup>2</sup>Service d'Aéronomie du Centre National de la Recherche Scientifique, Verrières le Buisson, France

<sup>3</sup>Environment Canada, Downsview, Ontario, Canada

<sup>4</sup>Space Science, Canadian Space Agency, Saint-Hubert, Québec, Canada

Received: 19 September 2006 – Accepted: 4 October 2006 – Published: 11 October 2006

Correspondence to: A. Fraser (amery@atmosp.physics.utoronto.ca)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Ground-based measurements of ozone and NO<sub>2</sub> slant columns by the SAOZ (Système d'Analyse par Observations Zénithales) and UT-GBS (University of Toronto Ground-Based Spectrometer) instruments during the MANTRA 2004 field campaign are presented herein. During the afternoon of 28 August, a thunderstorm passed over the instruments, which were installed at Vanscoy, Saskatchewan (52° N, 107° W). Enhanced slant columns of ozone and NO<sub>2</sub> were observed by both instruments during the storm, with maximum values of two and 25 times the expected columns, respectively. The enhanced ozone slant columns are primarily due to the longer path traversed by the solar radiation caused by multiple scattering inside the thick cloud layer associated with the thunderstorm. The enhanced NO<sub>2</sub> columns are attributed to NO<sub>x</sub> production by lightning. Two different methods are used to separate the NO<sub>2</sub> enhancements into contributions from the longer path length and production by lightning. Combining the observed excess NO<sub>2</sub> with lightning flash data from the Canadian Lightning Detection Network and Environment Canada Doppler radar measurements, the production of NO<sub>2</sub> molecules per lightning flash is determined. Using these two methods, the best estimate of the production rate is found to be  $(7.45 \pm 1.38) \times 10^{26}$  molecules NO<sub>2</sub>/flash from the UT-GBS and  $(6.58 \pm 1.21) \times 10^{26}$  molecules NO<sub>2</sub>/flash from SAOZ. These results are consistent with previous estimates which range from  $10^{26}$  molecules NO<sub>2</sub>/flash (Noxon, 1976) to  $(7 \pm 3) \times 10^{26}$  molecules NO<sub>2</sub>/flash (Franzblau and Popp, 1989).

## 1 Introduction

Active nitrogen oxides (NO<sub>x</sub>=NO+NO<sub>2</sub>) play an important role in ozone photochemistry, reacting catalytically with ozone in both the troposphere and the stratosphere (Crutzen, 1970). In the upper stratosphere NO<sub>x</sub> is one of the dominant sinks of ozone, while in the lower stratosphere, it mitigates ozone loss by converting other ozone-

ACPD

6, 10063–10086, 2006

### Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

depleting chemicals ( $\text{HO}_x$ ,  $\text{BrO}_x$ , and  $\text{ClO}_x$ ) into their reservoir species (McElroy et al., 1992). In the troposphere, in areas of high  $\text{NO}_x$  concentration, ozone is produced through reactions with volatile organic compounds (VOCs), while in areas of low  $\text{NO}_x$  concentration, ozone is catalytically destroyed by  $\text{NO}_x$ . The source of  $\text{NO}_x$  in the stratosphere is dominated by the transport of  $\text{N}_2\text{O}$  from the troposphere. Sources of  $\text{NO}_x$  in the troposphere are more numerous, and include ground-based fossil fuel burning ( $\sim 24 \text{ Tg N/year}$ ), biomass burning ( $\sim 8 \text{ Tg N/year}$ ), soil emissions ( $\sim 12 \text{ Tg N/year}$ ),  $\text{NH}_3$  oxidation ( $\sim 3 \text{ Tg N/year}$ ), aircraft emissions ( $\sim 0.4 \text{ Tg N/year}$ ), and transport from the stratosphere ( $\leq 0.4 \text{ Tg N/year}$ ) (Price et al., 1997, and references therein). Another source of  $\text{NO}_x$  in the middle and upper troposphere, with by far the largest uncertainty, is lightning. The large amounts of energy released by lightning flashes can break apart  $\text{N}_2$  and  $\text{O}_2$  molecules, which then recombine to form  $\text{NO}$ . This newly formed  $\text{NO}$  can then react with ozone to form  $\text{NO}_2$  (Zel'dovitch and Raizer, 1966). Recent estimates of the global annual production rate due to lightning lie between 2 and 20  $\text{Tg N/year}$  (Huntrieser et al., 1998; Price et al., 1997; Bond et al., 2001).

The MANTRA (Middle Atmosphere Nitrogen TRend Assessment) series of balloon campaigns was conducted in Vanscoy, Saskatchewan ( $52^\circ \text{N}$ ,  $107^\circ \text{W}$ ) in late summer during the reversal of the stratospheric flow, known as turnaround, to measure stratospheric constituents that impact ozone chemistry (Strong et al., 2005). The turnaround condition provides a scenario when the atmosphere is the closest to photochemical equilibrium and dynamics do not have a large influence (Wunch et al., 2005, and references therein). The MANTRA field campaigns were held biennially from 1998 to 2004. As a complement to the balloon measurements, a suite of ground-based instruments were operated throughout each campaign, measuring the day-to-day variability of some of the trace gases of interest. Ozonesondes were also launched regularly throughout each campaign. In 2004, ground-based measurements were accumulated on 43 days, from 3 August to 15 September.

The late afternoon of 28 August saw a heavy thunderstorm over Vanscoy, with lightning and thunder observed for three hours. Ozone and  $\text{NO}_2$  slant column densities

## Lightning-produced $\text{NO}_2$ observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

measured during the storm by two of the ground-based instruments are discussed herein. The observed NO<sub>2</sub> is separated into a contribution from path enhancement due to multiple scattering and production by lightning. The production of NO<sub>2</sub> per lightning flash is also calculated.

## 2 Instruments

The University of Toronto's Ground-Based Spectrometer (UT-GBS) was assembled in 1998 and has participated in all four MANTRA campaigns (Bassford et al., 2001, 2005; Farahani, 2006). It consists of a triple-grating spectrometer with a thermo-electrically cooled, two-dimensional CCD (charged-coupled device) array detector. Sunlight from the zenith-sky is gathered by a fused silica lens with a two-degree field of view, and focused on a liquid light guide, which minimizes the effects of polarization. Spectra are recorded continuously throughout the day, with varying exposure times to maximize the signal on the CCD. Shortly before the 2004 campaign, the CCD detector began malfunctioning. A replacement was obtained from the manufacturer, a liquid-nitrogen-cooled 1024×128 pixel front-illuminated CCD. Spectra were recorded between 345 and 555 nm, with a resolution of approximately 0.5 nm in the NO<sub>2</sub> region and 1.0 nm in the ozone region. The loaned CCD's sensitivity to UV-Vis radiation was lower than that of the original detector, meaning that longer exposure times were required to maximize the signal, leading to fewer measurements over twilight. A low signal-to-noise ratio was obtained during the MANTRA 2004 campaign due to an error in the data acquisition software, which has since been corrected. Despite these issues, good data was obtained, as seen in the spectral fits shown in Fig. 1.

The SAOZ (Système d'Analyse par Observations Zénithales) instrument was constructed in the late 1980s, and is now deployed in a global network for measurements of stratospheric concentrations of trace gases important to ozone loss (Pommereau and Goutail, 1998). SAOZ records spectra between 270 and 620 nm, with a resolution of 1.0 nm. The detector is an uncooled 1024-pixel linear diode array. SAOZ records

### Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

zenith-sky spectra with a 10° field-of-view. Spectra are recorded every thirty minutes throughout the day, and continuously during twilight, defined as when the solar zenith angle (SZA) is between 80° and 95°.

The DOAS technique (e.g., Solomon et al., 1987; Platt, 1994) is used for the analysis of spectra from both instruments with absorption cross-sections of ozone (Burrows et al., 1999), NO<sub>2</sub> (Vandaele et al., 1998), H<sub>2</sub>O (Rothman et al., 2003), and O<sub>4</sub> (Greenblatt et al., 1990) fitted using a Marquardt-Levenberg non-linear least-squares technique. Slant column densities (SCDs) of ozone are retrieved between 450 and 550 nm, and NO<sub>2</sub> SCDs are retrieved between 425 and 450 nm. The program WinDOAS, developed by the IASB (Belgian Institute for Space Aeronomy, Fayt and Van Roozendaal, 2001), has been used to analyze data from both instruments. A single reference spectrum was chosen for each instrument for the entire campaign (3 September, SZA=44.8°), and the two were selected to be as close as possible in time. Figure 1 shows typical ozone and NO<sub>2</sub> fits from the two instruments, taken during the thunderstorm.

Measurement errors for both instruments are calculated from the root-sum-square of individual sources of error, after Bassford et al. (2005) and references therein. The individual errors are the same for both instruments unless otherwise noted and include random noise on the spectra (2.5% for UT-GBS ozone, 1% for SAOZ ozone, 2% for NO<sub>2</sub>), instrument error arising from uncertainty in the dark current, bias, and slit function (2% for the UT-GBS, 1% for SAOZ), pseudo-random errors resulting from unaccounted-for structure in the spectra (1–2% for ozone, 4–6% for NO<sub>2</sub>), errors in the absorption cross sections (2.6% for ozone, 5% for NO<sub>2</sub>), the temperature dependence of the NO<sub>2</sub> cross-section (≤8%), and the effects of multiple Raman scattering, which acts to fill in absorption features (1% for ozone, 5% for NO<sub>2</sub>). The total error for the UT-GBS is 4.6% for ozone and 11.9% for NO<sub>2</sub>, while the errors for SAOZ are 3.5% for ozone and 11.8% for NO<sub>2</sub>.

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

### 3 Thunderstorm observations

Environment Canada (EC) weather observations from nearby Saskatoon (30 km North of Vanscoy) show a thunderstorm occurring between 17:00 and 20:00 LT (SZA 63°–90°). Figure 2 shows the hourly total cloud opacity and cloud base height recorded by Environment Canada. During the thunderstorm, the sky was completely overcast by a thick cloud with a base varying from 750 to 1350 m. EC radar observations from Radisson, Saskatchewan (60 km North-West of Vanscoy) also show thunderstorm and thick cloud activity occurring over Vanscoy. Figure 3 shows the precipitation occurring at an altitude of 1.5 km at 17:00 LT (23:00 UTC, SZA 63°). A cell of heavy rain (pink colour) was observed over Vanscoy (indicated by a red arrow on Fig. 3).

Lightning flash data was obtained from the Canadian Lightning Detection Network (CLDN), a series of ground-based detectors that sense lightning using the time-of-arrival of radio pulses generated by lightning (Burrows et al., 2002). Figure 4 shows the lightning detected over the Canadian Prairies for the one-hour period from 17:00 LT to 18:00 LT on 28 August (23:00 UTC to 00:00 UTC, SZA 63° to 72°). In the 50-km radius surrounding Vanscoy (marked by an X on Fig. 4), an average flash rate of 2.87 flashes/min was observed by the CLDN for the duration of the storm. Figure 4 also shows the visible GOES (Geostationary Operational Environmental Satellite) image of the storm, which extends over Saskatchewan and into Alberta and the northern United States.

### 4 Slant column measurements

Figure 5 shows the ozone, NO<sub>2</sub>, and O<sub>4</sub> SCDs measured by both instruments on 28 August. That afternoon, both instruments observed a maximum in all three species centered at a solar zenith angle of 77° (18:31 LT), which is consistent with the occurrence of the thunderstorm. At the ozone maximum, slant column densities measured by both instruments have increased by roughly a factor of two. At the peak of NO<sub>2</sub>,

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the slant column densities have increased by a factor of 25 as observed by the UT-GBS and 20 as observed by the SAOZ instrument 15 min before the peak. Because SAOZ had not switched to continuous measurements at the peak of the storm, which occurred before twilight, it did not capture the true maximum in NO<sub>2</sub> columns. However, since the NO<sub>2</sub> columns between the two instruments are in agreement throughout the campaign, it can be assumed that had SAOZ been measuring, it would have observed a maximum NO<sub>2</sub> SCD similar to that seen by the UT-GBS.

The increase in ozone is consistent with similar observations of ozone during thick cloud events (Pfeilsticker et al., 1999) and thunderstorms (Winterrath et al., 1999), both in Southern France. The increase in NO<sub>2</sub> is roughly double that seen by Langford et al. (2004) during a Colorado thunderstorm, though less than the 35-fold increase seen by Winterrath et al. (1999). The observed enhancements in both ozone and NO<sub>2</sub> are partly caused by increased path length through the atmosphere. In the case of NO<sub>2</sub>, the increase could also be due to lightning-produced NO<sub>x</sub>.

There are two processes that combine to increase the path length, or air mass factor (AMF), in the presence of thick clouds, described in detail in Pfeilsticker et al. (1998). Optical paths can be enhanced by multiple reflections between layers of clouds (the “ping-pong” effect) or by photon diffusion inside a thick cloud. Other processes could account for the observed increase of ozone, including transport, in-cloud production, convection, intrusion of stratospheric air, or reaction with NO (Winterrath et al., 1999). One way to verify the influence of multiple scattering is to look at the ratios of ozone and NO<sub>2</sub> to O<sub>4</sub>. O<sub>4</sub> concentrations are related to concentrations of oxygen, and hence a maximum in O<sub>4</sub> such as the one observed in Fig. 5 is evidence of multiple scattering through the atmosphere, in this case due to the thick clouds associated with the thunderstorm. The ratios of ozone and NO<sub>2</sub> to O<sub>4</sub> for the afternoon of 28 August are shown in Fig. 6. As seen in Fig. 6a, the O<sub>3</sub> to O<sub>4</sub> ratio is fairly constant until the start of twilight, indicating that the excess ozone is a result of increased scattering through clouds. In the case of NO<sub>2</sub>, in Fig. 6b, there is a clear maximum in the ratio at 77°, the peak of the thunderstorm, indicating additional NO<sub>2</sub> in the atmosphere, which is attributed here to

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

production by lightning.

## 5 Derivation of lightning-produced NO<sub>2</sub>

Two methods have been used to separate the observed enhancement of NO<sub>2</sub> into a portion due to path-enhancement and a portion due to lightning production. These methods are discussed in this section.

### 5.1 Method one: NO<sub>2</sub> to O<sub>4</sub> ratio

The behaviour of the NO<sub>2</sub>/O<sub>4</sub> ratios is fairly consistent between clear-sky days, however it can vary significantly between cloudy days, due to varying cloud cover. It is therefore difficult to predict what the NO<sub>2</sub>/O<sub>4</sub> ratio would have been on 28 August if no NO<sub>2</sub> was formed by lightning. However, the ratio generally increases exponentially with solar zenith angle. An exponential fit was made to the observed NO<sub>2</sub>/O<sub>4</sub> ratio, omitting the points between 65° and 82°, which correspond to the beginning and end of the observed maximum in the ratio. These fits for both the SAOZ and UT-GBS instruments are shown in Fig. 6b, with the measured value of the ratio used before and after the peak (before 65° and after 82°). From these fitted values of the NO<sub>2</sub>/O<sub>4</sub> ratio during the storm, an estimate of the portion of the NO<sub>2</sub> slant column due to enhanced path length can be derived:

$$\text{NO}_{2,\text{PE}}(\text{SZA}) = \left[ \frac{\text{NO}_2}{\text{O}_4}(\text{SZA}) \right]_{\text{fit}} \times \text{O}_{4,\text{meas}}(\text{SZA}) \quad (1)$$

where NO<sub>2,PE</sub> is the path-enhanced NO<sub>2</sub>, the subscript “fit” indicates the fitted NO<sub>2</sub>/O<sub>4</sub> ratio, and O<sub>4,meas</sub> the measured O<sub>4</sub> column.

The resulting NO<sub>2</sub> slant columns are shown in Fig. 7a. The difference between the observed NO<sub>2</sub> slant column and the slant column calculated from the interpolated NO<sub>2</sub>/O<sub>4</sub> ratio is the amount of NO<sub>2</sub> attributed to production by lightning, shown in Fig. 7b. Values from both instruments are similar.

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 5.2 Method two: derived air mass factors

A second method of deriving the amount of NO<sub>2</sub> caused by path enhancement is to use the measurements of ozone to derive the enhanced air mass factor, or path length, for ozone and NO<sub>2</sub>. Generally, measured slant column densities of both ozone and NO<sub>2</sub> are related to the air mass factor by:

$$\text{SCD}(\text{SZA}) = \text{VCD} \times \text{AMF}(\text{SZA}) - \text{RCD} \quad (2)$$

where VCD is the vertical column density, RCD is the reference column density, the amount of absorber in the reference spectrum used in the DOAS analysis, and SZA is the solar zenith angle. With no path enhancement, the afternoon ozone SCDs would follow the same trend as the morning SCDs, increasing with SZA. Assuming no change in the VCD, the path-enhanced SCD' will be related to the enhanced AMF' by:

$$\text{SCD}'(\text{SZA}) = \text{VCD} \times \text{AMF}'(\text{SZA}) - \text{RCD} \quad (3)$$

where the primes indicate path-enhancement. The enhanced AMF' can be found by combining these two equations:

$$\text{AMF}'(\text{SZA}) = \text{AMF}(\text{SZA}) \frac{\text{SCD}'(\text{SZA}) + \text{RCD}}{\text{SCD}(\text{SZA}) + \text{RCD}} \quad (4)$$

The enhanced AMF' is an upper limit on the AMF, and assumes that all of the excess ozone is due to multiple scattering in and between clouds. This assumption is justified by the behaviour of the ozone to O<sub>4</sub> ratio shown in Fig. 6a. The non-path-enhanced AMF was found using a radiative transfer model (McLinden et al., 2002) initialized with temperature, pressure, and ozone profiles taken from the average of all ozonesondes launched during the campaign. The RCDs for ozone and NO<sub>2</sub> were found from Langley plots, of SCD versus AMF, for the morning of 28 August. Measured ozone slant columns from the morning and afternoon were used as SCD and SCD', respectively, in Eq. (4). Figure 8a shows the enhanced ozone AMF' for both instruments, derived

using Eq. (4), as well as the AMF calculated with the radiative transfer model assuming no clouds and assuming a thick cumulus cloud near the surface, of optical depth 70, extending between 1 and 5 km (Bassford et al., 2001).

The air mass factors for ozone and NO<sub>2</sub> differ since they are retrieved in different wavelength regions. In the radiative transfer model, the ozone AMF is calculated at 500 nm, while the NO<sub>2</sub> AMF is calculated at 425 nm. The AMFs for the two species are affected by the cloud differently. Figure 8b shows the ratio of the NO<sub>2</sub> AMF to the O<sub>3</sub> AMF for the cloudless and thick cloud calculations. This ratio is a maximum for the cloud-free scenario. Without more detailed observations of the clouds, it is difficult to predict what the ratio of the AMFs would have been during the storm, and even more difficult to quantify how it would change with SZA. Given this, a maximum NO<sub>2</sub> AMF can be found by assuming there are no clouds during the storm. Although this assumption is certainly false, it allows a calculation of an upper limit on the NO<sub>2</sub> due to path enhancement, and a lower limit on the NO<sub>2</sub> produced by lightning. Taking this cloud-free ratio, estimated NO<sub>2</sub> (AMF')s are found from:

$$AMF'_{NO_2}(SZA) = AMF'_{O_3}(SZA) \frac{AMF_{NO_2}(SZA)}{AMF_{O_3}(SZA)} \quad (5)$$

Figure 8c shows this estimated NO<sub>2</sub> AMF for each instrument, as well as the calculated AMFs for the cloudless and cloudy scenarios.

Figure 7a shows the resulting upper limit for NO<sub>2</sub> SCDs due to path enhancement calculated from Eq. (4). The NO<sub>2</sub> slant columns measured in the morning were used as SCDs, increased to account for the expected increase due to the diurnal variation of NO<sub>2</sub>. The upper limit of NO<sub>2</sub> from both SAOZ and UT-GBS is similar. Prior to 67°, the upper limit exceeds the measured NO<sub>2</sub> SCD, an indication that an assumption made in calculating the NO<sub>2</sub> AMFs is incorrect – either there is additional ozone produced or transported into the optical path, or the ratio of the air mass factors is smaller than assumed, as expected given Fig. 8b. At SZAs between 75° and 85°, this maximum NO<sub>2</sub> is exceeded by the NO<sub>2</sub> found using the NO<sub>2</sub> to O<sub>4</sub> ratio method. This is perhaps an

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

indication that the interpolated NO<sub>2</sub> to O<sub>4</sub> ratio is too large in this region, a result of the difficulty in predicting the NO<sub>2</sub>/O<sub>4</sub> ratio. However, since both methods result in similar amounts of NO<sub>2</sub> due to path-enhancement they can be believed to be approximately correct.

## 6 NO<sub>2</sub> flash production rate

The difference between the measured NO<sub>2</sub> SCD and the derived path-enhanced NO<sub>2</sub> in Fig. 7a is the minimum amount of NO<sub>2</sub> produced by lightning, which is shown in Fig. 7b. Using the ratio method, the amount of excess NO<sub>2</sub>, found by integrating under the curves between 60° and 85° for method one in Fig. 7b is  $(1.28 \pm 0.20) \times 10^{18}$  molecules NO<sub>2</sub>/cm<sup>2</sup> for the UT-GBS and  $(1.13 \pm 0.18) \times 10^{18}$  molecules NO<sub>2</sub>/cm<sup>2</sup> for SAOZ. The error estimates on the total excess NO<sub>2</sub> are the root-sum-square of the SCD error discussed in Sect. 2 and the estimated integration error (10%). From Fig. 3, the size of the heavy-precipitation cell is  $(30 \pm 3)$  km<sup>2</sup>. Using the average flash rate during the storm from the CLDN (2.87 flashes/min), the storm-averaged NO<sub>2</sub> production per flash of lightning can be calculated from:

$$\text{NO}_2 \text{ produced} = \frac{E \times A}{F \times D} \quad (6)$$

where  $E$  is the excess NO<sub>2</sub> measured by the instrument,  $A$  is the area of the storm,  $F$  is the lightning flash rate, and  $D$  is the duration of the storm (three hours) (Noxon, 1976). The production amounts from this method are  $(7.45 \pm 1.38) \times 10^{26}$  molecules NO<sub>2</sub>/flash from the UT-GBS and  $(6.58 \pm 1.21) \times 10^{26}$  molecules NO<sub>2</sub>/flash from SAOZ. The values derived from both instruments agree within error bars.

Using the air mass factor method, the UT-GBS observed  $(1.18 \pm 0.83) \times 10^{18}$  excess molecules NO<sub>2</sub>/cm<sup>2</sup> and SAOZ observed  $(1.06 \pm 0.16) \times 10^{17}$  excess molecules NO<sub>2</sub>/cm<sup>2</sup>. The minimum production amounts are found to be  $(6.83 \pm 1.26) \times 10^{26}$  molecules NO<sub>2</sub>/flash from the UT-GBS, and  $(6.18 \pm 1.14) \times 10^{26}$  molecules NO<sub>2</sub>/flash

from SAOZ. Again, the two instruments agree within their combined error bars. The values from the two methods are also in agreement for the two instruments. However, it should be noted that the values calculated are slightly different quantities. The values found using the air mass factor method of limiting the NO<sub>2</sub> due to lightning production are minimum values for the amount of NO<sub>2</sub> production by lightning, whereas the values found using the NO<sub>2</sub> to O<sub>4</sub> ratio are best estimates of the actual NO<sub>2</sub> production rate.

Noxon (1976) estimated an order of magnitude value of 10<sup>26</sup> molecules NO<sub>2</sub>/flash based on DOAS slant column measurements of NO<sub>2</sub> made during a thunderstorm in Colorado. Franzblau and Popp (1989) derived a rate of (7±3)×10<sup>26</sup> molecules NO<sub>2</sub>/flash using a similar method in New Mexico. Langford et al. (2004) measured (5.8±2.9)×10<sup>26</sup> molecules NO<sub>2</sub>/flash from a zenith-viewing UV-Visible spectrometer. All the values derived here are in agreement within the combined error bars of these previously published values.

The detection efficiency of the CLDN is approximately 85–90% for cloud-to-ground flashes and only 1–4% for cloud-to-cloud flashes, due to the lower amounts of energy released during the latter (Burrows et al., 2002). The majority (90%) of lightning-produced NO<sub>2</sub> is produced by the more energetic cloud-to-ground flashes (Price et al., 1997). No correction has been made to the flash frequency observed by the CLDN. Taking the detection efficiency into account would slightly increase the flash rate, and therefore decrease the calculated NO<sub>2</sub> production value.

## 7 Conclusions

Elevated columns of ozone and NO<sub>2</sub> were observed by both SAOZ and UT-GBS during a thunderstorm that occurred the afternoon of 28 August 2004 in Vanscoy, Saskatchewan during the 2004 MANTRA campaign. In the case of ozone, the enhancement is most likely due to increased path length in and between clouds. In the case of NO<sub>2</sub> the enhancement is due to increased path length and the production of NO<sub>x</sub> by lightning. The amount of excess NO<sub>2</sub> due to lightning has been calculated, us-

### Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ing two methods to estimate the excess NO<sub>2</sub> due to path enhancement within the cloud. Lightning flash data from the CLDN and radar data from a nearby Environment Canada radar station were used to determine the amount of NO<sub>2</sub> produced per lightning flash. The production values in the range of (6.18–7.45×10<sup>26</sup>) molecules NO<sub>2</sub>/flash calculated from both instruments agree with each other, and are also in agreement with previously published values.

*Acknowledgements.* MANTRA 2004 was supported by the Canadian Space Agency, the Meteorological Service of Canada, and the Natural Sciences and Engineering Research Council of Canada. The UT-GBS instrument was funded by the Natural Sciences and Engineering Research Council of Canada. The loaned CCD was provided by the manufacturer, J. Y. Horiba of Edison, NJ. The participation of the French SAOZ in the campaign was supported by the Centre National d'Études Spatiales. WinDOAS was provided by C. Fayt and M. Van Roozendael of the Belgian Institute for Space Aeronomy (IASB-BIRA). Cloud parameter data was provided by Y. Rochon of Environment Canada. Canadian Lightning Detection Network data was provided by W. Burrows of the Hydrometeorology and Arctic Lab, Meteorological Service of Canada, Edmonton. The archived radar data was provided by T. Ostry of the Meteorological Service of Canada and was analysed by P. Rodriguez of the Science and Technology Branch of Environment Canada.

## References

- Bassford, M. R., Strong, K., and McLinden, C. A.: Zenith-sky observations of stratospheric gases: The sensitivity of air mass factors to geophysical parameters and the influence of tropospheric clouds, *J. Quant. Spectrosc. Radiat. Transfer*, 68, 657–677, 2001. [10066](#), [10072](#)
- Bassford, M. R., Strong, K., McLinden, C. A., and McElroy, C. T.: Ground-based measurements of ozone and NO<sub>2</sub> during MANTRA 1998 using a zenith-sky spectrometer, *Atmos.-Ocean*, 43, 325–338, 2005. [10066](#), [10067](#)
- Bond, D. W., Zhang, R., Tie, X., Brasseur, G., Huffman, G. R., Orville, R. E., and Boppcippio, D. J.: NO<sub>x</sub> production by lightning over the continental United States, *J. Geophys. Res.*, 106, 27 701–27 710, 2001. [10065](#)

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Burrows, J. P., Richter, A., Dehn, A., Deters, B., Himmelmann, S., Voight, S., and Orphal, J.: Atmospheric remote-sensing reference data from GOME – 2. Temperature dependent absorption cross-sections of O<sub>3</sub> in the 231–794 nm range, J. Quant. Spectrosc. Radiat. Transfer, 61, 509–517, 1999. [10067](#)
- 5 Burrows, W. R., Lewis, P. J., Kochtubajda, B., Snyder, B., and Turcotte, V.: Lightning occurrence patterns over Canada and adjacent United States from lightning detection network observations, Atmos. Ocean, 40, 59–80, 2002. [10068](#), [10074](#)
- Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, Quart. J. Roy. Meteorol. Soc., 96, 320–325, 1970. [10064](#)
- 10 Farahani, E.: Stratospheric composition measurements in the Arctic and at mid-latitudes and comparison with chemical fields from atmospheric models, Ph.D. Thesis, University of Toronto, Toronto, 2006. [10066](#)
- Fayt, C. and Van Roozendael, M.: WinDOAS 2.1 – Software user manual, Uccle, Belgium, BIRA-IASB, 2001. [10067](#)
- 15 Franzblau, E. and Popp, C. J.: Nitrogen oxides produced from lightning, J. Geophys. Res., 94, 11 089–11 104, 1989. [10064](#), [10074](#)
- Greenblatt, G. F., Orlando, J. J., Burkholder, J. B., and Ravishankara, A. R.: Absorption measurements of oxygen between 330 and 1140 nm, J. Geophys. Res., 95, 18 577–18 582, 1990. [10067](#)
- 20 Huntrieser, H., Schlager, H., Feigl, C., and Höller, H.: Transport and production of NO<sub>x</sub> in electrified thunderstorms: Survey of previous studies and new observations at midlatitudes, J. Geophys. Res., 103, 28 247–28 264, 1998. [10065](#)
- Langford, A. O., Portmann, R. W., Daniel, J. S., Miller, H. L., and Solomon, S.: Spectroscopic measurements of NO<sub>2</sub> in a Colorado thunderstorm: Determination of the mean production by cloud-to-ground lightning flashes, J. Geophys. Res., 109, D11304, doi:10.1029/2003JD004158, 2004. [10069](#), [10074](#)
- 25 McElroy, M. B., Salawitch, R. J., and Minschwaner, K.: The changing stratosphere, Planet. Space Sci., 40, 373–401, 1992. [10065](#)
- McLinden, C. A., McConnell, J. C., Griffioen, E., and McElroy, C. T.: A vector radiative-transfer model for the Odin/OSIRIS project, Can. J. Phys., 80, 375–393, 2002. [10071](#)
- 30 Noxon, J. F.: Atmospheric nitrogen fixation by lightning, Geophys. Res. Lett., 3, 463–465, 1976. [10064](#), [10073](#), [10074](#)
- Pfeilsticker, K., Erle, F., Funk, O., Marquard, L., Wagner, T., and Platt, U.: Optical path modifi-

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cations for zenith-sky measurements of stratospheric gases, J. Geophys. Res., 103, 25 323–25 335, 1998. [10069](#)

Peilstickler, K., Arlander, D. W., Burrows, J. P., Erle, F., Gil, M., Goutail, F., Hermans, C., Lambert, J.-C., Platt, U., Pommereau, J.-P., Richter, A., Sarkissian, A., Van Roozendaal, M., Wagner, T., and Winterrath, T.: Intercomparison of the influence of tropospheric clouds on UV-visible absorptions detected during the NDSC intercomparison campaign at OHP in June 1996, Geophys. Res. Lett., 26, 1169–1172, 1999. [10069](#)

Platt, U.: Differential optical absorption spectroscopy (DOAS), in: Air monitoring by spectroscopic techniques, edited by: Sigrist, M. W., pp. 27–84, John Wiley, Hoboken, NJ, 1994. [10067](#)

Pommereau, J. P. and Goutail, F.: O<sub>3</sub> and NO<sub>2</sub> ground-based measurements by visible spectrometry during Arctic winter and spring 1988, Geophys. Res. Lett., 15, 891–894, 1988. [10066](#)

Price, C., Penner, J., and Prather, M.: NO<sub>x</sub> from lightning 1. Global distribution based on lightning physics, J. Geophys. Res., 102, 5929–5941, 1997. [10065](#), [10074](#)

Rothman, L. S., Barbe, A., Benner, D. C., et al.: The HITRAN molecular spectroscopic database: edition of 2000 including updates through 2001, J. Quant. Spectrosc. Radiat. Transfer, 82, 5–44, 2003. [10067](#)

Solomon, S., Schmeltekopf, A. L., and Sanders, R. W.: On the interpretation of zenith sky absorption measurements, J. Geophys. Res., 92, 8311–8319, 1987. [10067](#)

Strong, K., Bailak, G., Barton, D., et al.: MANTRA – A balloon mission to study the odd-nitrogen budget of the stratosphere, Atmos.-Ocean, 43, 283–299, 2005. [10065](#)

Vandaele, A. C., Hermans, C., Simon, P. C., Carleer, M., Colin, R., Fally, S., Mérienne, M. -F., Jenouvrier, A., and Coquart, B.: Measurements of the NO<sub>2</sub> absorption cross-section from 42 000 cm<sup>-1</sup> to 10 000 cm<sup>-1</sup> (238–1000 nm) at 220 K and 294 K, J. Quant. Spectrosc. Radiat. Transfer, 59, 171–184, 1998. [10067](#)

Winterrath, T., Kurosu, T. P., Richter, A., and Burrows, J. P.: Enhanced O<sub>3</sub> and NO<sub>2</sub> in thunderstorm clouds: convection or production?, Geophys. Res. Lett., 26, 1291–1294, 1999. [10069](#)

Wunch, D., Tingley, M. P., Shepherd, T. G., Drummond, J. R., Moore, G. W. K., and Strong, K.: Climatology and predictability of the late summer stratospheric zonal wind turnaround over Vanscoy, Saskatchewan, Atmos.-Ocean, 43, 301–313, 2005. [10065](#)

Zel'dovitch, Y. B. and Raizer, Y. P.: Physics of shock waves and high-temperature hydrodynamic

ACPD

6, 10063–10086, 2006

## Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.




**Fig. 1.** Typical differential optical depth (DOD) fits for **(a)** ozone, UT-GBS, **(b)** ozone, SAOZ, **(c)** NO<sub>2</sub>, UT-GBS, and **(d)** NO<sub>2</sub>, SAOZ. All fits are for spectra taken during the thunderstorm, at a SZA of approximately 76° (18:25 LT). In all fits, the grey line is the data while the black line is the fit to the data.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.




**Fig. 2.** Cloud base heights and total cloud opacity recorded by Environment Canada at Saskatoon on 28 August 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.



**Fig. 3.** Precipitation rate at 17:00 LT (23:00 UTC, SZA 63°) observed by the Environment Canada radar in Radisson, Saskatchewan on 28 August. Vanscoy is indicated by the red arrow.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 4.** Lightning flash data for the one hour period beginning 28 August at 17:00 LT (23:00 UTC, SZA 63°) in Vanscoy (X) from the Canadian Lightning Detection Network (CLDN). Each plus (+) and minus (-) represents an individual lightning flash.

# Lightning-produced NO<sub>2</sub> observed at Vanscoy, SK

A. Fraser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.




**Fig. 5.** Ozone, NO<sub>2</sub>, and O<sub>4</sub> slant column densities measured by UT-GBS and SAOZ on 28 August 2004.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.




**Fig. 6.** (a) Measured  $O_3/O_4$  ratios, and (b) measured and fitted  $NO_2/O_4$  ratios for 28 August. The measured value of the ratio is plotted before and after the peak (before  $65^\circ$  and after  $82^\circ$ ) in the  $NO_2/O_4$  ratio (see text).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.




**Fig. 7.** (a) Measured total NO<sub>2</sub> SCDs as well as the derived contribution from path-enhancement (P-E) for methods one (ratios to O<sub>4</sub>) and two (derived AMFs) discussed in the text. (b) Residual NO<sub>2</sub> SCDs attributed to lightning.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Lightning-produced  
NO<sub>2</sub> observed at  
Vanscoy, SK**

A. Fraser et al.



**Fig. 8.** (a) Ozone air mass factors calculated for the case of no clouds and for a cloud of OD=70 from 1–5 km using a radiative transfer model, and derived from the measurements using Eq. (4). (b) Ratio of NO<sub>2</sub> to ozone AMFs using the radiative transfer model. (c) Same as (a), but for NO<sub>2</sub> and using Eq. (5).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)