Comparative bioaccumulation of trace elements between *Nautilus pompilius* and *N. macromphalus* (Cephalopoda: Nautiloidea) from Vanuatu and New-Caledonia

Mathieu Pernice, Julien Boucher, Renata Boucher-Rodoni, Pascale Joannot,
Paco Bustamante

► To cite this version:

Mathieu Pernice, Julien Boucher, Renata Boucher-Rodoni, Pascale Joannot, Paco Bustamante. Comparative bioaccumulation of trace elements between *Nautilus pompilius* and *N. macromphalus* (Cephalopoda: Nautiloidea) from Vanuatu and New-Caledonia. *Ecotoxicology and Environmental Safety*, 2009, 72 (2), pp.365- 371. 10.1016/j.ecoenv.2008.04.019 . hal-00336127

HAL Id: hal-00336127

<https://hal.science/hal-00336127>

Submitted on 1 Nov 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparative bioaccumulation of trace elements between *Nautilus pompilius* and *N. macromphalus* (Cephalopoda: Nautiloidea) from Vanuatu and New-Caledonia

Running title: influence of the environment on bioaccumulation of trace elements in *Nautilus*

PERNICE Mathieu¹, BOUCHER Julien², BOUCHER-RODONI Renata¹, JOANNOT Pascale³, BUSTAMANTE Paco^{4#}

¹ UMR 5178 Biologie des Organismes Marins et Ecosystèmes, DMPA, Muséum National d'Histoire Naturelle, 55 rue Buffon, 75005 Paris, France.

² Ecole Polytechnique Fédérale de Lausanne, LGCB-ISIC-SB-EPFL1015 Lausanne, Switzerland.

³ Direction des Collections, Muséum National d'Histoire Naturelle, Cour Fagon, 57 rue Cuvier, 75005 Paris, France.

⁴ Littoral Environnement et Sociétés (LIENSs), UMR 6250 CNRS-Université La Rochelle, 2 Rue Olympe de Gouges, 17042 La Rochelle Cedex 01, France

Correspondence to: Dr. Paco BUSTAMANTE
Littoral Environnement et Sociétés (LIENSs),
2 Rue Olympe de Gouges
F-17042 La Rochelle Cedex 01
France
Phone : +33 5 465 076 25
Fax : +33 5 464 582 64
E-mail : pbustama@univ-lr.fr

ABSTRACT: The concentrations of sixteen trace elements were investigated and compared for the first time in the digestive and excreting tissues of two *Nautilus* species (Cephalopoda: Nautiloidea) from two geologically contrasted areas: 1) *N. macromphalus* from New Caledonia, a region characterised by its richness in nickel ores and its lack of tectonic activities and 2) *N. pompilius* from the Vanuatu archipelago showing high volcanic and tectonic activities. In both *Nautilus* species, results clearly highlighted that the digestive gland played a key role in the bioaccumulation and storage of Ag, Cd, Ce, Co, Cu, Fe, La, Nd, V and Zn whereas As, Cr, Mn, Ni, Pb and Se were accumulated in a greater extent in the excreting tissues (i.e. pericardial and renal appendages). Despite contrasting environments, no significant difference ($P < 0.05$) was found between the two *Nautilus* species in the concentrations of most of the essential and non essential elements, including Ni and associated metals in Ni ores (i.e. Co and Mn). As nautilus lives on the outer shelf of barrier reefs, these results strongly support the hypothesis that the New Caledonian lagoon traps the major amount of the trace elements derived from natural erosion and the intense mining activities conducted on land. In contrast, the concentrations of the rare earth elements (Ce, La, and Nd) were significantly higher in *N. pompilius* than in *N. macromphalus*, probably as a result of the local enrichment of Vanuatu waters by specific environmental processes, such as volcanism or upwelling.

Keywords: metals, metalloids, cephalopods, detoxification, excretion, South Pacific

1. INTRODUCTION

Marine organisms are exposed to trace elements that are present in their diet and dissolved in seawater. This double exposure results in element accumulation in their different tissues whether or not these elements are essential to the metabolism (Rainbow, 2002). This process, widely described as bioaccumulation can vary greatly according to the specificities of the organisms such as age/size, sex and lifestyle, and according to the relative bioavailability of the metal in diet and seawater (Rainbow and Wang, 2001). Comparative analysis of trace element concentrations between closely related species living in different geographical areas can therefore allow at assessing the influence of the environment on their bioaccumulation.

Among the marine species, cephalopods belong to a molluscan group comprising ca. seven hundred species (Boyle and Rodhouse, 2005), in which high concentrations of trace elements such as Ag, Cd, Cu or Zn have generally been recorded (Martin and Flegal, 1975; Finger and Smith, 1987; Miramand and Bentley, 1992; Yamada et al., 1997; Bustamante et al., 1998; Dorneles et al., 2007). Indeed, several studies have demonstrated that the digestive and the excretory organs of cephalopods play a major role in the bioaccumulation of trace elements, as these organs are deeply involved in the assimilation processes, detoxification and storage of both essential and non-essential metals (Miramand and Guary, 1980; Miramand and Bentley, 1992; Bustamante et al., 2000, 2002a,b, 2004).

In the present work, we focused on the bioaccumulation of trace elements in the *Nautilus* genus. Nautiluses are scavenger cephalopods, inhabiting the outer shelf of barrier reefs in the Indo-Pacific area, generally at a depth of 300 to 500 meters (O'Dor et al., 1993; Norman, 2000). In spite of its evolutionary interest (*Nautilus* is the last representative of ectocochleate cephalopods, i.e. considered as a “living fossil”; Boyle and Rodhouse, 2005) and its physiological specificities (slow digestion and most of the excretory processes taking place in

the pericardial appendages; Schipp et al., 1985; Westermann et al., 2002), only very few studies deal with the bioaccumulation of trace elements in this ancestral genus. High concentrations of several metals and metalloids were reported in the digestive and excretory tissues of *Nautilus macromphalus* (Bustamante et al., 2000), a species endemic to the New Caledonian waters. These authors suggested that the high metal concentrations may be due to a high exposure in the metal-rich conditions in the surrounding New Caledonian waters. Indeed, the largest resources of Ni as laterites in the world are present in New Caledonia where open-cast mining is conducted since the 20th century, leading to a dramatic increase of metal concentrations in coastal waters and, subsequently, the metal burdens in marine biota (Labrosse et al., 2000; Metian et al., 2008). However, it is not clear whether metal accumulation observed in *N. macromphalus* reflected the impact of metal contamination through human activities on the outer shelf waters of New Caledonia or either if it is more due to a natural progressive bioaccumulation during nautilus long life span (*viz.* 10-15 years).

The aim of the present study was therefore to determine the bioaccumulation of a large range of elements in the genus *Nautilus* comparing two species from geologically contrasted areas: 1) *N. macromphalus* from New Caledonia which is characterised by its richness in nickel ores and its lack of tectonic activities and 2) *N. pompilius* from the Vanuatu archipelago which shows high volcanic and tectonic activities (Harrison et al, 1996). The analysed elements comprised 11 metals (Ag, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, V and Zn), 2 metalloids (As and Se) and 3 rare earth elements (Ce, La and Nd). These elements were selected for comparison with other species and other published works concerning Nautilus (e.g. Miramand et al. 1992, Bustamante et al. 2000, Ichihashi et al. 2001). The concentrations and relative contents of these different trace elements were determined in the digestive and the excretory tissues (i.e. pericardial and renal appendages) and compared between both nautilus species. The

specificities and the potential influence of the environment on bioaccumulation of these trace elements in the *Nautilus* genus are discussed.

2. MATERIALS AND METHODS

a) Biological material

Three *Nautilus pompilius* and five *Nautilus macromphalus* specimens were collected respectively off Santo island coast (Vanuatu, 15° 40' 522 S, 167° 02' 838 E, depth: 334m) and on the outer shelf of New-Caledonia (22° 20' 500 S, 166° 13' 400 E, depth: 300m) in May 2006 (Figure 1). Each individual was measured and weighted prior to dissection. The main features of the eight specimens sampled are reported in Table 1.

All specimens were dissected aseptically and for each specimen one pericardial appendage, one renal appendage and a fraction of the digestive gland were immediately frozen in liquid nitrogen and stored at -80°C for metal analyses.

b) Analytical procedure

Tissue samples were freeze-dried. Aliquots of the samples ranging from 20 to 300 mg were digested using a 3:1 v:v nitric-hydrochloric acid mixture with 65% ultrapure HNO₃ and ultrapure 70% HCl. The acidic digestion was performed overnight under ambient temperature and then heated in a microwave during 30 minutes, with increasing temperature until 105°C, and 15 minutes at 105°C (1200 Watts). After the mineralization process, each sample was diluted to 30 ml or 50 ml with milli-Q quality water, according to the volume of acid added to the mineralization (3 ml and 4.5 ml. respectively).

Ag, As, Cd, Ce, Co, Cr, Cu, Fe, La, Mn, Nd, Ni, Pb, Se, V and Zn were analysed either by ICP-OES (Varian® Vista-Pro) and ICP-MS (Varian® Ultra Mass 700). Reference tissues, dogfish liver DOLT-3 (NRCC), lobster hepatopancreas TORT-2 (NRCC) and Bush Branches

and Leaves (GBW 07602) were treated and analysed in the same way as the samples. Results were in good agreement with the certified values, and the standard deviations were low, proving good repeatability of the method. The results for standard reference materials displayed recoveries of the elements ranging from 81% to 119% (n=10). For each set of analyses, blanks were included in each analytical batch. The detection limits ($\mu\text{g.g}^{-1}$ dry wt) for ICP-OES were 8.3 (As, Fe, Zn), 3.3 (Ag, Se), 1.67 (Pb, V), 0.83 (Cd, Co, Cr, Cu, Mn, Ni), and for ICP-MS, they were 0.150 (Ni, V), 0.065 (Cd, Co, Cr, Cu, Mn, Pb), 0.033 (Ag), 0.017 (Nd), 0.008 (Ce, La). Average element concentrations are presented with standard deviations. All trace element concentrations are given on a dry weight basis ($\mu\text{g.g}^{-1}$ dry wt).

Data analyses

Statistical analyses were performed using MINITAB 13.2 Software. Comparison of trace element concentrations in nautilus tissues between sites was tested by 1-way ANOVA. Hypothesis of normal distribution was tested using the Anderson-Darling test and equality of variance by the Bartlett test. Pearson coefficients were calculated between trace elements in each tissue. The significance level for statistical analyses was always set at $\alpha = 0.05$.

3. RESULTS

Trace element concentrations in the tissues

Metal concentrations in the digestive gland, the renal appendages and the pericardial appendages of *N. macromphalus* and *N. pompilius* are reported in Table 2. La, Ce, and Nd concentrations fell below the detection limit in the renal appendages. Among the three tissues analysed, the digestive gland was the major site of accumulation for Ag, Cd, Ce, Co, Cu, Fe, La, Nd, V and Zn (Table 2). The digestive gland also concentrated As and Ni at

concentrations close to the highest ones recorded in the other tissues. As, Cr, Mn, Ni, Pb, and Se concentrations were remarkable in excreting tissues. More specifically, the renal appendages exhibited the highest Mn and Se concentrations, whereas As, Ni, and Pb were concentrated at similar concentrations within both renal and pericardial appendages. Finally, the pericardial appendages were the major site of accumulation for Cr.

Comparison between species

Generally, the pattern of trace element concentrations was the same in both *N. macromphalus* and *N. pompilius* tissues. However, there were few exceptions and more particularly in the digestive gland: firstly, Ce, Fe, La, and Nd concentrations were significantly ($P < 0.05$) higher in *N. pompilius*; secondly, Cd and Cr concentrations were significantly ($P < 0.05$) higher in *N. macromphalus* (Table 3).

In the case of the pericardial appendages, As, La, and Nd concentrations were significantly higher ($P < 0.05$) in *N. pompilius* (Table 3). For the renal appendages, Cu was the only element for which a significant difference was detected, *N. pompilius* having significantly higher concentrations than *N. macromphalus* (Table 3).

Relative proportion of trace elements between tissues

Among the three tissues considered, the digestive gland contained the largest proportions of most of the trace elements with 73 to 81% of Ag, 90 to 91% of Cd, 61 to 79% of Ce, 80 to 86% of Co, 67 to 72% of Cu, 90 to 93% of Fe, 72 to 81% of La, 55 to 75% of Nd, 56 to 67 of V, 81% to 82% of Zn in both *Nautilus* species (Fig. 2). The only exception were As, Cr, Mn, Ni, Se, and Pb which were mainly contained in the excreting tissues, i.e. renal and pericardial appendages (Fig. 2). The renal appendages were the major site of accumulation of Mn, Ni, and Se (i.e. 74 to 89% for Mn, 49 to 54% for Ni; 61 to 73% for Se) while the pericardial

appendages accumulated the highest relative proportion of Cr (i.e. 63 to 74%). Although the concentrations of some of the trace elements were high in the excretory organs, these tissues contained in fact low amounts of metals because of their small masses.

Correlations between metals

Table 4 shows the correlation between trace elements within the three tissues. It is noteworthy that the rare earth elements were correlated between each other within the digestive gland ($r > 0.985$ and $P < 0.001$ in all cases) and the pericardial appendages ($r > 0.922$ and $P < 0.003$ in all cases). In the digestive gland, rare earth elements varied positively with Fe and Mn ($r > 0.918$ and $r > 0.807$ and $P < 0.004$ and $P < 0.028$, respectively) but negatively with As ($r > -0.726$ and $P = 0.047$). Cu correlated positively with Zn ($r = 0.834$ and $P = 0.020$ in all cases) but negatively with Se in this tissue ($r = -0.809$ and $P = 0.028$ in all cases).

In the renal appendages, correlations were weak, Mn varied positively with Cd ($r = 0.833$, $P = 0.010$) and Zn ($r = 0.735$, $P = 0.038$), Fe with Zn ($r = 0.833$, $P = 0.010$) and Pb with Cr ($r = 0.936$, $P = 0.001$) but negatively with V ($r = -0.753$, $P = 0.031$; Table 3).

In the pericardial appendages, Cd concentrations correlated positively with Cr ($r = 0.897$, $P = 0.006$), Ni ($r = 0.965$, $P < 0.001$), and V ($r = 0.931$, $P = 0.002$).

4. DISCUSSION

The present extensive analysis of sixteen trace element concentrations in *N. macromphalus* and *N. pompilius* digestive and excreting tissues clearly highlighted that these organs constitute important tissues for the bioaccumulation of rare earth elements, metals and metalloids. Such results support the hypothesis of efficient regulation processes for non

196 essential metals in Nautilidae (Bustamante et al., 2000) even if, among the eight studied
197 specimens, the variability of metal concentrations was relatively high (Table 2).

198 In cephalopods, the digestive system plays a major role in both the digestive and the
199 detoxification processes: on the one hand, the digestive gland synthesizes the digestive
200 enzymes and is a place of absorption and assimilation of nutrients (Boucaud-Camou and
201 Boucher-Rodoni, 1983; Boucher-Rodoni and Boucaud-Camou, 1987); on the other hand, this
202 organ is involved in the storage and detoxification of trace elements (Bustamante et al.,
203 2002ab, 2004). Our investigations on trace element concentrations also confirmed the key role
204 of the digestive gland of both *N. macromphalus* and *N. pompilius* in the metabolism of most
205 of the analysed trace elements (i.e. Ag, Cd, Ce, Co, Cu, Fe, La, Nd, V and Zn), as it is also the
206 case in more recent cephalopods, i.e. belonging to the Coleoidae (e.g. Ghiretti-Magaldi et al.,
207 1958; Martin and Flegel, 1975; Miramand and Guary, 1980; Smith et al., 1984; Finger and
208 Smith, 1987; Miramand and Bentley, 1992; Seixas et al., 2005a; Pierce et al., 2008).

209 Our results also stressed the potential of excreting tissues (i.e. renal and pericardial
210 appendages) to accumulate several elements such as As, Cr, Mn, Ni, Pb, and Se. In contrast to
211 the Coleoidae, in nautiluses the renal appendages are specialized in the storage of calcium and
212 the pericardial appendages play a major role in the evacuation of the metabolic wastes
213 (Schippe and Martin, 1987). These organs are known to harbour a dual bacterial symbiosis
214 (Pernice et al., 2007) involving spirochaetes and betaproteobacteria. The exact mechanisms
215 by which these bacterial symbionts are involved in nautilus excretion are still enigmatic.

216 Nevertheless, it is widely admitted that micro-organisms can play an important role in the
217 incorporation of trace elements (Worms et al., 2006). Accumulation of Co, Fe and Ni by
218 betaproteobacterial symbionts of the pericardial appendages of *N. macromphalus* has already
219 been observed by electronic microscopy (Pernice et al., in preparation). However, it remains
220 unclear if this accumulative process is only passive (viz. through adsorption processes) or

active (viz. effective absorption of trace elements). Indeed, the betaproteobacterial symbionts present in nautilus tissues could play a role in the detoxification and storage of some metals such as Co, Fe and Ni (Table 2). Such processes could involve enzymatic reduction and/or complexation with organic compounds (Pernice et al., 2006), leading to the storage of these metals at high concentrations in the excreting tissues, independently of the nautilus species. The role of this bacterial symbiosis on trace element bioaccumulation and detoxification in nautilus excreting tissues clearly deserves further investigations.

Although Nautilidae have been previously reported for their special abilities to accumulate heavy metals at relatively elevated concentrations in their tissues (Bustamante et al., 2000), this study is the first comparative approach aiming to compare trace element concentrations in two *Nautilus* species from two geographically close areas (viz. distant from less than 1000 kms). Both species potentially share common trophic features but, at the same time, they live in geologically different environments (richness in nickel ores and lack of tectonic activities in New Caledonia and high volcanic and tectonic activities in Vanuatu), potentially resulting in a different degree of “natural” contamination. New Caledonia is currently the third largest producer of Ni in the world (Dalvi et al., 2004) and mining activities associated to a strong natural erosion have resulted in the enrichment of Ni and associated metals in Ni ores (i.e. Co, Cr, Cu, Fe and Mn) and some other extracted metals (i.e. Ag and Pb) in the coastal waters and consequently in the lagoon food webs (Monniot et al., 1994; Fernandez et al., 2006; Hédouin et al., 2006; Metian et al., 2008). Conversely, it is likely that the seawater concentrations of most of these metals are very low in Santo islands waters and around Vanuatu in general according to the absence of industrial and mining activities in this area (Harrison et al., 1996). Surprisingly, in the present study, very little difference appeared in the sixteen trace element concentrations between *N. macromphalus* from New Caledonia and *N. pompilius* from Vanuatu. Interestingly, there was no significant difference between both *Nautilus* species for

Ag, Co, Mn, Ni, Pb, Se, V and Zn (Table 3) indicating that the geographical origin does not seem to be a major factor contributing to interspecific variations of these trace element concentrations. As nautilus lives on the outer shelf of barrier reefs and rarely penetrates into lagoons (Saunders and Ward, 1987; Norman, 2000), our results strongly supported the hypothesis that the New Caledonian lagoon traps the major amount of the trace elements derived from the island (Ambatsian et al., 1997). To the best of our knowledge, there are no published data in the open literature concerning trace elements in organisms from the oceanic waters off New Caledonia, except for two marine mammal species (Bustamante et al., 2003). These authors reported very low concentrations of Co, Cr and Ni in the marine mammal tissues which suggested that mining activity in New Caledonia does not seem to be a significant source of contamination in the oceanic domain. Overall, the global geologic contagion due to the mining activity in New Caledonia seems to be limited to the shoreline waters and to the inner lagoon (Fernandez et al., 2006). More interestingly, the concentrations of rare earth elements were significantly different between the digestive gland and the pericardial appendages of *N. macromphalus* and *N. pompilius*. F-transition elements of rare earths have similar physicochemical properties, and it is therefore not surprising that they correlate to each other (Table 4). Our results suggested that rare earth elements would reflect the characteristics of the surrounding environment. Very little is known about the concentrations of Ce, La and Nd in cephalopods, which varied from 0.045 to 1.95 $\mu\text{g.g}^{-1}$ dwt in the digestive gland of the squid *Stenoteuthis oualaniensis* (Ichihashi et al., 2001). In fact, rare earth elements have chemistry and sorption/complexation characteristics analogous to transuranic elements such as americium or plutonium (Choppin Torres and Chopin, 1984; Kim et al., 1989; Buckau et al., 1992). In Coleoid cephalopods, transuranic elements also accumulate in the digestive and excreting tissues, in particular in the branchial heart and their appendages (Guary et al., 1981; Guary and Fowler, 1982;

Bustamante et al., 2006). It is therefore not surprising that they accumulate at relatively high concentrations in nautilus digestive gland and pericardial appendage.

Besides Ce, La and Nd, the concentrations of As, Cu and Fe were also higher in *N. pompilius* than in *N. macromphalus*. It is difficult to ascertain the reasons of such a difference but this might also be linked to geological features reflecting a potential enrichment of Vanuatu waters by local environmental processes, such as volcanism or upwelling (Malinovsky and Markevich, 2007). In fact, only two trace elements, Cd and Cr, displayed significantly higher concentrations in *N. macromphalus* than in *N. pompilius* tissues (Table 2). Therefore, the reasons of such interspecific differences in bioaccumulation, i.e. higher concentrations of trace elements in one species, remain unclear whereas an absence of difference could be explained by the similar physiology of these two species (Ward, 1988; Norman, 2000).

As for other marine organisms, in cephalopods, trace element concentrations may vary with biological and environmental factors such as temperature, salinity, season, age (size), sex, lifestyle and geographical origin (e.g. Monteiro et al., 1992; Bustamante et al., 1998; Raimundo et al., 2004; Seixas et al., 2005b; Pierce et al., 2008). Further investigations are needed to understand how exactly these factors affect their concentrations in the studied species. However, among these factors, individual size/age is known to be of primary importance in invertebrates (e.g. Warnau et al., 1995; Bustamante and Miramand, 2005; Hédouin et al., 2007). In the present study, the size as well as the two others main features measured (sex and weight) were globally homogenous among the eight studied specimens. Because nautilus are long lived animals, it would be striking to determine the influence of the size/age on trace elements bioaccumulation.

Sources of trace elements to nautilus can be (1) seawater, as it passes through the skin and through the gills during respiration, and (2) food, which is now considered as a major route for metal accumulation in marine animals, including both invertebrates and fish (Koyama et

al., 2000; Wang, 2002; Bustamante et al., 2002a, 2004). Knowledge of trace element speciation in the direct environment of nautilus is a key factor to ascertain the relative importance of each uptake pathways. Concerning the first pathway (i.e. seawater), further aquatic toxicity studies are needed to determine the fraction of total metal dissolved in seawater (the 'toxic fraction') that react with, and is transported across biological membranes such as nautilus gills. Concerning the second pathway (i.e. food), our comparative analysis suggested that the metal discharges due to intense mining activities conducted on land in New Caledonia are not easily transferred in the food web of nautilus. Therefore, to assess the impact of mining activities on cephalopods in the region of New Caledonia, further studies should be conducted on more coastal species known to be resident in the impacted areas, such as Sepiidae, Sepiolidae or Octopodidae spending all of their life in the waters of the New Caledonian lagoon.

5. ACKNOWLEDGMENTS

We thank the Aquarium and the centre IRD of Nouméa (New Caledonia) as well as the Maritime college of Luganville (Vanuatu) for their help in providing *Nautilus macromphalus* and *N. pompilius* specimens, respectively. This work was supported by the Muséum National d'Histoire Naturelle (pluriformations programme: "Ecologie fonctionnelle et développement durable des écosystèmes naturels et anthropisés"), the University Pierre-et-Marie Curie, the CNRS (UMR 5178 and UMR 6250 LIENSs) and a research grant from the Government of New Caledonia awarded by the Pacific Fund.

6. REFERENCES

321 Ambatsian, P., Fernex, F., Bernat, M., Parron, C., Lecolle, J. 1997. High metal inputs to
 322 closed seas: the New Caledonian lagoon. J. Geochem. Explor. 59, 59-74.
 323
 324 Boucaud-Camou, E., Boucher-Rodoni, R. 1983. Feeding and digestion in Cephalopods. In:
 325 The Mollusca, Part 2, physiology (Wilbur, K M ed). Academic press, New York, 1983, pp.
 326 149-180.
 327
 328 Boucher-Rodoni, R., Boucaud-Camou, E. 1987. Fine structure and absorption of ferritin in the
 329 digestive organs of *Loligo vulgaris* and *L. forbesi* (Cephalopoda, Teuthoidea). J. Morph. 193,
 330 173-184.
 331
 332 Boyle, P., Rodhouse, P. 2005. Cephalopods: Ecology and Fisheries Blackwell publishing,
 333 Ames. 452 pp.
 334
 335 Buckeau, G., Kim, J.I., Klenze, R., Rhee, D.S., Wimmer, H. 1992. A comparative study of the
 336 fulvate complexation of trivalent transuranic ions. Radiochim. Acta 57, 105-112.
 337
 338 Bustamante, P., Cherel, Y., Caurant, F., Miramand, P. 1998. Cadmium, copper and zinc in
 339 octopuses from Kerguelen Islands, southern Indian Ocean. Polar Biol. 19, 264-271.
 340
 341 Bustamante, P., Grigioni, S., Boucher-Rodoni, R., Caurant, F., Miramand, P. 2000.
 342 Bioaccumulation of 12 trace elements in the tissues of the nautilus *Nautilus macromphalus*
 343 from New Caledonia. Mar. Pollut. Bull. 40, 688-696.
 344

345 Bustamante, P., Cosson, R.P., Gallien, I., Caurant, F., Miramand, P. 2002a. Cadmium
 346 detoxification processes in the digestive gland of cephalopods in relation to accumulated
 347 cadmium concentrations. *Mar. Environ. Res.* 53, 227-241.
 348
 349 Bustamante, P., Teyssié, J-L., Fowler, S.W., Cotret, O., Danis, B., Miramand, P., Warnau, M.
 350 2002b. Biokinetics of zinc and cadmium accumulation and depuration at different stages in
 351 the life cycle of the cuttlefish *Sepia officinalis*. *Mar. Ecol. Prog. Ser.* 231, 167–177.
 352
 353 Bustamante, P., Garrigue, C., Breau, L., Caurant, F., Dabin, W., Greaves, J., Dodemont, R.
 354 2003. Trace elements in two odontocetes species (*Kogia breviceps* and *Globicephala*
 355 *macrorhynchus*) stranded in New Caledonia (South Pacific). *Environ. Pollut.* 124, 263-271.
 356
 357 Bustamante, P., Teyssié, J-L., Danis, B., Fowler, S.W., Miramand, P., Cotret, O., Miramand,
 358 P., Warnau, M. 2004. Uptake, transfer and distribution of silver and cobalt in tissues of the
 359 common cuttlefish *Sepia officinalis* at different stages of its life cycle. *Mar. Ecol. Prog. Ser.*
 360 269, 185-195.
 361
 362 Bustamante, P., Miramand, P. 2005. Evaluation of the variegated scallop *Chlamys varia* as a
 363 biomonitor of temporal trends of Cd, Cu, and Zn in the field. *Environ. Pollut.* 138, 109-120.
 364
 365 Bustamante, P., Teyssié, J-L., Fowler, S.W., Warnau, M. 2006. Assessment of the exposure
 366 pathway in the uptake and distribution of americium and cesium in cuttlefish (*Sepia*
 367 *officinalis*) at different stages of its life cycle. *J. Exp. Mar. Biol. Ecol.* 331, 198-207.
 368
 369 Choppin Torres, R.A., Choppin, G.R. 1984. Europium (III) and americium (III) stability
 370 constants with humic acid. *Radiochim. Acta* 34, 143-148.

371

372 Dalvi, A.D., Bacon, W.G., Osborne, R.C. 2004. The past and the future of nickel laterites.

373 PDAC International Convention, Trade Show and Investors Exchange. March 7–10, 27 pp.

374

375 Dorneles, P.R., Lailson-Brito, J., Aguiar dos Santos, R., Silva da Costa, P.A., Malm, O.,

376 Freitas Azevedo, A., Machado Torres, J.P. 2007. Cephalopods and cetaceans as indicators of

377 offshore bioavailability of cadmium off Central South Brazil Bight. Environ. Pollut. 148, 352–

378 359.

379

380 Eisler, R. 1981. Trace metals concentration in marine organisms. Pergamon Press, New York,

381 685 pp.

382

383 Fernandez, J.M., Ouillon, S., Chevillon, C., Douillet, P., Fichez, R., Legendre, R. 2006. A

384 combined modelling and geochemical study of the fate of terrigenous inputs from mixed

385 natural and mining sources in a coral reef lagoon (New Caledonia) Mar. Pollut. Bull. 52, 320–

386 331.

387

388 Finger, J.M., Smith, J.D. 1987. Molecular association of Cu, Zn, Cd and ^{210}Po in the digestive

389 gland of the squid *Nototodarus gouldi*. Mar. Biol. 95, 87-91.

390

391 Ghiretti-Magaldi, A., Giuditta, A., Ghiretti, F. 1958. Pathways of terminal respiration in

392 marine invertebrates. I. The respiratory system in cephalopods. J. Cell. Comp. Physiol. 52,

393 389-429.

394

395 Guary, J-C., Higgo, J.J.W., Cherry, R.D., Heyraud, M. 1981. High concentrations of
 396 transuranic and natural radioactive elements in the branchial hearts of the cephalopods
 397 *Octopus vulgaris*. Mar. Ecol. Prog. Ser. 4, 123-126.
 398

399 Guary, J-C., Fowler, S.W. 1982. Experimental studies on the biokinetics of plutonium and
 400 americium in the cephalopod *Octopus vulgaris*. Mar. Ecol. Prog. Ser. 7, 327-335.
 401

402 Harrison, N., Gangaiya, P., Morrison, R.J. 1996. Organochlorines in the Coastal Marine
 403 Environment of Vanuatu and Tonga. Mar. Pollut. Bull. 32, 575-579.
 404

405 Hédouin, L., Metian, M., Teyssié, J-L., Fowler, S.W., Fichez, R., Warnau, M. 2006.
 406 Allometric relationships in the bioconcentration of heavy metals by the edible tropical clam
 407 *Gafrarium tumidum*. Sci. Total Environ. 366, 154–163.
 408

409 Hédouin, L., Pringault, O., Metian, M., Bustamante, P., Warnau, M. 2007. Nickel
 410 bioaccumulation in bivalves from the New Caledonia lagoon: Seawater and food exposure.
 411 Chemosphere 66, 1449-1457.
 412

413 Hirano, S., Suzuki, K.T. 1996. Exposure, metabolism, and toxicity of rare earth and related
 414 compounds. Environ. Health Perspect. 104, 85-95.
 415

416 Kim, J.I., Buckeau, G., Bryant, E., Klenze, R. 1989. Complexation of americium (III) with
 417 humic acid. Radiochim. Acta 48, 135-143.
 418

419 Koyama, J., Nanamori, N., Segawa, S. 2000. Bioaccumulation of waterborne and dietary
 420 cadmium by oval squid *Sepioteuthis lessoniana*, and its distribution among organs. Mar.
 421 Pollut. Bull. 40, 961-996.
 422

Ichihashi, H., Kohno, H., Kannan, K., Tsumura, A., Yamasaki, S. 2001. Multielemental analysis of purpleback flying squid using high resolution inductively coupled plasma-mass spectrometry (HR ICP-MS). Environ. Sci. Technol. 35, 3103-3108.

Labrosse, P., Letourneur, Y., Kulbicki, M., Paddon, J.R. 2000. Fish stock assessment of the Northern New Caledonian lagoons: fishing pressure, potential yields and impact on management options. Aquat. Living Res. 13, 91-98.

Norman, M.D. 2000. *Cephalopods: a world guide*. IKAN publishing, Frankfurt, 320 pp.

Malinovsky, A.I., Markevich, P.V. 2007. Heavy Clastic Minerals from the Far East Island-Arc Complexes. Russian J. Pacific Geol. 1, 71-81.

Martin, J.H., Flegal, A.R. 1975. High copper concentrations in squid livers in association with elevated levels of silver, cadmium, and zinc. Mar. Biol. 30, 51-55.

Metian, M., Bustamante, P., Hédouin, L., Warnau, M. 2008. Accumulation of trace elements in the tropical scallop *Comptopallium radula* from coral reefs in New Caledonia. Environ. Pollut. 152: 543-552.

Miramand, P., Guary, J-C. 1980. High concentrations of some heavy metals in tissues of the Mediterranean octopus. Bull. Environ. Contam. Toxicol. 24, 783-788.

446 Miramand, P., Bentley, D. 1992. Concentration and distribution of heavy metals in tissues of
 447 two cephalopods, *Eledone cirrhosa* and *Sepia officinalis*, from the French coast of the English
 448 Channel. Mar. Biol. 114, 407-414.

449

450 Monniot, F., Martoja, R., Monniot, C. 1994. Cellular sites of iron and nickel accumulation in
 451 ascidians related to the naturally and anthropic enriched New Caledonian environment.
 452 Annal. Inst. Océanogr. 70, 205-216.

453

454 Monteiro, L.R., Porteiro, F.M., Gonçalves, J.M. 1992. Inter- and intra-specific variation of
 455 mercury levels in muscle of cephalopods from the Azores. Arquipelago 10, 13-22.

456

457 O'Dor, R.K., Forsythe, J., Webber, D.M., Wells, J., Wells, M.J. 1993. Activity levels of
 458 Nautilus in the wild. Nature 362, 626-627.

459

460 Pernice, M., Wetzel, S., Gros, O., Boucher-Rodoni, R., Dubilier, N. 2007. Enigmatic dual
 461 symbiosis in the excretory organs of *Nautilus macromphalus* (Cephalopoda: Nautiloidea).
 462 Proc. Royal Soc. Lond. Ser. Biol. Sci. 274, 1143–1152.

463

464 Pernice, M., Boucher, J., Joannot, P., Boucher-Rodoni, R. 2006. Comparative study of
 465 bacteria diversity and bioaccumulation of trace elements in the excretory organs of Nautilus.
 466 In: CIAC symposium 2006, February 6-10, Hobart, Australia, 120 pp.

467

468 Pierce, G.J., Stowasser, G., Hastie, L.C., Bustamante, P. 2008. Geographic, seasonal and
 469 ontogenetic variation in cadmium and mercury concentrations in squid (Cephalopoda:

470 Teuthoidea) from UK waters. Ecotox. Environ. Safety in press
471 ([doi:10.1016/j.ecoenv.2007.07.007](https://doi.org/10.1016/j.ecoenv.2007.07.007)).

472

473 Rainbow, P.S., Wang, W-X. 2001. Comparative assimilation of Cr, Cr, Se, and Zn by the
474 barnacle *Elminius modestus* from phytoplankton and zooplankton diets. Mar. Ecol. Prog. Ser.
475 218, 239–248.

476

477 Rainbow, P.S. 2002. Trace metal concentrations in aquatic invertebrates: why and so what?
478 Environ. Pollut. 120, 497–507.

479

480 Raimundo, J., Caetano, M., Vale, C. 2004. Geographical variation and partition of metals in
481 tissues of *Octopus vulgaris* along the Portuguese coast. Sci. Total Environ. 325, 71-81.

482

483 Saunders, W., Ward, P.D. 1987. Ecology, distribution and population characteristics of
484 Nautilus. In: Nautilus, the biology and paleobiology of a living fossil Saunders, WB &
485 Landman, NH. (eds). New-York & London, Plenum press, 137-162 pp.

486

487 Schipp, R., Martin, A.W., Liebermann, H., Magnier, Y. 1985. Cytomorphology and function
488 of pericardial appendages of Nautilus (Cephalopoda, Tetrabranchiata). Zoomorph. 105, 16-29.

489

490 Schipp, R., Martin, A.W. 1987. The excretory system of Nautilus. In: *Nautilus, the biology*
491 *and Paleobiology of a living fossil* (WB Sauders and NH. Landman, eds). Plenum Press, New
492 York, 281-304 pp.

493

494 Seixas, S., Bustamante, P., Pierce, G.J. 2005a. Interannual patterns of variation in
 495 concentrations of trace elements in arms of *Octopus vulgaris*. *Chemosphere* 59, 1113-1124.
 496
 497 Seixas, S., Bustamante, P., Pierce, G.J. 2005b. Accumulation of mercury in the tissues of the
 498 common octopus *Octopus vulgaris* (L.) in two localities on the Portuguese coast. *Sci. Total*
 499 *Environ.* 340, 113-122.
 500
 501 Smith, J.D., Plues, L., Heyraud, M., Cherry, R.D. 1984. Concentrations of the elements Ag,
 502 Al, Ca, Cd, Cu, Fe, Mg., Pb and Zn, and the radionuclides ²¹⁰Pb and ²¹⁰Po in the digestive
 503 gland of the squid *Nototodarus gouldi*. *Mar. Environ. Res.* 13, 55-68.
 504
 505 Wang, W-X. 2002. Cd and Se aqueous uptake and exposure of green mussels *Perna viridis*:
 506 influences of seston quantity. *Mar. Ecol. Prog. Ser.* 226, 211–221.
 507
 508 Ward, P.D. 1988. In search of Nautilus “A New York Academy of Sciences book”, Simon &
 509 Schuster eds, New-York.
 510
 511 Warnau, M., Ledent, G., Temara, A. 1995. Allometry of heavy metal bioconcentration in the
 512 echinoid *Paracentrotus lividus*. *Arch. Environ. Contam. Toxicol.* 29, 393–399.
 513
 514 Westermann, B., Ruth, P., Litzbauer, H.D., Beck, I., Beuerlin, K., Schimdtberg, H., Kleta,
 515 E.F., Schipp, R. 2002. The digestive tract of *Nautilus pompilius* (Cephalopoda,
 516 Tetrabranchiata): an X-ray analytical and computational tomography study on the living
 517 animal. *J. Exp. Biol.* 205, 1617-1624.
 518

519 Worms, I., Simona, D.F., Hasslera, C.S., Wilkinson, K.J. 2006. Bioavailability of trace metals
520 to aquatic microorganisms: importance of chemical, biological and physical processes on
521 biouptake. *Biochimie* 88, 1721–1731.
522
523 Yamada, H., Takayanagi, K., Tateishi, M., Tagata, H., Ikeda, K. 1997. Organotin compounds
524 and polychlorinated biphenyls of livers in squid collected from coastal waters and open ocean.
525 *Environ. Pollut.* 96(2), 217-226.

526 Table 1. Characteristics of the sampled specimens of *Nautilus pompilius* and *N.*
527 *macromphalus*.

528

Species	Sex	Length (mm)	Weight (g)
<i>N. pompilius</i>			
1	Male	177	825
2	Male	151	660
3	Male	171	796
<i>N. macromphalus</i>			
1	Male	164	766
2	Female	162	676
3	Male	173	870
4	Male	160	758
5	Male	140	480

529 Table 2: Trace elements concentrations (Mean $\pm$ SE; $\mu\text{g g}^{-1}$ dry wt) in the tissues of Nautiloidae from this study and from the literature. The
530 highest values are in bold. dl: detection limit; NA: not analysed; # only one value.

531

Tissue	Rare earth elements			Non essential metals			Essential metals and metalloids									
	La	Ce	Nd	Ag	Cd	Pb	As	Co	Cr	Cu	Fe	Mn	Ni	Se	V	Zn
<i>Nautilus macromphalus</i> (Bustamante et al. 2000)																
Digestive gland	NA	NA	NA	4.45 $\pm$ 2.68	45.1$\pm$13.2	0.46 $\pm$ 0.17	166 $\pm$ 26	7.8$\pm$3.6	4.4 $\pm$ 1.1	106$\pm$46	666$\pm$231	10.1 $\pm$ 1.7	16.3 $\pm$ 7.8	NA	8.8$\pm$2.0	672$\pm$208
Pericardial appendages	NA	NA	NA	20.3$\pm$6.2	2.16 $\pm$ 1.63	1.00 $\pm$ 0.82	260$\pm$111	1.31 $\pm$ 0.80	8.25$\pm$1.55	27.2 $\pm$ 33.4	78 $\pm$ 46	25.1 $\pm$ 23.0	5.64 $\pm$ 3.54	NA	0.18 $\pm$ 0.09	22.4 $\pm$ 26.6
Renal appendages	NA	NA	NA	1.80 $\pm$ 2.22	1.87 $\pm$ 1.23	1.06$\pm$0.88	112 $\pm$ 65	0.43 $\pm$ 0.37	2.54 $\pm$ 2.88	7.1 $\pm$ 10.2	343 $\pm$ 160	77.1$\pm$54.1	22.7$\pm$13.9	NA	1.02 $\pm$ 0.72	11.0 $\pm$ 16.8
<i>N. macromphalus</i> (Present study)																
Digestive gland	0.30$\pm$0.07	0.25$\pm$0.11	0.19$\pm$0.07	9.42$\pm$3.91	51.5$\pm$3.5	0.98 $\pm$ 0.15	257$\pm$19	4.69$\pm$1.51	2.95 $\pm$ 0.35	263$\pm$78	353$\pm$63	10.9 $\pm$ 1.3	14.0 $\pm$ 1.04	2.65 $\pm$ 0.90	14.8$\pm$2.28	388$\pm$61
Pericardial appendages	0.11 $\pm$ 0.02	0.17 $\pm$ 0.05	0.15 $\pm$ 0.05	1.82 $\pm$ 0.54	1.93 $\pm$ 0.76	1.57 $\pm$ 0.11	169 $\pm$ 19	0.65 $\pm$ 0.09	9.59$\pm$3.35	73.7 $\pm$ 9.9	19 $\pm$ 1	12.7 $\pm$ 2.9	6.93 $\pm$ 1.37	3.72 #	5.62 $\pm$ 2.26	56.5 $\pm$ 1.8
Renal appendages	<dl	<dl	<dl	0.44 $\pm$ 0.12	3.39 $\pm$ 1.16	2.11$\pm$0.51	152 $\pm$ 17	0.18 $\pm$ 0.03	2.57 $\pm$ 0.36	27.6 $\pm$ 4.2	20 $\pm$ 3	68.2$\pm$6.2	19.7$\pm$2.39	9.84$\pm$0.94	1.54 $\pm$ 0.13	35.2 $\pm$ 2.8
<i>N. pompilius</i> (Present study)																
Digestive gland	1.00$\pm$0.14	1.58$\pm$0.32	0.92$\pm$0.10	5.65$\pm$1.04	74.3$\pm$6.8	0.62 $\pm$ 0.07	210 $\pm$ 16	3.85$\pm$1.04	1.37 $\pm$ 0.24	311$\pm$61	625$\pm$51	15.4 $\pm$ 1.7	9.77 $\pm$ 0.07	2.35 $\pm$ 0.84	8.42$\pm$0.57	470$\pm$52
Pericardial appendages	0.24 $\pm$ 0.03	0.42 $\pm$ 0.05	0.31 $\pm$ 0.01	1.75 $\pm$ 0.07	2.39 $\pm$ 0.37	1.24$\pm$0.24	323$\pm$53	0.79 $\pm$ 0.07	9.30$\pm$3.87	99.7 $\pm$ 2.4	21 $\pm$ 3	24.1 $\pm$ 4.9	7.02 $\pm$ 0.79	2.49 #	4.94 $\pm$ 0.74	58.2 $\pm$ 2.7
Renal appendages	<dl	<dl	<dl	0.35 $\pm$ 0.03	5.93 $\pm$ 0.31	0.99 $\pm$ 0.21	292 $\pm$ 80	0.18 $\pm$ 0.03	1.98 $\pm$ 0.08	53.3 $\pm$ 4.6	24 $\pm$ 3	307$\pm$83	19.6$\pm$0.97	13.3$\pm$6.12	1.82 $\pm$ 0.02	42.7 $\pm$ 1.5

532 Table 3. One-way ANOVA comparison of trace element concentrations in *Nautilus pompilius*
533 (Np) and in *N. macromphalus* (Nm). NS: not significant.
534

Element	Digestive gland	Pericardial appendages	Renal appendages
Rare earth elements			
Ce	Np > Nm (F= 19.58 ; p= 0.007)	NS	-
La	Np > Nm (F= 23.40; p= 0.005)	Np > Nm (F= 0.609; p= 0.025)	-
Nd	Np > Nm (F= 25.13 ; p= 0.004)	Np > Nm (F= 6.54 ; p= 0.051)	-
Essential elements			
As	NS	Np > Nm (F= 9.58 ; p= 0.027)	NS
Co	NS	NS	NS
Cr	Nm > Np (F= 11.34 ; p= 0.020)	NS	NS
Cu	NS	NS	Np > Nm (F= 15.58; p=0.008)
Fe	Np > Nm (F= 10.09 ; p= 0.025)	NS	NS
Mn	NS	NS	NS
Ni	NS	NS	NS
Se	NS	NS	NS
V	NS	NS	NS
Zn	NS	NS	NS
Non essential elements			
Ag	NS	NS	NS
Cd	Nm > Np (F= 0.740 ; p= 0.023)	NS	NS
Pb	NS	NS	NS

535 Table 4: Correlations between trace elements within the different tissues of nautilus.

536 Not underlined: $p < 0.05$; underlined: $p < 0.001$.

537

Elements	Digestive gland	Renal appendages	Pericardial appendages
La	<u>+ Ce, + Nd, - As, + Fe, + Mn</u>	<dl	<u>+ Ce, + Nd, + Mn</u>
Ce	<u>+ Nd, - As, + Fe, + Mn,</u>	<dl	<u>+ Nd, + Co, + La</u>
Nd	<u>+ Fe, + Mn</u>	<dl	<u>+ Ce, + Co, + La</u>
Cu	+ Zn, -Se	-	-
Fe	+ Mn, -V, + <u>La, + Ce, + Nd</u>	+ Zn	+ Ag, <u>+ Mn</u> , + Cr
Pb	+ V	- V, <u>+ Cr</u>	-
Mn	+ La, + Ce, + Nd, + Fe	+ Cd, +Zn	+ La, <u>+ Fe</u>
Cd	-	<u>+Mn</u>	<u>+ Ni, + V, + Cr</u>

538

539 <dl: concentrations below the detection limit.

Caption to figures

Figure 1: Location of *Nautilus pompilius* and *N. macromphalus* specimens' collection.

Figure 2: Relative distribution of the trace elements between the digestive gland (DG), pericardial appendages (PA) and renal appendages (RA) of *Nautilus pompilius* from Vanuatu (black) and *N. macromphalus* from New Caledonia (white). Scale bars represent 1 standard error.

Figure 1.