

Orbital elements of double stars: ADS 7307 and ADS 17030

Marco Scardia, Jean-Louis Prieur, Laurent Koechlin, Eric Aristidi

▶ To cite this version:

Marco Scardia, Jean-Louis Prieur, Laurent Koechlin, Eric Aristidi. Orbital elements of double stars: ADS 7307 and ADS 17030. Information circular - IAU Commission 26. Double stars, 2002, 147, pp.1. hal-00339987

HAL Id: hal-00339987

https://hal.science/hal-00339987

Submitted on 23 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTERNATIONAL ASTRONOMICAL UNION COMMISSION 26 $({\small DOUBLE~STARS})$

INFORMATION CIRCULAR No. 147 (JUNE 2002)

NEW ORBITS								
$\mathbf{ADS} \\ \alpha 2000 \delta$	Name n	P a	T i	e ω	$\Omega(2000)$ Last ob.	2002 2003	${f Author(s)}$	
197 00153+4412	A 1256 AB 2°6095	$137^{y}96$ $0''153$	1997.78 62°9	0.724 118°1	68°4 1991.9	254°7 0"060 259.6 0.064	OLEVIC	
$440 \\ 00321 + 6715$	GJ 22 AC 22.1675	$16.24 \\ 0.535$	2000.39 46.6	0.178 91.3	179.0 2001.841	330.2 0.414 350.9 0.493	DOCOBO et al. (*)	
00427-6537	I 440 1.3246	271.78 0.457	1948.45 122.4	0.645 352.9	63.8 1991.73	272.3 0.431 271.8 0.439	LING	
$ \begin{array}{c} 650 \\ 00470 + 2315 \end{array} $	HU 413 1.0246	$351.7 \\ 0.623$	2074.21 52.5	0.424 25.9	44.9 1994.9	304.7 0.405 306.0 0.402	OLEVIC	
705 $00520+3154$	A 924 2.1727	$165.70 \\ 0.274$	1987.03 153.9	0.639 335.0	55.2 1996.7	331.6 0.183 328.4 0.190	OLEVIC	
01036+6341	MLR 87 6.3241	56.93 0.237	2011.83 131.4	$0.515 \\ 130.5$	62.1 1996.0	56.7 0.239 53.0 0.223	OLEVIC	
$993 \\ 01131 + 2942$	A 1260 2.8714	125.38 0.357	2013.12 82.9	0.736 88.5	59.8 1995.9	54.9 0.227 55.5 0.222	OLEVIC	
$1016\\01158\!+\!0947$	A 2102 1.9307	$186.46 \\ 0.592$	1934.61 97.6	0.398 255.6	147.8 1994.97	138.4 0.495 138.0 0.487	OLEVIC	
$1087 \\ 01200-1549$	HJ 2036 0.2494	1443.34 3.431	2674.91 114.1	0.092 288.9	124.8 1999.8	341.1 2.253 340.8 2.261	OLEVIC	
$2051 \\ 02423 + 4925$	HU 539 1.7549	$205.14 \\ 0.238$	2010.42 134.4	0.631 124.8	29.3 1995.9	343.3 0.094 336.6 0.096	OLEVIC	
2177 02512+0141	A 2338 2.2209	162.10 0.471	1975.49 73.0	0.610 114.0	127.0 1996.0	338.4 0.247 340.3 0.243	OLEVIC	

NEW ORBITS (continuation)

$\mathbf{ADS} \\ \alpha 2000 \delta$	Name n	P a	T i	e ω	$\Omega(2000)$ Last ob.		${f Author(s)}$
03496+6318	CAR 1 35.3634	10.18 0.102	1998.10 37.9		10.4 2000.7676	91.7 0.112 111.5 0.117	
04271+2542			1981.34 135.1		16.2 1998.775		DOCOBO & TAMAZIAN
04325+1732	GG TAU Aa 1.8311		2060.00 142.9	0.317 241.4	96.2 2001.110	347.4 0.245 345.9 0.244	
06253+0130			2001.03 129.5	0.069 220.6	177.8 1993.1	326.7 0.148 324.0 0.144	OLEVIC
	FIN 334 Aa 1.6880		1974.87 118.0	0.250 56.0	125.3 1993.0897	337.1 0.094 335.1 0.098	MANTE
6552 08047+4717	A 2050 3.1348	114.84 0.152	1986.22 22.7	0.753 40.3	89.4 1996.8638		ZIRM
7307 09210+3811	STF 1338 AB 1.1871	303.27 1.336	2023.25 29.9	0.254 191.9	137.3 1999.99		SCARDIA et al. (***) I
7307 09210+3811	STF 1338 AB 0.8103		1983.69 33.4	0.247 83.6	177.4 1999.99		SCARDIA et al. (***) II
11441-0448	RST 5524 6.6055		1984.34 48.3			154.5 0.159 157.6 0.165	ZIRM
17075+3810	COU 1291 7.0547		2011.66 62.6		121.2 1998.779	3297.7 0.170 301.0 0.164	
12961 19487+1504	A 1658 4.0863		1991.60 169.8	0.087 228.2	$51.5 \\ 2000.5$	133.4 0.207 128.7 0.208	DOCOBO & LING
13894 20290+0710	A 610 2.1100	170.62 0.472	1984.25 30.3	0.232 285.2	72.0 2000.519	57.3 0.393 59.8 0.398	COSTADO
20329+1142	J 1 0.6813	528.39 1.545	2396.22 58.5	0.745 228.4	38.8 1991.2	53.4 2.055 53.5 2.059	POPOVIC & OLEVIC
14749 21118+5959	MCA 67 Aa 4.4280	81.30 0.071	2034.547 112.7	0.347 82.7	75.8 1997.798	137.8 0.040 132.8 0.041	MANTE

NEW ORBITS (continuation)

$\mathbf{ADS} \\ \alpha 2000 \delta$	Name n	P a	T i	e ω	$\Omega(2000)$ Last ob.	$2002 \\ 2003$	$\operatorname{Author}(\operatorname{s})$
17030	A 424	173.05	1934.26	0.489	59.3	144.0 0.15	SCARDIA
23498 + 2741	2.0803	0.255	65.2	283.5	1998.663	146.0 0.15	et al. (***)

- (*) DOCOBO, TAMAZIAN, WOITAS and LEINERT
- (**) DOCOBO, TAMAZIAN and WOITAS
- (***) SCARDIA, PRIEUR, KOECHLIN and ARISTIDI

ANNOUNCEMENT

Changes to the WDS:

- The WDS currently consists of 585,254 mean positions of 98,084 systems.
- Duplicate Discovery Designations Removed:

One of the more difficult issues in dealing with the WDS as a database is the presence of duplicate discovery designations; that is, different systems assigned the same 3-character and 4-digit designation. These generally fall into one of two categories: systems given the same numbers but published in different lists, and those given some additional designation appended to the original one.

Examples of the first are the binaries first resolved by W. Herschel and both F.G.W. Struve and O. Struve. William Herschel published seven lists (I - VI, plus "new", or N), with stars of each list starting at number 1. In addition to their original discovery lists, each of the Struve's published an appendix, as well as a list of "rejected" doubles. These multiple lists were completely spelled out in the Aitken Double Star Catalogue (e.g., H IV 48), but when the Index (IDS) Catalogue was compiled at Lick all of these other designators were dropped for lack of space. To uniquely identify a system then required both the discovery designation and the position. As a result there are, for example, five components with the designation H 48! The source Herschel list was given in the notes file to the IDS. Appended and rejected stars from the lists of the Struve's were handled with an "a" or "r" towards the end of the WDS data line in most cases.

In the second (and fortunately rare) case, systems found quite near to known ones were given the same designation plus trailing character(s) (e.g., ES 1293a or BU 885 1/2). Sometimes both components were assigned these additional characters,

sometimes only one; occasionally two pairs in an entirely different section of the sky were given the same designation by the author (probably by mistake).

Each of these cases is being handled in a different manner. For the William Herschel discoveries, a list identifier is added to column three of each designation. For example:

H 19 (at 16 hours) was originally H II 19 and is now known as H 2 19, H 7 (at 18 hours) was originally H V 7 and is now known as H 5 7, and H 111 (at 06 hours) was originally H N 111 and is now known as H N 111.

In the case of the O. Struve appendix an A is added following STT in the name. For F. Struve, he provided two appendices. Those from the shorter list (Appendix II) are designated STFB. For example:

STF 11 (appendix I) is now STFA 11. STF 11 (appendix II) is now STFB 11. STT 252 (appendix) is now STTA252.

Stars of the second type are given the same 3-letter discovery designation but a new number, starting with 9001, to indicate that they originally had a different designation. For example:

BAL2356b is now BAL9001. BU 885 1/2 is now BU 9001.

A complete list of stars of the second type is provided in the error correction file. All changes in designation are described in the notes file. In addition to these, designations for 271 W. Herschel (H), 110 F. Struve (STF) and 227 O. Struve (STT) systems have been changed. Note that for some of these systems, the former three character, four digit reference (a3i4) has been replaced by a four character, three digit reference (a4i3). Although, for all USNO applications (e.g., data or observing list request) an a7 read will see no difference.

• Arcsecond Precise Coordinates:

Coordinates which are sixty times more precise than the WDS identifier are now provided for the majority of WDS systems (i.e., tenths of a second of time and seconds of arc). Coordinates are obtained from Hipparcos, Tycho-2, the Tycho Double Star Catalogue, or through manual inspection. It is expected that the next significant improvement in this area will be when either the 2MASS or UCAC data are cross-referenced with the WDS. Incremental changes will continue to be made as well. The ten-digit WDS identifier is retained and will continue to be the same for all components in hierarchical systems. However, the precise coordinate will be for the primary of the subsystem. For example, in the case of a system made up of A-BC and BC pairs, they will both have the same ten-digit WDS identifier, however, the fourteen digit precise coordinate would be of the A and B component, respectively.

So far this matching has been made for 80% of the WDS (i.e., 78,655 systems). These coordinates are found close to the end of the WDS summary line.

• Secondary Proper Motions added:

The most common note (N = 1421) in the WDS notes file was the proper motion of the secondary, when known. The above matching with Tycho-2 has allowed the determination of secondary proper motion for 37% of the WDS (i.e., 36,042 systems). The secondary proper motion is found following the precise position. This is expected to grow considerably when the UCAC is complete.

Brian D. Mason & William I. Hartkopf U.S. Naval Observatory

The deadline for contributions to Information Circular No. 148 is:

October 15th 2002

J. A. Docobo (oadoco@usc.es)

J. F. Ling (oafana@usc.es)

Tel. +34 981592747 Fax: +34 981597054

Observatorio Astronómico R. M. Aller P. O. Box 197 http://www.usc.es/astro Universidade de Santiago de Compostela SPAIN

ISSN: 1024-7769