Delineation of Pb contamination pathways in two Pectinidae: the variegated scallop

Chlamys varia* and the king scallop *Pecten maximus

Marc Metian^{1-2†}, Michel Warnau^{1‡}, François Oberhänsli¹, Paco Bustamante^{2*}

¹ International Atomic Energy Agency – Marine Environment Laboratories, 4 Quai Antoine
Ier, MC-98000 Principality of Monaco

² Littoral Environnement et Sociétés (LIENSs), UMR 6250 CNRS-Université de La Rochelle,
2 rue Olympe de Gouges, F-17042 La Rochelle Cedex 01, France

Correspondence to: Prof. Paco Bustamante

LIENSs, UMR 6250, CNRS-Université de La Rochelle

2 rue Olympe de Gouges

F-17042 La Rochelle Cedex 01

France

Phone : +33 5 46507625

Fax : +33 5 46458264

E-mail : pbustama@univ-lr.fr

* Corresponding author.

[†] Present address : Hawaii Institute of Marine Biology, University of Hawaii at Manoa,
Kaneohe, HI 96744 (USA) E-mail : metian@hawaii.com

[‡] Present address : LIENSs, UMR 6250, CNRS-Université de La Rochelle, 2 rue Olympe de
Gouges, F-17042 La Rochelle Cedex 01 (France). E-mail : warnaumichel@yahoo.com

Abstract: Bioaccumulation of Pb was determined in *Chlamys varia* and *Pecten maximus* exposed to ^{210}Pb via seawater, food and sediment. Both scallops readily concentrated dissolved Pb with whole-body 7-d concentration factors of 250 ± 40 and 170 ± 70 , respectively. In both species, more than 70% of Pb taken up from seawater was strongly retained within tissues (biological half-life > 1.5 month) whereas Pb ingested with phytoplankton was poorly assimilated ($< 20\%$). As *P. maximus* lives buried in the sediment, this exposure pathway was assessed and showed low bioaccumulation efficiency for sediment-bound Pb (transfer factor < 0.015). Despite the poor transfer efficiency of Pb from food and sediment, the use of a global bioaccumulation model indicated that the particulate pathway (food and/or sediment) constituted the major bioaccumulation route of Pb in both scallops. Whatever the exposure pathway, the digestive gland and kidneys always played a major role in Pb accumulation. In scallop tissues, Pb was predominantly associated with the insoluble subcellular fraction, suggesting a low bioavailability of Pb for scallop consumers.

Keywords: Metal; Uptake kinetics; Retention; Bioavailability; Mollusc

1. Introduction

Lead (Pb) is widely reported as a contaminant of concern in the marine environment (Laws, 2000; Clark, 2001). Marine contamination by this non-essential metal is mainly due to releases from anthropogenic activities and the atmosphere constitutes the principal transport vector towards the Oceans (Laws, 2000). Pb is bioaccumulated by living organisms and more particularly by marine invertebrates (e.g., Temara et al., 1998; Neff, 2002).

Among filter-feeders, scallops have demonstrated their ability to concentrate elevated levels of various metals in their tissues (e.g., Bryan, 1973; Bustamante and Miramand 2005a,b; Metian et al., 2007, 2008a,b), even in areas far from anthropogenic sources such as the Antarctic Ocean (e.g., Berkman and Nigro, 1992). In particular, scallops were reported to display higher bioaccumulation capacity for Pb than other filter-feeders such as oysters and mussels occurring in the same areas (Brooks and Rumsby, 1965; Segar et al., 1971). Moreover, Pb concentrations in *Chlamys varia* tissues were shown to reflect the contamination level of their environment (Bustamante and Miramand, 2005a). However, little is known about the metabolism of Pb in pectinids and available information is mainly restricted to baseline Pb concentrations in tissues and organs of scallops from different geographic areas, viz. the Antarctic Ocean (Berkman and Nigro, 1992), the Greenland waters (Johansen et al., 2000) and the coastal waters of France, United Kingdom and Canada (Bryan, 1973; Ray et al., 1984; Bustamante and Miramand, 2005b).

Therefore, the aim of the present work was to investigate the pathway-specific bioaccumulation of Pb in pectinids (viz., water, food and/or sediment pathways). Two pectinid species were considered (the variegated scallop *Chlamys varia* and the king scallop *Pecten maximus*) and bioaccumulation was investigated using highly sensitive radiotracer techniques (^{210}Pb) (Warnau and Bustamante, 2007). The bioaccumulation of ^{210}Pb was

studied at the levels of 1) whole individual, 2) tissues and organs and 3) cells of the different tissues, in order to determine the biokinetic parameters of Pb uptake and retention, and the body and subcellular distribution of Pb. In addition, the relative contribution of each exposure pathway to the global Pb bioaccumulation was assessed for both species using a dynamic bioaccumulation model (Landrum et al., 1992; Thomann et al., 1995; Wang et al., 1996; Metian et al., 2008b).

2. Materials and methods

2.1. Sampling

During spring 2005, one hundred variegated scallops, *Chlamys varia*, and seventy king scallops, *Pecten maximus*, were collected on the French Atlantic coast (Pertuis Breton, Charente-Maritime) by SCUBA diving. They were cautiously transported to IAEA-MEL premises in Monaco and were acclimated to laboratory conditions (constantly aerated open circuit aquarium; flux: 50 l h⁻¹, salinity: 36 p.s.u.; temperature: 17 ± 0.5 °C; pH: 8.0 ± 0.1; light/dark cycle: 12 h/12 h) for 6 weeks prior to experiments. During this period, scallops were fed daily an algal mixed diet (5 10⁴ cell ml⁻¹) composed of *Isochrysis galbana* and *Skeletonema costatum*.

2.2. Radiotracer and counting

The ²¹⁰Pb radiotracer of high specific activity was purchased from CERCA LEA, France (²¹⁰Pb as Pb(NO₃)₂ in 1M HNO₃, T_{1/2} = 22.3 years). The tracer was γ-counted (46.5 KeV; 4.05% intensity) using two well-type NaI detectors connected to a multi-channel analyzer and a computer equipped with a spectra analysis software (Interwinner[®] 6). The absolute detection efficiency was 3.4 ± 0.2% under the conditions selected. The radioactivity was determined by

comparison with standards of known activity and of appropriate geometries. Measurements were corrected for counting efficiency and physical radioactive decay. Activities of ^{210}Pb in the samples were much over the detection limits but the counting time was adjusted to obtain a propagated counting error less than 5% (Rodriguez y Baena et al., 2006a).

2.3. Seawater exposure

Fifteen *C. varia* and 15 *P. maximus* (average weight $\pm$ SD: 29 ± 6 g and 73 ± 6 g, respectively) were placed in a 70-l glass aquarium (0.45- μm filtered seawater, constantly aerated closed circuit aquarium; salinity: 36 p.s.u.; temperature: 17 ± 0.5 °C; pH: 8.0 ± 0.1 ; light/dark cycle: 12 h/12 h) and exposed for 7 d to ^{210}Pb dissolved in seawater (0.5 kBq l $^{-1}$). No change in pH was detectable after the tracer addition. ^{210}Pb spike and seawater were renewed twice a day the first two days and then daily in order to keep radioactivity constant in seawater. During each renewal of seawater and spike, the scallops were fed briefly (30 min) *S. costatum* and *I. galbana* (5×10^4 cells ml $^{-1}$) in clean seawater. ^{210}Pb activity in seawater was checked before and after each spike renewal, yielding a time-integrated activity of ^{210}Pb (0.33 ± 0.14 kBq l $^{-1}$) (Rodriguez y Baena et al., 2006b).

Nine tag-identified scallops of each species were removed at different time intervals to be radioanalyzed alive and then returned to their aquarium. At the end of the 7-d exposure period, 5 individuals of *C. varia* and *P. maximus* (not belonging to the tag-identified batches) were sacrificed and dissected. Shell, digestive gland, kidneys, gills, gonad, mantle, intestine, adductor muscle and the remaining soft tissues were separated and radioanalyzed to assess ^{210}Pb body distribution. The 10 remaining scallops were placed in non-contaminating conditions (constantly aerated open circuit; flux: 50 l h $^{-1}$; salinity: 36 p.s.u.; temperature: 17 ± 0.5 °C; pH: 8.0 ± 0.1 ; light/dark cycle: 12 h/12 h; daily feeding on *S. costatum* and *I. galbana*, 5×10^4 cells ml $^{-1}$) for 36 d (*P. maximus*) or 91 d (*C. varia*). The 9 tag-identified individuals of

each species were regularly radioanalyzed to determine the depuration kinetics of ^{210}Pb . At the end of the depuration period, 4 individuals of *C. varia* and *P. maximus* were collected and dissected into several body compartments as previously described.

When dissections were carried out, subcellular distribution of ^{210}Pb between soluble and insoluble fractions was also investigated, according to the method described in Bustamante and Miramand (2005b). Briefly, a part of digestive gland, gills and adductor muscle was homogenized individually with a mortar and pestle on ice with 10 ml of 0.02 M Tris-HCl buffer, 0.25 M sucrose, 1 mM phenylmethylsulfonylfluoride (PMSF, as protease inhibitor), at pH 8.6 and in presence of dithiothreitol as antioxidant agent. The homogenates were centrifuged at 80,000 G for 1 h at 5°C in a Sorvall RC28S ultracentrifuge to separate particle-free supernatant (cytosol; soluble fraction) from the pellet (insoluble fraction). Homogenate aliquots, cytosols, and pellets were then γ -counted.

2.4. Food exposure

The Haptophyceae *Isochrysis galbana* was used to study ^{210}Pb dietary transfer to scallops. Phytoplankton cells were exposed to 1.5 kBq l⁻¹ of ^{210}Pb during their exponential growth phase (7 d). The cells were then filtrated (1- μm mesh size Osmonic filters) when haemocytometer counting determined a concentration of 10⁷ cell ml⁻¹. Phytoplankton cells were then resuspended in a 70-l aquarium where 6 individuals of both *C. varia* and *P. maximus* (average weight $\pm$ SD: 22 $\pm$ 4 g and 91 $\pm$ 4 g, respectively) had been placed previously for one week. The ^{210}Pb content of the radiolabelled *I. galbana* and exposure medium was γ -counted before and after the filtration. Scallops were allowed feeding for 2 h on the radiolabelled *I. galbana* (average concentration of 5 10⁴ cell ml⁻¹ among this time). After the feeding period, all scallops were γ -counted and flowing seawater conditions (50 l h⁻¹) were restored in the aquarium, with daily feeding on *S. costatum* and *I. galbana* (5 10⁴ cells

ml⁻¹). Over the next 8 d, all individuals were regularly radioanalyzed to follow the whole-body depuration kinetics of ²¹⁰Pb. All individuals were collected after 8 d of depuration and sacrificed to determine the body distribution of the remaining ²¹⁰Pb (shell, digestive gland, kidneys, gills, gonad, mantle, intestine, adductor muscle and the remaining soft tissues).

2.5. Sediment exposure

Sediment was collected in Wimereux (North-Atlantic coast of France). Sediment grain size distribution was determined using a Mastersizer micro (Malvern) and the dry/wet wt ratio was calculated after freeze-drying using a LABCONCO Freezone18. Aerated sediment (8 kg) was dispatched in two 5-l plastic bottles, spiked with ²¹⁰Pb (275 kBq in each bottle) and then constantly agitated for 6 d according to the method described in Danis et al. (2003) and adapted by Metian et al. (2007). After this period, part of the sediment was laid in a 20-l aquarium to form a continuous, 4-cm height layer. Weakly bound ²¹⁰Pb was allowed to leach overnight under flowing seawater (50 l h⁻¹). Twelve *P. maximus* (average weight $\pm$ SD: 143 $\pm$ 3 g) were then placed in the aquarium for 7 d (open circuit, seawater flux: 50 l h⁻¹), during which 8 tag-identified individuals were regularly radioanalyzed alive. During the exposure period, sediment samples were regularly collected and γ -counted to verify that the ²¹⁰Pb activity in sediment remained constant. At the end of the uptake period, 4 *P. maximus* were collected, weighed and dissected. Separated tissues and organs (shell, digestive gland, kidneys, gills, gonad, adductor muscle and remaining soft tissues) were radioanalyzed to determine the body distribution of ²¹⁰Pb. The remaining individuals were transferred for 16 d to a new aquarium (non contaminating conditions; open circuit, seawater flux: 50 l h⁻¹) to follow the depuration kinetics of ²¹⁰Pb. At the end of the depuration period, 4 scallops were collected and dissected as described above to determine the ²¹⁰Pb body distribution. The 4

remaining organisms were dissected grossly (separation of the whole soft parts from the shells) and γ -counted.

2.6. Data analysis

Uptake and depuration kinetics for all exposure pathways were fitted using kinetic models and statistical methods as described by Warnau et al. (1996a,b) and Metian et al. (2007). The level of significance for statistics and modelling was always set at $\alpha = 0.05$.

3. Results

3.1. Seawater exposure

Uptake of ^{210}Pb in whole-body *C. varia* and *P. maximus* followed linear kinetics ($R^2 = 0.81$ and 0.93 , respectively; Fig. 1A). ^{210}Pb was accumulated more rapidly in *C. varia* than in *P. maximus* ($k_u = 36$ vs. 24 d^{-1} , respectively; Fig. 1A). Consequently, the whole-body concentration factor (CF) of ^{210}Pb measured at the end of the uptake period (CF_{7d}) was higher in *C. varia* (250 ± 40) than in *P. maximus* (170 ± 70). The CF_{7d} calculated for the different organs and tissues (Table 1) clearly show that ^{210}Pb was concentrated differently by the body compartments of each species. For *C. varia*, the highest CF_{7d} were found in the digestive gland and kidneys reached ($1,490 \pm 240$ and $1,780 \pm 770$, respectively). These CF_{7d} were not significantly different from each other ($p > 0.05$) but were higher than those of the other body compartments. In the case of *P. maximus*, the kidneys displayed the highest CF_{7d} ($1,890 \pm 1,050$).

In terms of body load, ^{210}Pb distribution among soft tissues also differed between the two species (Table 1). In *C. varia*, ^{210}Pb was mainly distributed in the digestive gland (50%) and only 15% of the total ^{210}Pb was found in the kidneys. In contrast, ^{210}Pb was distributed

193 similarly ($p_{\text{Tukey after arcsin transformation}} > 0.05$) among the different organs of *P. maximus* at the
194 end of the exposure period: digestive gland, kidneys, gills, gonad, mantle, intestine, adductor
195 muscle all accounted for ~12 to ~30% of the total ^{210}Pb body load.

196 After the 7-d exposure experiment, non-contaminating conditions were restored and
197 depuration kinetics were followed for 91 and 36 d for *C. varia* and *P. maximus*, respectively.
198 The whole-body depuration kinetics of ^{210}Pb were best described by a double exponential
199 model ($R^2 \geq 0.85$; Fig. 1) displaying similar ^{210}Pb depuration kinetic parameters for both
200 species. The major part of ^{210}Pb ($A_{01} > 70\%$) was efficiently retained ($T_{b/21} > 1.5$ month; Fig.
201 1A).

202 At the end of the depuration period, the ^{210}Pb body distribution differed from that observed at
203 the end of the exposure period (Table 1). This difference was limited in *C. varia*: the digestive
204 gland remained the organ containing the highest ^{210}Pb proportion ($54 \pm 14\%$ of the total body
205 load), and the only slight but significant difference ($p = 0.028$) was found in the gonad (7% at
206 the end of exposure period vs. 4% at the end of the depuration period). In contrast, difference
207 in ^{210}Pb body distribution between exposure and depuration periods was more pronounced in
208 *P. maximus*: the percentage in the kidney increased from $15 \pm 5\%$ at the end of the uptake
209 period to $70 \pm 19\%$ at the end the depuration period. This increase was due to (1) a significant
210 increase in ^{210}Pb activity in the kidneys between the two periods (from 620 ± 350 up to $1,390$
211 $\pm 1,110 \text{ Bq g}^{-1}$ wet wt); and (2) a simultaneous decrease in ^{210}Pb activity in the other tissues
212 (i.e., 76 to 96% according to the body compartment; data not shown).

213 In Figure 2, we show the subcellular distribution of ^{210}Pb in the digestive gland, gills and
214 adductor muscle of both species at the end of the uptake and depuration periods of the
215 seawater experiment. ^{210}Pb was always mainly present in the insoluble fraction of the cells of
216 the considered tissues ($> 50\%$) and increased between the two sampling times (end of uptake
217 and end depuration period), except in the adductor muscle of *C. varia*.

218

219 3.2. Dietary exposure

220 The depuration kinetics of ^{210}Pb ingested with food (the Haptophyceae phytoplankton
221 *Isochrysis galbana*) were best fitted using a double exponential model in both *C. varia* and *P.*
222 *maximus* ($R^2 = 0.94$ and 0.99 , respectively; Fig. 1 B). Both species were characterized by a
223 rather low assimilation efficiency for ^{210}Pb ingested with food ($\text{AE} = 6 - 18\%$). However,
224 once incorporated, ^{210}Pb was very strongly retained in the tissues of both species, with
225 biological half-lives that were virtually infinite (depuration rate constants k_{el} not significantly
226 different from 0; Fig 1B).

227

228 3.3. Sediment exposure

229 Sediment used in our experiment was mainly composed (95.8%) of grains with a diameter
230 ranging from 76 to 302 μm and was characterized by a dry/wet wt ratio of approximately
231 0.80. The ^{210}Pb activity in spiked sediment remained constant all along the exposure period
232 (viz., $47 \pm 3 \text{ Bq g}^{-1}$ wet wt, $n = 21$).

233 The whole-body uptake kinetics of sediment-bound ^{210}Pb in *P. maximus* was best fitted by a
234 first-order saturation model ($R^2 = 0.62$; Fig. 3) that rapidly reached a plateau (estimated $\text{TF}_{ss} =$
235 0.011 ± 0.001).

236 Among the different body compartments of *P. maximus*, the kidneys displayed the highest
237 TF_{7d} which values were characterized by a rather high inter-individual variation ($\text{TF}_{7d \text{ kidneys}} =$
238 0.20 ± 0.20 ; Table 2). The digestive gland showed the second highest TF_{7d} (0.17 ± 0.07) and
239 contained the highest proportion of the total ^{210}Pb body load (i.e. 46%) at the end of the
240 exposure period (Table 2).

241 The ^{210}Pb whole-body depuration kinetics after exposure via the sediment were best described
242 by a bi-exponential model ($R^2 = 0.54$; Fig. 3). A large fraction of the ^{210}Pb taken up ($A_{0s} =$

47%) was rapidly lost whereas the remaining ^{210}Pb fraction ($A_{01} = 53\%$) was strongly retained in *P. maximus* tissues, with an estimated $T_{b1/2}$ of 43 d (Fig. 3).

When the different tissues were considered separately, the activity levels of ^{210}Pb were near or under the detection limit. Nevertheless, ^{210}Pb activity could be measured accurately in the whole soft tissues and compared to that determined at the end of the uptake period. A significant decrease of ^{210}Pb activity was observed at the end of the depuration period (from 0.88 ± 0.19 down to $0.14 \pm 0.02 \text{ Bq g}^{-1}$ wet wt). After 16 d in non contaminated conditions, only 15% of the initial ^{210}Pb activity remained associated with the whole soft parts (data not shown).

4. Discussion

Pectinids are well known to accumulate high levels of metals in their tissues (e.g., Brooks and Rumsby, 1965; Bryan, 1973; Bustamante and Miramand, 2005b; Metian et al., 2007, 2008a,b). However, only few works have focused on Pb levels in scallops (see Table 3). These studies have shown that the concentrations of this metal in the whole soft parts ranged between 0.65 and $16 \mu\text{g g}^{-1}$ dry wt, which are generally higher than those found in other filter-feeders such as oysters and mussels inhabiting the same area (e.g., Brooks and Rumsby, 1965; Segar et al., 1971). It has also been showed that Pb body concentrations in *Chlamys varia* were directly related to the degree of Pb contamination in the surrounding environment (Bustamante and Miramand, 2005b) and that, in a given area, Pb concentrations varied according to the scallop species considered (Bryan, 1973). Therefore it appeared important to investigate Pb uptake and depuration kinetics in scallops exposed via different contamination pathways in order to better understand/characterize the processes driving Pb bioaccumulation in scallops.

267 When exposed to dissolved Pb, both the variegated scallop *C. varia* and the king scallop
 268 *Pecten maximus* displayed efficient bioconcentration capacities. *C. varia* bioconcentrated Pb
 269 ca. 50% faster than *P. maximus*. After 7 days of exposure, uptake kinetics were still in a linear
 270 phase and reached relatively elevated concentration factors (CF) for both species ($CF_{7d} = 250$
 271 ± 40 for *C. varia* and 170 ± 70 for *P. maximus*) compared to other organisms such as the
 272 mussel *Mytilus galloprovincialis* (CF_{7d} : 150 ± 20 ; Boudjenoun et al., 2007). The high
 273 bioconcentration capacity of the scallops is probably related to their biological characteristics
 274 as they have a more efficient filtration system and/or a larger gill surface area than mussels
 275 (Beninger and St-Jean, 1997, Ward et al., 1998). Indeed, gills are generally considered as the
 276 main entry of dissolved metals in marine filter-feeder organisms (Marigómez et al., 2002).
 277 Once taken up from the dissolved phase, most of the Pb ($A_{01} > 70\%$) was strongly retained
 278 within the tissues of both species ($T_{b/2l} > 1.5$ month). These observations are in accordance
 279 with previous studies that reported elevated absorption efficiencies ($A_{01} \geq 60\%$; Fisher et al.,
 280 1996) and retention ($T_{b/2l} \geq 23$ d; Boudjenoun et al., 2007) of dissolved Pb in mussels. Such
 281 efficient bioconcentration and retention of Pb in *C. varia* and *P. maximus* suggests that
 282 seawater could play an important role in the global Pb bioaccumulation in scallops.
 283 When scallops were exposed to Pb via the food pathway, the metal was poorly assimilated by
 284 the two species ($AE \leq 18\%$). The major part of the Pb ingested with phytoplankton ($> 80\%$)
 285 was lost within less than 2 days. The Pb AE determined in scallops is much lower than in the
 286 mussel *Mytilus galloprovincialis* fed the phytoplankton *Thalassiosira pseudonana* ($AE =$
 287 50% , Fisher et al., 1996). Metian et al. (2008b) have shown that the AE of Ag ingested with
 288 food was influenced substantially by the phytoplankton species used as food. However,
 289 according to Fisher et al. (1983; 1987), this should not be the case for Pb as these authors
 290 have shown that Pb bioaccumulation is rather non specific and invariant with physiological
 291 condition of the phytoplankton cells. Therefore the low Pb AE found for both scallop species

would not be due to the particular phytoplanktonic species used as food but rather to some specificities of the scallops (e.g., differences in digestive processes). Interestingly, even though AE ranged between only 6 and 18%, we found out that dietary transfer may play a major role in the global accumulation of Pb in the scallops (see below).

In contrast to *C. varia* which is living on rocky substrata, *P. maximus* lives buried into the sediments. When exposed to radiolabelled sediment, uptake of Pb by *P. maximus* was quite poor ($TF_{ss} = 0.011$). However the assimilated fraction of the sediment-bound metal was retained with a similar biological half-life ($T_{b/2l} = 43$ d) than when taken up via the food (52 d) or seawater (47 d). The granulometry of the sediment used in our study (mostly composed of grain of size from 76 to 302 μm) allows assuming that no ingestion of particles did occur, as the gills, the peribuccal system or the stomach were reported to be efficient in selecting the size of the particles that are actually ingested by the scallops (e.g., Beninger et al., 1991; Brillant et al., 2000). If this hypothesis was to be true, this would suggest that sediment-related contamination would have occurred through Pb bioaccumulation from porewater. However, two facts are supporting the counter hypothesis that particles could actually have been ingested. First, some studies have reported the occurrence of large particles (up to 950 μm) in the stomachs of scallops (Mikulich and Tsikhon-Lukanina, 1981; Shumway et al., 1987) and, second, the digestive gland contained 45% of the whole Pb load when scallops were exposed to contaminated sediment. As a porewater exposure pathway should be equivalent to a seawater exposure, one would have expected to find a rather more homogeneous distribution among digestive gland, gills and kidneys, as was observed at the end of the exposure period in our seawater experiment (see Table 1). Hence, although requiring confirmation, ingestion of sediment grains could actually represent a source of Pb exposure for *P. maximus*.

316 In order to assess the contribution of the different routes of exposure in global Pb
 317 bioaccumulation, the biokinetic parameters determined from our experiments were used along
 318 with parameters taken from the literature (distribution coefficient in sediment, concentration
 319 factor in phytoplankton, scallops ingestion rate; IAEA, 2004; Metian et al., 2008b) to feed the
 320 global bioaccumulation model originally described Thomann (1981). This model was further
 321 developed and commonly used by several authors (e.g., Thomann et al., 1995; Wang et al.,
 322 1996; Reinfelder et al., 1998; Metian et al., 2008b). Assuming that Pb is distributed in the
 323 different compartments of the marine environment (seawater, phytoplankton and sediment)
 324 according to the distribution coefficient in sediment ($K_{d,Pb} = 10^5$; ocean margin sediments)
 325 and the range of concentration factors in phytoplankton ($CF_{phyto,Pb} = 10^4 - 10^5$) available from
 326 the literature (IAEA, 2004), the model computations indicated that the relative contributions
 327 of each exposure route were substantially influenced by the $CF_{phyto,Pb}$ values. In particular, in
 328 *C. varia* for which only seawater and food exposure pathways were considered, the dietary
 329 pathway was found to contribute for 86% of the global Pb bioaccumulation when considering
 330 the recommended $CF_{phyto,Pb}$ value (10^5 ; IAEA, 2004), whereas the dissolved pathway was the
 331 dominant pathway (62%) when considering the lowest $CF_{phyto,Pb}$ value (10^4). Lower food
 332 contribution when $CF_{phyto,Pb}$ increase could be related to a stronger retention of Pb on the
 333 phytoplanktonic cell, thus reducing the bioavailability of the metal. For the sediment-dweller
 334 *P. maximus*, three different exposure pathways were taken into account (seawater, food and
 335 sediment). Model computations showed that the particulate pathway (food and sediment) was
 336 always the major source (95 - 97%) of Pb bioaccumulation for *P. maximus*. However,
 337 depending on the $CF_{phyto,Pb}$ value, the contributions of food vs sediment pathways shifted from
 338 5 vs 90% (for $CF_{phyto,Pb} = 10^4$) to 33 vs 64% (for $CF_{phyto,Pb} = 10^5$). As available values for
 339 $CF_{phyto,Pb}$ and $K_{d,Pb}$ parameters are essentially average, non-specific values published by the
 340 IAEA (2004), it is recommended to further refine these two parameters for environmental

conditions close to those prevailing in the scallop habitats before further interpreting and extrapolating the results presented here.

The kidneys and the digestive gland appeared to play an important role in the uptake and retention of Pb in the tissues of both scallop species. However, these organs were involved differently according to the exposure pathway and the species considered. In scallops collected in the field, the highest Pb concentrations are always reported in the kidneys, when those organs were analysed separately (Table 3). In addition, the kidneys contained more than 50% of the total Pb body burden in *Aequipecten opercularis*, *C. varia* and *P. maximus* (e.g., Bryan, 1973; Bustamante and Miramand, 2005b). As concentrations recorded from the field are supposed to represent a steady-state situation, kidneys may be reasonably considered as organs playing a key role in detoxification and storage of Pb. This is also supported by our experimental observations in *P. maximus*. Indeed, differences in body distribution observed between the end of the seawater uptake and depuration periods suggest that translocation of Pb from different tissues and organs towards the kidneys and/or its storage in these organs occurred in *P. maximus* (which contained 70% of total whole soft parts Pb body burden one month after the exposure period). In contrast, if any translocations of Pb in *C. varia*, target organ(s) of those processes would be different than in *P. maximus* as Pb remained mainly present in the digestive gland all along both exposure and depuration periods. It is known that bivalves take up dissolved Pb mainly through the gills and that the metal is then translocated towards the different organs and tissues via blood granulocytes and high molecular weight plasma ligands (Marigómez et al., 2002). Interspecific variations in Pb concentrations observed in scallops in the field (see Table 3) could thus be related to interspecific differences in translocation mechanisms. In this respect, waterborne Pb would be mainly translocated towards the kidneys (followed by storage) in *P. maximus* whereas it would be mainly translocated towards the digestive gland in *C. varia*. It is noteworthy that in field studies,

renal Pb levels are always much higher in *P. maximus* than in *C. varia*, even for organisms displaying similar whole soft parts Pb concentrations (see Table 3). Nevertheless, it would be worth developing studies on translocation routes to and storage mechanisms in scallop digestive gland and kidneys as it would help greatly in better understanding how Pb as well as other metals are handled and detoxified by these organisms. The results of the subcellular fractioning, in both *C. varia* and *P. maximus*, indicated that Pb was mainly associated with the insoluble cellular fraction, which is consistent with similar fractioning carried out with *C. varia* from the field (Bustamante and Miramand, 2005b). As this insoluble subcellular fraction includes the haemocyte lysosomes which are well-known to play an important role in metal detoxification and storage in bivalves (e.g., Ballan-Dufrançais et al., 1985), further studies should not neglect the role of these free-circulating cells in addressing Pb translocation targetting.

Acknowledgements

We thank the Aquarium of La Rochelle for providing the scallops and J.L. Teyssié (IAEA-MEL) for his valuable technical advices. MW is an Honorary Senior Research Associate of the National Fund for Scientific Research (NFSR, Belgium) and holds a 2008 Invited Expert position at LIENSs (CNRS-Université de La Rochelle), supported by the Conseil Régional de Poitou-Charentes. This work was supported by the IAEA, the French GIP Seine-Aval Programme, LIENSs and the Conseil Général de Charente-Maritime. The IAEA is grateful for the support provided to its Marine Environment Laboratories by the Government of Monaco.

References

389 Ballan-Dufrançais, C., Jeantet, A.Y., Feghali, C., Halpern, S. Physiological features of heavy
390 metal storage in bivalve digestive cells and amoebocytes: EPMA and factor analysis
391 of correspondences. *Biol Cell* 1985;53:283-292.

392 Beninger, P.G., Le Pennec, M. Functional anatomy of scallops. In: Shumway, S.E. (ed)
393 Scallops: biology, ecology and aquaculture. Elsevier, pp. 133-224; 1991

394 Beninger, P.G., St-Jean, S.D. Particle-processing on the labial palps of *Mytilus edulis* and
395 *Placopecten magellanicus* (Mollusca: Bivalvia). *Mar Ecol Prog Ser* 1997;147:117-
396 127.

397 Berkman, P.A., Nigro, M. Trace metal concentrations in scallops around Antarctica:
398 Extending the Mussel Watch Programme to the Southern Ocean. *Mar Pollut Bull*
399 1992;24:322-323.

400 Boudjenoun, R., Teyssié, J.L., Rodriguez y Baena, A., Fowler, S.W., Warnau, M. Lead
401 bioaccumulation in the Mediterranean mussel *Mytilus galloprovincialis*. *Rapp Com*
402 *Intern Mer Méd* 2007;38:237.

403 Brooks, R.R., Rumsby, M.G. The biogeochemistry of trace element uptake by some New
404 Zealand bivalves. *Limnol Oceanogr* 1965;10:521-527.

405 Bryan, G.W. The occurrence and seasonal variation of trace metals in the scallops *Pecten*
406 *maximus* (L.) and *Chlamys opercularis* (L.). *J Mar Biol Assoc UK* 1973;53:145-166.

407 Brilliant, M.G.S., MacDonald, B.A. Postingestive selection in the sea scallop, *Placopecten*
408 *magellanicus* (Gmelin): the role of particle size and density. *J Exp Mar Biol Ecol*
409 2000;253:211-227.

410 Bustamante, P., Miramand, P. Evaluation of the variegated scallop *Chlamys varia* as a
411 biomonitor of temporal trends of Cd, Cu, and Zn in the field. *Environ Pollut*
412 2005a;138:109-120.

413 Bustamante, P., Miramand, P. Subcellular and body distributions of 17 trace elements in the
 414 variegated scallop *Chlamys varia* from the French coast of the Bay of Biscay. Sci Tot
 415 Environ 2005b;337:59-73.

416 Clark, R.B. Marine Pollution. Fifth edition. Oxford University Press, 184 p; 2001

417 Cossa, D., Auger, D., Averty, B., Lucon, M., Masselin, P., Noel, J., Sanjuan, J. Niveaux de
 418 concentration en métaux, métalloïdes et composés organochlorés dans les produits de
 419 la pêche côtière française, IFREMER, Plouzane, France, 24 p ; 1990

420 Danis, B., Cotret, O., Teyssié, J.L., Fowler, S.W., Bustamante, P., Warnau M. Delineation of
 421 PCB uptake pathways in a benthic sea star using a radiolabelled congener. Mar Ecol
 422 Prog Ser 2003;253:155-163.

423 Fang, Z.Q., Cheung, R.Y., Wong, M.H. Heavy metal concentrations in edible bivalves and
 424 gastropods available in major markets of the Pearl River Delta. J Environ Sci (China)
 425 2001;13:210-217.

426 Fisher, N.S., Burns, K.A., Cherry, R.D., Heyraud, M. Accumulation and cellular distribution
 427 of ²⁴¹Am, ²¹⁰Po, and ²¹⁰Pb in two marine algae. Mar Ecol Prog Ser 1983;11:233-237.

428 Fisher, N.S., Teyssié, J.-L., Fowler, S.W., Wang, W.-X. Accumulation and retention of metals
 429 in mussels from food and water: A comparison under field and laboratory conditions.
 430 Environ Sci Technol 1996;30:3232-3242.

431 Fisher, N.S., Teyssié, J.-L., Krishnaswami, S., Baskaran, M. Accumulation of Th, Pb, U, and
 432 Ra in marine phytoplankton and its geochemical significance. Limnol Oceanogr
 433 1987;32:131-142.

434 Greig, R.A., Wenzloff, D.R., Mackenzie, C.L., Merrill, A.S., Zdanowicz, V.S. Trace metals in
 435 sea scallops *Placopecten magellanicus*, from eastern United States. Bull Environ
 436 Contam Toxicol 1978;19:326-333.

437 IAEA. Sediments distribution coefficients and concentration factors for biota in the marine
 438 environment. Technical Report Series (TRS) No. 422. International Atomic Energy
 439 Agency, Vienna, Austria, 95 p; 2004

440 Johansen, P., Pars, T., Bjerregaard, P. Lead, cadmium and selenium intake by Greenlanders
 441 from local marine food. *Sci Tot Environ* 2000;245:187-194.

442 Julshamn, K., Duinker, A., Frantzen, S., Torkildsen, L., Maage, A. Organ distribution and
 443 food safety aspects of cadmium and lead in great scallops, *Pecten maximus* L., and
 444 horse mussels, *Modiolus modiolus* L., from Norwegian Waters. *Bull Environ Contam*
 445 *Toxicol* 2008;80:385-389.

446 Landrum, P.F., Lee, H., Lydy, M.J. Toxicokinetics in aquatic systems: model comparisons
 447 and use in hazard assessment. *Environ Toxicol Chem* 1992;11:1709-1725.

448 Laws, .E.A. Metals. In: *Aquatic Pollution: An Introductory Text*, 3rd ed. John Wiley & Sons,
 449 New York, 639 p; 2000

450 Marigómez, I., Soto, M., Cajaraville, M.P., Angulo, E., Giamberini, L. Cellular and
 451 subcellular distribution of metals in molluscs. *Microscop Res Tech* 2002;56:358-392.

452 Metian, M., Warnau, M., Oberhansli, F., Teyssié, J.L., Bustamante, P. Interspecific
 453 comparison of Cd bioaccumulation in European Pectinidae (*Chlamys varia* & *Pecten*
 454 *maximus*). *J Exp Mar Biol Ecol* 2007;353:58-67.

455 Metian, M., Bustamante, P., Hédouin, L., Warnau, M. Accumulation of trace elements in the
 456 tropical scallop *Comptopallium radula* from coral reefs in New Caledonia. *Environ*
 457 *Pollut* 2008a;152:543-552.

458 Metian, M., Bustamante, P., Cosson, R.P., Hédouin, L., Warnau, M. Investigation of Ag in
 459 the king scallop *Pecten maximus* using field and laboratory approaches. *J Exp Mar*
 460 *Biol Ecol* 2008b;367:53-60.

461 Mikulich, L.V., Tsikhon-Lukamina, A. Food of the scallop. *Oceanol* 1981;21:633-635.

462 Neff, J.M. Bioaccumulation in marine organisms: effect of contaminants from oil well
 463 produced water, Elsevier Science, Oxford, UK, 460 p; 2002

464 Nielsen, S.A., Nathan, A. Heavy Metal Levels in New Zealand Molluscs. New Zealand J Mar
 465 Freshwat Res 1975;9:467-481.

466 Pesch, G.G., Renolds, B., Rogerson, P. Trace metals in scallops from within and around two
 467 ocean disposal sites. Mar Pollut Bull 1977;8:224-228.

468 Ray, S., Woodside, M., Jerome, V.E., Akagi, H. Copper, zinc, cadmium and lead in scallop
 469 (*Placopecten magellanicus*) from Georges and Browns Banks. Chemosphere.
 470 1984;13:1247-1254.

471 Reinfelder, J.R., Fisher, N.S., Luoma, S.N., Nichols, J.W., Wang, W.X. Trace element trophic
 472 transfer in aquatic organisms: A critique of the kinetic model approach. Sci Tot
 473 Environ 1998;219:117-135.

474 Rodriguez y Baena, A.M., Miquel, J.C., Masqué, P., Povinec, P., La Rosa, J. A single vs.
 475 double spike approach to improve the accuracy of ^{234}Th measurements in small-volume
 476 seawater samples. Mar Chem 2006a;100:269-281.

477 Rodriguez y Baena, A.M., Metian, M., Teyssié, J.L., De Broyer, C., Warnau, M.
 478 Experimental evidence for ^{234}Th bioaccumulation in three Antarctic crustaceans:
 479 potential implications in particle flux studies. Mar Chem 2006b;100:354-365.

480 Segar, D.A., Collins, J.D., Riley, J.P. The distribution of the major and some minor elements
 481 in marine animals. Part II. Molluscs. J Mar Biol Assoc UK 1971;51:131-136.

482 Shumway, S.E., Selvin, R., Schick, D.F. Food resources related to habitat in the scallop
 483 *Placopecten magellanicus* (Gmelin, 1791): A qualitative study. J Shellfish Res
 484 1987;6:89-95.

485 Storelli, M.M., Marcotrigiano, G.O. Consumption of bivalve molluscs in Italy: estimated
 486 intake of cadmium and lead. Food Additives Contam 2001;18:303-307.

487 Temara, A., Aboutboul, P., Warnau, M., Jangoux, M., Dubois, P. Uptake and fate of lead in
 488 the common asteroid *Asterias rubens* (Echinoderm). Water, Air, Soil Pollut
 489 1998;102:201-208.

490 Thomann, R.V. Equilibrium model of fate of microcontaminants in diverse aquatic food
 491 chains. Can J Fish Aquatic Sci 1981;38:280-296.

492 Thomann, R.V., Mahony, J.D., Mueller, R. Steady-state model of biota sediment
 493 accumulation factor for metals in two marine bivalves. Environ Toxicol Chem
 494 1995;14:1989-1998.

495 Topping, G. Heavy metals in shellfish from Scottish waters. Aquaculture 1973;1:379-384.

496 Wang, W.-X., Fisher, N.S., Luoma, S.N. Kinetic determinations of trace element
 497 bioaccumulation in the mussel *Mytilus edulis*. Mar Ecol Prog Ser 1996;140:91-113.

498 Ward, J.E., Levinton, J.S., Shumway, S.E., Cucci, T. Particle sorting in bivalves: *in vivo*
 499 determination of the pallial organs of selection. Mar Biol 1998;131:283-292.

500 Warnau, M., Bustamante, P. Radiotracer techniques: a unique tool in marine ecotoxicological
 501 studies. Environ Bioindicators 2007;2:217-218.

502 Warnau, M., Fowler, S.W., Teyssie, J.L. Biokinetics of selected heavy metals and
 503 radionuclides in two marine macrophytes: the seagrass *Posidonia oceanica* and the
 504 alga *Caulerpa taxifolia*. Mar Environ Res 1996a;41:343-362.

505 Warnau, M., Teyssié, J.L., Fowler, S.W. Biokinetics of selected heavy metals and
 506 radionuclides in the common Mediterranean echinoid *Paracentrotus lividus*: sea water
 507 and food exposures. Mar Ecol Prog Ser 1996b;141:83-94.

508 Wu, R., Groves, A. Cadmium and lead in tissues of scallops from Port Phillip Bay, Australia.
 509 Wat Sci Technol 1995;31:473-483.

A. Seawater experiment

A1. Uptake kinetics

A2. Depuration kinetics

B. Food experiment

Figure 1. Uptake and depuration biokinetics of ^{210}Pb in the scallops *Chlamys varia* and *Pecten maximus* (A) exposed for 7 d to the radiotracer via seawater (A1. Concentration Factors; mean $\pm$ SD; $n = 9$), then maintained for 36 or 91 d in non contaminated conditions (A2. Remaining activity, %; mean $\pm$ SD; $n = 9$) and (B) fed for 2 h radiolabelled *Isochrysis galbana* (Remaining activity, %; mean $\pm$ SD; $n = 6$).

Estimated parameters. k_u : uptake rate constant (d^{-1}); A_{0s} and A_{0l} : activity (%) lost according to the short- and the long-lived exponential component, respectively; $T_{b/2}$: biological half-life (d). ns: parameter not significantly different ($p > 0.05$) from 0 (k_{el}) or from the infinite ($T_{b/2l}$).

A. *Chlamys varia*

B. *Pecten maximus*

Figure 2. Subcellular distribution (%) of ^{210}Pb in selected tissues (digestive gland, gills and muscle) of *Chlamys varia* (A) and *Pecten maximus* (B) at the end of the 7-d seawater exposure ($n = 4$ for both species) and of the depuration period (36 d, $n = 4$ for *P. maximus* and 91 d, $n = 4$ for *C. varia*) periods.

1

2

3

4 Figure 3. Uptake and depuration biokinetics of ^{210}Pb in the scallop *Pecten maximus* exposed for 7 d via the sediment (Transfer Factor ; mean $\pm$

5 SD; n = 8), then maintained for 16 d in non contaminated conditions (Remaining activity, %; mean $\pm$ SD; n = 8).

6 Estimated parameters. TF_{ss} : transfer factor at steady state; other parameters as in Fig. 1.

7

Table 1. Concentration Factors (mean $\pm$ SD) and body distribution among soft parts (mean % $\pm$ SD) of ^{210}Pb in the scallops *Chlamys varia* and *Pecten maximus* exposed to the metal via seawater.

Compartment	Seawater exposure		
	End of 7-d uptake period (n = 5)		End of depuration period (91 d for <i>C. varia</i> , 36 d for <i>P. maximus</i> , n = 4)
	Concentration Factor	Distribution (%)	Distribution (%)
<i>Chlamys varia</i>			
Digestive gland	1490 $\pm$ 240	50 $\pm$ 6	54 $\pm$ 14
Gills	250 $\pm$ 240	10 $\pm$ 3	4 $\pm$ 3
Kidneys	1780 $\pm$ 550	15 $\pm$ 6	17 $\pm$ 16
Intestine	490 $\pm$ 680	1 $\pm$ 0	3 $\pm$ 4
Gonad	410 $\pm$ 140	7 $\pm$ 2	4 $\pm$ 1
Foot	550 $\pm$ 360	3 $\pm$ 3	3 $\pm$ 4
Mantle	45 $\pm$ 11	5 $\pm$ 1	9 $\pm$ 8
Adductor muscle	56 $\pm$ 26	8 $\pm$ 3	4 $\pm$ 2
Remaining	230 $\pm$ 80	3 $\pm$ 2	2 $\pm$ 1
Whole body	250 $\pm$ 40		
<i>Pecten maximus</i>			
Digestive gland	570 $\pm$ 250	28 $\pm$ 5	12 $\pm$ 8
Gills	110 $\pm$ 30	22 $\pm$ 7	7 $\pm$ 6
Kidneys	1890 $\pm$ 1050	15 $\pm$ 5	70 $\pm$ 19
Intestine	340 $\pm$ 130	2 $\pm$ 1	< 1
Gonad	300 $\pm$ 230	4 $\pm$ 2	5 $\pm$ 4
Foot	150 $\pm$ 60	1 $\pm$ 1	1 $\pm$ 0
Mantle	35 $\pm$ 8	14 $\pm$ 4	1 $\pm$ 1
Adductor muscle	48 $\pm$ 35	12 $\pm$ 5	4 $\pm$ 2
Remaining	170 $\pm$ 40	4 $\pm$ 2	1 $\pm$ 0
Whole body	170 $\pm$ 70		

Table 2. Transfer Factors (mean $\pm$ SD, n = 4) and body distribution (mean % $\pm$ SD, n = 4) of ^{210}Pb in *Pecten maximus* after 7 d of exposure via the sediment.

Compartment	Transfer Factor	Distribution (%)
Digestive gland	0.17 ± 0.07	46 ± 4
Gills	0.008 ± 0.006	5 ± 5
Kidneys	0.20 ± 0.20	15 ± 13
Gonad	0.05 ± 0.06	7 ± 7
Adductor muscle	0.003 ± 0.003	8 ± 6
Remaining tissues	0.009 ± 0.005	18 ± 5
Whole soft parts	0.017 ± 0.004	100

Table 3. Concentration of Pb ($\mu\text{g g}^{-1}$ dry wt) in different scallop species from various locations.

Scallop species	Location	Whole soft parts concentration ($\mu\text{g g}^{-1}$ dry wt)	Organ displaying highest concentration ($\mu\text{g g}^{-1}$ dry wt)	References
<i>Adamussium colbecki</i>	Antarctica (Explorers cove)	2.6 ± 0.4	-	Berkman and Nigro, 1992
<i>A. colbecki</i>	Antarctica (Swoya station)	from 1.0 to 3.2	-	Berkman and Nigro, 1992
<i>Aequipecten opercularis</i>	U.K. (English channel)	12.0 ± 5.2	kidneys (827 ± 236)	Bryan, 1973
<i>Chlamys islandica</i>	Greenland	0.65 ± 0.12^a	-	Johansen et al., 2000
<i>C. nobilis</i>	Hong Kong markets	2.05 ± 1.35^a	-	Fang et al., 2001
<i>C. nobilis</i>	Guangdong markets	1.0 ± 0.2^a	-	Fang et al., 2001
<i>C. varia</i>	W France (Ré island)	1.91 ± 0.07	kidneys (28.4 ± 7.0)	Bustamante and Miramand, 2005b
<i>C. varia</i>	W France (La Rochelle)	2.66 ± 0.21	kidneys (41.6 ± 3.7)	Bustamante and Miramand, 2005b
<i>Pecten alba</i>	Australia (Port Phillip bay)	$14.06 \pm 0.72^*$	gonad (18.8 ± 3.9) **	Wu and Groves, 1995
<i>P. jacobaeus</i>	Italy (Middle Adriatic Sea)	0.75 ± 0.30^a	-	Storelli and Marcotrigiano, 2001
<i>P. novae-zelandiae</i>	New Zealand (Tasman bay)	16	kidneys (137)	Brooks and Rumsby, 1965
<i>P. novae-zelandiae</i>	New Zealand	6.70^a	stomach (7^a)	Nielsen and Nathan, 1975
<i>P. maximus</i>	NW France (W. English channel)	-	$0.57 \pm 0.14^{\dagger\dagger\dagger}$	Cossa et al., 1990
<i>P. maximus</i>	NW France (E. English channel)	-	$0.69 \pm 0.38^{\dagger\dagger\dagger}$	Cossa et al., 1990
<i>P. maximus</i>	U.K. (English channel)	2.0 ± 0.2	kidneys (159 ± 38)	Bryan, 1973
<i>P. maximus</i>	Scotland (different locations)	from < 0.5 to 5.0^a	gonad (from < 1.0 to 8.5^a) **	Topping, 1973
<i>P. maximus</i>	Irish Sea	8.3	gonad (31) **	Segar et al., 1971
<i>P. maximus</i>	Norway (Frøya)	0.6 ± 0.1^a	kidneys (41 ± 13^a)	Julshamn et al., 2008
<i>Placopecten magellanicus</i>	Canada (George & Browns Banks)	$2.0^\dagger$	gills (2.1) ††	Ray et al., 1984
<i>P. magellanicus</i>	U.S.A. (mid-Atlantic coast)	3.59	-	Pesch et al., 1977
<i>P. magellanicus</i>	U.S.A. (Atlantic coast)	-	gonad ($< 19.5^a$) **	Greig et al., 1978

^a Data on a wet wt basis were converted on a dry wt basis, using a conversion factor of 5 (Greig et al., 1978).

* whole soft parts without gonad and adductor muscle.

** only adductor muscle and gonad were analyzed separately.

† whole soft parts without adductor muscle, mantle and gills.

†† only adductor muscle, mantle, gills and viscera were analyzed separately.

††† only adductor muscle were analyzed separately.