

Connecting walks and connecting dart sequences for n-D combinatorial pyramids

Sébastien Fourey, Luc Brun

► **To cite this version:**

Sébastien Fourey, Luc Brun. Connecting walks and connecting dart sequences for n-D combinatorial pyramids. GREYC-TR-2009-2. 2009. <hal-00408202>

HAL Id: hal-00408202

<https://hal.archives-ouvertes.fr/hal-00408202>

Submitted on 29 Jul 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



GREYC Laboratory
Groupe de Recherche en Informatique, Image, Automatique
et Instrumentation de Caen
CNRS UMR-6072
6, bd maréchal Juin
F-14050 Caen CEDEX 4 – France

Technical report GREYC-TR-2009-02

*Connecing walks and connecting dart
sequences for n -D combinatorial
pyramids*

Sébastien Fourey and Luc Brun
{sebastien.fourey,luc.brun}@greyc.ensicaen.fr

July 2009

Connecting walks and connecting dart sequences for n -D combinatorial pyramids

Sébastien Fourey Luc Brun

*GREYC, UMR6072 – ENSICAEN, 6 bd maréchal Juin 14050 Caen CEDEX,
France*

Abstract

Combinatorial maps define a general framework which allows to encode any subdivision of an n -D orientable quasi-manifold with or without boundaries. Combinatorial pyramids are defined as stacks of successively reduced combinatorial maps. Such pyramids provide a rich framework which allows to encode fine properties of objects (either shapes or partitions). Combinatorial pyramids have first been defined in 2D. This first work has later been extended to pyramids of n -D generalized combinatorial maps. Such pyramids allow to encode stacks of non orientable partitions but at the price of a twice bigger pyramid. These pyramids are also not designed to capture efficiently the properties connected with orientation. This work presents the design of pyramids of n -D combinatorial maps and important notions for their encoding and processing.

Contents

1	Introduction	2
2	Combinatorial maps	3
2.1	Basic definitions	3
2.2	Cells in maps	6
3	Cells removal and contraction in G-maps and maps	10
3.1	Cells removal	10
3.2	Cells contraction	13
4	Connecting walks	14

Email addresses: Sebastien.Fourey@greyc.ensicaen.fr (Sébastien Fourey),
Luc.Brun@greyc.ensicaen.fr (Luc Brun).

4.1	In generalized maps	15
4.2	In combinatorial maps	22
5	Pyramids of maps	29
5.1	Definition	29
6	Connecting dart sequences	30
6.1	Definition and properties	30
6.2	An iterative definition	34
7	Conclusion and perspectives	43
	References	44

1 Introduction

Pyramids of combinatorial maps have first been defined in 2D [2], and later extended to pyramids of n -dimensional generalized maps by Grasset et al. [12]. Generalized maps model subdivisions of orientable but also non-orientable quasi-manifolds [13] at the expense of twice the data size of the one required for combinatorial maps. For practical use (for example in image segmentation), this may have an impact on the efficiency of the associated algorithms or may even prevent their use. Furthermore, properties and constraints linked to the notion of orientation may be expressed in a more natural way with the formalism of combinatorial maps. For these reasons, we are interested here in the definition of pyramids of n -dimensional combinatorial maps.

The key notion for the definition of pyramids of maps is the operation of simultaneous removal or contraction of cells. These two notions have been defined in [9] (see also [10]) where the definitions have been related to the ones given in [7] for generalized maps. Their validity was indeed proved using the link between maps and generalized maps established by Lienhardt [13].

After recalling some preliminaries about combinatorial maps and the main results obtained in [9], we present in this paper two important notions in the design of combinatorial pyramids: connecting walks and connecting darts sequences. These two notions correspond respectively to the notions of *reduction window* and *receptive field* within the regular pyramid framework. They also have interesting properties that should allow us to derive, in future works, efficient encoding schemes and operations on pyramids of n -D maps the same way Brun and Kropatsch did for 2-dimensional combinatorial pyramids [3].

Connecting walks which are introduced in Section 4, somehow fill the gap between two consecutive levels of the pyramid. We first provide a definition of connecting walks in generalized maps and establish a link (Proposition 13) with the definition we give for such walks in combinatorial maps. On the other hand, connecting dart sequences (Section 5) link a level of a pyramid of maps to any of its lower levels. The definition of the latter sequence as well as a discussion of its expected use are given in Section 5.

2 Combinatorial maps

2.1 Basic definitions

We recall here the definitions of n -dimensional generalized maps (n -G-maps for short) and n -dimensional combinatorial maps (n -maps).

Definition 1 (n -G-map [13]) *Let $n \geq 0$, an n -G-map is defined as an $n+2$ -tuple $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ where:*

- \mathcal{D} is a finite non-empty set of darts;
- $\alpha_0, \dots, \alpha_n$ are involutions on \mathcal{D} (i.e. $\forall i \in \{0, \dots, n\}, \alpha_i^2(b) = b$) such that:
 - $\forall i \in \{0, \dots, n-1\}, \alpha_i$ is an involution without fixed point (i.e. $\forall b \in \mathcal{D}, \alpha_i(b) \neq b$);
 - $\forall i \in \{0, \dots, n-2\}, \forall j \in \{i+2, \dots, n\}, \alpha_i \alpha_j$ is an involution.

If α_n is an involution without fixed point, G is said to be without boundaries or closed. In the following we only consider closed n -G-maps with $n \geq 2$.

Remark 1 *For any $i, j \in \{0, \dots, n\}$ such that $j \geq i+2$, the permutation $\alpha_i \alpha_j$ is an involution. Therefore, in any n -G-map we have:*

$$\forall i \in \{0, \dots, n-2\}, \forall j \in \{i+2, \dots, n\}, \alpha_i \alpha_j = \alpha_j \alpha_i$$

Indeed, if $\alpha_i \alpha_j \alpha_i \alpha_j = 1_{\mathcal{D}}$, then $\alpha_i \alpha_j \alpha_i = \alpha_j^{-1} = \alpha_j$ and $\alpha_i \alpha_j = \alpha_j \alpha_i^{-1} = \alpha_j \alpha_i$.

Definition 2 (n -map [13]) *An n -map ($n \geq 1$) is defined as an $(n+1)$ -tuple $M = (\mathcal{D}, \delta_0, \dots, \delta_{n-1})$ such that:*

- \mathcal{D} is a finite non-empty set of darts;
- $\delta_0, \dots, \delta_{n-2}$ are involutions on \mathcal{D} and δ_{n-1} is a permutation on \mathcal{D} such that
 - $\forall i \in \{0, \dots, n-2\}, \forall j \in \{i+2, \dots, n\}, \delta_i \delta_j$ is an involution.

Definition 3 (Orbit) *Let $\Phi = \{\phi_1, \dots, \phi_k\}$ be a set of permutations on \mathcal{D} (a set of darts). We denote by $\langle \Phi \rangle$ the permutation group generated by*

Φ , i.e. the set of permutations obtained by any composition and inversion of permutations contained in Φ . The orbit of a dart $d \in \mathcal{D}$ relatively to Φ is defined by $\langle \Phi \rangle (d) = \{ \phi(d) \mid \phi \in \langle \Phi \rangle \}$. Furthermore, we extend this notation to the empty set by defining $\langle \emptyset \rangle$ as the identity map.

Definition 4 (Connected component) Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map (resp. $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map). A subset \mathcal{D}' of \mathcal{D} is called a connected component of M (resp. G) if $\mathcal{D}' = \langle \gamma_0, \dots, \gamma_{n-1} \rangle (d)$ (resp. $\mathcal{D}' = \langle \alpha_0, \dots, \alpha_{n-1} \rangle (d)$) for some dart $d \in \mathcal{D}'$.

An n -map may be associated to an n -G-map, as stated by the next definition. In this paper, we use this direct link between the two structures to show that the removal operation we introduce for maps is properly defined (Section 3.1). For that purpose, we notably use the fact that a removal operation (as defined by Damiand and Lienhardt) in a G-map has a counterpart (according to our definition) in its associated map and vice versa.

Definition 5 (Map of the hypervolumes) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map ($n \geq 1$). The n -map $HV = (\mathcal{D}, \delta_0 = \alpha_n \alpha_0, \dots, \delta_{n-1} = \alpha_n \alpha_{n-1})$ is called the map of the hypervolumes of G .

Lienhardt [14] proved that if the n -G-map G is orientable, HV has two connected components. In the following we only consider orientable n -G-maps and assume that an arbitrary component of the map HV is chosen.

Definition 6 (Dual and inverse) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map and let $M = (\mathcal{D}, \delta_0, \dots, \delta_{n-1})$ be an n -map.

- The dual of G , denoted by \overline{G} , is defined by:

$$\overline{G} = (\mathcal{D}, \alpha_n, \alpha_{n-1}, \dots, \alpha_0)$$

- The dual and the inverse of M are respectively defined by:

$$\begin{aligned} \overline{M} &= (\mathcal{D}, \delta_0^{-1}, \delta_0^{-1} \delta_{n-1}, \dots, \delta_0^{-1} \delta_1) \\ &= (\mathcal{D}, \delta_0, \delta_0 \delta_{n-1}, \dots, \delta_0 \delta_1) \\ M^{-1} &= (\mathcal{D}, \delta_0, \dots, \delta_{n-2}, \delta_{n-1}^{-1}) \end{aligned}$$

Note that we also have $M^{-1} = (\mathcal{D}, \delta_0^{-1}, \dots, \delta_{n-2}^{-1}, \delta_{n-1}^{-1})$ since $(\delta_i)_{i \in \{0, \dots, n-2\}}$ is an involution.

Proposition 1 ([10]) If we consider the function HV which maps each n -G-map G to an n -map of the hypervolumes $HV(G)$. We have if $n > 1$:

$$\overline{HV(G)} = HV(\overline{G}).$$

In other words the following diagram commutes:

$$\begin{array}{ccc} G & \xrightarrow{\text{dual}} & \overline{G} \\ HV \downarrow & & \downarrow HV \\ M & \xrightarrow{\text{dual}} & \overline{M} \end{array}$$

Proposition 2 (Associated maps of an n -G-map [13]) *The two n -maps associated to an n -G-map $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ are defined as:*

$$M_1 = HV(G) = (\mathcal{D}, \alpha_n \alpha_0, \alpha_n \alpha_1, \dots, \alpha_n \alpha_{n-1}) \stackrel{\text{not.}}{=} (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$$

$$M_2 = \overline{M_1} = HV(\overline{G}) = (\mathcal{D}, \alpha_0 \alpha_n, \alpha_0 \alpha_{n-1}, \dots, \alpha_0 \alpha_1) \stackrel{\text{not.}}{=} (\mathcal{D}, \overline{\gamma}_0, \dots, \overline{\gamma}_{n-1})$$

Since there is a one-to-one correspondence between i -cells¹ of M_1 and $(n-i)$ -cells of $M_2 = \overline{M_1}$, and since there is a direct link between the subscripts of the permutations of a map and the way cells are defined, it is convenient to denote² $(\mathcal{D}, \overline{\gamma}_0, \dots, \overline{\gamma}_{n-1}) \stackrel{\text{not.}}{=} (\mathcal{D}, \beta_n, \dots, \beta_1)$.

Moreover the permutations $(\beta_i)_{i \in \{1, \dots, n\}}$ and $(\gamma_i)_{i \in \{0, \dots, n-1\}}$ are related by the following relationships ($n \geq 2$):

$$\gamma_0 = \beta_n \text{ and } \forall i \in \{1, \dots, n-1\} \begin{cases} \gamma_i = \beta_n \beta_i \\ \beta_i = \gamma_0 \gamma_i \end{cases}$$

The two maps M_1 and M_2 associated to an n -G-map G are respectively defined as the maps of the hypervolumes of G and \overline{G} . However, whenever the reference to the n -G-map will not be required we will simply consider that we have two dual maps describing a same partition of the space. The equivalence between these two representations is illustrated in Fig. 1 where the 2-G-map G is the triple $(\mathcal{D} = \mathcal{D}_1 \cup \mathcal{D}_2, \gamma_0, \gamma_1)$ with

$$\begin{aligned} \mathcal{D}_1 &= \{1, 2, 3, 4, 5, 6\}, \mathcal{D}_2 = \{-1, -2, -3, -4, -5, -6\} \\ \alpha_0 &= (1, -1)(2, -2)(3, -3)(4, -4)(5, -5)(6, -6) \\ \alpha_1 &= (-1, -2)(2, -1)(3, -4)(4, -3)(5, -6)(6, -5) \\ \alpha_2 &= (1, -6)(2, -3)(3, -2)(4, -5)(5, -4)(6, -1) \end{aligned}$$

¹ Cells are formally defined in subsection 2.2.

² These notation are also used for example in [4].




Fig. 1. A closed 2-G-map G (left). The 2-maps $HV(G)$ and $HV(\overline{G})$ (right).
The 2-map $HV(G)$ is $(\mathcal{D}_1, \gamma_0, \gamma_1)$ where

$$\begin{aligned}\gamma_0 &= (1, 6)(2, 3)(4, 5)(-1, -6)(-2, -3)(-4, -5) \\ \gamma_1 &= (1, 5, 3)(2, 4, 6)(-2, -4, -6)(-1, -5, -3)\end{aligned}$$

The 2-map $HV(\overline{G})$ is $(\mathcal{D}_2, \beta_2, \beta_1)$ where

$$\begin{aligned}\beta_2 &= \gamma_0 \\ \beta_1 &= (1, 2)(3, 4)(5, 6)(-1, -2)(-3, -4)(-5, -6)\end{aligned}$$

2.2 Cells in maps

Definition 7 (*n*-G-maps and cells [13]) *Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map ($n \geq 1$). Let us consider $d \in \mathcal{D}$. The i -cell, or cell of dimension i , which contains d is denoted by $\mathcal{C}_i(d)$ and defined by the orbit:*

$$\mathcal{C}_i(d) = \langle \alpha_0, \dots, \hat{\alpha}_i, \dots, \alpha_n \rangle (d)$$

where $\hat{\alpha}_i$ denotes the absence of the involution α_i .

Definition 8 (*n*-maps and cells [13]) *Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map. The i -cell or cell of dimension i (vertex, edges, ...) of M which owns a*




Fig. 2. A part of a closed 2-G-map G (left) and a part of a connected component of the associated n -map $M_2 = HV(\overline{G})$ (right).

given dart $d \in \mathcal{D}$ is denoted by $\mathcal{C}_i(d)$ and defined by the orbits:

$$\begin{aligned} \forall i \in \{0, \dots, n-1\} \quad \mathcal{C}_i(d) &= \langle \gamma_0, \dots, \hat{\gamma}_i, \dots, \gamma_{n-1} \rangle (d) \\ \text{For } i = n \quad \mathcal{C}_n(d) &= \langle \gamma_0 \gamma_1, \dots, \gamma_0 \gamma_{n-1} \rangle (d) \end{aligned}$$

In both an n -map and an n -G-map, two cells \mathcal{C} and \mathcal{C}' with different dimensions will be called *incident* if $\mathcal{C} \cap \mathcal{C}' \neq \emptyset$.

Proposition 3 ([10]) *The cells of the map $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ may be equivalently expressed according to the permutations of $\overline{M} = (\mathcal{D}, \beta_n, \dots, \beta_1)$ using the following equations:*

$$\begin{aligned} \text{For } i = 0 \quad \mathcal{C}_0(d) &= \langle \beta_1^{-1} \beta_2, \dots, \beta_1^{-1} \beta_n \rangle (d) \\ \forall i \in \{1, \dots, n\} \quad \mathcal{C}_i(d) &= \langle \beta_0, \dots, \hat{\beta}_i, \dots, \beta_n \rangle (d) \end{aligned}$$

Definition 9 (Degree and dual degree of a cell) *Let \mathcal{C} be an i -cell in an n -(G-)map, $0 \leq i \leq n$.*

- The degree of \mathcal{C} is the number of $(i+1)$ -cells incident to \mathcal{C} .
- The dual degree of \mathcal{C} is the number of $(i-1)$ -cells incident to \mathcal{C} .

Property 1 ([10]) *Following Definition 9, the degree of a cell \mathcal{C} in an n -G-map $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ is precisely the number of sets in the set $\Delta = \{ \langle \hat{\alpha}_{i+1} \rangle (d) \mid d \in \mathcal{C} \}$.*

Notation 1 *Let $(\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map. For $d \in \mathcal{D}$, we denote by $\langle \hat{\alpha}_{k_1}, \hat{\alpha}_{k_2}, \dots, \hat{\alpha}_{k_p} \rangle (d)$, where all the involutions are excluded, the orbit $\langle \Phi \rangle (d)$ where $\Phi = \{ \alpha_0, \dots, \alpha_n \} \setminus \{ \alpha_{k_1}, \alpha_{k_2}, \dots, \alpha_{k_p} \}$.*




Fig. 3. Degree and local degree: The cell v_2 has a degree 2 and a local degree 3. Dual degree and dual local degree: The cell e_1 has a dual degree and a dual local degree 2, whereas e_2 has a dual degree 1 and a dual local degree 2.

Definition 10 (Local degree in G-maps) Let \mathcal{C} be an i -cell in an n -G-map.

- For $i \in \{0, \dots, n-1\}$, the local degree of \mathcal{C} is the number

$$|\{\langle \hat{\alpha}_i, \hat{\alpha}_{i+1} \rangle(b) \mid b \in \mathcal{C}\}|$$

- For $i \in \{1, \dots, n\}$, the dual local degree of \mathcal{C} is the number

$$|\{\langle \hat{\alpha}_{i-1}, \hat{\alpha}_i \rangle(b) \mid b \in \mathcal{C}\}|$$

The local degree (resp. dual local degree) of an n -cell (resp. a 0-cell) is 0.

More intuitively, the local degree of an i -cell \mathcal{C} is the number of $i+1$ -cells that locally appear to be incident to \mathcal{C} . It is called *local* because it may be different from the degree since an $i+1$ -cell may be incident more than once to an i -cell, as illustrated in Figure 3 where the 1-cell e_2 is multi-incident to the 0-cell v_2 .

On the other hand, the dual local degree of an i -cell \mathcal{C} is the number of $(i-1)$ -cells that appear to be incident to \mathcal{C} . As in the example given in Figure 3 where the edge e_2 locally appears to be bounded by two vertices, as it is always the case for a 1-cell, whereas the darts involved by the orbits considered in Definition 10 all belong to a unique vertex (v_2).

Property 2 ([10]) The degree of a cell in an n -G-map is greater than zero and less than or equal to its local degree.

Property 3 (Cell with local degree 1 [10]) An i -cell \mathcal{C} in an n -G-map is of local degree 1 if and only if for all $d \in \mathcal{C}$, $d\alpha_{i+1} \in \langle \hat{\alpha}_i, \hat{\alpha}_{i+1} \rangle(d)$.

It is known since [5,7] that cells that may be removed or contracted in a G-map must satisfy a criterion which, although correct, was mistakenly called “having a local degree 2”. In [9,10], the notion of *regularity*, recalled below, was introduced in order to state a new criterion based on the correct definition of the local degree (Definitions 10 and 12).

Definition 11 (Regular cell) An i -cell ($i \leq n - 2$) in an n -G-map is said to be regular if it satisfies the two following conditions:

- a) $\forall d \in \mathcal{C}$, $d\alpha_{i+1}\alpha_{i+2} = d\alpha_{i+2}\alpha_{i+1}$ or $d\alpha_{i+1}\alpha_{i+2} \notin \langle \hat{\alpha}_i, \hat{\alpha}_{i+1} \rangle (d\alpha_{i+2}\alpha_{i+1})$,
and
- b) $\forall b \in \mathcal{C}$, $b\alpha_{i+1} \notin \langle \hat{\alpha}_i, \hat{\alpha}_{i+1} \rangle (b)$

Any $(n - 1)$ -cell is said to be regular too.

The following theorem provides a characterization of cells that may be removed from a G-map.

Theorem 1 ([9]) For any $i \in \{0, \dots, n - 2\}$, an i -cell \mathcal{C} is a regular cell with local degree 2 if and only if

- i) $\exists d \in \mathcal{C}$, $d\alpha_{i+1} \notin \langle \hat{\alpha}_i, \hat{\alpha}_{i+1} \rangle (d)$, and
- ii) $\forall d \in \mathcal{C}$, $d\alpha_{i+1}\alpha_{i+2} = d\alpha_{i+2}\alpha_{i+1}$

Definition 12 (Local degree in maps [9]) Let \mathcal{C} be an i -cell in an n -map.

- The local degree of \mathcal{C} is the number

$$\begin{cases} \left| \langle \hat{\gamma}_i, \hat{\gamma}_{i+1} \rangle (b) \mid b \in \mathcal{C} \right| & \text{if } i \in \{0, \dots, n - 2\} \\ \left| \langle \gamma_0\gamma_1, \dots, \gamma_0\gamma_{n-2} \rangle (b) \mid b \in \mathcal{C} \right| & \text{if } i = n - 1 \end{cases}$$

- The dual local degree of \mathcal{C} is the number

$$\begin{cases} \left| \langle \hat{\gamma}_i, \hat{\gamma}_{i-1} \rangle (b) \mid b \in \mathcal{C} \right| & \text{for } i \in \{1, \dots, n - 1\} \\ \left| \langle \gamma_0\gamma_1, \dots, \gamma_0\gamma_{n-2} \rangle (b) \mid b \in \mathcal{C} \right| & \text{for } i = n \end{cases}$$

The local degree (resp. dual local degree) of an n -cell (resp. a 0-cell) is 0.

Let us justify the orbits considered in the definition of the local degree when $i \in \{0, \dots, n - 2\}$. As for G-maps, we consider the darts that may be reached from d while allowing no change of i -cell (hence $\hat{\gamma}_i$) and no change of $(i + 1)$ -cell (hence $\hat{\gamma}_{i+1}$). In the case when $i = n - 1$, preventing any change of $(n - 1)$ -cell means that the allowed involutions are in $\{\gamma_0, \dots, \gamma_{n-2}\}$, and preventing any change of n -cell means, according to Definition 8, that the set of allowed involutions is $\{\gamma_0\gamma_1, \dots, \gamma_0\gamma_{n-1}\}$. Overall, only the involution $\gamma_0\gamma_{n-1}$ of the latter set is not allowed by the first one, therefore we obtain the orbit that must be considered, i.e. $\langle \gamma_0\gamma_1, \dots, \gamma_0\gamma_{n-2} \rangle$.

Definition 13 (Regular cell in n -maps [9]) An i -cell in an n -map, $0 \leq i < n - 1$ is said to be regular if it satisfies the two conditions a) and b) below:

a) If $i < n - 3$, for all $d \in \mathcal{C}$ we have:

$$d\gamma_{i+1}\gamma_{i+2} = d\gamma_{i+2}\gamma_{i+1} \text{ or } d\gamma_{i+1}\gamma_{i+2} \notin \langle \hat{\gamma}_i, \hat{\gamma}_{i+1} \rangle (d\gamma_{i+2}\gamma_{i+1})$$

If $i = n - 3$, for all $d \in \mathcal{C}$ we have:

$$d\gamma_{n-2}\gamma_{n-1} = d\gamma_{n-1}^{-1}\gamma_{n-2} \text{ and } d\gamma_{n-2}\gamma_{n-1}^{-1} = d\gamma_{n-1}\gamma_{n-2}, \text{ or}$$

$$d\gamma_{n-2} \notin \left(\langle \hat{\gamma}_{n-3}, \hat{\gamma}_{n-2} \rangle (d\gamma_{n-1}^{-1}\gamma_{n-2})\gamma_{n-1}^{-1} \cup \langle \hat{\gamma}_{n-3}, \hat{\gamma}_{n-2} \rangle (d\gamma_{n-1}\gamma_{n-2})\gamma_{n-1} \right)$$

If $i = n - 2$, for all $d \in \mathcal{C}$ we have:

$$d\gamma_{n-1}^{-1} = d\gamma_{n-1}, \text{ or}$$

$$d\gamma_{n-1}^{-1} \notin \langle \gamma_1, \dots, \gamma_{n-3} \rangle (d\gamma_{n-1})$$

b) If $i < n - 2$, for all $b \in \mathcal{C}$ we have:

$$b\gamma_{i+1} \notin \langle \hat{\gamma}_i, \hat{\gamma}_{i+1} \rangle (b)$$

If $i = n - 2$, for all $b \in \mathcal{C}$ we have:

$$\{b\gamma_{n-1}, b\gamma_{n-1}^{-1}\} \cap \langle \gamma_0, \dots, \gamma_{n-3} \rangle (b) = \emptyset$$

Any $(n - 1)$ -cell is said to be regular too.

Notation 2 If \mathcal{D} and \mathcal{D}' are sets, $\sigma : \mathcal{D} \longrightarrow \mathcal{D}'$, and $S \subset \mathcal{D}$, we denote by $S\sigma$ the image of S by σ , i.e.

$$S\sigma = \{ \sigma(s) \mid s \in S \}$$

3 Cells removal and contraction in G-maps and maps

3.1 Cells removal

Notation 3 If $S = \{E_i\}_{0 \leq i \leq N}$ is a set of sets for $N \in \mathbb{N}$, we denote by S^* the union of sets in S , i.e.

$$S^* = \bigcup_{0 \leq i \leq N} E_i$$

Definition 14 (Removal set) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map (resp. $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map) and $S_r = \{R_i\}_{0 \leq i \leq n}$ be sets of i -cells with $R_n = \emptyset$. The family of sets S_r is called a removal set in G (resp. in M). Furthermore, for such a family we will denote $R = \bigcup_{i=0}^n R_i$, the set of all cells of S_r , so that R^* is the set of all darts in S_r .

Definition 15 (Removal kernel) Let G be an n - G -map. A removal kernel $K_r = \{R_i\}_{0 \leq i \leq n}$ in G is a removal set such that all cells of R are disjoint (i.e. $\forall \mathcal{C}, \mathcal{C}' \in R, \mathcal{C} \cap \mathcal{C}' = \emptyset$) and all of them are regular cells with local degree 2 (Definitions 11 and 10). A removal kernel is defined the same way for an n -map M using Definitions 13 and 12 for the notions of regularity and local degree, respectively.

The following definition for cells removal is adapted from [11] where a definition combining removals and contractions is given. The definition below is obtained by considering that no cell is to be contracted.

In her definition, Grasset required that cells of the removal kernel should have a “local degree two” according to her definition. However, this definition is both too restrictive to be a valid definition for the local degree 2, and it also does not exclude cells with local degree 1.

Excluding cells with local degree 1 from Grasset’s definition yields the notion of *regular cells* with local degree 2 (Definition 11). Both notions being equivalent, as shown by Theorem 1. Therefore, the definition we present below as a note, although not exactly the one given by Grasset, is just more restrictive in the sense that no cell with a local degree one should be in the removal kernel.

Note 1 The operation of cells removal in n - G -maps is defined as follows in [11]: Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $K_r = \{R_i\}_{0 \leq i \leq n-1}$ a removal kernel in G . Let $BV_i = R_i^* \alpha_i \setminus R_i^*, \forall i, 0 \leq i \leq n$. The set BV_i is called the set of surviving darts which are neighbors of an i -cell to be removed. The n - G -map resulting of the removal of the cells of R is $G' = (\mathcal{D}', \alpha'_0, \dots, \alpha'_n)$ defined by:

- (1) $\mathcal{D}' = \mathcal{D} \setminus R^*$;
- (2) $\forall i, 0 \leq i \leq n, \forall b \in \mathcal{D}' \setminus BV_i, b\alpha'_i = b\alpha_i$;
- (3) $\forall i, 0 \leq i < n, \forall b \in BV_i, b\alpha'_i = b' = b(\alpha_i \alpha_{i+1})^k \alpha_i$ where k is the smallest integer such that $b' \in BV_i$.

In this report, we use the following definition for the removal of cells, definition which is slightly simpler and proved to be equivalent to the one used in [11].

Definition 16 (Cells removal in n - G -maps [11,9]) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $K_r = \{R_i\}_{0 \leq i \leq n-1}$ be a removal kernel in G . The n - G -map resulting of the removal of the cells of R is $G' = (\mathcal{D}', \alpha'_0, \dots, \alpha'_n)$ where:

- (1) $\mathcal{D}' = \mathcal{D} \setminus R^*$;
- (2) $\forall d \in \mathcal{D}', d\alpha'_n = d\alpha_n$;
- (3) $\forall i, 0 \leq i < n, \forall d \in \mathcal{D}', d\alpha'_i = d' = d(\alpha_i \alpha_{i+1})^k \alpha_i$ where k is the smallest integer such that $d' \in \mathcal{D}'$.

Notation 4 The n - G -map obtained after removal of a kernel $K_r = \{R_i\}_{0 \leq i \leq n}$

from an n -G-map G will be denoted either by $G \setminus K_r$, or by $G \setminus R^*$.

As stated by the next proposition, the involution α_n remains unchanged after the removal operation.

Proposition 4 *Let G, G' be n -G-maps and $K_r = \{R_i\}_{0 \leq i \leq n}$ be a removal kernel as in Definition 16. Since $R_n = \emptyset$, then $d\alpha_n \in \mathcal{D}'$ for all $d \in \mathcal{D}'$.*

Remark 2 *Let G, G' be n -G-maps and $K_r = \{R_i\}_{0 \leq i \leq n}$ be a removal kernel as in Definition 16. If a dart d belongs to an i -cell \mathcal{C} of R_i , then $d\alpha_n \in \mathcal{C}$. Indeed, since there are no n -cell in K_r , $i < n$ so that $d\alpha_n \in \langle \hat{\alpha}_i \rangle(d) = \mathcal{C}$.*

The equivalence between Definition 10 of [11] (see Note 1) and Definition 16 is stated by the following proposition.

Proposition 5 ([9]) *Definition 16 and the one given in Note 1 are equivalent.*

The following definition for the operation of simultaneous removal of cells in an n -map was given in [9] (see also [10]).

Definition 17 (Cells removal in n -maps) *Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map and $S_r = \{R_i\}_{0 \leq i \leq n-1}$ a removal kernel in M . The n -map $M \setminus S_r = (\mathcal{D}', \gamma'_0, \dots, \gamma'_{n-1})$ obtained after removal of the cells of S_r is defined by:*

- $\mathcal{D}' = \mathcal{D} \setminus R^*$;
- $\forall i \in \{0, \dots, n-2\}, \forall d \in \mathcal{D}', d\gamma'_i = d(\gamma_i \gamma_{i+1}^{-1})^k \gamma_i$, where k is the smallest integer such that $d(\gamma_i \gamma_{i+1}^{-1})^k \gamma_i \in \mathcal{D}'$.
- For $i = n-1, \forall d \in \mathcal{D}', d\gamma'_{n-1} = d\gamma_{n-1}^{k+1}$ where k is the smallest integer such that $d\gamma_{n-1}^{k+1} \in \mathcal{D}'$.

It was proved ([9, Theorem 2], see also [10]) that the such defined $(n-1)$ -tuple $M \setminus S_r$ is actually an n -map, this by establishing the link between the removal operation in n -maps and the same operation in n -G-maps. This link required the following definition for the removal kernel in the map of the hypervolumes of an n -G-map, associated with a removal kernel in the latter G-map.

Definition 18 *Let G be an n -G-map, $S_r = \{R_i\}_{0 \leq i \leq n}$ be a removal set in G and $M = HV(G)$. We define the set $HV(S_r) = \{R'_i\}_{0 \leq i \leq n}$ as follows:*

- $\forall i \in \{0, \dots, n-1\}, R'_i = \{\langle \alpha_n \alpha_0, \dots, \alpha_n \hat{\alpha}_i, \dots, \alpha_n \alpha_{n-1} \rangle(d) \mid d \in R_i^*\}$
- $R'_n = \{\langle \alpha_0 \alpha_1, \dots, \alpha_0 \alpha_{n-1} \rangle(d) \mid \exists \mathcal{C} \in R_n, d \in \mathcal{C}\}$

3.2 Cells contraction

Definition 19 (Contraction kernel) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $K_c = \{C_i\}_{0 \leq i \leq n}$ be sets of i -cells with $C_0 = \emptyset$. Let $C = \cup_{i=0}^n C_i$. Furthermore, we suppose that the cells of C are disjoint (i.e. $\forall c, c' \in C, c \cap c' = \emptyset$), have a dual local degree 2, and are regular cells in \overline{G} . The family of sets K_c is then called a contraction kernel in G .

We also denote:

$$C_i^* = \bigcup_{c \in C_i} c \text{ and } C^* = \bigcup_{i \in \{0, \dots, n\}} C_i^*$$

A contraction kernel is defined in a similar way for an n -map M .

Remark 3 If G is an n - G -map, then from the very definition of cells (Definition 7) an i -cell in G is an $(n - i)$ -cell in \overline{G} .

The following definition is adapted from [11] where a definition combining removals and contractions was given. The definition below is obtained by considering that no cell is to be removed.

Note 2 (Definition of cells contraction by [11]) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $K_c = \{C_i\}_{1 \leq i \leq n}$ a contraction kernel in G . Let $BV_i = C_i^* \alpha_i \setminus C_i^*$, $\forall i, 0 \leq i \leq n$. The set BV_i is called the set of surviving darts which are neighbors of an i -cell to be contracted. The n - G -map resulting of the contraction of the cells of C is $G' = (\mathcal{D}', \alpha'_0, \dots, \alpha'_n)$ defined by:

- (1) $\mathcal{D}' = \mathcal{D} \setminus C^*$;
- (2) $\forall i, 0 \leq i \leq n, \forall b \in \mathcal{D}' \setminus BV_i, b\alpha'_i = b\alpha_i$;
- (3) $\forall i, 0 < i \leq n, \forall b \in BV_i, b\alpha'_i = b' = b(\alpha_i \alpha_{i-1})^k \alpha_i$ where k is the smallest integer such that $b' \in BV_i$.

In [9], the contraction operation in G -maps is defined as a removal operation in the dual map (Definition 20 below). The equivalence between this definition and the one (Note 2) given by Grasset in [11] or by Damiand and Lienhardt in [7] will be stated by Proposition 7.

Definition 20 (Cells contraction in n - G -maps [9]) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $K_c = \{C_i\}_{1 \leq i \leq n}$ be a contraction kernel. The n - G -map resulting of the contraction of the cells of K_c is $G' = \overline{\overline{G} \setminus K_c}$.

Notation 5 The n - G -map obtained after the contraction of a kernel $K_c = \{C_i\}_{0 \leq i \leq n}$ from an n - G -map G will be denoted either by G/K_c , or by G/C^* .

Proposition 6 ([9]) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $K_c =$

$\{C_i\}_{1 \leq i \leq n}$ be a contraction kernel. The n - G -map resulting of the contraction of the cells of C according to Definition 20 is $G' = (\mathcal{D}', \alpha'_0, \dots, \alpha'_n)$ defined by:

- (1) $\mathcal{D}' = \mathcal{D} \setminus C$;
- (2) $\forall d \in \mathcal{D}', d\alpha'_0 = d\alpha_0$;
- (3) $\forall i, 0 < i \leq n, \forall d \in \mathcal{D}', d\alpha'_i = d' = d(\alpha_i\alpha_{i-1})^k\alpha_i$ where k is the smallest integer such that $d' \in \mathcal{D}'$.

Proposition 7 ([9]) *Definition 20 and the one given by Grasset (Note 2, see also [11]) are equivalent.*

Definition 21 (Cells contraction in n -maps) *Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map and let $K_c = \{C_i\}_{1 \leq i \leq n}$ be a contraction kernel. The n -map resulting of the contraction of the cells of K_c , which we denote M/K_c is the n -map $\overline{M} \setminus K_c$.*

Proposition 8 ([9]) *Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map. Let $K_c = \{C_i\}_{1 \leq i \leq n}$ be a contraction kernel. The n -map obtained after contraction of the cells of K_c , $M' = (\mathcal{D}', \gamma'_0, \dots, \gamma'_{n-1})$ is defined by:*

- $\mathcal{D}' = \mathcal{D} \setminus C$;
- $\forall d \in \mathcal{D}', d\gamma'_0 = d\gamma_{n-1}^k\gamma_0$ where k is the smallest integer such that $d\gamma_{n-1}^k\gamma_0 \in \mathcal{D}'$;
- $\forall i \in \{1, \dots, n-1\}, \forall d \in \mathcal{D}', d\gamma'_i = d\gamma_{n-1}^k(\gamma_i\gamma_{i-1}^{-1})^{k'}\gamma_i$, where k is the smallest integer such that $d\gamma_{n-1}^k \in \mathcal{D}'$ and k' is the smallest integer such that $d\gamma_{n-1}^k(\gamma_i\gamma_{i-1}^{-1})^{k'}\gamma_i \in \mathcal{D}'$.

4 Connecting walks

The permutations or involutions which define the map resulting from a removal operation are obtained by somehow following a path in the original map until a surviving dart has been found (Definitions 16 and 17). This leads to the notion of the so called *connecting walks* which we define here and whose main properties are described.

Notation 6 *If $S = (d_1, d_2, \dots, d_p)$ and $S' = (b_1, b_2, \dots, b_q)$, $p, q \in \mathbb{N}$, are sequences of darts in a (G -)map then we denote:*

- $S^\times = (d_2, \dots, d_p)$, i.e. S without its first dart,
- $S^\kappa = (d_1, \dots, d_{p-1})$, i.e. S without its last dart,
- $\text{reverse}(S) = (d_p, d_{p-1}, \dots, d_1)$, and
- $S \cdot S' = (d_1, \dots, d_p, b_1, \dots, b_q)$.

We also denote by S^* the set $\{d_1, d_2, \dots, d_p\}$ and by $\text{last}(S)$ the last dart d_p

of S .

4.1 In generalized maps

4.1.1 Definition and properties

Definition 22 (Connecting walk in n -G-maps) Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map and $K_r = \{R_i\}_{0 \leq i \leq n}$ be a removal kernel in G . Let $G' = G \setminus K_r = (\mathcal{D}', \alpha'_0, \dots, \alpha'_n)$. The i -connecting walk associated to a dart $d \in \mathcal{D}'$ for $i \in \{0, \dots, n\}$, denoted by $\text{CW}_{G,G'}^i(d)$, is the sequence of darts of \mathcal{D} defined by:

$$\text{CW}_{G,G'}^i(d) = (d_0, d_1, \dots, d_p)$$

where

- $d_0 = d$,
- $\forall u, 0 \leq u \leq p, d_u = d(\alpha_i \alpha_{i+1})^u$,
- $p = \text{Min}\{k \in \mathbb{N} \mid d_k \alpha_i \in \mathcal{D}'\}$.

The above definition is clearly linked to the one of the removal operation (Definition 16). To make this link explicit, we first prove the following property which states that darts of an i -connecting walk are, except for the first one, darts of i -cells that have been removed. This property as well as the next one is illustrated by Figure 4(b), in the 2D case for the ease of visualization.

Property 4 With the notations of Definition 22, for all $d \in \mathcal{D}'$ such that $\text{CW}_{G,G'}^i(d) = (d_0, d_1, \dots, d_p)$ we have:

$$\forall k \in \{1, \dots, p\}, d_{k-1} \alpha_i \in R_i^* \text{ and } d_k \in R_i^*$$

Proof: We prove the property by recurrence on k . From the definition of connecting walks, we know that if $p \geq 1$ we have $d_0 \alpha_i \notin \mathcal{D}'$ (i.e. $d_0 \alpha_i \in R_i^*$). Suppose that $d_0 \alpha_i \in R_j$ for $j \neq i$, in other words that $d_0 \alpha_i \in \langle \hat{\alpha}_j \rangle (d')$ for some $d' \in R_j^*$. Since $i \neq j$ we have $d_0 = d_0 \alpha_i \alpha_i \in \langle \hat{\alpha}_j \rangle (d') \in R_j$, which contradicts the fact that $d_0 \in \mathcal{D}'$; therefore $d_0 \alpha_i \in R_i^*$. Since $d_0 \alpha_i$ and $d_0 \alpha_i \alpha_{i+1}$ belong to the same i -cell, we also obtain $d_1 = d_0 \alpha_i \alpha_{i+1} \in R_i$. Eventually, the property is true for $k = 1$.

Now, suppose that $d_k \in R_i^*$ for $k < p$. Since $k < p$ and p is the smallest integer such that $d_p \alpha_i \in \mathcal{D}'$, then $d_k \alpha_i \in R_i^*$. If we suppose that $d_k \alpha_i \in R_j$ with $j \neq i$, since $d_k \alpha_i$ and $d_k \alpha_i \alpha_i = d_k$ belong to the same j -cell, we would obtain that d_k belongs to both an i -cell and a j -cell of the removal kernel K_r , which is not allowed by Definition 15. Therefore, we have $d_k \alpha_i = d_{(k+1)-1} \alpha_i \in R_i$. Again,

since $d_k\alpha_i$ and $d_k\alpha_i\alpha_{i+1}$ belong to the same i -cell we also have $d_k\alpha_i\alpha_{i+1} = d_{k+1} \in R_i$. Thus, the property is true for $k + 1$. \square




Fig. 4. A 2-G-map G (top row) from which the two white vertices are to be removed, yielding a map G' (bottom row). The connecting walks $CW_{G,G'}^0(b) = (b = b_0, b_1, b_2)$ (second row) and $CW_{G,G'}^0(d) = (d = d_0, d_1, d_2)$ (third row).

In [11], Grasset defines connecting walks in G-maps in a slightly different way than what is done by Definition 22. A first difference is that in Grasset's definition, d does not appear at the beginning of the sequence that defines $CW_{G',G'}^i(d)$, whereas the dart $d_p\alpha_i$ of Definition 22 is added at the end of the sequence. On the other hand, consecutive darts in a connecting walk as defined by Grasset are linked by alternately either an α_i or an α_{i+1} involution when they are always linked by the permutation $\alpha_i\alpha_{i+1}$ in our definition. Thus, our connecting walk for a given dart and dimension counts $((k - 1)/2) + 1$ darts when the corresponding one with Grasset's definition has k ones.

Following the definition of [11], connecting walks that are distinct (up to reverse ordering and after removal of their last dart) are always disjoint [11, Proposition 22]. With our definition the property simply becomes that connecting walks are either equal or disjoint. In other words, a removed dart belongs to at most one connecting walk for some $i \in \{0, \dots, n\}$. This property, stated by the following proposition, induces a father-child relationship between darts of consecutive levels similar to the *reduction windows* in the context of regular pyramids.

Proposition 9 *Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n -G-map, K_r be a removal kernel in G , and d be a dart of R_i^* for $0 \leq i < n$. The dart d belongs to at most one connecting walk. In other words, the two following properties hold:*

- i) $d \in \bigcup_{b \in \mathcal{D}'} CW_{G,G \setminus K_r}^i(b)^* \Rightarrow \exists! b \in \mathcal{D}', d \in CW_{G,G \setminus K_r}^i(b)^{*\ast}$
- ii) $\forall j \in \{0, \dots, n\} \setminus \{i\}, \forall b \in \mathcal{D}', d \notin CW_{G,G \setminus K_r}^j(b)^{*\ast}$

Proof: First note that Figure 5(b) shows an example of a removal kernel for which some darts are contained in no connecting walk.




Fig. 5. (a) Removal of a 0-cells: Each dart of the removed vertex belongs to exactly one 0-connecting walk and no i -connecting walk for $i \in \{1, 2\}$. (b) Removal of two edges from a 2-G-map: darts of the vertex v belong to no connecting walk. (c) Four regular edges with local degree 2 (dashed circles) which may be removed from a 2-G-map. All the removed darts belong to exactly one 1-connecting walk and to no i -connecting walk for $i \in \{0, 2\}$.

From Property 4, $\text{CW}_{G,G'}^j(d)^{\times*} \subset R_j^*$ for all $j \in \{0, \dots, n\} \setminus \{i\}$. Since all cells of a removal kernel are disjoint (Definition 15) and $d \in R_i^*$, d may only belong to some i -connecting walks, hence property *ii*).

Now, let us suppose that there exists two darts a and b such that:

- $\text{CW}_{G,G'}^i(a) = (a_0 = a, a_1, \dots, a_p)$ and $d = a_k$ for $0 < k \leq p$;
- $\text{CW}_{G,G'}^i(b) = (b_0 = b, b_1, \dots, b_q)$ and $d = b_h$ for $0 < h \leq q$.

Thus, by Definition 22 we have $d = a(\alpha_i \alpha_{i+1})^k$ and $d = b(\alpha_i \alpha_{i+1})^h$. If $a \neq b$, it is clear that $k \neq h$ and we may suppose without loss of generality that $h < k$. Then, from

$$a(\alpha_i \alpha_{i+1})^k = b(\alpha_i \alpha_{i+1})^h$$

we deduce that

$$a(\alpha_i \alpha_{i+1})^{k-h} = b$$

Note that $0 < k - h < k$, so by Property 4 we have $a(\alpha_i \alpha_{i+1})^{k-h} \in R_i^*$. This contradicts the fact that $b \in \mathcal{D}'$. It follows that $a = b$. \square

The following property establishes a one-to-one correspondence between connecting walks, as any i -connecting walk, associated with a dart $d \in \mathcal{D}'$, may

be built from the connecting walk associated with $d\alpha'_i$ (with the notations of Definition 16). This is illustrated on Figure 4. In fact, the above mentioned correspondence coincides with the application of an involution; it is therefore itself an involution on the set of connecting walks.

Property 5 *Let G be an n - G -map and K_r be a removal kernel in G . Let $G' = G \setminus K_r = (\mathcal{D}', \alpha'_0, \dots, \alpha'_n)$. For $i \in \{0, \dots, n-1\}$, $d \in \mathcal{D}'$, and $b = d\alpha'_i$; if $\text{CW}_{G,G'}^i(d) = (d_0 = d, d_1, \dots, d_p)$ we have:*

$$\text{CW}_{G,G'}^i(b) = (b_0 = b, b_1, \dots, b_p) \text{ where } b_k = d_{p-k}\alpha_i \text{ for } 0 \leq k \leq p$$

Proof: From the definition of connecting walks, we have:

$$\text{CW}_{G,G'}^i(d) = (d, d(\alpha_i\alpha_{i+1}), d(\alpha_i\alpha_{i+1})^2, \dots, d(\alpha_i\alpha_{i+1})^p)$$

where $p = \text{Min}\{h \in \mathbb{N} \mid d(\alpha_i\alpha_{i+1})^h\alpha_i \in \mathcal{D}'\}$.

With $b = d\alpha'_i$, we have:

$$\text{CW}_{G,G'}^i(b) = (b, b(\alpha_i\alpha_{i+1}), b(\alpha_i\alpha_{i+1})^2, \dots, b(\alpha_i\alpha_{i+1})^q)$$

where $q = \text{Min}\{h \in \mathbb{N} \mid b(\alpha_i\alpha_{i+1})^h\alpha_i \in \mathcal{D}'\}$.

If $p = 0$, we have $\text{CW}_{G,G'}^i(d) = (d)$ and $b = d\alpha_i$ so we immediately obtain $b\alpha_i = d$, therefore $\text{CW}_{G,G'}^i(b) = (b)$. Thus, we suppose in the sequel that $p > 0$.

Let us prove by recurrence on k that for $0 \leq k \leq p$, we have:

$$b(\alpha_i\alpha_{i+1})^k\alpha_i = d(\alpha_i\alpha_{i+1})^{p-k} \quad (1)$$

From Definition 22, we have $d_p\alpha_i \in \mathcal{D}'$ and (by Property 4) for all $k \in \{0, \dots, p-1\}$, $d_k = d(\alpha_i\alpha_{i+1})^k \notin \mathcal{D}'$. It follows from the very definition of α'_i (Item (3) of Definition 16) that $d_p\alpha_i = d\alpha'_i$, i.e. $b = d_p\alpha_i$. Since $d_p = d(\alpha_i\alpha_{i+1})^p$ we obtain $b = d(\alpha_i\alpha_{i+1})^p\alpha_i$ which is (1) for $k = 0$.

Now, suppose that (1) holds for $k < p$, we have:

$$\begin{aligned} b(\alpha_i\alpha_{i+1})^k\alpha_i &= d(\alpha_i\alpha_{i+1})^{p-k} \\ b(\alpha_i\alpha_{i+1})^k\alpha_i\alpha_{i+1}\alpha_i &= d(\alpha_i\alpha_{i+1})^{p-k}\alpha_{i+1}\alpha_i \\ b(\alpha_i\alpha_{i+1})^{k+1}\alpha_i &= d(\alpha_i\alpha_{i+1})^{p-(k+1)} \end{aligned}$$

The latter equality is precisely (1) for $k+1$.

Since $p > 0$, from Property 4 we know that for all $k \in \{0, \dots, p-1\}$, $d(\alpha_i \alpha_{i+1})^{p-k} \notin \mathcal{D}'$ which by (1) means that $b(\alpha_i \alpha_{i+1})^k \alpha_i \notin \mathcal{D}'$. Furthermore, and again from (1) with $k = p$, we obtain $d = b(\alpha_i \alpha_{i+1})^p \alpha_i$ so that $b(\alpha_i \alpha_{i+1})^p \alpha_i \in \mathcal{D}'$ (since $d \in \mathcal{D}'$). It follows that p is the smallest integer such that $b(\alpha_i \alpha_{i+1})^p \alpha_i \in \mathcal{D}'$. This shows that $q = p$ and by (1) that $b_k = d_{p-k} \alpha_i$ for all $k \in \{0, \dots, p\}$. \square

4.1.2 Additional properties

In Section 5, which is dedicated to the notion of a pyramid of G-maps or maps, we will use the following notion of a

Since Property 9 does not guarantee that a dart always belong to a connecting walk, all darts that have been removed may not be traversed by following all the connecting walks. Hence, we define the notion of a *simple removal kernel* which allows us to state

Definition 23 *A removal kernel K_r is called simple if*

$$\forall i \in \{0, \dots, n-1\}, \forall d \in R_i, \exists s \in \mathcal{D}' \mid d \in \text{CW}_{G,G'}^i(s)^{\times*}$$

(Note that by Proposition 9 the dart s is necessarily unique.)

We derive the following property after the suppression of a simple removal kernel from an n -G-map.

Property 6 *If G is an n -G-map and K_r is a simple removal kernel in G , then we have:*

$$\mathcal{D} = \mathcal{D}' \sqcup \left[\bigsqcup_{d \in \mathcal{D}', 0 \leq i \leq n-1} \text{CW}_{G,G'}^i(d)^{\times*} \right] \quad (2)$$

where \sqcup denotes the union of disjoint sets.

Proof: As a consequence of Proposition 9, $\text{CW}_{G,G'}^i(d_1)^{\times*} \cap \text{CW}_{G,G'}^j(d_2)^{\times*} = \emptyset$ when either $d_1 \neq d_2$ or $i \neq j$. Thus, the union within the brackets in (2) is indeed a union of disjoint sets; and the leftmost \sqcup comes from the fact that $\text{CW}_{G,G'}^i(d)^{\times*} \subset R = \mathcal{D} \setminus \mathcal{D}'$.

The \supseteq part is immediate, so we only need to prove the \subseteq part of equality (2), and we denote by U the set on the right side of the equality. Let $b \in \mathcal{D}$. If $b \in \mathcal{D}'$ we immediately obtain $b \in U$, so we suppose that $b \notin \mathcal{D}'$. In this case, there exists $i \in \{0, \dots, n-1\}$ such that $b \in R_i$. The removal kernel being simple, following Definition 23 there exists $d \in \mathcal{D}'$ such that $b \in \text{CW}_{G,G'}^i(d)^{\times*}$, and again $b \in U$. \square

When Property 6 applies, the traversal of the connecting walks of all the darts of \mathcal{D}' is guaranteed to visit all darts of \mathcal{D} . In a pyramid, this means that a level may be rebuilt with no hole from the level above it; in other words there is no loss of information when reducing a map using a simple kernel.

Note that the removed vertex in Figure 5(a) constitutes a simple removal kernel, whereas the one made of the two edges in Figure 5(b) does not. As a preliminary remark, we may observe that the edge resulting from the removal of vertex v in Figure 5(b) is a simple removal kernel by itself, thus that the removal of the two edges depicted in this figure may be achieved after the suppression of two simple removal kernels. In fact, not all removal kernel may be decomposed into simple ones. However, using Proposition 10 (below) some removal operations may be delayed in order to obtain a simple kernel between two specified levels. Indeed, simple removal kernels may be characterized, in a computationally more efficient way, using Proposition 10. The proof of this proposition will use the following Lemma.

Lemma 1 *Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map, K_r be a removal kernel in G , and \mathcal{D}' be the set of darts of $G \setminus K_r$. Let $i \in \{0, \dots, n-1\}$. If a sequence of darts $S = (d_0, d_1, \dots, d_k)$, $k \in \mathbb{N}^*$, satisfies*

- i) $d_0 \in \mathcal{D}'$,*
- ii) $\forall h \in \{0, \dots, k\}$, $d_h = d_0(\alpha_i \alpha_{i+1})^h$,*
- iii) $\forall h \in \{1, \dots, k\}$, $d_h \notin \mathcal{D}'$,*
- iv) $d_k \in R_i$.*

then, the two following properties are also satisfied:

- (a) $\forall h \in \{1, \dots, k\}$ we have $d_h \in R_i$,*
- (b) $\forall h \in \{0, \dots, k-1\}$ we have $d_h \alpha_i \in R_i$.*

Proof: We prove this lemma by recurrence on the “length” k of the sequence (d_0, \dots, d_k) . For $k = 1$ the sequence is $(d_0, d_1 = d_0 \alpha_i \alpha_{i+1})$. Item *iv)* is $d_1 \in R_i$, hence (a). Now, we note that d_1 and $d_1 \alpha_{i+1}$ belong to the the same i -cell of G . Since $d_1 \alpha_{i+1} = d_0 \alpha_i \alpha_{i+1} \alpha_{i+1} = d_0 \alpha_i$, we thus deduce that $d_0 \alpha_i$ also belongs to R_i , hence (b). Therefore, the property is valid for any sequence (d_0, d_1) , i.e. for $k = 1$.

Now, we assume that the lemma is valid for a sequence (d_0, \dots, d_k) , for a given $k > 0$, and we consider a sequence S' which satisfies items *i)* to *iv)*.

$$S' = (d_0, d_1, \dots, d_k, d_{k+1})$$

Since $d_{k+1} \in R_i$, and because d_{k+1} and $d_{k+1} \alpha_{i+1}$ belong to the same i -cell of G , we obtain that $d_{k+1} \alpha_{i+1} = d_k \alpha_i \alpha_{i+1} \alpha_{i+1} = d_k \alpha_i$ also belongs to R_i . This

makes (b) valid in S' for $h = k + 1$. Now, by item *iii*), we have $d_k \in R$. Let us suppose that $d_k \in R_j$ with $j \neq i$. In this case, d_k and $d_k\alpha_i$ belong to the same j -cell of R_j . However, $d_k\alpha_i$ cannot belong to both an i -cell of R_i , which has been proved to be true, and a j -cell of R_j (Definition 15 of a removal kernel). It follows that $d_k \in R_i$ which is (a) for $h = k$.

Now, if $d_k \in R_i$ we obtain that the subsequence $S'' = (d_0, \dots, d_k)$ of S' is a sequence which satisfies all the conditions of the lemma. Indeed, beside item *iv*) which is precisely $d_k \in R_i$, items *i*) to *iii*) are true for S'' , as a subsequence of S' . By the recurrence hypothesis, the lemma may be applied to S'' and we obtain:

- $\forall h \in \{1, \dots, k\}$ we have $d_h \in R_i$,
- $\forall h \in \{0, \dots, k - 1\}$ we have $d_h\alpha_i \in R_i$.

Since we have $d_{k+1} \in R_i$ and we also proved that $d_k\alpha_i \in R_i$, we deduce that properties (a) and (b) are both valid for the the sequence S' , and the lemma holds with $k + 1$. \square

Proposition 10 *A removal kernel $K_r = \{R_i\}_{i=0, \dots, n}$ in an n -G-map G is simple if and only if:*

$$\forall i \in \{0, \dots, n - 1\}, \forall d \in R_i^*, \langle \alpha_i\alpha_{i+1} \rangle(d) \cap \mathcal{D}' \neq \emptyset \quad (3)$$

where \mathcal{D}' is the set of darts of $G \setminus K_r$.

Proof: (\Leftarrow) Since $\alpha_i\alpha_{i+1}$ is a permutation, for all $d \in R_i^*$ there exists $p \in \mathbb{N}$ such that

$$\langle \alpha_i\alpha_{i+1} \rangle(d) = \{d, d(\alpha_i\alpha_{i+1}), d(\alpha_i\alpha_{i+1})^2, \dots, d(\alpha_i\alpha_{i+1})^p\}$$

with $d(\alpha_i\alpha_{i+1})^{p+1} = d$.

By (3), $\langle \alpha_i\alpha_{i+1} \rangle(d)$ contains at least one dart of \mathcal{D}' , so $p > 0$ (since $d \notin \mathcal{D}'$) and $d(\alpha_i\alpha_{i+1})^h \in \mathcal{D}'$ for at least one $h \in \{1, \dots, p\}$. Let k be the greatest such integer h , and let us denote $b_k = d(\alpha_i\alpha_{i+1})^k$. Thus we may consider the sequence:

$$\left(d(\alpha_i\alpha_{i+1})^k, d(\alpha_i\alpha_{i+1})^{k+1}, \dots, d(\alpha_i\alpha_{i+1})^{k+(p-k)}, d(\alpha_i\alpha_{i+1})^{k+(p-k)+1} \right)$$

where

- $b_k = d(\alpha_i\alpha_{i+1})^k \in \mathcal{D}'$,
- $\left\{ d(\alpha_i\alpha_{i+1})^{k+1}, \dots, d(\alpha_i\alpha_{i+1})^{k+(p-k)}, d(\alpha_i\alpha_{i+1})^{k+(p-k)+1} = d \right\} \subset R_i^*$,
- $d(\alpha_i\alpha_{i+1})^{k+(p-k)+1} = d \in R_i^*$ (by hypothesis).

By Lemma 1, we deduce that

- (a) $d(\alpha_i \alpha_{i+1})^{k+h+1} \in R_i^*$ for all $h \in \{0, \dots, (p-k)\}$, and
- (b) $d(\alpha_i \alpha_{i+1})^{k+l} \alpha_i \in R_i^*$, for all $l \in \{0, \dots, (p-k) - 1\}$.

From (b) and the very definition of the connecting walk $CW_{G,G'}^i(b_k)$, it follows that $d(\alpha_i \alpha_{i+1})^{k+h+1} \in CW_{G,G'}^i(b_k)^*$ for all $h \in \{0, \dots, (p-k)\}$. In particular, $d(\alpha_i \alpha_{i+1})^{k+(p-k)+1} = d \in CW_{G,G'}^i(b_k)^*$.

(\Rightarrow) Suppose that the removal kernel K_r is simple, and let d be a dart of R_i^* for $i \in \{0, \dots, n-1\}$. From Definition 23, there exists $s \in \mathcal{D}'$ such that $d \in CW_{G,G'}^i(s)^{x*}$. It follows that $d = s(\alpha_i \alpha_{i+1})^k$ for $k \in \mathbb{N}^*$ \square

4.2 In combinatorial maps

Definition 24 (Connecting walk in n -maps) Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map and $K_r = \{R_i\}_{0 \leq i \leq n}$ be a removal kernel in M . Let $M = M \setminus K_r = (\mathcal{D}', \gamma'_0, \dots, \gamma'_{n-1})$. The i -connecting walk associated to a dart $d \in \mathcal{D}'$ for $i \in \{0, \dots, n-1\}$, denoted by $CW_{M,M'}^i(d)$, is the sequence of darts of \mathcal{D} defined by

$$CW_{M,M'}^i(d) = (d_0, d_1, \dots, d_p) \text{ with } p = \text{Min}\{k \in \mathbb{N} \mid d_k \gamma_i \in \mathcal{D}'\}$$

where

- For $i \in \{0, \dots, n-2\}$, $\forall u, 0 \leq u \leq p$, $d_u = d(\gamma_i \gamma_{i+1}^{-1})^u$
- For $i = n-1$, $\forall u, 0 \leq u \leq p$, $d_u = d\gamma_{n-1}^u$


Property 7 With the notations of Definition 24, for all $d \in \mathcal{D}'$ such that $CW_{M,M'}^i(d) = (d_0, d_1, \dots, d_p)$ we have:

$$\forall k \in \{1, \dots, p\}, d_{k-1} \gamma_i \in R_i^* \text{ and } d_k \in R_i^*$$

Proof:

• (If $i < n-1$) We prove the property by recurrence on k . From the definition of connecting walks, we know that if $p \geq 1$ we have $d_0 \gamma_i \notin \mathcal{D}'$ (i.e. $d_0 \gamma_i \in R^*$). Suppose that $d_0 \gamma_i \in R_j$ for $j \neq i$, in other words that $d_0 \gamma_i \in \langle \hat{\gamma}_j \rangle(d')$ for some $d' \in R_j^*$. Since $i \neq j$ we have $d_0 = d_0 \gamma_i \gamma_i \in \langle \hat{\gamma}_j \rangle(d') \in R_j$, which contradicts the fact that $d_0 \in \mathcal{D}'$; therefore $d_0 \gamma_i \in R_i^*$. Since $d_0 \gamma_i$ and $d_0 \gamma_i \gamma_{i+1}^{-1}$ belong to the same i -cell, we also obtain $d_1 = d_0 \gamma_i \gamma_{i+1}^{-1} \in R_i$. Eventually, the property is true for $k = 1$.

Now, suppose that $d_k \in R_i^*$ for $k < p$. Since $k < p$ and p is the smallest integer such that $d_p \gamma_i \in \mathcal{D}'$, then $d_k \gamma_i \in R^*$. If we suppose that $d_k \gamma_i \in R_j$ with $j \neq i$,



(a) Removal of two 1-cells in a map. The two darts of vertex v do not belong to any connecting walk.

Fig. 6. Removed darts which do not belong to any connecting walk.

since $d_k \gamma_i$ and $d_k \gamma_i \gamma_i = d_k$ belong to the same j -cell, we would obtain that d_k belongs to both an i -cell and a j -cell of the removal kernel K_r , which is not allowed by Definition 15. Therefore, we have $d_k \gamma_i = d_{(k+1)-1} \gamma_i \in R_i$. Again, since $d_k \gamma_i$ and $d_k \gamma_i \gamma_{i+1}$ belong to the same i -cell we also have $d_k \gamma_i \gamma_{i+1} = d_{k+1} \in R_i^*$. Thus, the property is true for $k + 1$.

- (If $i = n - 1$) We prove the property by recurrence on k . From the definition of connecting walks, we know that if $p \geq 1$ we have $d_0 \gamma_{n-1} \notin \mathcal{D}'$ (i.e. $d_0 \gamma_{n-1} \in R^*$). Suppose that $d_0 \gamma_{n-1} \in R_j$ for $j \neq n - 1$, in other words that $d_0 \gamma_{n-1} \in \langle \hat{\gamma}_j \rangle (d')$ for some $d' \in R_j^*$. Since $j \neq n - 1$ we have $d_0 = d_0 \gamma_{n-1} \gamma_{n-1}^{-1} \in \langle \hat{\gamma}_j \rangle (d') \in R_j$, which contradicts the fact that $d_0 \in \mathcal{D}'$; therefore $d_1 = d_0 \gamma_{n-1} \in R_{n-1}^*$. Eventually, the property is true for $k = 1$.

Now, suppose that $d_k \in R_{n-1}^*$ for $k < p$. Since $k < p$ and p is the smallest integer such that $d_p \gamma_{n-1} \in \mathcal{D}'$, then $d_k \gamma_{n-1} \in R^*$. If we suppose that $d_k \gamma_{n-1} \in R_j$ with $j \neq n - 1$, since $d_k \gamma_{n-1}$ and $d_k \gamma_{n-1} \gamma_{n-1}^{-1} = d_k$ belong to the same j -cell, we would obtain that d_k belongs to both an $(n - 1)$ -cell and a j -cell of the removal kernel K_r , which is not allowed by Definition 15. Therefore, we have $d_k \gamma_{n-1} = d_{(k+1)-1} \gamma_{n-1} \in R_{n-1}$. Thus, the property is true for $k + 1$. \square

As for G-map, connecting walks within maps also provide a father-child relationship, as stated by the following proposition:

Proposition 11 *Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map, K_r be a removal kernel in M , and d be a dart of R_i^* for $0 \leq i < n$. The dart d belongs to at most one connecting walk. In other words, the two following properties hold:*

- i) $d \in \bigcup_{b \in \mathcal{D}'} \text{CW}_{M, M \setminus K_r}^i(b)^* \Rightarrow \exists! b \in \mathcal{D}', d \in \text{CW}_{M, M \setminus K_r}^i(b)^{**}$
- ii) $\forall j \in \{0, \dots, n\} \setminus \{i\}, \forall b \in \mathcal{D}', d \notin \text{CW}_{M, M \setminus K_r}^j(b)^{**}$

Where $\bigcup_{b \in \mathcal{D}'} \text{CW}_{M, M \setminus K_r}^i(b)^*$ represents the set of darts belonging to at least one connecting walk.

Proof: First note that Figure 6(a) shows an example of a removal kernel for which some darts are contained in no connecting walk.

From Property 7, $CW_{M,M'}^j(d)^{\times*} \subset R_j^*$ for all $j \in \{0, \dots, n\} \setminus \{i\}$. Since all cells of a removal kernel are disjoint (Definition 15) and $d \in R_i^*$, d may only belong to some i -connecting walks, hence property *ii*).

Now, let us suppose that there exists two darts a and b , with $a \neq b$, such that:

- $CW_{M,M'}^i(a) = (a_0 = a, a_1, \dots, a_p)$ and $d = a_k$ for $0 < k \leq p$;
- $CW_{M,M'}^i(b) = (b_0 = b, b_1, \dots, b_q)$ and $d = b_h$ for $0 < h \leq q$.

– If $i < n-1$, by Definition 24 we have $d = a(\gamma_i \gamma_{i+1}^{-1})^k$ and $d = b(\gamma_i \gamma_{i+1}^{-1})^h$. Since $a \neq b$, if $a(\gamma_i \gamma_{i+1}^{-1})^k = b(\gamma_i \gamma_{i+1}^{-1})^h$ we necessarily have $k \neq h$ (the composition of permutations being a one-to-one map). We may then suppose without loss of generality that $h < k$. Then, from

$$a(\gamma_i \gamma_{i+1}^{-1})^k = b(\gamma_i \gamma_{i+1}^{-1})^h$$

we deduce that

$$a(\gamma_i \gamma_{i+1}^{-1})^{k-h} = b$$

Note that $0 < k-h < k$, so by Property 7 applied to $CW_{M,M'}(a)$ we have $a(\gamma_i \gamma_{i+1}^{-1})^{k-h} \in R_i^*$. This contradicts the fact that $b \in \mathcal{D}'$. It follows that $a = b$.

– If $i = n-1$, by Definition 24 we have $d = a\gamma_{n-1}^k$ and $d = b\gamma_{n-1}^h$. Since $a \neq b$, if $a\gamma_{n-1}^k = b\gamma_{n-1}^h$ we necessarily have $k \neq h$. Thus, we may suppose without loss of generality that $h < k$. Then, from $a\gamma_{n-1}^k = b\gamma_{n-1}^h$ we deduce that $a(\gamma_{n-1})^{k-h} = b$. Since $0 < k-h < k$, by Property 7 applied to $CW_{M,M'}(a)$ we obtain $a\gamma_{n-1}^{k-h} \in R_{n-1}^*$. This contradicts the fact that $b \in \mathcal{D}'$. It follows that $a = b$. \square

Using Property 4, it is clear from Definition 16 that we also have the following property, which relates i -connecting walks to the corresponding involution α'_i in the resulting map.

Property 8 *Let $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ be an n -map, K_r be a removal kernel in M , $M' = M \setminus K_r = (\mathcal{D}', \gamma'_0, \dots, \gamma'_{n-1})$ and $d \in \mathcal{D}'$. For all $i \in \{0, \dots, n\}$ we have:*

$$d\gamma'_i = \text{last}(CW_{M, M \setminus K_r}^i(d))\gamma_i$$

Proof: This result is a direct consequence of the definition of a connecting walk (Definition 24), considering Property 7. \square

As for G-maps, the following property establishes a one-to-one correspondence between connecting walks, as any i -connecting walk ($i < n-1$) associated with

a dart $d \in \mathcal{D}'$, may be built from the connecting walk associated with $d\gamma'_i$ (with the notations of Definition 17).

Lemma 2 *If $M = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$ is an n -map and d is a dart of a regular $(n-2)$ -cell \mathcal{C} with local degree 2, then $d\gamma_{n-1} = d\gamma_{n-1}^{-1}$.*

Proof: Since \mathcal{C} is a regular cell, from Definition 13 we have:

$$d\gamma_{n-1}^{-1} = d\gamma_{n-1} \text{ or } d\gamma_{n-1}^{-1} \notin \langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d\gamma_{n-1})$$

Let us suppose that $d\gamma_{n-1}^{-1} \notin \langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d\gamma_{n-1})$, which implies that $\langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d\gamma_{n-1}) \cap \langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1}^{-1} \rangle (d\gamma_{n-1}) = \emptyset$. By property *b*) of Definition 13, we also have $\{d\gamma_{n-1}, d\gamma_{n-1}^{-1}\} \cap \langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d) = \emptyset$. It follows that the three sets $\langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d)$, $\langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d\gamma_{n-1})$, and $\langle \hat{\gamma}_{n-2}, \hat{\gamma}_{n-1} \rangle (d\gamma_{n-1}^{-1})$ are disjoint sets. Since $d\gamma_{n-1}$ and $d\gamma_{n-1}^{-1}$ both belong to \mathcal{C} (as $n-1 \neq n-2$), we conclude that

$$\left| \left\{ \langle \hat{\gamma}_{n-1}, \hat{\gamma}_{n-1} \rangle (b) \mid b \in \mathcal{C} \right\} \right| \geq 3$$

which contradicts the fact that \mathcal{C} has local degree 2 (Definition 12).

Therefore we necessarily have $d\gamma_{n-1}^{-1} = d\gamma_{n-1}$. \square

As we claimed in our introduction, generalized maps do not allow to manipulate easily notions related with the orientation over the underlying quasimanifold, when the latter is orientable. This is due, in part, to the fact that in this case a G-map, by using twice as many darts as really needed, actually encodes the two possible orientations at the same time. A connecting walk in a G-map G , as defined in this paper, uses a fixed orientation by skipping darts. Indeed, all darts of the walk thus belong to a single connected component of the map of the hypervolumes $HV(G)$ associated to the G-map G (Definition 5). It is therefore consistent with respect to the orientation property since each component of $HV(G)$ corresponds to one orientation of G . These remarks are based on Proposition 13, for which Proposition 12 is an intermediary result.

Proposition 12 *With the notations of Definition 24, for all $d \in \mathcal{D}'$ and $i \in \{0, \dots, n-2\}$, if $CW_{M, M'}^i(d) = (d_0, d_1, \dots, d_p)$ we have:*

$$\forall r \in \{0, \dots, p\}, d_r = d_0(\gamma_i \gamma_{i+1})^r$$

Proof: If $i < n-2$, γ_{i+1} is an involution so that $d\gamma_{i+1} = d\gamma_{i+1}^{-1}$ for all $b \in \mathcal{D}$. The proposition is then straightforward following Definition 24.

If $i = n-2$, for all $r \in \{0, \dots, p-1\}$ in Definition 24, by Property 7 we have $d_r \gamma_{n-2} \in R_{n-2}^*$. Therefore, from the very definition of a removal kernel $d_r \gamma_{n-2}$

belongs to a regular $(n - 2)$ -cell with local degree 2 so that, by Lemma 2, $d_r(\gamma_{n-2}\gamma_{n-1}^{-1}) = d_r(\gamma_{n-2}\gamma_{n-1})$. \square

As shown by the the next property also based on Proposition 12, an involution may also be defined on the set of i -connecting walks in a map, for $i < n - 1$.

Property 9 *Let M be an n -map and K_r be a removal kernel in M . Let $M' = M \setminus K_r = (\mathcal{D}', \gamma'_0, \dots, \gamma'_{n-1})$. For $i \in \{0, \dots, n - 2\}$, $d \in \mathcal{D}'$, and $b = d\gamma'_i$; if $\text{CW}_{M,M'}^i(d) = (d_0 = d, d_1, \dots, d_p)$ we have:*

$$\text{CW}_{M,M'}^i(b) = (b_0 = b, b_1, \dots, b_p) \text{ where } b_k = d_{p-k}\gamma_i \text{ for } 0 \leq k \leq p$$

Proof: From the definition of connecting walks for $i < n - 1$, we have:

$$\text{CW}_{M,M'}^i(d) = (d, d(\gamma_i\alpha_{i+1}^{-1}), d(\gamma_i\gamma_{i+1}^{-1})^2, \dots, d(\gamma_i\gamma_{i+1}^{-1})^p)$$

where $p = \text{Min}\{h \in \mathbb{N} \mid d(\gamma_i\gamma_{i+1}^{-1})^h\gamma_i \in \mathcal{D}'\}$.

With $b = d\gamma'_i$, we have:

$$\text{CW}_{M,M'}^i(b) = (b, b(\gamma_i\gamma_{i+1}^{-1}), b(\gamma_i\gamma_{i+1}^{-1})^2, \dots, b(\gamma_i\gamma_{i+1}^{-1})^q)$$

where $q = \text{Min}\{h \in \mathbb{N} \mid b(\gamma_i\gamma_{i+1}^{-1})^h\gamma_i \in \mathcal{D}'\}$.

If $p = 0$, we have $\text{CW}_{M,M'}^i(d) = (d)$ and $b = d\gamma_i$ so we immediately obtain $b\gamma_i = d$, therefore $\text{CW}_{M,M'}^i(b) = (b)$. Thus, we suppose in the sequel that $p > 0$.

Let us prove by recurrence on k that for $0 \leq k \leq p$, we have:

$$b(\gamma_i\gamma_{i+1})^k\gamma_i = d(\gamma_i\gamma_{i+1})^{p-k} \quad (4)$$

From Definition 24, we have $d_p\gamma_i \in \mathcal{D}'$ and (by Property 7) for all $k \in \{0, \dots, p - 1\}$, $d_k = d(\gamma_i\gamma_{i+1}^{-1})^k \notin \mathcal{D}'$. It follows from the very definition of γ'_i (Definition 17) that $d_p\gamma_i = d\gamma'_i$, i.e. $b = d_p\gamma_i$. Since $d_p = d(\gamma_i\gamma_{i+1}^{-1})^p$ we obtain $b = d(\gamma_i\gamma_{i+1}^{-1})^p\gamma_i$ which is (4) for $k = 0$.

Now, suppose that (4) holds for $k < p$, we have:

$$\begin{aligned} b(\gamma_i\gamma_{i+1}^{-1})^k\gamma_i &= d(\gamma_i\gamma_{i+1}^{-1})^{p-k} \\ b(\gamma_i\gamma_{i+1})^k\gamma_i &= d(\gamma_i\gamma_{i+1}^{-1})^{p-k} \quad (\text{by Proposition 12}) \\ b(\gamma_i\gamma_{i+1})^k\gamma_i\gamma_{i+1}\gamma_i &= d(\gamma_i\gamma_{i+1}^{-1})^{p-k}\gamma_{i+1}\gamma_i \\ b(\gamma_i\gamma_{i+1})^{k+1}\gamma_i &= d(\gamma_i\gamma_{i+1}^{-1})^{p-(k+1)} \quad (i < n - 1 \Rightarrow \gamma_i^2 = 1_{\mathcal{D}}) \end{aligned}$$

The latter equality is precisely (4) for $k + 1$.

Since $p > 0$, from Property 7 we know that for all $k \in \{0, \dots, p - 1\}$, $d(\gamma_i \gamma_{i+1}^{-1})^{p-k} \notin \mathcal{D}'$ which by (4) means that $b(\gamma_i \gamma_{i+1}^{-1})^k \gamma_i \notin \mathcal{D}'$. Furthermore, and again from (4) with $k = p$, we obtain $d = b(\gamma_i \gamma_{i+1}^{-1})^p \gamma_i$ so that $b(\gamma_i \gamma_{i+1}^{-1})^p \gamma_i \in \mathcal{D}'$ (since $d \in \mathcal{D}'$). It follows that p is the smallest integer such that $b(\gamma_i \gamma_{i+1}^{-1})^p \gamma_i \in \mathcal{D}'$. This shows that $q = p$ and by (4) that $b_k = d_{p-k} \gamma_i$ for all $k \in \{0, \dots, p\}$. \square

Proposition 13 *Let $G = (\mathcal{D}, \alpha_0, \dots, \alpha_n)$ be an n - G -map and $M = HV(G)$ be its n -map of the hypervolumes. Let K_r be a removal kernel in G , let $G' = G \setminus K_r$ and $M' = M \setminus HV(K_r) = (\mathcal{D}', \gamma'_0, \dots, \gamma'_{n-1})$. The i -connecting walks of d respectively in G and M (with respect to K_r and $HV(K_r)$) satisfy*

$$\forall d \in \mathcal{D}, \forall i \in \{0, \dots, n - 2\}, \text{CW}_{G,G'}^i(d) = \text{CW}_{M,M'}^i(d)$$

Furthermore, we have:

$$\forall d \in \mathcal{D}, \text{CW}_{G,G'}^{(n-1)}(d)^\times = \text{reverse}(\text{CW}_{M,M'}^{(n-1)}(d\gamma'_{n-1})^\times)$$

A first part of the proof of this proposition is achieved through the two following lemmas.

Lemma 3 *With the notation of Proposition 13 we have:*

$$\forall d \in \mathcal{D}', \text{CW}_{G,G'}^{n-2}(d) = \text{CW}_{M,M'}^{n-2}(d)$$

Proof: Let $d \in \mathcal{D}'$ and $HV(K_r) = \{R'_i\}$ following Definition 18; we have:

$$\text{CW}_{M,M'}^{n-2}(d) = (d, d(\gamma_{n-2} \gamma_{n-1}^{-1}), \dots, d(\gamma_{n-2} \gamma_{n-2}^{-1})^p)$$

where $d(\gamma_{n-2} \gamma_{n-1}^{-1})^p \gamma_{n-2} \in \mathcal{D}'$, and for all $k \in \{1, \dots, p\}$, $d(\gamma_{n-2} \gamma_{n-1}^{-1})^{k-1} \gamma_{n-2} \in R'_{n-2}$ and $d(\gamma_{n-2} \gamma_{n-1}^{-1})^k \in R'_{n-2}$ (Property 7). In the sequel, we denote $d_0 = d$ and $d_k = d(\gamma_{n-2} \gamma_{n-1}^{-1})^k$.

By Proposition 12, we have $\text{CW}_{M,M'}^{n-2}(d) = (d, d(\gamma_{n-2} \gamma_{n-1}), \dots, d(\gamma_{n-2} \gamma_{n-1})^p)$ and from the definition of the map of the hypervolumes M we may write:

$$\begin{aligned} \text{CW}_{M,M'}^{n-2}(d) &= (d, d(\alpha_n \alpha_{n-2} \alpha_n \alpha_{n-1}), \dots, d(\alpha_n \alpha_{n-2} \alpha_n \alpha_{n-1})^p) \\ &= (d, d(\alpha_{n-2} \alpha_n \alpha_n \alpha_{n-1}), \dots, d(\alpha_{n-2} \alpha_n \alpha_n \alpha_{n-1})^p) \\ &= (d, d(\alpha_{n-2} \alpha_{n-1}), \dots, d(\alpha_{n-2} \alpha_{n-1})^p) \end{aligned}$$

In other words, we have $d_k = d(\alpha_{n-2} \alpha_{n-1})^k = d_{k-1}(\alpha_{n-2} \alpha_{n-1})$ for $k \in \{1, \dots, p\}$.

From Definition 18 and for all $k \in \{1, \dots, p\}$, if $d_{k-1}\gamma_{n-2} \in R_{n-2}^*$ there exists $b \in R_{n-2}^*$ such that $d_{k-1}\gamma_{n-2} \in \langle \alpha_n\alpha_0, \dots, \alpha_n\hat{\alpha}_{n-2}, \alpha_n\alpha_{n-1} \rangle (b)$. Since $\langle \alpha_n\alpha_0, \dots, \alpha_n\hat{\alpha}_{n-2}, \alpha_n\alpha_{n-1} \rangle (b) \subset \langle \alpha_{n-2} \rangle (b)$, we deduce that the darts $d_{k-1}\gamma_{n-2}$ and b belong to the same $(n-2)$ -cell, thus $d_{k-1}\gamma_{n-2} \in R_{n-2}^*$. We may prove in a similar way that $d_k \in R_{n-2}^*$. Eventually, we have:

$$\forall k \in \{1, \dots, p\}, d_{k-1}\gamma_{n-2} \in R_{n-2}^* \text{ and } d_k \in R_{n-2}^* \quad (5)$$

So far, we have proved that $d_k = d(\alpha_{n-2}\alpha_{n-1})^k \in R_{n-2}^*$ for all $k \in \{1, \dots, p\}$ and $d_k(\alpha_n\alpha_{n-2}) \in R_{n-2}^*$ for all $k \in \{0, \dots, p-1\}$. We now prove that $d_k\alpha_{n-2} \in R_{n-2}^*$ for all $k \in \{0, \dots, p-1\}$. Indeed, from the conditions on the involutions of a G-map, we have $d_k\alpha_n\alpha_{n-2} = d_k\alpha_{n-2}\alpha_n$. Since $d_k\alpha_{n-2}\alpha_n$ and $d_k\alpha_{n-2}\alpha_n\alpha_n = d_k\alpha_{n-2}$ belong to the same $(n-2)$ -cell, from $d_k(\alpha_n\alpha_{n-2}) \in R_{n-2}^*$ we deduce that $d_k\alpha_{n-2} \in R_{n-2}^*$.

We also have $d_p\gamma_{n-2} = d_p\alpha_n\alpha_{n-2} = d_p\alpha_{n-2}\alpha_n \in \mathcal{D}'$. For all $i \in \{0, \dots, n-1\}$, $d_p\alpha_{n-2}\alpha_n$ and $d_p\alpha_{n-2}$ belong to the same i -cell so that $d_p\alpha_{n-2}$ may not belong to an i -cell to be removed. Since $R_n = \emptyset$ it follows that $d_p\alpha_{n-2} \in \mathcal{D}'$.

Eventually, we have proved that $(d = d_0, d_1, \dots, d_p)$ is the connecting walk associated to the dart d in G according to Definition 22 since $d_k\alpha_{n-2} \in R_{n-2}^*$ for all $k \in \{0, \dots, p-1\}$ and $d_p\alpha_{n-2} \in \mathcal{D}'$. \square

Lemma 4 *With the notations of Proposition 13 we have:*

$$\text{CW}_{G,G'}^{(n-1)}(d)^\times = \text{reverse}(\text{CW}_{M,M'}^{(n-1)}(d\gamma_{n-1}'^{-1})^\times)$$

Proof: Following Definition 22 and the one of the map of the hypervolumes M we have:

$$\text{CW}_{G,G'}^{(n-1)}(d) = (d, d\alpha_{n-1}\alpha_n, \dots, d(\alpha_{n-1}\alpha_n)^p) \quad (6)$$

where p is the smallest integer such that $d(\alpha_{n-1}\alpha_n)^p\alpha_{n-1} \in \mathcal{D}'$. Following Remark 2 we deduce that $d(\alpha_{n-1}\alpha_n)^p\alpha_{n-1}\alpha_n = d(\alpha_{n-1}\alpha_n)^{p+1} = d(\gamma_{n-1}^{-1})^{p+1} \in \mathcal{D}'$. Furthermore, from the definition of connecting walks in G-maps we have $d(\alpha_{n-1}\alpha_n)^k\alpha_{n-1} \notin \mathcal{D}'$, for $k < p$ and from Property 4 we also know that $d(\alpha_{n-1}\alpha_n)^k = d\gamma_{n-1}^{-k} \notin \mathcal{D}'$, for $k \in \{1, \dots, p\}$. It follows from the definition of cells removal in n -maps (Definition 17) that $d(\alpha_{n-1}\alpha_n)^{p+1} = d(\gamma_{n-1}^{-1})^{p+1} = d\gamma_{n-1}'^{-1}$.

Now, for $h \in \{1, \dots, p\}$, we obtain:

$$d\gamma_{n-1}'^{-1}(\alpha_n\alpha_{n-1})^h = d(\alpha_{n-1}\alpha_n)^{p+1}(\alpha_n\alpha_{n-1})^h = d(\alpha_{n-1}\alpha_n)^{p+1-h}$$

where $0 < p+1-h \leq p$, so that $d\gamma_{n-1}'^{-1}(\alpha_n\alpha_{n-1})^h \notin \mathcal{D}'$ ((6) and Property 4). On the other hand, $d\gamma_{n-1}'^{-1}(\alpha_n\alpha_{n-1})^{p+1} = d \in \mathcal{D}'$, therefore from the definition of $\text{CW}_{M,M'}^{n-1}(d\gamma_{n-1}'^{-1})$ we have:

$$\begin{aligned}
\text{CW}_{M,M'}^{n-1}(d\gamma'_{n-1}^{-1}) &= (d\gamma'_{n-1}^{-1}, d\gamma'_{n-1}^{-1}(\alpha_n\alpha_{n-1}), \dots, d\gamma'_{n-1}^{-1}(\alpha_n\alpha_{n-1})^p) & (7) \\
&= (d\gamma'_{n-1}^{-1}, d(\alpha_{n-1}\alpha_n)^p, \dots, d(\alpha_{n-1}\alpha_n)) & (8)
\end{aligned}$$

The result comes from (6) and (8). \square

Proof of Proposition 13:

If $i < n - 2$:

$$\begin{aligned}
\text{CW}_{M,M'}^i(d) &= (d, d(\gamma_i\gamma_{i+1}^{-1}), \dots, d(\gamma_i\gamma_{i+1}^{-1})^p) \\
&= (d, d(\alpha_n\alpha_i\alpha_{i+1}\alpha_n), \dots, d(\alpha_n\alpha_i\alpha_{i+1}\alpha_n)^p) \\
&= (d, d(\alpha_i\alpha_n\alpha_n\alpha_{i+1}), \dots, d(\alpha_i\alpha_n\alpha_n\alpha_{i+1})^p) \\
&= (d, d(\alpha_i\alpha_{i+1}), \dots, d(\alpha_i\alpha_{i+1})^p)
\end{aligned}$$

The cases $i = n - 2$ and $i = n - 1$ are respectively given by Lemma 3 and Lemma 4. \square

5 Pyramids of maps

In this section we define pyramids of combinatorial n -maps and introduce the connecting dart sequences which will be used to derive a concise encoding of pyramids.

5.1 Definition

Definition 25 (Pyramid of n -maps) *A pyramid of n -maps with height h is an h -tuple (M_0, K_1, \dots, K_h) where M_0 is an n -map and K_l , $l \in \{1, \dots, h\}$ is a removal kernel for the map M_{l-1} , which is defined by $M_l = M_{l-1} \setminus K_l$ for $l \in \{1, \dots, h\}$.*

Notation 7 *When dealing with a pyramid of n -maps (M_0, K_1, \dots, K_h) , $h \in \mathbb{N}^*$, we usually denote $M_l = (\mathcal{D}_l, \gamma_{l,0}, \dots, \gamma_{l,n-1})$, $0 \leq l \leq h$, and when no confusion may arise we simply refer to a permutation of M_l as $\gamma_{l,i}$ for $i \in \{0, \dots, n\}$ without mentioning the map M_l . We also shorten $\gamma_{0,i}$ as γ_i for all $i \in \{0, \dots, n-1\}$. Eventually, we denote $K_l = \{R_{l,i}\}_{i=1, \dots, n}$.*

6 Connecting dart sequences

We may now give the definition of a connected dart sequence which makes the link, as shown by two propositions given further on, between any two levels of a pyramid the same way a connecting walk does between two consecutive levels.

6.1 Definition and properties

Definition 26 (Connecting dart sequence) *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps and d be a dart of \mathcal{D}_l for $l \in \{0, \dots, h\}$. If $CW_{M_{l-1}, M_l}^i(d) = (d = d_0, \dots, d_p)$ for $i \in \{0, \dots, n-1\}$, we define the i -connecting dart sequence associated to d at level l , denoted by $CDS_l^i(d)$, as follows:*

- For $l = 0$, $CDS_0^i(d) = (d)$, and
- for $l \in \{1, \dots, h\}$
 - If $i \leq n - 2$, $CDS_l^i(d) = GL_{l-1}^i(d_0) \cdot GL_{l-1}^i(d_1) \cdot \dots \cdot GL_{l-1}^i(d_p)$
where:

$$\begin{cases} \forall r \in \{0, \dots, p-1\}, GL_{l-1}^i(d_r) = CDS_{l-1}^i(d_r) \cdot CDS_{l-1}^{i+1}(d_r \gamma_{l-1, i}) \\ GL_{l-1}^i(d_p) = CDS_{l-1}^i(d_p) \end{cases}$$

- If $i = n - 1$, $CDS_l^{n-1}(d) = CDS_{l-1}^{n-1}(d_0) \cdot CDS_{l-1}^{n-1}(d_1) \cdot \dots \cdot CDS_{l-1}^{n-1}(d_p)$.

Notation 8 *Given a pyramid of n -maps (M_0, K_1, \dots, K_h) , $h \in \mathbb{N}$, we may define a set of removed darts at level k , $1 \leq k \leq h$, denoted $K_{0\dots k}^*$, as follows:*

$$K_{0\dots k}^* = \bigsqcup_{l=0, \dots, k} \bigsqcup_{R \in K_l} R^*$$

The above unions are disjoint because of the very definition of the removal kernels K_l , $1 \leq l \leq h$ and the fact that cells of a removal kernel are disjoint.

Note that we trivially have $CDS_1^i(d) = CW_{M_0, M_1}(d)$ for any $d \in D_1$. Connecting dart sequences correspond thus to a natural extension of connecting walks.

The such defined connecting dart sequences also induce a father-child relationship between darts of any two levels in a pyramid, as the transitive closure of the one provided by connecting walks for consecutive levels.

One may obviously not expect the darts of a such defined connecting dart sequence to belong to removed cells of a single dimension, as it is the case for

connecting walks (Propositions 4 and 7). For example, darts of the connecting dart sequence $\text{CDS}_2^0(b)$ in Figure 7 belong to both 1-cells and 0-cells which have been removed from M_0 and M_1 , respectively. Still, the first dart of a connecting dart sequence at level l is the only dart belonging to \mathcal{D}_l . Indeed, we have the following proposition.




Fig. 7. A 3D combinatorial pyramid (M_0, K_1, K_2) . (a) The 3-map M_0 . (b) The 3-map M_1 obtained after removing the edges e_1 and e_2 from M_0 . (c) The map M_2 obtained after removing the vertices v_1 , v_2 , and v_3 from M_1 . Four involutions γ_0 are materialized by two dotted lines. (d) The connecting walk $\text{CW}^1_{M_0, M_1}(b')$ (black darts). (e) The connecting dart sequence $\text{CDS}_2^0(b)$ (black darts).

Proposition 14 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps and $l \in \{1, \dots, h\}$. For all dart $d \in \mathcal{D}_l$ and $i \in \{0, \dots, n-1\}$ we have $\text{CDS}_l^i(d)^{\times*} \cap \mathcal{D}_l = \emptyset$.*

Proof: We prove the property by recurrence on the level l . For $l = 1$, it is readily seen that we have $\text{CDS}_1^i(d) = \text{CW}^i_{M_0, M_1}(d)$ and the result holds by Property 7.

Thus, we suppose that the property is valid for $l \in \{1, \dots, h-1\}$.

– If $i \leq n-2$. Following Definition 26:

$$\begin{aligned} \text{CDS}_{l+1}^i(d) &= \text{CDS}_l^i(d_0) \cdot \text{CDS}_l^{i+1}(d_0 \gamma_{l,i}) \cdot \dots \\ &\quad \dots \cdot \text{CDS}_l^i(d_{p-1}) \cdot \text{CDS}_l^{i+1}(d_{p-1} \gamma_{l,i}) \cdot \text{CDS}_l^i(d_p) \end{aligned}$$

where $(d_0, \dots, d_p) = \text{CW}^i_{M_l, M_{l+1}}(d)$.

From the recurrence hypothesis, $\text{CDS}_l^{i+1}(d_r \gamma_{l,i})^{\times*} \cap \mathcal{D}_l = \emptyset$ for all $r \in \{0, \dots, p-1\}$, and $\text{CDS}_l^i(d_r)^{\times*} \cap \mathcal{D}_l = \emptyset$, so that $\text{CDS}_l^i(d_r) \cap \mathcal{D}_l = d_r$, for all $r \in \{1, \dots, p\}$. From Property 7, d_0 is the only dart of $\text{CW}^i_{M_l, M_{l+1}}(d)$ that belongs to \mathcal{D}_{l+1} . Since it is also the first dart of $\text{CDS}_l^i(d_0)$, hence the first dart of $\text{CDS}_{l+1}^i(d)$, we eventually obtain $\text{CDS}_{l+1}^i(d)^{\times*} \cap \mathcal{D}_l = \emptyset$.

– If $i = n - 1$.

$$\text{CDS}_{l+1}^{n-1}(d) = \text{CDS}_l^{n-1}(d_0) \cdot \dots \cdot \text{CDS}_l^{n-1}(d_p)$$

where $(d_0, \dots, d_p) = \text{CW}_{M_l, M_{l+1}}^{n-1}(d)$. The result is again immediate after application of the recurrence hypothesis to the sequences $\text{CDS}_l^{n-1}(d_r)$, $0 \leq r \leq p$, and using the fact that $\text{CW}_{M_l, M_{l+1}}^{n-1}(d)^\times \cap \mathcal{D}_l = \emptyset$ (Property 7). \square

Lemma 5 *With the notation of Definition 26, for all $l \in \{0, \dots, h\}$ and all $d \in \mathcal{D}_l$ we have $\text{first}(\text{CDS}_l^i(d)) = d$. In particular, $\text{CDS}_l^i(d)$ contains at least one dart: d itself.*

Proof: We proceed by recurrence on l . First, if $l = 0$ we have by Definition 26 $\text{CDS}_0^i(d) = (d)$ and the property is verified.

Thus we suppose that the result holds for $l \in \{0, \dots, h-1\}$. At level $l+1$, by Definition 26 we have:

- If $i < n - 1$,

$$\begin{aligned} \text{CDS}_{l+1}^i(d) &= \text{CDS}_l^i(d_0) \cdot \text{CDS}_l^{i+1}(d_0 \gamma_{l,i}) \cdot \dots \\ &\quad \dots \cdot \text{CDS}_l^i(d_{p-1}) \cdot \text{CDS}_l^{i+1}(d_{p-1} \gamma_{l,i}) \cdot \text{CDS}_l^i(d_p) \end{aligned}$$

- If $i = n - 1$, $\text{CDS}_{l+1}^{n-1}(d) = \text{CDS}_{l-1}^{n-1}(d_0) \cdot \text{CDS}_{l-1}^{n-1}(d_1) \cdot \dots \cdot \text{CDS}_{l-1}^{n-1}(d_p)$

where $(d_0, \dots, d_p) = \text{CW}_{M_l, M_{l+1}}^i(d)$.

In both cases, and by the recurrence hypothesis, $\text{CDS}_l^i(d_0)$ is not an empty sequence and we have $\text{first}(\text{CDS}_l^i(d_0)) = d_0$. From the very definition of the connecting walk $\text{CW}_{M_l, M_{l+1}}^i(d)$ we have $d_0 = d$. It follows immediately from the above equalities that $\text{first}(\text{CDS}_{l+1}^i(d)) = d$, hence the property is verified at level $l+1$. \square

Lemma 6 *With the notations of Definition 26, a connecting dart sequence CDS_l^i always counts an odd number of darts when $i < n - 1$.*

Proof: We prove the result by recurrence on l for $l \in \{0, \dots, h\}$. The property is true for $n = 0$ since $\text{CDS}_0^i(d) = (d)$. Thus we may assume that the property is valid for some $l \in \{0, \dots, h-1\}$.

Now, following Definition 26, if $\text{CW}_{M_l, M_{l+1}}^i(d) = (d_0, d_1, \dots, d_p)$ we have

$$\text{CDS}_{l+1}^i(d) = \text{GL}_l^i(d_0) \cdot \text{GL}_l^i(d_1) \cdot \dots \cdot \text{GL}_l^i(d_p)$$

If $p = 0$, the property holds immediately by the recurrence hypothesis. Thus, we now consider the case when $p > 0$.

For $h \in \{0, \dots, p-1\}$ we have $\text{GL}_l^i(d_h) = \text{CDS}_l^i(d_h) \cdot \text{CDS}_l^{i+1}(d_h \gamma_{l,i})$. From the recurrence hypothesis applied to both $\text{CDS}_l^i(d_h)$ and $\text{CDS}_l^{i+1}(d_h \gamma_{l,i})$, we deduce that the sequence $\text{GL}_l^i(d_h)$ counts an even number of darts. It follows that the sequence $\text{GL}_l^i(d_0) \cdot \dots \cdot \text{GL}_l^i(d_{p-1})$ also counts an even number of darts.

Eventually, $\text{GL}_l^i(d_p) = \text{CDS}_l^i(d_p)$, which, from the recurrence hypothesis counts an odd number of darts, hence the result. \square

Connecting dart sequences also share with connecting walks the property that the last dart of an i -connecting dart sequence associated with a dart d at level l is linked with the dart $d\gamma_{l,i}$ by the permutation γ_i .

Proposition 15 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$, with the notations of Definition 26. Let $d \in \mathcal{D}_l$ for $l \in \{1, \dots, h\}$. We have*

$$\text{last}(\text{CDS}_l^i(d))\gamma_{0,i} = d\gamma_{l,i}$$

Proof: We prove the proposition by recurrence on l . If $l = 0$, from Definition 26 we have $\text{CDS}_0^i(d) = (d)$, hence the property is obviously valid. Thus, we suppose that the property is satisfied for $l \in \{0, \dots, h-1\}$.

Let us consider $\text{CDS}_{l+1}^i(d)$ for some dart d of \mathcal{D}_l .

$$\begin{aligned} \text{CDS}_{l+1}^i(d) &= \text{GL}_l^i(d_0) \cdot \dots \cdot \text{GL}_l^i(d_p) \\ &= \text{CDS}_l^i(d_0) \cdot \text{CDS}_l^{i+1}(d_0 \gamma_i) \cdot \dots \\ &\quad \dots \cdot \text{CDS}_l^i(d_{p-1}) \cdot \text{CDS}_l^{i+1}(d_{p-1} \gamma_i) \cdot \text{CDS}_l^i(d_p) \end{aligned} \tag{9}$$

where $\text{CW}_{M_l, M_{l+1}}^i(d) = (d_0 = d, \dots, d_p)$.

In (9), since $\text{CDS}_l^i(d_p)$ has d_p as its first dart (Lemma 5), it is in particular not empty and it is therefore readily seen that

$$\text{last}(\text{CDS}_{l+1}^i(d)) = \text{last}(\text{CDS}_l^i(d_p))$$

By the recurrence hypothesis applied to $\text{CDS}_l^i(d_p)$, we have:

$$\text{last}(\text{CDS}_l^i(d_p))\gamma_{0,i} = d_p \gamma_{l,i}$$

From Property 8 applied to $\text{CW}_{M_l, M_{l+1}}^i(d)$ we know that $d_0 \gamma_{l+1,i} = d_p \gamma_{l,i}$. It follows that

$$\text{last}(\text{CDS}_{l+1}^i(d))\gamma_{0,i} = d_0 \gamma_{l+1,i}$$

The property is thus valid at level $l + 1$. \square

6.2 An iterative definition

In this section, we introduce a non-recursive definition of connecting walks, stated by Proposition 17. Thanks to Proposition 15, this definition allows to retrieve the image of an involution at any level of a pyramid using a traversal of the base level map directed by the knowledge of the level at which a dart disappears and the degree of the corresponding cells that has been removed. These two values (level and degree) are formally described by the membership of a dart to a set $\text{AR}_{l,i}$ according to the following notation.

Notation 9 *If (M_0, K_1, \dots, K_h) is a pyramid of n -maps and l is an integer in $\{1, \dots, h\}$ and $0 \leq i \leq n - 1$, then we denote*

$$\text{AR}_{l,i} = \bigcup_{k=0}^{l-1} R_{k,i}^* \quad (10)$$

where “AR” stands for “aggregated removal set”, i.e., the union of the i -cells to be removed from every levels of the pyramid below level l (excluded).

We now state several properties of these sets.

Lemma 7 *For all $l \in \{0, \dots, h\}$ and all $i, j \in \{0, \dots, n - 1\}$ with $i \neq j$, we have $\text{AR}_{l,i} \cap \text{AR}_{l,j} = \emptyset$.*

Proof: We prove this result by a recurrence on l . If $l = 1$, we have $\text{AR}_{1,i} = R_{1,i}^*$ and $\text{AR}_{1,j} = R_{1,j}^*$, therefore the result comes from the very definition of a removal kernel (i.e., cells to be removed are disjoint). We thus suppose that the property holds for $l \in \{1, \dots, h - 1\}$.

Following (10), we have $\text{AR}_{l+1,i} = \text{AR}_{l,i} \cup R_{l,i}^*$ and $\text{AR}_{l+1,j} = \text{AR}_{l,j} \cup R_{l,j}^*$. It follows that

$$\text{AR}_{l+1,i} \cap \text{AR}_{l+1,j} = (\text{AR}_{l,i} \cup R_{l,i}^*) \cap (\text{AR}_{l,j} \cup R_{l,j}^*)$$

By (10) we also deduce that $R_{l,i}^* \cap \text{AR}_{l,j} = \emptyset$ since $R_{l,i}^* \subset \mathcal{D}_l$ and $\text{AR}_{l,j} \cap \mathcal{D}_l = \emptyset$. Similarly, we have $\text{AR}_{l,i} \cap R_{l,j}^* = \emptyset$. Furthermore, from the very definition of a removal kernel we know that $R_{l,i}^* \cap R_{l,j}^* = \emptyset$. Eventually, from the recurrence hypothesis we have $\text{AR}_{l,i} \cap \text{AR}_{l,j} = \emptyset$. We thus deduce from the above equation that $\text{AR}_{l+1,i} \cap \text{AR}_{l+1,j} = \emptyset$, hence the property is valid at level $l + 1$. \square

We now prove several useful lemmas which [...]

Lemma 8 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ and let d_1, d_2 be two darts of \mathcal{D}_l , $0 \leq l \leq h$. If $d_1\gamma_i = d_2$ for $i \in \{0, \dots, n-1\}$, then $d_1\gamma_{l,i} = d_2$.*

Proof: This may be proved by a simple recurrence on k , $0 \leq k \leq l$, using the very definition of cells removal in an n -map (Definition 17).

Let $d_1, d_2 \in \mathcal{D}_l$ and $i \in \{0, \dots, n-1\}$ be such that $d_1\gamma_i = d_2$. Using the notation introduced before (Notation 7), this may be written $d_1\gamma_{0,i} = d_2$, hence the property is valid for $k = 0$.

Now we suppose that $d_1\gamma_{k,i} = d_2$ for $k \in \{0, \dots, l-1\}$. As $d_2 = d_1\gamma_{k,i}$ belongs to \mathcal{D}_l , it necessarily belongs to \mathcal{D}_{k+1} since $k < l$. It follows from Definition 17 that $d_1\gamma_{k+1} = d_1\gamma_k$, i.e. $d_1\gamma_{k+1} = d_2$. The property is thus satisfied at level $k+1$; it is therefore true for all $k \in \{0, \dots, l\}$ and in particular for l . We obtain $d_1\gamma_{l,i} = d_2$. \square

Lemma 9 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ and d be a dart of \mathcal{D}_l . If $d\gamma_j \notin \mathcal{D}_l$ for $j \in \{0, \dots, n-1\}$ then $d\gamma_j \in \text{AR}_{l,j}$.*

Proof: If $d\gamma_j \notin \mathcal{D}_l$ then $d\gamma_j \in R_{u,k}$ for $u \in \{0, \dots, l-1\}$ and $k \in \{0, \dots, n-1\}$. Since $d \in \mathcal{D}_l \subset \mathcal{D}_u$ and $d\gamma_j \in \mathcal{D}_u$, from Lemma 8 we deduce that $d\gamma_j = d\gamma_{u,j}$. If we suppose that $k \neq j$, as $k < n$ we obtain that d and $d\gamma_{u,j}$ belong to the same k -cell of R_u ; a contradiction with the fact that $d \in \mathcal{D}_l$ ($l > u$). Necessarily, we have $k = j$ and therefore $d\gamma_j \in \text{AR}_{l,j}$. \square

Lemma 10 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ and d be a dart of \mathcal{D}_0 , such that $d \in \text{AR}_{l,i}$ for $l \in \{1, \dots, h\}$ and $i \in \{0, \dots, n-1\}$. If $d\gamma_j \in \text{AR}_{l,k}$ for $j \in \{0, \dots, n-1\} \setminus \{i\}$ and $k \in \{0, \dots, n-1\}$, then either $k = i$ or $k = j$.*

Proof: We prove this result by recurrence on the level l .

If $l = 1$, we have $\text{AR}_{1,i} = R_{0,i}^*$ and $\text{AR}_{1,k} = R_{0,k}^*$. If $d \in R_{0,i}^*$ and $d\gamma_j \in R_{0,k}^*$ with $i \neq j$, since d and $d\gamma_j = d\gamma_{0,j}$ belong to the same i -cell of M_0 ($j < n$), we necessarily have $k = i$. Otherwise, $d\gamma_j$ would belong to two cells to be removed at level 0, which is not allowed by the definition of a removal kernel. If $i = j$ and $d\gamma_i \notin \mathcal{D}_1$ then $d\gamma_i \in R_{0,k}$ for some $k \in \{0, \dots, n-1\}$. Let us suppose that $k \neq i$, since $R_{0,n} = \emptyset$ it follows that d and $d\gamma_i$ belong to the same k -cell of $R_{0,k}$. Again, this would mean that d belongs to both an i -cell of $R_{0,i}^*$ and a k -cell of $R_{0,k}^*$, which is not allowed from the definition of the removal kernel K_0 . Therefore $k = i$, i.e. $d\gamma_i \in \text{AR}_{1,i}$. Hence the property is satisfied when $l = 1$.

We consider the property as satisfied for $l \in \{1, \dots, h-1\}$ and we prove that it is also satisfied at level $l+1$. Thus, suppose that $d \in \text{AR}_{l+1,i}$ and $d\gamma_j \in \text{AR}_{l+1,k}$ for $i, j, k \in \{0, \dots, n-1\}$ with $i \neq j$. We have $\text{AR}_{l+1,i} = \text{AR}_{l,i} \cup R_{l,i}^*$ and $\text{AR}_{l+1,k} = \text{AR}_{l,k} \cup R_{l,k}^*$. Therefore, we may distinguish the following cases:

- If $d \in \text{AR}_{l,i}$ we have $d \in R_{u,i}^*$ for some $u \in \{0, \dots, l-1\}$.

Let us suppose that $d\gamma_j \in R_{l,k}^*$. In this case $d\gamma_j \in \mathcal{D}_l \subset \mathcal{D}_u$. It follows from Lemma 8 that $d\gamma_j = d\gamma_{u,j}$. As $j \neq i$ and $i < n$ we obtain that d and $d\gamma_{u,j}$ belong to the same i -cell of $R_{u,i}$, which is a contradiction with the fact that $d\gamma_{u,j} \in \mathcal{D}_l$ with $l > u$; hence $d\gamma_j \notin R_{l,k}^*$, i.e. $d\gamma_j \in \text{AR}_{l,k}$. Eventually, since $d \in \text{AR}_{l,i}$ and $d\gamma_j \in \text{AR}_{l,k}$, by the recurrence hypothesis we immediately have $k \in \{i, j\}$.

- If $d \in R_{l,i}^*$ and $d\gamma_j \in R_{l,k}^*$. Since in this case both d and $d\gamma_j$ belong to \mathcal{D}_l , we deduce from Lemma 8 that $d\gamma_j = d\gamma_{l,j}$.

Suppose that $k = j$. As $i \neq j$ (i.e. $k \neq i$), d and $d\gamma_{l,j}$ would belong to the same i -cell of $R_{l,i}^*$ which contradicts the fact that $R_{l,i}^* \cap R_{l,k}^* = \emptyset$. Hence $k \neq j$. It follows that d and $d\gamma_{l,j}$ belong to the same k -cell of $R_{l,k}^*$. Therefore, $k \neq i$ would contradict the fact that K_l is a removal kernel since d would belong to both an i -cell and a k -cell to be removed; eventually we necessarily have $k = i$.

- If $d \in R_{l,i}^*$ and $d\gamma_j \in \text{AR}_{l,k}$, we have $d\gamma_j \in R_{u,k}^*$ for $u \in \{0, \dots, l-1\}$. Since $d \in R_{l,i}^*$, in particular $d \in \mathcal{D}_v$ for all $v \in \{0, \dots, l\}$ so that $d \in \mathcal{D}_u$. Thus, as d and $d\gamma_j$ both belong to \mathcal{D}_u , by Lemma 8 we obtain $d\gamma_j = d\gamma_{u,j}$. Now, suppose that $j \neq k$. We deduce that d and $d\gamma_{u,j}$ belong to the same k -cell of K_u . In other words, d is to be removed from M_u , which is a contradiction with the fact that $d \in \mathcal{D}_l$ with $l > u$. Hence, we necessarily have $k = j$.

We conclude that the property is valid at level $l+1$, and therefore for all $l \in \{1, \dots, h\}$. \square

Proposition 16 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ with $M_0 = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$. If d is a dart of M_l , with $1 \leq l \leq h$, we have:*

- $\text{CDS}_l^{n-1}(d)^{\times*} \subset \text{AR}_{l+1, n-1}$
- If $\text{CDS}_l^{n-1}(d) = (d_0 = d, d_1, \dots, d_p)$ for $p \in \mathbb{N}^*$, then $d_{u+1} = d_u \gamma_{n-1}$ for all $u \in \{0, \dots, p-1\}$.

Proof: We prove the property by recurrence on l . If $l = 0$ we have $\text{CDS}_l^{n-1}(d) = (d)$ and there is nothing to be proved. Thus, we suppose that the property holds for $l \in \{0, \dots, h-1\}$.

According to Definition 26, we have:

$$\text{CDS}_{l+1}^{n-1}(d) = \text{CDS}_l^{n-1}(d_0) \cdot \text{CDS}_l^{n-1}(d_1) \cdot \dots \cdot \text{CDS}_l^{n-1}(d_p)$$

with $\text{CW}_{M_l, M_{l+1}}^{n-1}(d) = (d_0 = d, d_1, \dots, d_p)$.

By Lemma 5, $\text{CDS}_l^{n-1}(d_h)$ contains at least d_h as its first element, thus it follows that

$$\text{CDS}_{l+1}^{n-1}(d)^{\times*} = \text{CDS}_{l+1}^{n-1}(d)^{\times*} \bigcup_{h \in \{1, \dots, p\}} \text{CDS}_{l+1}^{n-1}(d_h)^* \quad (11)$$

From Property 7 applied to $\text{CW}_{l+1, n-1}$ we know that $d_k \in R_l^{n-1} \subset \text{AR}_{l+1, n-1}$ for all $k \in \{1, \dots, p\}$, and by the recurrence hypothesis we have $\text{CDS}_l^{n-1}(d_h)^{\times*} \subset \text{AR}_{l, n-1}$ for all $h \in \{0, \dots, p\}$. Since $\text{AR}_{l, n-1} \subset \text{AR}_{l+1, n-1}$, from (11) we deduce that $\text{CDS}_{l+1}^{n-1}(d)^{\times*} \subset \text{AR}_{l+1, n-1}$. Property *i*) is therefore satisfied by $\text{CDS}_{l+1}^{n-1}(d)$.

Now, let us write $\text{CDS}_{l+1}^{n-1}(d) = (d'_0, \dots, d'_q)$. For all $v \in \{0, \dots, q-1\}$ we may distinguish two cases:

- If $d'_v = \text{last}(\text{CDS}_l^{n-1}(d_k))$ for some $k \in \{0, \dots, p-1\}$, then by Proposition 15 we have $\text{last}(\text{CDS}_l^{n-1}(d_k))\gamma_{n-1} = d_k\gamma_{l, n-1}$, hence $d'_v\gamma_{n-1} = d_k\gamma_{l, n-1}$. Now, from the definition of the connecting walk $\text{CW}_{M_l, M_{l+1}}^{n-1}(d)$, since $k < p$ we have $d_k\gamma_{l, n-1} = d_{k+1}$. As in this case $d_{k+1} = d'_{v+1}$ we obtain $d'_{v+1} = d'_v\gamma_{n-1}$.
- If $d'_v \in \text{CDS}_l^{n-1}(d_k)^{\times*}$ we may apply the recurrence hypothesis to d'_v in $\text{CDS}_l^{n-1}(d_k)$ and immediately obtain that $d'_{v+1} = d'_v\gamma_{n-1}$.

We conclude that property *ii*) is satisfied for $l+1$.

Eventually, both properties are therefore satisfied for all $l \in \{0, \dots, h\}$. \square

Lemma 11 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ with $M_0 = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$. If $\text{CDS}_l^i(d) = (d_0 = d, d_1, \dots, d_p)$, $p \in \mathbb{N}^*$, is the i -connecting dart sequence associated to a dart d of M_l , with $1 \leq l \leq h$ and $0 \leq i \leq n$; then $d_p \in \text{AR}_{l, i}$ and $d_1 \in \text{AR}_{l, i}$.*

Proof: We prove the lemma by recurrence on l . If $l = 0$ there is nothing to prove since from Definition 26 we have $\text{CDS}_l^i(d) = (d)$, hence $p = 0$. If $l = 1$, we have $\text{CDS}_l^i(d) = \text{CW}_{M_0, M_1}^i(d)$ and from Property 7, if $\text{CW}_{M_0, M_1}^i(d)$ counts more than one dart we have $\text{CW}_{M_0, M_1}^i(d)^{\times*} \subset R_{0, i}^*$. Thus, in this case $d_1 = d_p \in \text{AR}_{1, i}$. We may then suppose that the property is valid for $l \in \{0, \dots, h-1\}$.

Let $\text{CW}_{M_l, M_{l+1}}(d) = (d = d'_0, \dots, d'_q)$. If $q = 0$, then from Definition 26 we have $\text{CDS}_{l+1}^i(d) = \text{CDS}_l^i(d)$ and the property holds by the recurrence hypothesis.

applied to $\text{CDS}_l^i(d)$. Thus, we suppose in the following that $q > 0$, which by Lemma 5 implies $p > 0$.

From Definition 26, the first sequence of darts defining $\text{CDS}_{l+1}^i(d)$ is $\text{CDS}_l^i(d)$.

- If $\text{CDS}_l^i(d)$ contains more than one dart, then its second dart is precisely d_1 , which by the recurrence hypothesis belongs to $\text{AR}_{l,i} \subset \text{AR}_{l+1,i}$.
- If $\text{CDS}_l^i(d) = (d)$, then following Definition 26 d_1 is the second dart of $\text{CW}_{M_l, M_{l+1}}(d)$. From Property 7 of the latter connecting walk we obtain $d_1 \in R_{l,i}^* \subset \text{AR}_{l+1,i}$.

On the other hand, still following Definition 26, the last sequence of darts defining $\text{CDS}_{l+1}^i(d)$ is $\text{CDS}_l^i(d'_q)$.

- If $\text{CDS}_l^i(d'_q) = (d'_q)$, then from Property 7 of $\text{CW}_{M_l, M_{l+1}}^i(d)$, since $q > 0$, we have $d'_q \in R_{l,i}^* \subset \text{AR}_{l+1,i}$. As $d'_q = d_p$ in this case, we obtain $d_p \in \text{AR}_{l+1,i}$.
- Otherwise, $\text{CDS}_l^i(d'_q)$ counts more than one dart and by the recurrence hypothesis its last dart belongs to $\text{AR}_{l,i}$, and therefore to $\text{AR}_{l+1,i}$ since $\text{AR}_{l,i} \subset \text{AR}_{l+1,i}$. Furthermore, it is also the last dart of $\text{CDS}_{l+1}^i(d)$, i.e. d_p . We obtain $d_p \in \text{AR}_{l+1,i}$.

In both cases, we thus have $d_1 \in \text{AR}_{l+1,i}$. Eventually, the property is valid for $l + 1$, and therefore satisfied for all $l \in \{0, \dots, h\}$. \square

Proposition 17 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ with $M_0 = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$. If $\text{CDS}_l^i(d) = (d_0 = d, d_1, \dots, d_p)$, $p \in \mathbb{N}^*$, is the i -connecting dart sequence associated to a dart d of M_l , with $1 \leq l \leq h$ and $0 \leq i \leq n$; then for all $u \in \{0, \dots, p-1\}$ we have $d_{u+1} = d_u \gamma_{t_u}$ where t_u , which is called the involution subscript associated with d_u in $\text{CDS}_l^i(d)$, is defined as follows:*

- $t_0 = i$
- For all $u \in \{1, \dots, p-1\}$

$$t_u = \begin{cases} t_{u-1} + 1 & \text{if } t_{u-1} < n - 1 \text{ and } d_u \in \text{AR}_{l, t_{u-1}} \\ t_{u-1} & \text{if } t_{u-1} = n - 1 \text{ and } d_u \in \text{AR}_{l, t_{u-1}} \\ t_{u-1} - 1 & \text{if } d_u \in \text{AR}_{l, t_{u-1}-1} \end{cases} \quad (12)$$

Moreover $t_{p-1} = n - 1$ if $i = n - 1$, otherwise $t_{p-1} = i + 1$. (Note that $p \neq 0$ implies that $p > 1$ by Lemma 6.)

Proof: If $i = n - 1$, from Proposition 16 we know that $d_u \in \text{AR}_{l+1, n-1}$ for all $u \in \{0, \dots, p\}$, and $d_{u+1} = d_u \gamma_{n-1}$ for all $u \in \{0, \dots, p-1\}$. It is therefore readily seen that the proposition holds since, in particular, only the second

condition of (12) applies for all $u \in \{0, \dots, p-1\}$. Thus, we suppose in the following that $i < n-1$.

We prove the proposition by recurrence on l . Let $\mathcal{H}_1(l)$ be the property of $\text{CDS}_l^i(d)$ described by the proposition. We first show that the property holds for $l=0$. Following Definition 26, if $l=0$ we have $\text{CDS}_l^i(d) = (d)$ so that $p=0$ and there is nothing left to prove. We may thus suppose that the property $\mathcal{H}_1(l)$ is satisfied and we show that it is also valid for $l+1$.

With $\text{CW}_{M_l, M_{l+1}}^i(d) = (b_0 = d, \dots, b_q)$ we have:

$$\begin{aligned} \text{CDS}_{l+1}^i(d) &= \text{GL}_l^i(b_0) \cdot \text{GL}_l^i(b_1) \cdot \dots \cdot \text{GL}_l^i(b_p) \\ &= \text{CDS}_l^i(b_0) \cdot \text{CDS}_l^{i+1}(b_0\gamma_{l,i}) \cdot \text{CDS}_l^i(b_1) \cdot \text{CDS}_l^{i+1}(b_1\gamma_{l,i}) \cdot \dots \\ &\quad \dots \cdot \text{CDS}_l^i(b_q) \end{aligned} \quad (13)$$

We may denote

$$\text{CDS}_{l+1}^i = (d'_0, \dots, d'_{p'})$$

If $q=0$, we immediately obtain $\text{CDS}_{l+1}^i(d) = \text{CDS}_l^i(d)$. In this case, the property is immediate since the recurrence hypothesis applies for $\text{CDS}_l^i(d)$ and $\text{AR}_{l,j} \subset \text{AR}_{l+1,j}$ for all $j \in \{0, \dots, n\}$. Therefore, we assume in the following that $q > 0$. Furthermore, as each subsequence in (13) is not empty, if $q > 0$ we necessarily have $p' > 0$.

Now, we prove by recurrence on v that the following property $\mathcal{H}_2(v)$, made of the two properties below, holds for all $v \in \{0, \dots, p'-1\}$:

Recurrence hypothesis $\mathcal{H}_2(v)$

i) $d'_{v+1} = d'_v \gamma_{t'_v}$ where $t'_0 = i$ and if $v > 0$ we have

$$t'_v = \begin{cases} t'_{v-1} + 1 & \text{if } t'_{v-1} < n-1 \text{ and } d'_v \in \text{AR}_{l+1, t'_{v-1}} \\ t'_{v-1} & \text{if } t'_{v-1} = n-1 \text{ and } d'_v \in \text{AR}_{l+1, t'_{v-1}} \\ t'_{v-1} - 1 & \text{if } d'_v \in \text{AR}_{l, t'_{v-1}-1} \end{cases} \quad (14)$$

ii) Exactly one of the following conditions holds:

- (1) $d'_v = \text{last}(\text{CDS}_l^i(d_k))$ for some $k \in \{0, \dots, q-1\}$ and $t'_v = i$;
- (2) $d'_v = \text{last}(\text{CDS}_l^{i+1}(d_p))$;
- (3) $d'_v = \text{last}(\text{CDS}_l^{i+1}(d_k\gamma_{l,i}))$ for some $k \in \{0, \dots, q-1\}$ and $t'_v = i+1$;
- (4) $d'_v \in \text{CDS}_l^i(d_h)^{\times*}$ for some $h \in \{0, \dots, q\}$ or $d'_v \in \text{CDS}_l^i(d_k\gamma_{l,i})^{\times*}$ for some $k \in \{0, \dots, q-1\}$, and t'_v is the involution subscript associated with d'_v in $\text{CDS}_l^i(d_h)$ or $\text{CDS}_l^i(d_k\gamma_{l,i})$, respectively.

If $v = 0$, we write $\text{CDS}_l^i(d) = (d''_0, \dots, d''_r)$ which is the first sequence of darts of $\text{CDS}_{l+1}^i(d)$ following Definition 26. We may thus distinguish two cases:

- If $r > 0$, then from the recurrence hypothesis $\mathcal{H}_1(l)$ applied to $\text{CDS}_l^i(d)$ we have $d''_1 = d''_0 \gamma_i$. Since in this case $d''_1 = d'_1$ and $d''_0 = d'_0$, condition i) of the property is valid for $v = 0$ with $t'_v = i$, and only condition (4) of ii) holds.
- If $r = 0$, we have $d'_0 = \text{last}(\text{CDS}_l^i(b_0))$. We know by Proposition 15 that $\text{last}(\text{CDS}_l^i(d_0)) \gamma_i = d_0 \gamma_{l,i}$. Since $q > 0$ we have $d_0 \gamma_{l,i} = b_1$ (from the definition of the connecting walk $\text{CW}_{M_l, M_{l+1}}^i(d)$) with $b_1 = d'_1$ as $r = 0$. We eventually obtain $d'_0 \gamma_i = d'_1$. Condition i) of the hypothesis is then valid for $v = 0$ with $t'_0 = i$, and condition (1) of ii) holds.

We have just shown that property $\mathcal{H}_2(0)$ holds. We may then suppose that $\mathcal{H}_2(w)$ holds for all $w \leq v$, with $v < p' - 1$; and we consider the dart d'_{v+2} for $v \in \{1, \dots, p' - 2\}$. We may distinguish the four following cases:

- (a) $d'_{v+1} = \text{last}(\text{CDS}_l^i(d_k))$ for some $k \in \{0, \dots, q - 1\}$ d'_ρ in (15)
- (b) $d'_{v+1} \in \text{CDS}_l^i(d_h)^{\times*}$ for some $h \in \{0, \dots, q\}$ d'_μ in (15)
- (c) $d'_{v+1} = \text{last}(\text{CDS}_l^{i+1}(d_k \gamma_{l,i}))$ for some $k \in \{0, \dots, q - 1\}$ d'_ϕ in (15)
- (d) $d'_{v+1} \in \text{CDS}_l^{i+1}(d_k \gamma_{l,i})^{\times*}$ for some $k \in \{0, \dots, q - 1\}$ d'_ω in (15)

as illustrated below

$$\begin{array}{cccccccc}
\text{CDS}_{l+1}^i(d) & = & \text{CDS}_l^i(b_0) & \cdot & \text{CDS}_l^{i+1}(b_0 \gamma_{l,i}) & \cdot & \text{CDS}_l^i(b_1) & \cdot & \dots & \cdot & \text{CDS}_l^i(b_q) \\
\text{CDS}_{l+1}^i(d) & = & (b_0, \dots) & \cdot & (b_0 \gamma_{l,i}, \dots, \boxtimes, \dots, \boxtimes) & \cdot & (b_1, \dots, \boxtimes, \dots, \boxtimes) & \cdot & \dots & \cdot & (b_q, \dots) \\
\text{CDS}_{l+1}^i(d) & = & (d'_0, d'_1, \dots) & & \dots, d'_\omega, \dots, d'_\phi, \dots & & \dots, d'_\mu, \dots, d'_\rho, \dots & & \dots, d'_{p'} & & \\
& & & & \updownarrow & \updownarrow & \updownarrow & \updownarrow & & & \\
& & & & \text{(d)} & \text{(c)} & \text{(b)} & \text{(a)} & & & \text{(15)}
\end{array}$$

- (a) If $d'_{v+1} = \text{last}(\text{CDS}_l^i(b_k))$ for some $k \in \{0, \dots, q - 1\}$.
Let us denote $\text{CDS}_l^i(d_k) = (d''_0 = b_k, \dots, d''_r)$, for $r \in \mathbb{N}$. We have $d'_{v+1} = d''_r$.

From Proposition 15, we know that $d'_{v+1} \gamma_i = d_k \gamma_{l,i}$. Since $k < q$, from the very definition of $\text{CDS}_{l+1}^i(d)$ (see (15)) we have $d_k \gamma_{l,i} = d'_{v+2}$, therefore $d'_{v+2} = d'_{v+1} \gamma_i$.

- If $r = 0$, since $d'_{v+1} = d''_0$ and $v > 0$, we necessarily have $k > 0$ (otherwise we would have $d'_{v+1} = d$, a contradiction with $v > 0$). It follows that $d'_v = \text{last}(\text{CDS}_l^{i+1}(b_{k-1} \gamma_{l,i}))$. By condition (3) of the recurrence hypothesis $\mathcal{H}_2(v)$, we have $t'_v = i + 1$. On the other hand, if $r = 0$ we have $d'_{v+1} = b_k$ and by Property 7 applied to $\text{CW}_{M_l, M_{l+1}}^i(d)$ we know that $b_k \in R_{l,i}^*$, thus $d'_{v+1} \in \text{AR}_{l+1,i}$. Since $t'_v = i + 1$ we eventually obtain that $d'_{v+1} \in \text{AR}_{l+1, t'_v - 1}$ and $d'_{v+2} = d'_{v+1} \gamma_{t'_v - 1}$, so that the property $\mathcal{H}_2(v + 1)$ is satisfied with $t'_{v+1} = t'_v - 1 = i$.
- If $r > 0$, we may apply the recurrence hypothesis $\mathcal{H}_1(l)$ to $\text{CDS}_l^i(b_k)$ and therefore denote by $(t''_0, \dots, t''_{r-1})$ the sequence of involution subscripts associated with $\text{CDS}_l^i(d_k)$. From the *moreover* part of $\mathcal{H}_1(l)$, since $r > 0$

we have $t''_{r-1} = i + 1$ (and $r > 1$ by Lemma 6). From the recurrence hypothesis $\mathcal{H}_2(v)$, we know that t'_v is equal to the involution subscript associated with d'_v (i.e. b_{r-1}) in $\text{CDS}_l^i(d_k)$, namely $t'_v = t''_{r-1} = i + 1$.

Furthermore, by Lemma 11, we know that $d'_{v+1} \in \text{AR}_{l,i}$, i.e. $d'_{v+1} \in \text{AR}_{l+1,i}$ since $\text{AR}_{l,i} \subset \text{AR}_{l+1,i}$. Eventually, we have $d'_{v+1} \in \text{AR}_{t'_v-1}$ and $d'_{v+2} = d'_{v+1}\gamma_{t'_v-1}$ so that the first part of the property \mathcal{H}_{v+1} is satisfied with $t'_{v+1} = t'_v - 1 = i$.

Furthermore, in all cases, condition (1) of $\mathcal{H}_2(v+1)$ holds since $t'_{v+1} = i$.

- (b) If $d'_{v+1} \in \text{CDS}_l^i(b_h)^{\times*}$ for some $h \in \{0, \dots, q\}$.

Let us denote $\text{CDS}_l^i(b_h) = (d''_0 = b_h, \dots, d''_r)$, for $r \in \mathbb{N}$. Since $d'_{v+1} \in \text{CDS}_l^i(b_h)^{\times*}$ we necessarily have $r > 0$, with $d'_{v+1} = d''_s$ for some $s \in \{0, \dots, r-1\}$. Since $s < r$, we also have $d'_{v+2} = d''_{s+1}$.

- If $s = 0$ we have $d'_{v+1} = b_h$ and from Property 7 applied to $\text{CDS}_{M_l, M_{l+1}}(d)$ we deduce that $d'_{v+1} \in R_{l,i}^* \subset \text{AR}_{l+1,i}$. By the recurrence hypothesis $\mathcal{H}_1(l)$ we know that $d''_1 = d''_0\gamma_i$, i.e. $d'_{v+2} = d'_{v+1}\gamma_i$. Now, since in this case $d'_{v+1} = \text{first}(\text{CDS}_l^i(b_h))$ we necessarily have $h > 0$ and $d'_v = \text{last}(\text{CDS}_l^{i+1}(b_{h-1}\gamma_{l,i}))$. From condition (3) of the recurrence hypothesis $\mathcal{H}_2(v)$ we deduce that $t'_v = i + 1$. On the other hand, by Lemma 11 we know that $d'_v = \text{last}(\text{CDS}_l^{i+1}(b_{h-1}\gamma_{l,i})) \in \text{AR}_{l,i+1}$. As $\text{AR}_{l,i+1} \subset \text{AR}_{l+1,i+1}$, since $d'_{v+2} = d'_{v+1}\gamma_i$ we conclude that the first part of the recurrence hypothesis $\mathcal{H}_1(l+1)$ is satisfied with $t'_{v+1} = t'_v - 1 = i$. Furthermore, condition (4) of the recurrence hypothesis $\mathcal{H}_2(v+1)$ is satisfied since by the recurrence hypothesis $\mathcal{H}_1(l)$ the involution subscript associated with b_h in $\text{CDS}_{l,i}(b_h)$ is i .

- If $s > 0$, we may apply the recurrence hypothesis $\mathcal{H}_1(l)$ to $\text{CDS}_l^i(b_h)$ and therefore denote by $(t''_0, \dots, t''_{r-1})$ the sequence of involution subscripts associated with $\text{CDS}_l^i(b_h)$. We thus have $d''_{s+1} = d''_s\gamma_{t''_s}$.

Since $s > 0$ we have $b''_s \notin \mathcal{D}_l$, and therefore obtain that $d''_s \in \text{AR}_{l,j}$ if and only if $d''_s = d'_{v+1} \in \text{AR}_{l+1,j}$, for all $j \in \{0, \dots, n-1\}$.

Furthermore, from condition (4) of the recurrence hypothesis $\mathcal{H}_2(v)$, we know that t'_v is equal to the involution subscript associated with d'_v (i.e. d''_{s-1}) in $\text{CDS}_l^i(b_h)$, namely $t'_v = t''_{s-1}$.

It follows immediately from the definition of t''_s given by $\mathcal{H}_1(l)$, that the property i of $\mathcal{H}_2(v+1)$ is satisfied with $t'_{v+1} = t''_s$, which is also precisely the condition (4) of property ii) in the latter hypothesis.

- (c) If $d'_{v+1} = \text{last}(\text{CDS}_l^{i+1}(b_k\gamma_{l,i}))$ for some $k \in \{0, \dots, p-1\}$.

Let us denote $\text{CDS}_l^{i+1}(b_k\gamma_{l,i}) = (d''_0 = b_k\gamma_{l,i}, \dots, d''_r)$, for $r \in \mathbb{N}$. We have $d'_{v+1} = d''_r$.

From Proposition 15, since $d'_{v+1} = \text{last}(\text{CDS}_l^{i+1}(b_k\gamma_{l,i}))$ we know that $d'_{v+1}\gamma_{i+1} = (b_k\gamma_{l,i})\gamma_{l,i+1}$. Furthermore, since $k < p$, from the very definition of $\text{CDS}_{l+1}^i(d)$ we have $d'_{v+2} = b_{k+1}$ (see (15)). Now, from the definition of $\text{CW}_{l,l+1}^i(d)$ we know that $b_{k+1} = b_k\gamma_{l,i}\gamma_{l,i+1}$, therefore $d'_{v+2} = d'_{v+1}\gamma_{i+1}$.

- If $r = 0$, from the definition of $\text{CDS}_{l+1}^i(d)$ we have $d'_v = \text{last}(\text{CDS}_l^i(b_{k-1}))$ (see (15)). By condition (1) of the recurrence hypothesis $\mathcal{H}_2(v)$, we have

$t'_v = i$. On the other hand, since $r = 0$ we have $d''_{v+1} = b_k \gamma_{l,i}$ and by Property 7 applied to $CW_{M_l, M_{l+1}}^i(d)$ we know that $b_k \gamma_{l,i} \in R_{l,i}^*$, thus $d''_{v+1} \in \text{AR}_{l+1,i}$. Since $t'_v = i$ we eventually obtain that $d''_{v+1} \in \text{AR}_{l+1,t'_v}$ and $d''_{v+2} = d''_{v+1} \gamma_{t'_v+1}$, so that the property i) of $\mathcal{H}_2(v+1)$ is satisfied with $t'_{v+1} = t'_v + 1 = i + 1$. Condition (3) of property ii) therefore holds too.

- If $r > 0$, we may apply the recurrence hypothesis $\mathcal{H}_1(l)$ to $\text{CDS}_l^{i+1}(b_k \gamma_{l,i})$ and therefore denote by $(t''_0, \dots, t''_{r-1})$ the sequence of involution subscripts associated with $\text{CDS}_l^{i+1}(b_k \gamma_{l,i})$. From the same hypothesis, since $r > 0$ we have $t''_{r-1} = i + 2$, and by Lemma 11 we have $d''_{v+1} \in \text{AR}_{i+1}$.

From the recurrence hypothesis $\mathcal{H}_2(v)$, we know that t'_v is equal to the involution subscript associated with d'_v (i.e. b_{r-1}) in $\text{CDS}_l^{i+1}(b_k \gamma_{l,i})$, namely $t'_v = t''_{r-1} = i + 2$.

Eventually, we have $d''_{v+1} \in \text{AR}_{t''_{r-1}}$ and $d''_{v+2} = d''_{v+1} \gamma_{t''_{r-1}}$ so that the property i) of the hypothesis $\mathcal{H}_2(v+1)$ is satisfied with $t'_{v+1} = t'_v - 1 = i + 1$; hence condition (3) of property ii) holds too.

- (d) If $d''_{v+1} \in \text{CDS}_l^{i+1}(b_k \gamma_{l,i})^{\times*}$ for some $k \in \{0, \dots, q-1\}$. This case is in all parts similar to case (b), considering a connecting dart sequence $\text{CDS}_l^{i+1}(b_k \gamma_{l,i})$ instead of $\text{CDS}_l^i(b_k)$ for some $k \in \{0, \dots, q-1\}$.

We conclude that the property $\mathcal{H}_2(v+1)$ is satisfied, hence it is valid for all $v \in \{0, \dots, p'-1\}$. This shows that the first part of $\mathcal{H}_1(l+1)$ is valid. Now, from the definition of $\text{CDS}_{l+1}^i(d)$ we have $d'_{p'} = \text{last}(\text{CDS}_l^i(b_q))$. As stated before, we only consider the case $q > 0$.

Let us denote $\text{CDS}_l^i(b_q) = (d''_0, \dots, d''_r)$ with $r \in \mathbb{N}$.

- If $r = 0$ we have $d'_{p'} = b_q$. Since $q > 0$ we deduce that $b_q \in CW_{M_l, M_{l+1}}^i(d)^*$ and by Property 7 we obtain $d'_{p'} \in R_l^i(d) \subset \text{AR}_{l+1,i}$. Furthermore, by condition (3) of hypothesis $\mathcal{H}_2(p'-1)$ we know that $t'_{p'-1} = i + 1$. We conclude that the *moreover* part of $\mathcal{H}_1(l+1)$ is satisfied by $\text{CDS}_{l+1}^i(d)$.
- If $r > 0$ (i.e. $r > 1$ by Lemma 6), from the *moreover* part of the recurrence hypothesis $\mathcal{H}_1(l)$ applied to $\text{CDS}_l^i(b_q)$ we deduce that $t''_{r-1} = i + 1$ (since $i < n-1$). By the recurrence hypothesis $\mathcal{H}_2(p'-1)$, t''_{r-1} is also the involution subscript associated with $d''_{p'-1}$ (i.e. d''_{r-1}) in $\text{CDS}_{l+1}^i(d)$. We conclude that $t'_{p'-1} = i + 1$ and the *moreover* part of the recurrence hypothesis $\mathcal{H}_1(l+1)$ is satisfied.

The overall conclusion is that the recurrence hypothesis $\mathcal{H}_1(l+1)$ is satisfied, it is therefore valid for all $l \in \{0, \dots, h\}$. \square

Eventually, the following corollary shows that the dimension of cells that are removed and which are traversed by a single connecting dart sequence cannot change radically for consecutive darts. Precisely, the difference is bounded by one, in absolute value.

Corollary 2 *Let (M_0, K_1, \dots, K_h) be a pyramid of n -maps for $h \in \mathbb{N}^*$ with $M_0 = (\mathcal{D}, \gamma_0, \dots, \gamma_{n-1})$. If $\text{CDS}_l^i(d) = (d_0 = d, d_1, \dots, d_p)$, $p \in \mathbb{N}^*$, is the i -connecting dart sequence associated to a dart d of M_l , with $1 \leq l \leq h$ and $0 \leq i \leq n$; then for all $u \in \{1, \dots, p\}$ we have $d_u \in \text{AR}_{\delta_u}$ where $\delta_u \in \{0, \dots, n-1\}$ satisfies:*

- $\delta_1 = i$
- For all $u \in \{1, \dots, p\}$, $\delta_u \in \{\delta_{u-1} - 1, \delta_{u-1}, \delta_{u-1} + 1\}$.

Proof: We prove the property by recurrence on u . If $u = 1$, by Lemma 11 we have $d_u \in \text{AR}_{l,i}$ hence $\delta_1 = i$. We may then suppose that the property holds for $u \in \{1, \dots, p-1\}$.

By Proposition 17, we necessarily have $d_u \in \text{AR}_{l,t_{u-1}} \cup \text{AR}_{l,t_{u-1}-1}$ where t_u is the involution subscript associated with d_u in $\text{CDS}_l^i(d)$, as defined by the same proposition.

- If $d_u \in \text{AR}_{l,t_{u-1}}$, since the recurrence hypothesis holds for d_u , following Lemma 7 we necessarily have $\delta_u = t_{u-1}$. Now, according to Proposition 17, in this case $t_u = t_{u-1} + 1$ (if $i < n-1$) or $t_u = t_{u-1}$ (if $i = n-1$). Again by Proposition 17 defining t_{u+1} we deduce that necessarily $d_{u+1} \in \text{AR}_{l,t_u}$ or $d_{u+1} \in \text{AR}_{l,t_{u-1}}$. If $i < n-1$ we obtain that either $d_{u+1} \in \text{AR}_{l,\delta_u+1}$ or $d_{u+1} \in \text{AR}_{l,t_{u-1}+1-1} = \text{AR}_{l,\delta_u}$; and if $i = n-1$ we obtain that either $d_{u+1} \in \text{AR}_{l,\delta_u}$ or $d_{u+1} \in \text{AR}_{l,\delta_u-1}$.
- If $d_u \in \text{AR}_{l,t_{u-1}-1}$, since the recurrence hypothesis holds for d_u , following Lemma 7 we necessarily have $\delta_u = t_{u-1} - 1$. Now, according to Proposition 17, in this case $t_u = t_{u-1} - 1 = \delta_u$. Again by Proposition 17 defining t_{u+1} we deduce that necessarily $d_{u+1} \in \text{AR}_{l,t_u} = \text{AR}_{l,\delta_u}$ or $d_{u+1} \in \text{AR}_{l,t_{u-1}} = \text{AR}_{l,\delta_u-1}$.

In the two cases, we obtain $d_{u+1} \in \text{AR}_{l,\delta_u} \cup \text{AR}_{l,\delta_u+1} \cup \text{AR}_{l,\delta_u-1}$ so that the property holds for $\delta_{u+1} \in \{\delta_u, \delta_u + 1, \delta_u - 1\}$. \square

7 Conclusion and perspectives

Using the definition given in [9] for the simultaneous removal of cells in an n -map, we have defined here n -dimensional combinatorial pyramids the way Brun and Kropatsch did in the two-dimensional case ([2]) and following the works of Grasset et al. about pyramids of generalized maps ([12]). We have defined connecting walks in both maps and G-maps, and established a link between the two definitions. Such walks are analogous to the *reduction windows* of regular pyramids. Connecting dart sequences, which are analogous to the *receptive fields* within regular pyramids, have also been defined.

The next step of this work consists in the definition of an implicit encoding of n -dimensional combinatorial pyramids (see [3]), based on the notion of a connecting dart sequence and Proposition 15, which yields a mean to retrieve the value of any permutation $\gamma_{l,i}$ and hence to build efficiently any map M_l .

References

- [1] L. Brun, W. Kropatsch, The construction of pyramids with combinatorial maps, Tech. Rep. 63, Institute of Computer Aided Design, Vienna University of Technology, lstr. 3/1832,A-1040 Vienna Austria (June 2000).
- [2] L. Brun, W. Kropatsch, Combinatorial pyramids, in: Suvisoft (ed.), IEEE International conference on Image Processing (ICIP), vol. II, IEEE, Barcelona, 2003.
- [3] L. Brun, W. Kropatsch, Implicit encoding of combinatorial pyramids, in: O. Drbohlav (ed.), Proceedings of the Computer Vision Winter Workshop, Valtice, Czech Republic, 2003.
- [4] G. Damiand, Définition et étude d'un modèle topologique minimal de représentation d'images 2D et 3D, Thèse de doctorat, Université Montpellier II (Décembre 2001).
- [5] G. Damiand, P. Lienhardt, Removal and contraction for n -dimensional generalized maps, in: Proceedings of the Computer Vision Winter Workshop, Bad Aussee, Austria, 2002.
- [6] G. Damiand, P. Lienhardt, Removal and contraction for n -dimensional generalized maps, in: Proceedings of the Computer Vision Winter Workshop, Bad Aussee, Austria, 2002.
- [7] G. Damiand, P. Lienhardt, Removal and contraction for n -dimensional generalized maps, in: Proceedings of 11th Discrete Geometry for Computer Imagery, vol. 2886 of Lecture Notes in Computer Science, Naples, Italy, 2003.
- [8] G. Damiand, P. Lienhardt, Removal and contraction for n -dimensional generalized maps, technical report (2003).
- [9] S. Fourey, L. Brun, A first step toward combinatorial pyramids in nD spaces, in: Graph-Based Representations in Pattern Recognition, vol. 5534 of Lecture Notes in Computer Science, Springer, 2009.
- [10] S. Fourey, L. Brun, A second step toward combinatorial pyramids in nd spaces, Tech. rep., GREYC (2009).
- [11] C. Grasset-Simon, Définition et étude des pyramides généralisées nD : application pour la segmentation multi-échelle d'images 3d, Thèse de doctorat, Université de Poitiers (2006).

- [12] C. Grasset-Simon, G. Damiand, P. Lienhardt, n D generalized map pyramids: Definition, representations and basic operations, *Pattern Recognition* 39 (4) (2006) 527–538.
- [13] P. Lienhardt, Topological models for boundary representation: a comparison with n -dimensional generalized maps, *Computer-Aided Design* 23 (1) (1991) 59–82.
- [14] P. Lienhardt, N -dimensional generalized combinatorial maps and cellular quasi-manifolds, *International Journal of Computational Geometry & Applications* 4 (3) (1994) 275–324.