

# New mathematical approach using fractional derivatives to take into account excess losses in magnetic materials

Marie-Ange Raulet, Benjamin Ducharne, Daniel Guyomar

► **To cite this version:**

Marie-Ange Raulet, Benjamin Ducharne, Daniel Guyomar. New mathematical approach using fractional derivatives to take into account excess losses in magnetic materials. 19th Soft Magnetic Materials Conference, Sep 2009, Turin, Italy. hal-00411166

**HAL Id: hal-00411166**

**<https://hal.archives-ouvertes.fr/hal-00411166>**

Submitted on 26 Aug 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# New mathematical approach using fractional derivatives to take into account excess losses in magnetic materials

Marie-Ange Raulet<sup>a</sup>, Benjamin Ducharne<sup>b</sup>, Daniel Guyomar<sup>b</sup>

<sup>a</sup> Université de Lyon, F-69622 Lyon, France, Université Lyon 1, F-69622 Villeurbanne, France, CNRS, UMR 5005, Laboratoire Ampère, F-69621 Villeurbanne France,

<sup>b</sup> Laboratoire de Génie Electrique et Ferroélectricité, Institut national des sciences appliquées de Lyon, 8, rue de la Physique, 69621 Villeurbanne, France

Accurate prediction of magnetic circuit dynamical behaviour constitutes a fundamental problem for the design and the optimization of electromagnetic devices. The representation of dynamical effects due to eddy currents with a macroscopic model [1] is usually well admitted and adopted by researchers and electrical engineers. On the other hand, macroscopic modelling of microscopic dynamical effects due to wall motions which induce the so called excess losses, always exhibits interests of various research workers [2]-[6].

This paper proposes a new mathematical approach for the modelling of the dynamical microscopic effects. This approach is based on a fractional derivative formulation. This fractional term is used to describe the excess field contribution (1):

$$Hexc(t) = \lambda \cdot \frac{d^\alpha B(t)}{dt^\alpha}. \quad (1)$$

The fractional derivative model has already been used with success to model hysteresis on a large frequency bandwidth of ferroelectric ceramics ( $1\text{mHz} < f < 10\text{Hz}$ ) [7]. In the case of magnetic materials the fractional formulation is required to describe excess losses.

The new model is implemented and tested by considering a magnetic sample magnetized by a sine flux. An analytical study is carried out and gives results which are in a good agreement with the well known Bertotti's results related to the frequency relation of the excess energy [2].

Finally, comparisons between simulations and large number of experimental results performed on a SiFe sample magnetized by a varying frequency sine flux will be considered as validation of the new model.

- 
- [1] G. Bertotti, J. Appl. Phys., **54** (1983), 5293-5305.
  - [2] G. Bertotti, IEEE Trans. Magn. **24**, N°1 (1988), 621-630.
  - [3] N. Sadowski, IEEE Trans. Magn. **38**, (2002), 797-800.
  - [4] S.E. Zirka, IEEE Trans. Magn. **42**, N°9 (2006), 2121-2130.
  - [5] M. Drouineau, JMMM. **320** (2008), e1070-e1073.
  - [6] MA Raulet, IEEE Trans. Magn. **40**, N°2 (2004), 872-875.
  - [7] D. Guyomar, IEEE J. Phys. D : Appl.Phys., (2007).