

A macroscopic crowd motion model of gradient flow type

Bertrand Maury, Aude Roudneff-Chupin, Filippo Santambrogio

► **To cite this version:**

Bertrand Maury, Aude Roudneff-Chupin, Filippo Santambrogio. A macroscopic crowd motion model of gradient flow type. *Mathematical Models and Methods in Applied Sciences*, World Scientific Publishing, 2010, à paraître. hal-00418511v2

HAL Id: hal-00418511

<https://hal.archives-ouvertes.fr/hal-00418511v2>

Submitted on 2 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A MACROSCOPIC CROWD MOTION MODEL OF GRADIENT FLOW TYPE

BERTRAND MAURY, AUDE ROUDNEFF-CHUPIN, AND FILIPPO SANTAMBROGIO

ABSTRACT. A simple model to handle the flow of people in emergency evacuation situations is considered: at every point x , the velocity $\mathbf{U}(x)$ that individuals at x would like to realize is given. Yet, the incompressibility constraint prevents this velocity field to be realized and the actual velocity is the projection of the desired one onto the set of admissible velocities. Instead of looking at a microscopic setting (where individuals are represented by rigid discs), here the macroscopic approach is investigated, where the unknown is a density $\rho(t, x)$. If a gradient structure is given, say $\mathbf{U} = -\nabla D$ where D is, for instance, the distance to the exit door, the problem is presented as a Gradient Flow in the Wasserstein space of probability measures. The functional which gives the Gradient Flow is neither finitely valued (since it takes into account the constraints on the density), nor geodesically convex, which requires for an *ad-hoc* study of the convergence of a discrete scheme.

Crowd motion; Gradient Flow; Wasserstein distance; Continuity equation.

1. INTRODUCTION

In the last two decades, several strategies have been proposed to model the motion of pedestrians. Most of them rely on a microscopic approach: the degrees of freedom are the positions of individuals, and their evolution depends on a balance between selfish behaviour, congestion constraints, and possibly social factors (politeness, gregariousness). Among those microscopic models, some are based on a stochastic description of the individual behaviour (see e.g. [27]), whereas others are purely deterministic (see [24, 28, 29]).

An essential ingredient in those models lies in the way interactions between individuals are handled, in particular in the case of high density (congestion phenomena). Following the classification which holds in the modelling of granular flows, one can differentiate the Molecular Dynamics (MD) approach (the non overlapping constraint between rigid grains is relaxed, and handled by a short range repulsive force) and the Contact Dynamics (CD) one (the collisions are explicitly taken into account). In the context of pedestrians, MD strategy has proved to be quite efficient to model congestion. In particular Helbing[22, 24, 26] introduced the concept of social forces, which are designed in such a way that individuals tend to repel each other when their distance drops below a certain value. The model proposed in [34] relies on the alternative strategy: individuals do not interact with each other as soon as they are not in contact, and the non overlapping constraint is treated in a strong (non relaxed) way. Although it is natural to expect some link between the two approaches (MD models are likely to converge in some way to their CD counterparts as the repulsive force stiffness goes to infinity), it is to be noticed that the mathematical structures of the two classes of models are quite different. In the first case, Cauchy-Lipschitz theory for ODE's applies, whereas CD models present some analogies with the so-called sweeping process introduced by Moreau[36] in the 70', for which a dedicated framework has been developed (see [20], [21], [35]).

In the case of macroscopic models, the first strategy (congestion is treated in a relaxed way) is favoured, as it allows to use classical methods for studying PDE. For example, crowd motion models inspired from traffic flow models have been developed (see Refs. [14, 12, 13]). They take the form of hyperbolic conservation laws, and they are essentially monodimensional in

space. In higher dimension, Bellomo and Dogbe[4, 19] proposed second order models, where a phenomenological relation describes how the crowd modifies its own speed:

$$\begin{cases} \partial_t \rho + \nabla_{\mathbf{x}} \cdot (\rho \mathbf{v}) = 0 \\ \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla_{\mathbf{x}}) \mathbf{v} = \mathbf{F}(\rho, \mathbf{v}). \end{cases}$$

Typically, the motion is governed by \mathbf{F} , which has two parts: a relaxation term toward a definite speed, and a repulsive term to take into account that pedestrians tend to avoid high density areas. Degond[18] uses the same approach to model sheep herds. In this model, the term \mathbf{F} depends on a pressure which blows up when the density approaches a given congestion density (barrier method). There also exist first order models, where the velocity field is directly defined as a function of the density (see e.g. [30, 31, 15]). Another class of models is described by Piccoli and Tosin in Refs. [38], [39]. They propose a time-evolving measures framework, where the velocity of the pedestrian is composed by two terms: a desired velocity and an interaction velocity. The last one models the reaction of the pedestrian to the other surrounding pedestrians (namely, people can deviate from their preferred path if they enter a crowded area).

To our knowledge, as the ones presented above, all macroscopic models rely on a relaxed expression of the congestion. Let us mention however the work of Buttazzo, Jimenez and Oudet in [9], where the optimal transportation between two given densities is computed under constraints (obstacles, congestion, ...) which can be strongly expressed. Yet, this approach is very different from the model we describe later, since its goal is to find an optimal transport between densities as in the work of Benamou and Brenier[5] (which is the classical reference for dynamical formulations of transport problems), whereas optimal transportation is in our case a very suitable tool. Moreover, we will mainly make use of the distance that optimal transport induces on probability measures rather than looking at the optimal maps themselves, as we will see after a brief description of the model we consider.

The macroscopic model we present here is based on a strong expression of the congestion constraint. It is a natural extension of the microscopic approach proposed in Refs. [34, 35, 41], which we describe here in its simpler form. The crowd configuration is represented by the position vector $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$. Each of the N individuals wishes to have a velocity \mathbf{U}_i which depends on its position only: $\mathbf{U}_i = \mathbf{U}(\mathbf{q}_i)$, where $\mathbf{U}(\cdot)$ is some given velocity field over \mathbb{R}^2 (typically $\mathbf{U} = -\nabla D$, where D is the geodesic distance to the exit). To account for non-overlapping, it is assumed that the actual velocity $\mathbf{u} = (\mathbf{u}_1, \dots, \mathbf{u}_N)$ is the ℓ^2 -projection of $\tilde{\mathbf{U}} = (\mathbf{U}_1, \dots, \mathbf{U}_N) = (\mathbf{U}(\mathbf{q}_1), \dots, \mathbf{U}(\mathbf{q}_N))$ onto the cone of feasible velocities $C_{\mathbf{q}}$ (i.e. the set of velocities which do not lead to a violation of the non-overlapping constraint). The model takes the form

$$(1) \quad \begin{cases} \frac{d\mathbf{q}}{dt} = \mathbf{u} \\ \mathbf{u} = P_{C_{\mathbf{q}}} \tilde{\mathbf{U}}. \end{cases}$$

In the spirit of this microscopic approach, the model we propose here rests on the two following principles

- (1) the pedestrian population is described by a density ρ which is subject to remain below a certain maximal value (equal to 1 in what follows), this density follows an advection equation,
- (2) the advecting field \mathbf{u} is the closest, among admissible fields (i.e. which do not lead to a violation of the constraint), to some spontaneous field \mathbf{U} , which corresponds to the strategy people would follow in the absence of others.

If we denote by C_{ρ} the cone of admissible velocities (i.e. set of velocities which do not increase density in already saturated zones, see next section for a proper definition), the model takes the

following form

$$(2) \quad \begin{cases} \partial_t \rho + \nabla \cdot \rho \mathbf{u} &= 0 \\ \mathbf{u} &= P_{C_\rho} \mathbf{U}, \end{cases}$$

where the projection is meant in the L^2 sense. As a matter of fact, in the same way as Cauchy-Lipschitz theory for ODE's no longer applies for CD in the microscopic case, we cannot use classical methods to study Equation (2), as well as most of the PDE's we could encounter in the CD macroscopic models. This is due in particular to the lack of regularity of the velocity \mathbf{u} (whose natural regularity is L^2), which prevents us to apply the characteristic method or even DiPerna-Lions theory. The non-continuous dependence of the operator P_{C_ρ} with respect to ρ is another source of problems.

Instead, we will see that this PDE corresponds to a Gradient Flow in the Wasserstein space (i.e. the space of probability measures endowed with the distance W_2 induced by the optimal transport under quadratic cost), provided that the spontaneous velocity field has a gradient structure: $\mathbf{U} = -\nabla D$. This means that we consider the functional

$$\Phi(\rho) = \begin{cases} \int_{\Omega} D(x)\rho(x)dx & \text{if } \rho \leq 1 \\ +\infty & \text{otherwise} \end{cases}$$

and we look for the curve of measures $\rho(t, \cdot)$ which follows the steepest descent direction of Φ starting from a given datum ρ^0 . This curve will happen to solve equation (2). This is a general and very efficient method to find solutions to certain evolution PDE's which been made possible by the theory of optimal transportation. This theory owes its origin to Kantorovich[33], but has been widely developed thereafter (see the books by Villani[42, 43]). Several equations have been approached by this method, for instance the classical heat equation, as well as the Fokker-Planck or the porous media equations (see Refs. [32, 37, 11]). Notice that, as the functional which is used to produce the porous media equation as a Gradient Flow is

$$\rho \mapsto \int \rho(x)^m dx,$$

our case can be considered as its formal limit when m tends to infinity. All the theory of Gradient Flow in Wasserstein Spaces is treated in the reference book by Ambrosio-Gigli-Savaré[2] and one of the key assumptions is the λ -convexity of the functional, which ensures better estimates. On the other hand, some existence results can be obtained without this assumption, but they have to be treated carefully by hand, as it happens in Ref. [7]. In our case, even if we suppose D to be λ -convex, we face the same kind of difficulties if we want to add the presence of an exit door on the boundary of Ω where the measure can concentrate (see Section 2).

The paper is organized as follows: In Section 2 we present the model in the Eulerian setting and a related discrete minimizing movement scheme (MMS). We explain how a straightforward use of a convergence theorem in [2] asserts a convergence of the trajectories for the discrete MMS to some continuous pathline. Identification of this limit with a solution to the initial problem can be done unformally. Yet some technical obstacles (in particular the handling of walls) prevent us from obtaining a fully rigorous proof based on this approach. The actual proof of convergence to a solution of the crowd motion model is based on alternative arguments. The end of this section describes this convergence results. As the presence of an exit raises some very specific technical difficulties, we propose in Section 3 a proof in the case there is no exit. The proof in the general case is given in Section 4. To illustrate the convergence theorem, we present in Section 5 an idealized (yet non trivial) situation where both eulerian solutions and discrete MMS trajectories can be described with accuracy. Finally, we discuss in Section 6 the limitations of this model and its possible extensions to other fields of natural sciences. In particular, we explain why

we developed the theory in any dimension although dimensions greater than two do not make clear sense as far as crowd motion is concerned.

2. THE EULERIAN MODEL AND ITS GRADIENT FLOW FORMULATION

2.1. Eulerian model. The model we propose is designed to handle emergency evacuation situations : the behaviour of individuals is based on optimizing their very own trajectory, regardless of others, but the fulfillment of individual strategies is made impossible because of congestion.

The model takes the following form: given a domain Ω (the building), whose boundary Γ is composed of Γ_{out} (the exit) and Γ_w (the walls), we describe the current distribution of people by a measure ρ of given mass (say 1 without loss of generality) supported within $\bar{\Omega}$. To model the fact that people getting through the door are out of danger, yet keeping a constant total mass without having to model the exterior of the building, we shall assume that ρ may concentrate on Γ_{out} . In this spirit, we denote by K the set of all those probability measures over \mathbb{R}^2 that are supported in $\bar{\Omega}$, and that are the sum of a diffuse part, with density between 0 and 1, in Ω , and a singular part carried by Γ_{out} .




FIGURE 1. Geometry.

We shall denote by \mathbf{U} the spontaneous velocity field: $\mathbf{U}(x)$ represents the velocity that an individual at x would have if he were alone. It is taken equal to 0 outside Ω . The set C_ρ of feasible velocities corresponds to all those fields which do not increase ρ on the saturated zone (unformally, $\nabla \cdot \mathbf{u} \geq 0$ in $[\rho = 1]$), and which account for walls (people do not walk through them). As we plan to define C_ρ as a closed convex set in $L^2(\Omega)$, those constraints do not make sense as they are, and we shall favor a dual definition of this set. Let us introduce the “pressure” space

$$H_\rho^1 = \{q \in H^1(\Omega), q \geq 0 \text{ a.e. in } \Omega, q(x) = 0 \text{ a.e. on } [\rho < 1], q|_{\Gamma_{out}} = 0\}.$$

The proper definition of C_ρ reads

$$(3) \quad C_\rho = \{\mathbf{v} \in L^2(\Omega)^2, \int_\Omega \mathbf{v} \cdot \nabla q \leq 0 \quad \forall q \in H_\rho^1\}.$$

The model is based on the assumption that the actual instantaneous velocity field is the feasible field which is the closest to \mathbf{U} in the least-square sense, i.e. it is defined as the L^2 -projection of \mathbf{U} onto the closed convex cone C_ρ . Finally the problem consists in finding a trajectory $t \mapsto \rho(t) \in K$ which is advected by \mathbf{u} , i.e. such that (ρ, \mathbf{u}) is a (weak) solution of the transport equation in \mathbb{R}^2

$$(4) \quad \partial_t \rho + \nabla \cdot (\rho \mathbf{u}) = 0,$$

where \mathbf{u} verifies, for almost every t ,

$$(5) \quad \mathbf{u} = P_{C_\rho} \mathbf{U}.$$

Remark 2.1. The fact that Γ_{out} is likely to carry some mass calls for some proper definition of the velocity on this zero-measure set. As the exit plays the role of a reservoir in our model, we shall actually consider that all feasible fields vanish on Γ_{out} , so that velocity \mathbf{u} given by (5) will be considered as defined Lebesgue-a.e. in Ω and vanishing on Γ_{out} .

Remark 2.2. Boundary conditions (walls and exit).

The unilateral divergence constraint and the behaviour at walls and exit are implicitly contained in the dual expression of C_ρ , as illustrated by the following considerations. We assume in this remark that $[\rho = 1] = \bar{\omega}$ where $\omega \subset \Omega$ is a smooth subdomain, and that all fields are smooth. First of all, by taking tests pressures which are smooth and compactly supported in ω , we obtain $\nabla \cdot \mathbf{u} \geq 0$ in the saturated zone. As the pressure vanishes on Γ_{out} , the velocity is free on that part of the boundary (free outlet condition, as in Darcy flows). Consider now a situation where the saturated zone covers the wall Γ_w . For any smooth function φ defined on Γ_w consider a sequence of extensions φ_ε supported within $\omega \cup \Gamma_w$, which converges to 0 in $L^2(\Omega)$. Then

$$\int_{\Omega} \mathbf{u} \cdot \nabla \varphi_\varepsilon \leq 0 \quad \forall \varepsilon > 0$$

implies

$$-\int_{\Omega} \varphi_\varepsilon \nabla \cdot \mathbf{u} + \int_{\Gamma_w} \varphi_\varepsilon \mathbf{u} \cdot \mathbf{n} \leq 0 \quad \forall \varepsilon > 0.$$

As the first term goes to 0 with ε we obtain that the velocity necessarily enters the domain on the saturated wall (what we adressed before as “people do not walk through walls”).

2.2. Gradient flow formulation. In this section we introduce a discrete evolution problem in the Wasserstein space, whose limit will be the gradient flow of a suitable functional, and we establish informally the link between this new problem and the crowd motion model. The formal equivalence, which will be proved rigorously in the following sections, will be satisfied in the case where $\mathbf{U} = -\nabla D$ is the opposite of a gradient.

Let us denote by \mathcal{P}_2 the set of probability measures over \mathbb{R}^2 endowed with the Wasserstein distance, and by

$$(6) \quad K = \{\rho \in \mathcal{P}_2, \text{supp}(\rho) \subset \bar{\Omega}, \rho = \rho_{out} + \rho_\Omega, \rho_\Omega(x) \leq 1 \text{ a.e.}, \text{supp}(\rho_{out}) \subset \Gamma_{out}\}$$

the set of feasible densities. Let an initial density ρ^0 be given, and $\tau > 0$ a time step. We build $\rho_\tau^0 = \rho^0, \rho_\tau^1, \dots$ as follows

$$(7) \quad \rho_\tau^k = \underset{\mathcal{P}_2(\mathbb{R}^d)}{\text{argmin}} \left\{ J(\rho) + \mathbf{I}_K(\rho) + \frac{1}{2\tau} W_2^2(\rho, \rho_\tau^{k-1}) \right\},$$

where W_2 is the Wasserstein distance, J is the dissatisfaction functional defined as

$$(8) \quad J(\rho) := \int_{\Omega} D(x) \rho(x) dx,$$

and \mathbf{I}_K is the indicatrix of K :

$$\mathbf{I}_K(\rho) = \begin{cases} 0 & \text{if } \rho \in K \\ +\infty & \text{if } \rho \notin K. \end{cases}$$

The function D is typically the distance to the door Γ_{out} , and to D we associate a vector field $\mathbf{U} = -\nabla D$. It is important in order to have vanishing velocities on the door that D is minimal and constant on Γ_{out} .

We admit here that under reasonable assumptions this process is indeed an algorithm (i.e. ρ_τ^{k+1} is uniquely defined as the minimizer of the function above), and we denote by ρ_τ the piecewise constant interpolate of $\rho_\tau^0, \rho_\tau^1, \dots$.

As τ goes to 0, by Prop. 2.2.3, Th. 2.3.1, and Th. 11.1.3 in [2], ρ_τ converges to some trajectory $t \mapsto \rho$ in K , which is a (weak) solution to

$$\partial_t \rho + \nabla \cdot (\rho \mathbf{u}) = 0,$$

where \mathbf{u} is such that, for almost every t ,

$$\mathbf{u} \in -\partial(J + I_K)(\rho),$$

where $\partial\Psi$ denotes the strong subdifferential of Ψ . Furthermore \mathbf{u} minimizes the L^2 norm among all those fields in the subdifferential above.

Let us now prove informally that this characterizes the instantaneous velocity as the projection of $\mathbf{U} = -\nabla D$ onto C_ρ . The subdifferential of a function Ψ at ρ in the Wasserstein setting is defined as the set of fields \mathbf{u} such that

$$\Psi(\rho) + \int_{\Omega} \langle \mathbf{u}, \mathbf{t}(x) - x \rangle d\rho(x) \leq \Psi(\mathbf{t}_{\#}\rho) + o(\|\mathbf{t} - \mathbf{i}\|),$$

where \mathbf{t} denotes a transport map acting on ρ . Note that the previous inequality does not provide any information as soon as $\mathbf{t}_{\#}\rho$ is not feasible (in that case the right-hand side is $+\infty$). Let us consider a feasible field $\mathbf{v} \in C_\rho$, and let us assume that, for ε small enough, $\mathbf{t}_\varepsilon = \mathbf{i} + \varepsilon \mathbf{v}$ pushes forward ρ onto a measure in K (this is not true in general, see Remark 2.3). Note that \mathbf{t}_ε is defined ρ -almost everywhere, with Γ_{out} carrying some mass, but as it vanishes on Γ_{out} (see Remark 2.1), the singular part of ρ remains unchanged. Having ε go to 0 in the subdifferential inequality, we obtain

$$\int \nabla D \cdot \mathbf{v} d\rho(x) + \int \mathbf{u} \cdot \mathbf{v} d\rho(x) \leq 0,$$

so that $\mathbf{u} + \nabla D = \mathbf{u} - \mathbf{U}$ belongs to C_ρ° , the polar cone to C_ρ . As \mathbf{u} minimizes the L^2 norm over $\mathbf{U} + C_\rho$, \mathbf{u} identifies with the projection of \mathbf{U} onto C_ρ , which ends this informal proof.

Remark 2.3. In general, there exist feasible densities $\rho \in K$ (defined by (6)) and fields $\mathbf{v} \in C_\rho$ (defined by (3)) such that $(\mathbf{i} + \varepsilon \mathbf{v})_{\#}\rho$ exits K for any $\varepsilon > 0$, this is why the considerations above do not make a rigorous proof. Consider for example ω a dense open subset in Ω , with a small measure, and define ρ as $\mathbf{1}_{\omega^c}$. The pressure space is $\{0\}$, and C_ρ is $L^2(\Omega)$: any field is feasible. If one considers now a strictly contractant field (with negative divergence), it is clear that $(\mathbf{i} + \varepsilon \mathbf{v})_{\#}\rho \notin K$ for any $\varepsilon > 0$. Notice also that this kind of paradox does not depend on the fact that we chose a “linear” perturbation $(\mathbf{i} + \varepsilon \mathbf{v})$, since the same would happen if one, instead, perturbs the identity by following the flow of the vector field \mathbf{v} for a time ε (which is classically a better choice in order to satisfy the density constraint).

As explained in the previous remark, the approach carried out in this section is not a rigorous proof that the advecting field is actually the projection of \mathbf{U} onto C_ρ . We conjecture that projecting $(\mathbf{i} + \varepsilon \mathbf{v})_{\#}\rho$ onto K (for the Wasserstein distance) introduces a perturbation which is negligible compared to ε , so that \mathbf{v} may actually be used as a test-function, but this conjecture raises some technical issues which we were not able to solve. In what follows we give an alternate proof which circumvents the necessity to characterize $\partial(J + I_K)$.

2.3. Notations and statement of the main result. We first recall some results on the continuity equation: let $(\rho(t, \cdot))_{t>0}$ be a family of density measures on \mathbb{R}^d , and $\mathbf{v} : (t, x) \in \mathbb{R}^+ \times \mathbb{R}^d \mapsto \mathbf{v}(t, x) \in \mathbb{R}^d$ be a Borel velocity field such that

$$(9) \quad \int_0^T \int_{\mathbb{R}^d} |\mathbf{v}(t, x)| \rho(t, x) dx < +\infty.$$

We say that (ρ, \mathbf{v}) satisfies the continuity equation with initial condition ρ^0

$$(10) \quad \begin{cases} \partial_t \rho + \nabla \cdot (\rho \mathbf{v}) &= 0 \\ \rho(0, \cdot) &= \rho^0 \end{cases}$$

if for all $\varphi \in \mathcal{C}_c^\infty([0, T] \times \mathbb{R}^d)$ we have

$$(11) \quad \int_0^T \int_{\mathbb{R}^d} (\partial_t \varphi(t, x) + \nabla \varphi(t, x) \cdot \mathbf{v}(t, x)) \rho(t, x) dx + \int_{\mathbb{R}^d} \varphi(0, x) \rho^0(x) dx = 0.$$

Let us recall that if $\rho(t, \cdot)$ is a solution of the continuity equation, there exists a narrowly continuous curve $\tilde{\rho}(t, \cdot)$ such that $\rho(t, \cdot) = \tilde{\rho}(t, \cdot)$ for a.e. t . In general, we will always focus on this continuous representation.

We now detail the construction of a discrete family of densities (ρ_τ^k) that approaches in a sense we will precise later the solution of the continuity equation we are interested with. For a fixed time step $\tau > 0$, we define the sequence (ρ_τ^k) of density measures on Ω using the recursive scheme:

$$(12) \quad \begin{cases} \rho_\tau^0 &= \rho^0 \\ \rho_\tau^k &\in \operatorname{argmin}_{\mathcal{P}_2(\mathbb{R}^d)} \left\{ J(\rho) + \mathbf{I}_K(\rho) + \frac{1}{2\tau} W_2^2(\rho, \rho_\tau^{k-1}) \right\}, \end{cases}$$

where W_2 is the Wasserstein distance, and J is the dissatisfaction functional defined in (8).

This construction is a minimizing movement scheme as described by DeGiorgi and Ambrosio in [17, 1] and then - in the framework of probability measures - in [2, 3] with functional $\Phi(\rho) = J(\rho) + \mathbf{I}_K(\rho)$.

We define on $\mathring{\Omega}$ the discrete velocities: $\mathbf{v}_\tau^k = \frac{\mathbf{i} - \mathbf{t}_\tau^k}{\tau}$, where \mathbf{t}_τ^k is the unique optimal transport function from ρ_τ^k to ρ_τ^{k-1} , which is well defined on $\mathring{\Omega}$ (but not necessarily on Γ_{out} , due to the singular part of ρ_τ^k). We also define $\mathbf{E}_\tau^k = \rho_\tau^k \mathbf{v}_\tau^k$ on $\mathring{\Omega}$ (by abuse of notation, we will write $\mathring{\Omega}$ instead of Ω when we want to stress that we are not considering the boundary). We can interpolate these discrete values $(\rho_\tau^k, \mathbf{v}_\tau^k, \mathbf{E}_\tau^k)_{k \geq 0}$ by the piecewise constant functions defined by:

$$(13) \quad \begin{cases} \rho_\tau(t, \cdot) = \rho_\tau^k \\ \mathbf{v}_\tau(t, \cdot) = \mathbf{v}_\tau^k \\ \mathbf{E}_\tau(t, \cdot) = \mathbf{E}_\tau^k \end{cases} \quad \text{if } t \in](k-1)\tau, k\tau].$$

Our goal is to prove that ρ_τ converges when $\tau \rightarrow 0$ to a solution of the continuity equation (10). Here is our main result:

Theorem 2.4. *Let Ω be a convex bounded set of \mathbb{R}^d , $D : \mathbb{R}^d \rightarrow \mathbb{R}$ a continuous λ -convex function, ρ^0 a probability density, and (ρ_τ^k) constructed following the recursive scheme (12).*

Then there exists a family of probability densities $(\rho(t, \cdot))_{t > 0}$, and a family of velocities $(\mathbf{u}(t, \cdot))_{t > 0}$ such that $(\rho_\tau(t, \cdot), \mathbf{E}_\tau(t, \cdot))$ narrowly converges to $(\rho(t, \cdot), \rho(t, \cdot) \mathbf{u}(t, \cdot))$ for a.e. t . Moreover, (ρ, \mathbf{u}) satisfies the continuity equation:

$$(14) \quad \begin{cases} \partial_t \rho + \nabla \cdot (\rho \mathbf{u}) &= 0 \\ \mathbf{u}(t, \cdot) &= P_{\mathcal{C}_{\rho(t, \cdot)}} \mathbf{U} \quad \text{for a.e. } t \\ \rho(0, \cdot) &= \rho^0 \end{cases}$$

where $\mathbf{U} = -\nabla D$, and $\mathcal{C}_{\rho(t, \cdot)}$ is defined in (3).

We will first prove this theorem in the particular case where there is no exit. In the following section, we thus assume that $\Gamma_{out} = \emptyset$, which will imply that all the measures are absolutely

continuous with respect to the Lebesgue measure. Then we will extend the proof to the general case.

Remark 2.5. We chose to assume here a λ -convexity hypothesis on D both in order to clarify some statements, which are easier to state and prove under this assumption (see for instance Lemma 3.1 and the subsequent Remark 3.2) and because the typical case we think of is $D = d(\cdot, \Gamma_{out})$, where Γ_{out} is a flat part of the boundary of the convex set Ω . This implies that D is convex as well. It would be interesting to study the case of non-convex domains Ω (for instance with obstacles), and use the geodesic distance for computing D , which would lead to a non- λ -convex function, but this is not yet possible by means of our techniques, since one should work with the Wasserstein distance W_2 computed w.r.t. the geodesic distance itself, which is not much studied.

Anyway, it can be checked that the only point throughout the paper where λ -convexity is used is the proof of Lemma 3.1, but Remark 3.2 explains how to get rid of this assumption: this means that, for existence purposes, this assumption may be withdrawn. On the other hand, the λ -convexity assumption is typical in this gradient flow problems, because it allows for uniqueness and stability results, and we think that similar results could be achieved in our case as well.

3. EXISTENCE RESULT IN A DOMAIN WITH NO EXIT

3.1. Technical lemmas. Since we will make a strong use of optimality conditions in terms of the dual problem in Monge-Kantorovitch theory, let us briefly recall what we need.

Given the two probabilities μ and ν on $\bar{\Omega}$ we always have

$$\frac{1}{2}W_2^2(\mu, \nu) = \max \left\{ \int_{\Omega} \varphi d\mu + \int_{\Omega} \psi d\nu, \varphi, \psi \in C^0(\bar{\Omega}) : \varphi(x) + \psi(y) \leq \frac{1}{2}|x - y|^2 \right\},$$

the maximum being always realized by a pair of c -concave conjugate functions (φ, ψ) with $\varphi = \psi^c$ and $\psi = \varphi^c$, where the c -transform of a function χ is defined through

$$\chi^c(y) = \inf_{x \in \Omega} \frac{1}{2}|x - y|^2 - \chi(x)$$

(with generalizations to other costs c rather than the square of the distance). We will call Kantorovitch potential from μ to ν (resp., from ν to μ) any c -concave function φ (resp., ψ) such that (φ, φ^c) (resp., (ψ^c, ψ)) realizes such a maximum. We have uniqueness of the optimal pair as soon as one of the support of one of the two measures is the whole domain $\bar{\Omega}$.

Lemma 3.1. *Let $D : \mathbb{R}^d \mapsto \mathbb{R}$ λ -convex, and $\bar{\rho} \in K$. Then, there exists τ^* such that for all $\tau < \tau^*$:*

- (i) *The functional $\phi(\rho) = \Phi(\rho) + \frac{1}{2\tau}W_2^2(\rho, \bar{\rho})$ admits a unique minimizer ρ_m .*
- (ii) *There exists a Kantorovitch potential $\bar{\varphi}$ from ρ_m to $\bar{\rho}$, such that:*

$$(15) \quad \int_{\Omega} \left(D + \frac{\bar{\varphi}}{\tau} \right) \rho \geq \int_{\Omega} \left(D + \frac{\bar{\varphi}}{\tau} \right) \rho_m \quad \text{for all } \rho \leq 1 \text{ a.e..}$$

Proof. (i) The existence of a minimizer can easily be proved using a minimizing sequence of $\phi(\rho)$. Let ρ_1, ρ_2 be two different minimizers, and \mathbf{r}_1 the optimal transport between $\bar{\rho}$ and ρ_1 . We define $\mathbf{r}_t := (1-t)\mathbf{r}_1 + t\mathbf{r}_2$ and $\rho_t := \mathbf{r}_{t\#}\rho_1$, for $t \in]0, 1[$. We know that $\rho_t = \frac{\rho}{|\det \nabla \mathbf{r}_t|} \circ (\mathbf{r}_t)^{-1}$.

As $M \mapsto (\det M)^{-1}$ is convex on the set of positive definite matrices S_d^{++} , and $\nabla \mathbf{r}_1 \in S_d^{++}$, we have :

$$\rho_t(x) \leq \left(\frac{1-t}{\det \nabla \mathbf{r}_1} + \frac{t}{\det \nabla \mathbf{r}_2} \right) \bar{\rho} \circ (\mathbf{r}_t)^{-1}(x).$$

We also know that ρ_1 and ρ_2 are admissible, therefore: $\frac{\bar{\rho}}{\det \nabla \mathbf{r}_1} \leq 1$ a.e., and we obtain: $\rho_t \leq 1$. We have then

$$\phi(\rho_t) = \int_{\Omega} D((1-t)\mathbf{r}_1(x) + t\mathbf{r}_2(x))\bar{\rho}(x) dx + \frac{1}{2\tau} W_2^2(\rho_t, \bar{\rho}).$$

Since D is λ -convex

$$D((1-t)\mathbf{r}_1(x) + t\mathbf{r}_2(x)) \leq (1-t)D(\mathbf{r}_1(x)) + tD(\mathbf{r}_2(x)) - \frac{\lambda}{2}t(1-t)|\mathbf{r}_1(x) - \mathbf{r}_2(x)|^2.$$

Moreover, $W_2^2(\cdot, \bar{\rho})$ is 1-convex along the interpolation ρ_t (see lemma 9.2.1 p. 206 in Ref [2]), therefore, for τ small enough, we have

$$\phi(\rho_t) < (1-t)\phi(\rho_1) + t\phi(\rho_2) = \inf_K \phi(\rho),$$

which is absurd.

(ii) We first assume that $\bar{\rho} > 0$ a.e., which implies that the Kantorovich potential $\bar{\varphi}$ from ρ_m to $\bar{\rho}$, satisfying $\bar{\varphi}(x_0) = 0$ (with x_0 any fixed point in Ω), is unique. Let us define a small perturbation of ρ_m : let $\rho \leq 1$ be a probability density, $\varepsilon > 0$ and $\rho_\varepsilon := \rho_m + \varepsilon(\rho - \rho_m)$. As ρ_m minimizes $\phi(\rho)$, we have:

$$(16) \quad J(\rho_\varepsilon) - J(\rho_m) + \frac{1}{2\tau}(W_2^2(\rho_\varepsilon, \bar{\rho}) - W_2^2(\rho_m, \bar{\rho})) \geq 0.$$

The first part of the left side of the inequality can easily be calculated:

$$J(\rho_\varepsilon) - J(\rho_m) = \int_{\Omega} D(x)(\rho_\varepsilon - \rho_m)(x) dx = \varepsilon \int_{\Omega} D(x)(\rho - \rho_m)(x) dx.$$

Let us estimate the second part: we denote by $(\varphi_\varepsilon, \psi_\varepsilon)$ some Kantorovich potentials associated to $\bar{\rho}$ and ρ_ε . We have

$$\begin{cases} \frac{1}{2}W_2^2(\rho_\varepsilon, \bar{\rho}) &= \int_{\Omega} \varphi_\varepsilon(x)\rho_\varepsilon(x) dx + \int_{\Omega} \psi_\varepsilon(y)\bar{\rho}(y) dy \\ \frac{1}{2}W_2^2(\rho_m, \bar{\rho}) &\geq \int_{\Omega} \varphi_\varepsilon(x)\rho_m(x) dx + \int_{\Omega} \psi_\varepsilon(y)\bar{\rho}(y) dy, \end{cases}$$

where φ_ε is a Kantorovitch potential from ρ_ε to $\bar{\rho}$. Thus:

$$\frac{1}{2}(W_2^2(\rho_\varepsilon, \bar{\rho}) - W_2^2(\rho_m, \bar{\rho})) \leq \int_{\Omega} \varphi_\varepsilon(x)(\rho_\varepsilon - \rho_m)(x) dx = \varepsilon \int_{\Omega} \varphi_\varepsilon(x)(\rho - \rho_m)(x) dx,$$

and we can deduce from inequality (16) that:

$$\int_{\Omega} D(x)(\rho - \rho_m)(x) dx + \frac{1}{\tau} \int_{\Omega} \varphi_\varepsilon(x)(\rho - \rho_m)(x) dx \geq 0 \quad \text{for all admissible } \rho.$$

Let ε tend to 0: φ_ε converges to the unique Kantorovich potential $\bar{\varphi}$ from ρ_m to $\bar{\rho}$. This gives

$$\int_{\Omega} D(x)(\rho - \rho_m)(x) dx + \frac{1}{\tau} \int_{\Omega} \bar{\varphi}(x)(\rho - \rho_m)(x) dx \geq 0 \quad \text{for all admissible } \rho.$$

We now prove the general case: let $\bar{\rho}_\delta > 0$ a.e., $\bar{\rho}_\delta \leq 1$ a.e., such that $\bar{\rho}_\delta$ converges to $\bar{\rho}$ when δ tends to 0. Using (i), there exists a unique minimizer $\rho_{m,\delta}$ of $\phi_\delta(\rho) := \int_{\Omega} D\rho + I_K + \frac{1}{2\tau}W_2^2(\rho, \bar{\rho}_\delta)$, and it converges to ρ_m as δ tends to 0. Moreover, we have proved that:

$$\int_{\Omega} D(x)(\rho - \rho_{m,\delta})(x) dx + \frac{1}{\tau} \int_{\Omega} \bar{\varphi}_\delta(x)(\rho - \rho_{m,\delta})(x) dx \geq 0 \quad \text{for all admissible } \rho,$$

with $\bar{\varphi}_\delta$ that converges to a Kantorovitch potential $\bar{\varphi}$. Taking the limit $\delta \rightarrow 0$, we obtain the desired inequality. For this kind of arguments concerning optimality for transport costs and other functionals, see for instance Ref. [10]. \square

Remark 3.2. if D is not λ -convex, we cannot prove uniqueness of the minimizer of ϕ . However, if ρ_m is a minimizer, it still satisfies inequality (ii). Indeed, in the second part of the proof of (ii), we can define $\rho_{m,\delta}$ as a minimizer of $\phi_\delta(\rho) + c_\delta W_2^2(\rho, \rho_m)$, where $c_\delta \rightarrow 0$ (so that the optimality condition that we see at the limit $\delta \rightarrow 0$ disregards this term), but slowly (so that it makes $\rho_{m,\delta}$ converge to ρ_m). Obviously this kind of argument was not necessary if one only wanted to prove this optimality condition for one minimizer ρ_m , and not for every minimizer.

Lemma 3.3. (*Decomposition of the spontaneous velocity*):

The spontaneous velocity $\mathbf{U} = -\nabla D$ can be written as follows:

$$(17) \quad \mathbf{U} = \mathbf{v}_\tau^k + \nabla p_\tau^k \quad \text{with } p_\tau^k \in H_{\rho_\tau^k}^1.$$

Proof. Using the previous lemma, we know that there exists a Kantorovich potential $\bar{\varphi}$ from ρ_τ^k to ρ_τ^{k-1} such that ρ_τ^k is a solution of the minimizing problem:

$$\rho_\tau^k \in \operatorname{argmin}_{\rho \in K} \left\{ \int_{\Omega} D(x)\rho(x) dx + \frac{1}{\tau} \int_{\Omega} \bar{\varphi}(x)\rho(x) dx \right\},$$

which imposes:

$$\begin{cases} \rho_\tau^k = 1 & \text{on } [F < l] \\ \rho_\tau^k \leq 1 & \text{on } [F = l] \\ \rho_\tau^k = 0 & \text{on } [F > l], \end{cases} \quad \text{with } F : \begin{cases} \Omega & \rightarrow \mathbb{R} \\ x & \mapsto D(x) + \frac{\bar{\varphi}(x)}{\tau}, \end{cases}$$

and $l \in \mathbb{R}$ chosen such that ρ_τ^k satisfies: $\int_{\Omega} \rho_\tau^k dx = 1$.

We can then define a pressure like function:

$$(18) \quad p_\tau^k(x) := (l - F(x))_+ = \left(l - D(x) - \frac{\bar{\varphi}(x)}{\tau} \right)_+$$

which satisfies: $p_\tau^k \geq 0$, and $p_\tau^k = 0$ on $[\rho_\tau^k < 1]$, therefore $p_\tau^k \in H_{\rho_\tau^k}^1$.

Moreover, on $[\rho_\tau^k > 0]$, we have: $\nabla p_\tau^k = -\nabla D - \frac{\nabla \bar{\varphi}}{\tau}$ (where the density vanishes \mathbf{v}_τ^k may be modified at will, so that we can keep the same formula). Since we have

$$\mathbf{v}_\tau^k = \frac{\mathbf{i} - \mathbf{t}_\tau^k}{\tau} = \frac{\nabla \bar{\varphi}}{\tau},$$

we get the desired decomposition for the spontaneous velocity : $\mathbf{U} = \mathbf{v}_\tau^k + \nabla p_\tau^k$. \square

Let us now define the densities $\tilde{\rho}_\tau(t)$ that interpolate the discrete values (ρ_τ^k) along geodesics:

$$(19) \quad \tilde{\rho}_\tau(t) = \left(\frac{t - (k-1)\tau}{\tau} (\mathbf{id} - \mathbf{t}_\tau^k) + \mathbf{t}_\tau^k \right)_{\#} \rho_\tau^k.$$

We also define $\tilde{\mathbf{v}}_\tau(t, \cdot)$ as the unique velocity field such that $\tilde{\mathbf{v}}_\tau(t, \cdot) \in \operatorname{Tan}_{\tilde{\rho}_\tau} \mathcal{P}_2(\mathbb{R}^d)$ and $(\tilde{\rho}_\tau, \tilde{\mathbf{v}}_\tau)$ satisfy the continuity equation. As before, we define: $\tilde{\mathbf{E}}_\tau = \tilde{\rho}_\tau \tilde{\mathbf{v}}_\tau$.

After these definitions we will give some a priori bounds on the curves, the pressures and the velocities that we defined. In order to get these bounds, we need to start from some estimates which are standard in the framework of Minimizing Movements. The sequence $(\rho_\tau^k)_k$ satisfies an estimate on its variation which gives a Hölder and H^1 behavior. From the minimality of ρ_τ^k , compared to ρ_τ^{k-1} , one gets

$$W_2^2(\rho_\tau^k, \rho_\tau^{k-1}) \leq 2\tau(\Phi(\rho_\tau^k) - \Phi(\rho_\tau^{k-1})).$$

Since Φ coincides with J , which is bounded, on the sequence $(\rho_\tau^k)_k$, then we have $W_2^2(\rho_\tau^k, \rho_\tau^{k-1}) \leq C\tau$ (discrete Hölder behavior), but we also have, if we sum up over k

$$(20) \quad \sum_k \tau \left(\frac{W_2(\rho_\tau^k, \rho_\tau^{k-1})}{\tau} \right)^2 \leq 2\Phi(\rho^0),$$

which is the discrete version of an H^1 estimate. As for $\tilde{\rho}_\tau(t)$, it is an absolutely continuous curve in the Wasserstein space and its velocity on the time interval $[(k-1)\tau, k\tau]$ is given by the ratio $W_2(\rho_\tau^{k-1}, \rho_\tau^k)/\tau$. Hence, the L^2 norm of its velocity on $[0, T]$ is given by

$$(21) \quad \int_0^T |\tilde{\rho}'_\tau|_{W_2}^2(t) dt = \sum_k \frac{W_2^2(\rho_\tau^k, \rho_\tau^{k-1})}{\tau},$$

and, thanks to (20), it admits a uniform bound independent of τ (here we use the notation $|\sigma'|$ for the metric derivative of a curve σ and $|\sigma'|_{W_2}(t)$ means that this metric derivative is computed according to the distance W_2). This gives compactness of the curves $\tilde{\rho}_\tau$, as well as an Hölder estimate on their variations (since $H^1 \subset C^{0,1/2}$).

Lemma 3.4. (*A priori estimates*):

We have the following a priori estimates:

- (i) \mathbf{v}_τ is τ -uniformly bounded in $L^2((0, T), L^2_{\rho_\tau}(\Omega))$.
- (ii) p_τ is τ -uniformly bounded in $L^2((0, T), H^1(\Omega))$.
- (iii) \mathbf{E}_τ and $\tilde{\mathbf{E}}_\tau$ are τ -uniformly bounded measures.

Proof. (i) We have the following equalities:

$$\begin{aligned} \int_0^T \int_\Omega \rho_\tau |\mathbf{v}_\tau|^2 &= \sum_k \int_{(k-1)\tau}^{k\tau} \int_\Omega \rho_\tau^k |\mathbf{v}_\tau^k|^2 \\ &= \sum_k \left(\int_{(k-1)\tau}^{k\tau} dt \right) \left(\int_\Omega \rho_\tau^k(x) \frac{|x - \mathbf{t}_\tau^k(x)|^2}{\tau^2} dx \right) \\ &= \sum_k \tau \frac{W_2^2(\rho_\tau^{k-1}, \rho_\tau^k)}{\tau^2} = \frac{1}{\tau} \sum_k W_2^2(\rho_\tau^{k-1}, \rho_\tau^k). \end{aligned}$$

Thanks to the general estimate (20) we get $\int_0^T \int_\Omega \rho_\tau |\mathbf{v}_\tau|^2 \leq 2\Phi(\rho^0)$.

(ii) Since we have shown the following decomposition: $\nabla p_\tau = -\nabla D - \mathbf{v}_\tau$, we have:

$$\int_0^T \int_\Omega \rho_\tau |\nabla p_\tau|^2 \leq 2 \int_0^T \int_\Omega \rho_\tau |\mathbf{v}_\tau|^2 + 2 \int_0^T \int_\Omega \rho_\tau |\nabla D|^2 \leq C.$$

But $p_\tau = 0$ on $[\rho_\tau < 1]$, therefore $\int_0^T \int_\Omega |\nabla p_\tau|^2 = \int_0^T \int_\Omega \rho_\tau |\nabla p_\tau|^2 \leq C$.

(iii) We look at $\tilde{\mathbf{E}}_\tau$ and we use the estimates (20) and (21).

$$\begin{aligned} \int_0^T \int_\Omega |\tilde{\mathbf{E}}_\tau| &= \int_0^T \int_\Omega \tilde{\rho}_\tau |\tilde{\mathbf{v}}_\tau| \leq \int_0^T \left(\int_\Omega \tilde{\rho}_\tau |\tilde{\mathbf{v}}_\tau|^2 \right)^{\frac{1}{2}} \underbrace{\left(\int_\Omega \rho_\tau \right)^{\frac{1}{2}}}_{=1} \leq \int_0^T \left(\int_\Omega \tilde{\rho}_\tau |\tilde{\mathbf{v}}_\tau|^2 \right)^{\frac{1}{2}} \\ &\leq \sqrt{T} \left(\int_0^T \int_\Omega \rho_\tau |\mathbf{v}_\tau|^2 \right)^{\frac{1}{2}} \leq C. \end{aligned}$$

Therefore, $\tilde{\mathbf{E}}_\tau$ is a τ -uniformly bounded measure. The proof for \mathbf{E}_τ is almost the same, estimating L^1 norms with L^2 norms by Cauchy-Schwartz. \square

Lemma 3.5. *Assume that μ and ν are absolutely continuous measures, whose densities are bounded by a same constant C . Then, for all function $f \in H^1(\Omega)$, we have the following inequality:*

$$(22) \quad \int_{\Omega} f d(\mu - \nu) \leq \sqrt{C} \|\nabla f\|_{L^2(\Omega)} W_2(\mu, \nu).$$

Proof. Let μ_t be the constant speed geodesic between μ and ν , and \mathbf{w}_t the velocity field such that (μ, \mathbf{w}) satisfies the continuity equation, and $\|\mathbf{w}_t\|_{L^2(\mu_t)} = W_2(\mu, \nu)$. For all t , μ_t is absolutely continuous, and its density is bounded by the same constant C a.e.. Therefore:

$$\begin{aligned} \int_{\Omega} f d(\mu - \nu) &= \int_0^1 \frac{d}{dt} \left(\int_{\Omega} f(x) d\mu_t(x) \right) dt = \int_0^1 \int_{\Omega} \nabla f \cdot \mathbf{w}_t d\mu_t dt \\ &\leq \left(\int_0^1 \int_{\Omega} |\nabla f|^2 d\mu_t dt \right)^{1/2} \left(\int_0^1 \int_{\Omega} |\mathbf{w}_t|^2 d\mu_t dt \right)^{1/2} \\ &\leq \sqrt{C} \|\nabla f\|_{L^2(\Omega)} W_2(\mu, \nu). \end{aligned}$$

□

Remark: With the same method, we can also prove:

$$\int_{\Omega} f d(\mu - \nu) \leq C^{\frac{1}{p}} \|\nabla f\|_{L^p(\Omega)} W_q(\mu, \nu)$$

for all $f \in W^{1,p}$ and q such that $\frac{1}{p} + \frac{1}{q} = 1$. More generally, if $\mu, \nu \in L^r(\Omega)$ and $\|\mu\|_{L^r}, \|\nu\|_{L^r} \leq C$, one has $\int_{\Omega} f d(\mu - \nu) \leq C^{\frac{1}{q'}} \|\nabla f\|_{L^p(\Omega)} W_q(\mu, \nu)$, provided $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1 + \frac{1}{q'}$.

3.2. Proof of the theorem in a domain with no exit.

Step 1: convergence of $(\tilde{\rho}_\tau, \tilde{\mathbf{E}}_\tau)$ and $(\rho_\tau, \mathbf{E}_\tau)$. We have proved that $\tilde{\rho}_\tau$ and $\tilde{\mathbf{E}}_\tau$ are τ -uniformly bounded measures, thus there exists (ρ, \mathbf{E}) such that $(\tilde{\rho}_\tau, \tilde{\mathbf{E}}_\tau)$ converges narrowly to (ρ, \mathbf{E}) . Let us prove that $(\rho_\tau, \mathbf{E}_\tau)$ converges to the same limit as $(\tilde{\rho}_\tau, \tilde{\mathbf{E}}_\tau)$.

We start from the ρ -part. The curves $\tilde{\rho}_\tau$ actually converge uniformly in $[0, T]$ with respect to the W_2 -distance. The curves ρ_τ and $\tilde{\rho}_\tau$ coincide on every time of the form $k\tau$. The former is constant on every interval $](k-1)\tau, k\tau]$, whereas the latter is uniformly Hölder continuous of exponent $1/2$, which implies $W_2(\tilde{\rho}_\tau(t), \rho_\tau(t)) \leq C\tau^{\frac{1}{2}}$. This proves that ρ_τ converges uniformly to the same limit as $\tilde{\rho}_\tau$.

We now consider a function $f \in C_c^\infty([0, T] \times \Omega)$, and prove that $\int_0^T \int_{\Omega} f(\tilde{\mathbf{E}}_\tau - \mathbf{E}_\tau)$ converges to 0 as τ tends to 0. We have: $\tilde{\rho}_\tau(t, \cdot) = \mathbf{T}_t \# \rho_\tau^k$ where

$$\mathbf{T}_t = (t - (k-1)\tau) \mathbf{v}_\tau^k + \mathbf{t}_\tau^k.$$

Therefore

$$\tilde{\rho}_\tau(t+h, \cdot) = (\mathbf{T}_t + h \mathbf{v}_\tau^k) \# \rho_\tau^k = ((\mathbf{id} + h \mathbf{v}_\tau^k \circ \mathbf{T}_t^{-1}) \circ \mathbf{T}_t) \# \rho_\tau^k = (\mathbf{id} + h \mathbf{v}_\tau^k \circ \mathbf{T}_t^{-1}) \# \rho_\tau(t, \cdot),$$

which implies that: $\mathbf{t}_{\tilde{\rho}_\tau(t, \cdot)}^{\tilde{\rho}_\tau(t+h, \cdot)} = \mathbf{id} + h \mathbf{v}_\tau^k \circ \mathbf{T}_t^{-1}$. We can then express $\tilde{\mathbf{v}}_\tau$ explicitly :

$$\tilde{\mathbf{v}}_\tau(t, \cdot) = \lim_{h \rightarrow 0} \frac{\mathbf{t}_{\tilde{\rho}_\tau(t, \cdot)}^{\tilde{\rho}_\tau(t+h, \cdot)} - \mathbf{id}}{h} = \lim_{h \rightarrow 0} \frac{h \mathbf{v}_\tau^k \circ \mathbf{T}_t^{-1}}{h} = \mathbf{v}_\tau^k \circ \mathbf{T}_t^{-1},$$

and obtain

$$\begin{aligned} \int_{\Omega} f(t, x) \tilde{\rho}_{\tau}(t, x) \tilde{\mathbf{v}}_{\tau}(t, x) dx &= \int_{\Omega} f(t, \mathbf{T}_{\mathbf{t}}(x)) \rho_{\tau}^k(x) \tilde{\mathbf{v}}_{\tau}(t, \mathbf{T}_{\mathbf{t}}(x)) dx \\ &= \int_{\Omega} f(t, \mathbf{T}_{\mathbf{t}}(x)) \rho_{\tau}^k(x) \mathbf{v}_{\tau}^k(x) dx. \end{aligned}$$

Hence

$$\begin{aligned} \int_0^T \int_{\Omega} f(\tilde{\mathbf{E}}_{\tau} - \mathbf{E}_{\tau}) &\leq \sum_k \int_{\tau_k}^{\tau_{k+1}} \int_{\Omega} |f(t, x) - f(t, \mathbf{T}_{\mathbf{t}}(x))| |\mathbf{v}_{\tau}^k(x)| \rho_{\tau}^k(x) dx dt \\ &\leq \sum_k \int_{\tau_k}^{\tau_{k+1}} \int_{\Omega} \text{Lip} f |x - \mathbf{T}_{\mathbf{t}}(x)| |\mathbf{v}_{\tau}^k(x)| \rho_{\tau}^k(x) dx dt \\ &\leq \sum_k \int_{\tau_k}^{\tau_{k+1}} \int_{\Omega} \text{Lip} f \tau |\mathbf{v}_{\tau}^k(x)|^2 \rho_{\tau}^k(x) dx dt \leq C \text{Lip} f \tau. \end{aligned}$$

Step 2: existence of the limit velocity. Let us prove that \mathbf{E} is absolutely continuous with respect to ρ . Let θ be a scalar measure, and \mathbf{F} a vectorial measure: the function

$$\Theta : (\theta, \mathbf{F}) \mapsto \begin{cases} \int_0^T \int_{\Omega} \frac{|\mathbf{F}|^2}{\theta} & \text{if } \mathbf{F} \ll \theta \text{ a.e. } t \in [0, T] \\ +\infty & \text{otherwise} \end{cases}$$

is l.s.c. for the weak- \star convergence of measures. Since we have shown the τ -uniform bound:

$$\int_0^T \int_{\Omega} \frac{|\mathbf{E}_{\tau}|^2}{\rho_{\tau}} \leq C,$$

we have $\Theta(\rho, \mathbf{E}) < +\infty$. Therefore \mathbf{E} is absolutely continuous with respect ρ , and there exists $\mathbf{u}(t, \cdot) \in L^2(\rho(t, \cdot))$ such that $\mathbf{E} = \rho \mathbf{u}$. Moreover, $(\rho, \rho \mathbf{u})$ satisfies the (linear) continuity equation, as limit of $(\tilde{\rho}_{\tau}, \tilde{\mathbf{E}}_{\tau})$.

Let us now prove that $\mathbf{u}(t) \in C_{\rho(t)}$. Let $t_0 \in (0, T)$, $h > 0$, and $q \in H^1_{\rho(t_0, \cdot)}$. By the continuity equation, we have

$$\int_{t_0}^{t_0+h} \int_{\Omega} \nabla q(x) \cdot \mathbf{u}_t(x) \rho(t, x) dx = \int_{\Omega} [\rho(t_0, x) - \rho(t_0 + h, x)] q(x) dx.$$

Since $\rho(t_0, \cdot) = 1$ wherever $q > 0$, and $\rho(t_0 + h, \cdot) \leq 1$ a.e., $\int_{\Omega} [\rho(t_0, x) - \rho(t_0 + h, x)] q(x) dx \leq 0$, and we have for a.e. t_0

$$\begin{aligned} 0 &\geq \frac{1}{h} \int_{t_0}^{t_0+h} \int_{\Omega} \nabla q(x) \cdot \mathbf{u}_t(x) \rho(t, x) dx \xrightarrow{h \rightarrow 0} \int_{\Omega} \nabla q(x) \cdot \mathbf{u}(t_0, x) \rho(t_0, \cdot)(x) dx \\ &= \int_{\Omega} \nabla q(x) \cdot \mathbf{u}(t_0, x) dx. \end{aligned}$$

Using the same method between $t_0 - h$ and t_0 , we also obtain the converse inequality. Finally, we have for a.e. t_0

$$(23) \quad \int_{\Omega} \nabla q(x) \cdot \mathbf{u}(t_0, x) dx = 0 \quad \text{for all } q \in H^1_{\rho(t_0, \cdot)}.$$

Step 3: the limit velocity satisfies: $\mathbf{u} = P_{C_p} \mathbf{U}$. We first prove the decomposition: $\mathbf{U} = \mathbf{u}(t, \cdot) + \nabla p(t, \cdot)$ for a.e. t . We have $\mathbf{E}_\tau = \rho_\tau \mathbf{v}_\tau = -\rho_\tau (\nabla D + \nabla p_\tau) = -\rho_\tau \nabla D - \nabla p_\tau$ since $p_\tau = 0$ on $[\rho_\tau < 1]$. Let us prove that p_τ converges to $p \in H^1_\rho$: as $p_\tau \in L^2([0, T], H^1(\Omega))$, there exists p such that p_τ weakly converges to p in $L^2([0, T], H^1(\Omega))$. We have obviously: $p \geq 0$ a.e., but it is more difficult to show that $p(t, \cdot) = 0$ on $[\rho(t) < 1]$. We consider the average functions:

$$p_\tau^{a,b} = \frac{1}{b-a} \int_a^b p_\tau(t, \cdot) dt \quad \text{and} \quad p^{a,b} = \frac{1}{b-a} \int_a^b p(t, \cdot) dt.$$

Since $p_\tau = 0$ on $[\rho_\tau < 1]$, we have

$$\begin{aligned} 0 = \int_a^b \int_\Omega p_\tau(t, x) (1 - \rho_\tau(t, x)) dx dt &= \frac{1}{b-a} \int_a^b \int_\Omega p_\tau(t, x) (1 - \rho_\tau(a, x)) dx dt \\ &\quad + \frac{1}{b-a} \int_a^b \int_\Omega p_\tau(t, x) (\rho_\tau(a, x) - \rho_\tau(t, x)) dx dt. \end{aligned}$$

The first integral reads: $\int_\Omega p_\tau^{a,b}(x) (1 - \rho_\tau(a, x)) dx \xrightarrow{\tau \rightarrow 0} \int_\Omega p^{a,b}(x) (1 - \rho(a, x)) dx$, as $p_\tau^{a,b}$ weakly converges in $H^1(\Omega)$ – therefore strongly in $L^2(\Omega)$ – to $p^{a,b}$, and $\rho_\tau(a, \cdot)$ weakly- \star converges in $L^\infty(\Omega)$ to $\rho(a, \cdot)$. Moreover, for every Lebesgue point a of $p(\cdot, x)$, we have: $p^{a,b} \xrightarrow{b \rightarrow a} p(a, \cdot)$, therefore, for all these a , we have

$$\int_\Omega p^{a,b}(x) (1 - \rho(a, x)) dx dt \xrightarrow{b \rightarrow a} \int_\Omega p(a, x) (1 - \rho(a, x)) dx.$$

Using lemma 3.5, we obtain for the second integral:

$$\begin{aligned} &\int_a^b \int_\Omega p_\tau(t, x) (\rho_\tau(a, x) - \rho_\tau(t, x)) dx dt \\ &\leq \int_a^b \|\nabla p_\tau(t, \cdot)\|_{L^2(\Omega)} \underbrace{W_2(\rho_\tau(a, \cdot), \rho_\tau(t, \cdot))}_{\leq C\sqrt{b-a}} dt \\ &\leq C\sqrt{b-a} \left(\int_a^b \|\nabla p_\tau(t, \cdot)\|_{L^2(\Omega)}^2 dt \right)^{\frac{1}{2}} \left(\int_a^b dt \right)^{\frac{1}{2}} \\ &\leq C(b-a) \left(\int_a^b \|\nabla p_\tau(t, \cdot)\|_{L^2(\Omega)}^2 dt \right)^{\frac{1}{2}}. \end{aligned}$$

As $\int_0^T \|\nabla p_\tau(t, \cdot)\|_{L^2(\Omega)}^2 dt$ is τ -uniformly bounded, $\|\nabla p_\tau(t, \cdot)\|_{L^2(\Omega)}$ weakly converges to a measure μ . Therefore, beyond a zero measure set of points a , we have

$$\lim_{\tau \rightarrow 0} \frac{1}{b-a} \int_a^b \int_\Omega p_\tau(t, x) (\rho_\tau(a, x) - \rho_\tau(t, x)) dx dt \leq C \sqrt{\mu([a, b])} \xrightarrow{b \rightarrow a} 0.$$

We finally obtain: $\int_\Omega p(a, x) (1 - \rho(a, x)) dx = 0$ for almost every a .

Hence $\mathbf{E} = -\rho \nabla D - \nabla p$, with $p = 0$ on $[\rho < 1]$, so: $\mathbf{E} = -\rho (\nabla D + \nabla p)$. Since: $\mathbf{E} = \rho \mathbf{u}$, we have shown the following decomposition:

$$\mathbf{u} = -\nabla D - \nabla p \quad \text{i.e.} \quad \mathbf{U} = \nabla p + \mathbf{u}.$$

Moreover, by Equality (23), \mathbf{u} and ∇p satisfy the complementarity relation

$$\int_{\Omega} \nabla p(t, x) \cdot \mathbf{u}(t, x) dx = 0 \quad \text{for a.e. } t,$$

which implies that we have exactly: $\mathbf{u}(t) = P_{C_{\rho(t)}} \mathbf{U}$.

4. PROOF OF THE THEOREM IN THE GENERAL CASE

We consider here the general case where $\Gamma_{out} \neq \emptyset$.

4.1. Lack of geodesic convexity. The main problem we encounter when we want to generalize the previous proof is the fact that the classical geodesics no longer belong to the admissible space K , which is no more a geodesically convex set. Indeed, if we consider a density ρ^0 which is constant equal to 1 on a subset of Ω , a measure ρ^1 which is concentrated on Γ_{out} , and the geodesic $\rho(t, \cdot)$ between them, the density of $\rho(t, \cdot)$ will be of the order of $1/(1-t)$ where it is positive, and therefore $\rho(t, \cdot) \notin K$ for all $t \in]0, 1[$.

This is one of the main reasons that prevent from using the standard theory of gradient flow for geodesically convex functionals in the Wasserstein space (see [2]).

In this section we will investigate the connectedness properties of the set K . For the sake of this work, we will see that we need to estimate the length to connect two measures in K at a very single point of the proof. Yet, we think that these estimates are interesting in themselves and this is why we try to present them so that they will be valid in any dimension d .

We define a new distance, coming from a minimal length approach, on K :

Proposition 4.1. *(Continuity of the length L) For $\mu, \nu \in K$, we define the length*

$$(24) \quad L(\mu, \nu) = \inf \left\{ \int_0^1 |\sigma'|_{W_2}(t) dt : \sigma(t) \in K, \sigma(0) = \mu, \sigma(1) = \nu \right\}.$$

This length is finite, and it is a distance on K which is continuous for the narrow convergence: if $(\mu_n), (\nu_n)$ are sequences that narrowly converge in K to μ and ν , then $L(\mu_n, \nu_n)$ converges to $L(\mu, \nu)$.

To prove this proposition, we will first analyze the case where the domain Ω is the unit cube and the door is one of the sides. We set $Q =]0, 1[^{d-1} \times]-1, 0[$, $\overline{Q} = [0, 1]^{d-1} \times [-1, 0]$ and $S = [0, 1]^{d-1} \times \{0\}$. We will still denote by K the set of admissible measures, i.e. those who are composed by a density less than 1 in Q and by a possibly singular part on S . We will denote by y the last component of a point $(x, y) \in \mathbb{R}^d = \mathbb{R}^{d-1} \times \mathbb{R}$. When integrating over S , we write dx instead of $\mathcal{H}^{d-1}(dx)$ or similar expressions.

Let us start from a simpler case.

A first useful lemma is the following:

Lemma 4.2. *Let ρ^0, ρ^1 be two probability measures on \overline{Q} of the form $\rho^i = \rho_Q^i + \rho_S^i$, where ρ_Q^i has a density on Q bounded by \overline{k} and ρ_S^i is concentrated on S . Set $\ell = W_1(\rho^0, \rho^1)$. Then, for any Lipschitz continuous function j we have*

$$\begin{aligned} \int_S j d(\rho_S^0 - \rho_S^1) &\leq \text{Lip}(j)\ell + c(\overline{k})\|j\|_{L^\infty}\ell^{1/2}, \\ \int_Q j(\rho_Q^0 - \rho_Q^1) &\leq 2\text{Lip}(j)\ell + c(\overline{k})\|j\|_{L^\infty}\ell^{1/2}. \end{aligned}$$

Proof. We start from the first estimate: consider a function $\chi_\delta : \overline{Q} \rightarrow [0, 1]$ such that $\chi_\delta = 1$ on S , $\chi_\delta = 0$ outside a strip of width δ from S , and $|\nabla \chi_\delta| \leq \delta^{-1}$ (as a matter of fact, it defines this

function as $\chi_\delta(x, y) = (1 + \delta^{-1}y)_+$. We may write

$$\int_S jd(\rho_S^0 - \rho_S^1) = \int_{\overline{Q}} j\chi_\delta d(\rho^0 - \rho^1) - \int_Q j\chi_\delta d(\rho_Q^0 - \rho_Q^1) \leq Lip(j\chi_\delta)\ell + \overline{k}\delta\|j\|_{L^\infty}.$$

Then we use $Lip(j\chi_\delta) \leq Lip(j) + \|j\|_{L^\infty}\delta^{-1}$ and we get

$$\int_S jd(\rho_S^0 - \rho_S^1) \leq \left(Lip(j) + \frac{\|j\|_{L^\infty}}{\delta} \right) \ell + \overline{k}\delta\|j\|_{L^\infty},$$

which implies, by choosing $\delta = \ell^{1/2}$,

$$\int_S jd(\rho_S^0 - \rho_S^1) \leq Lip(j)\ell + c(\overline{k})\|j\|_{L^\infty}\ell^{1/2}.$$

As far as the second estimate is concerned, just write

$$\int_Q j(\rho_Q^0 - \rho_Q^1) = \int_{\overline{Q}} j(\rho^0 - \rho^1) - \int_S jd(\rho_S^0 - \rho_S^1)$$

and use $\int_{\overline{Q}} j(\rho^0 - \rho^1) \leq Lip(j)\ell$ and the previous inequality. \square

It is important to notice in the above inequality that, once we fix ρ_Q^i or ρ_S^i , the two estimates separately make ℓ appear, where ℓ may be the W_1 distance between any pair of measures, satisfying the constraints, having ρ_Q^i or ρ_S^i as an internal or boundary part, respectively. The pair of measures we use need not to be the same in the two estimates.

Lemma 4.3. *Let $\rho^0, \rho^1 \in K$ be two admissible probability measures on \overline{Q} and $L, M \geq 1$. Suppose that ρ^0 and ρ^1 are of the following form:*

$$\rho^i = \rho_Q^i + \rho_S^i, \quad \rho_Q^i \ll \mathcal{L}^d, \quad \rho_S^i = h_i \cdot \mathcal{H}^{d-1}, \quad h_i \leq M, \quad Lip(h_i) \leq L, \quad i = 0, 1.$$

Then, there exists a curve ρ^t from ρ^0 to ρ^1 , contained in K (the set of admissible measures) and such that its W_2 -length does not exceed $C(d)M^{1/2}\sqrt{L\ell + M\ell^{1/2}}$, where we set $\ell := W_1(\rho^0, \rho^1)$.

Moreover, the same stays true if ℓ stands for a number such that there exist ‘‘extensions’’ of ρ_Q^i on S and of ρ_S^i on Q that belong to K and such that for both extensions the new W_1 -distance is smaller than ℓ (but the two extensions may be different). If instead of staying in K the constraint on the density in Q is relaxed to ‘‘being smaller than \overline{k} ’’ with $\overline{k} > 1$, the constant will also depend on \overline{k} , as in Lemma 4.2.

Proof. It is possible to replace the two probabilities on \overline{Q} with probabilities $\tilde{\rho}^i$ on $R = [0, 1]^{d-1} \times [-1, M]$ so that $\tilde{\rho}^i$ is absolutely continuous with density less than 1 and $(\pi_Q)_\# \tilde{\rho}^i = \rho^i$ (where π_Q is the projection on Q). We will take

$$\tilde{\rho}^i = \rho_Q^i + \mathbf{1}_{y < h_i(x)} \cdot \mathcal{L}^d.$$

Consider the geodesic $\tilde{\rho}^t$ from $\tilde{\rho}^0$ to $\tilde{\rho}^1$. It is a curve of measure whose length is exactly $W_2(\tilde{\rho}^0, \tilde{\rho}^1)$. Moreover, if one projects on Q all the trajectories of the particles of this curve, one gets the curve $(\pi_Q)_\# \tilde{\rho}^t$, which connects ρ^0 to ρ^1 but stays in K (since the only effect of the projection is to send all the mass on $R \setminus Q$ on S , while the densities inside Q are not affected). And we know that the densities of $\tilde{\rho}^t$ will not be larger than 1, since this is the case for a geodesic between two measures with densities bounded by 1.

Hence we only need to estimate $W_2(\tilde{\rho}^0, \tilde{\rho}^1)$. For simplicity, let us estimate W_1 instead of W_2 . We will conclude by the inequality $W_2 \leq (\text{diam } R)^{1/2}W_1^{1/2}$. Notice that the diameter of R is $\sqrt{(M+1)^2 + d-1} \leq C(d)M$.

To estimate W_1 , take a function $f \in Lip_1(R)$. What follows will be easier to justify in case f is regular but everything will work (by density, or instance), for any f whose Lipschitz constant does not exceed 1. Let us define, for $x \in [0, 1]^{d-1}$ and $a, b \in [0, M]$,

$$g(x, a, b) = \frac{\int_a^b f(x, t) dt}{b - a}.$$

We denote by g_x , g_a and g_b the partial derivatives of g . We can verify that

$$|g_x(x, a, b)| = \frac{\left| \int_a^b f_x(x, t) dt \right|}{|b - a|} \leq Lip(f) = 1,$$

then we compute g_b and we get

$$|g_b(x, a, b)| = \left| \frac{f(x, b)}{b - a} - \frac{g(x, a, b)}{b - a} \right| \leq \frac{Lip(f)}{|b - a|} |b - a| = 1,$$

and, analogously, $|g_a| \leq 1$.

In particular, if one takes two Lipschitz functions $a(x)$ and $b(x)$, one has $Lip(g(x, a(x), b(x))) \leq 1 + Lip(a) + Lip(b)$.

Now we write

$$\int_R f d(\tilde{\rho}^0 - \tilde{\rho}^1) = \int_Q f d(\rho_Q^0 - \rho_Q^1) + \int_S \left(\int_0^{h_0(x)} f(x, t) dt - \int_0^{h_1(x)} f(x, t) dt \right) dx.$$

We estimate both terms thanks to the previous lemma. The first term in the right hand side is less than ℓ , while for the second we may write

$$\int_S \left(\int_0^{h_0(x)} f(x, t) dt - \int_0^{h_1(x)} f(x, t) dt \right) dx = \int_S g(x, h_0(x), h_1(x)) (h_0 - h_1)(x) dx.$$

Hence we are in the case of the previous lemma with $j(x) = g(x, h_0(x), h_1(x))$, and hence $Lip(j) \leq 1 + 2L$ and $\|j\|_{L^\infty} \leq M + \sqrt{d}$ (the first estimate comes from our study of g , for the second just suppose that f vanishes somewhere on S).

Hence we get, using the arbitrariness of the function f

$$W_1(\tilde{\rho}^0, \tilde{\rho}^1) \leq \ell + (1 + 2L)\ell + 2M\ell^{1/2}.$$

To simplify the computations we use $1 \leq M, L$ and get

$$W_2(\tilde{\rho}^0, \tilde{\rho}^1) \leq C(d)M^{1/2}W_1(\tilde{\rho}^0, \tilde{\rho}^1)^{1/2} \leq C(d)M^{1/2}\sqrt{L\ell + M\ell^{1/2}}.$$

The last part of the statement is an easy consequence of the technique we used and of Lemma 4.2. \square

Theorem 4.4. *Let μ_0 and μ_1 be two probabilities in K . Then there exists a curve $(\mu_t)_t$ connecting μ_0 to μ_1 , such that its W_2 -length does not exceed $C(d)W_1(\mu_0, \mu_1)^{1/(4d)}$ and that $\mu_t \in K$ for every t .*

Proof. Take $\varepsilon > 0$ and modify μ^i into a new measure $\rho^i \in K$ by regularizing in the direction of x : it is sufficient to take the convolution of μ_S^i with a kernel of the form $C(\varepsilon^{1-d} - \varepsilon^{-d}|x|)_+$. This ensures that the W_1 distance has not increased and that the new measures on S will have Lipschitz and bounded densities on S , with $M \leq C\varepsilon^{1-d}$ and $L \leq C\varepsilon^{-d}$, and on Q the constraint is kept as well. Yet, there is a problem: these measures may exit the domain. There are two possible ways for solving this problem, and both will be useful.

One possibility is rescaling of a factor $(1 + 2\varepsilon)^{-1}$, so that all the mass is pushed again into the domain. This does not change significantly the values of M and L but the densities inside will be no more bounded by 1. They will be bounded by a constant \bar{k} close to 1. In this case too the Wasserstein distance has not increased, since the rescaling was a contraction.

The other possibility is composing with a contracting transport $T : \overline{Q}_\varepsilon \rightarrow \overline{Q}$ (\overline{Q}_ε being the ε -neighborhood of \overline{Q}), which is chosen so that the convolution of the constant function 1 is sent onto the constant function 1 (this is possible thanks to the fact the convolution keeps the mass unchanged). This construction ensures that the constraint inside Q will be satisfied but unluckily, since the inverse of T is not Lipschitz continuous (due to the fact that the densities vanished on the boundary of \overline{Q}_ε), it is not suitable for S . Anyway, in this case too, the Wasserstein distance was not increased.

Hence we do a mixed procedure: we use the second possibility in Q and the first on S . It is clear that in this way we have good densities both in Q and on S , and we can apply Lemma 4.2 and the last statement of Lemma 4.3. Notice that the W_1 -distance between the two measures $\rho^i \in K$ that we constructed could be larger than ℓ . It is easily estimated by something like $\ell + \varepsilon$ but this is not sufficient for the following estimates.

Now, to connect μ^0 to μ^1 , one can first connect each μ^i to ρ^i , and the cost is no more than ε , since it is sufficient to spread every particle on a ball of radius ε (i.e. the radius of the support of the previous kernel) when we do convolution, and then to move it no more than ε when we compose with a contraction. After that, one uses the previous Lemma to estimate the length for connecting ρ^0 to ρ^1 and gets

$$\min \text{length} \leq 2\varepsilon + C(d)\varepsilon^{(1-d)/2} \sqrt{\varepsilon^{-d}\ell + \varepsilon^{1-d}\ell^{1/2}} = 2\varepsilon + C(d)\varepsilon^{\frac{3}{2}-d} \sqrt{\frac{\ell}{\varepsilon^2} + \frac{\ell^{1/2}}{\varepsilon}},$$

where ℓ denotes the W_1 distance between μ^0 and μ^1 . If one supposes that ε is chosen so that $\ell\varepsilon^{-2}$ is smaller than 1, one can estimate the last sum in the square root and get

$$\min \text{length} \leq 2\varepsilon + C(d)\varepsilon^{1-d}\ell^{1/4}.$$

Choosing $\varepsilon = \ell^{1/(4d)}$ gives at the same time that $\ell\varepsilon^{-2}$ is small and that the minimal length may be estimated by $\ell^{1/(4d)}$. \square

To approach the general case one needs to use the following theorem, which has already been used in a transport-related setting with density constraints in the variational theory of incompressible Euler equations by Y. Brenier and provides a useful tool for reducing to the cube (see Refs [16] and [40] for the applications to fluid mechanics).

Theorem 4.5. *For any sufficiently good domain $\Omega \subset \mathbb{R}^d$ which is homeomorphic to the cube, there exists a bi-lipschitz homeomorphism $\phi : \overline{\Omega} \rightarrow \overline{Q}$ such that $\phi_{\#}(\mathcal{L}_{|\Omega}^d) = c\mathcal{L}_{|Q}^d$. Moreover, the behavior of ϕ on the boundary may be prescribed at will.*

4.2. Generalization of the technical lemmas. In this section, we briefly explain how to generalize the technical lemmas we used in the first proof (with $\Gamma_{out} = \emptyset$). Conditions on the minimizer for the discrete problem.

- First of all, in lemma 3.1, we can't prove the uniqueness of the minimizer ρ_m with the same method. Indeed, exactly as we explained in the previous section for geodesics, the interpolation ρ_t between two possible minimizers does not necessarily belong to K . Therefore, we will have to apply the "selection" method explained in Remark 3.2 in order to prove inequality (15). More precisely, the case where $\bar{\rho} > 0$ remains unchanged, but in the general case, we fix a minimizer ρ_m of ϕ , and we define $\rho_{m,\delta}$ as a minimizer of

$$\phi_\delta(\rho) := \int_{\Omega} D\rho + I_K(\rho) + \frac{1}{2\tau} W_2^2(\rho, \bar{\rho}_\delta) + c_\delta W_2^2(\rho, \rho_m),$$

with c_δ that converges to 0 slower than $W_2(\bar{\rho}, \bar{\rho}_\delta)$. Since ρ_m and $\rho_{m,\delta}$ are minimizers of J and ϕ_δ , we have the following inequalities:

$$\begin{cases} \int_{\Omega} D\rho_{m,\delta} + \frac{1}{2\tau} W_2^2(\rho_{m,\delta}, \bar{\rho}_\delta) + c_\delta W_2^2(\rho, \rho_m) & \leq \int_{\Omega} D\rho_m + \frac{1}{2\tau} W_2^2(\rho_m, \bar{\rho}_\delta) \\ \int_{\Omega} D\rho_m + \frac{1}{2\tau} W_2^2(\rho_m, \bar{\rho}) & \leq \int_{\Omega} D\rho_{m,\delta} + \frac{1}{2\tau} W_2^2(\rho_{m,\delta}, \bar{\rho}) \end{cases}$$

which implies, using the triangular inequality:

$$\begin{aligned} W_2^2(\rho_{m,\delta}, \rho_m) &\leq \frac{1}{2\tau c_\delta} [W_2^2(\rho_{m,\delta}, \bar{\rho}) - W_2^2(\rho_{m,\delta}, \bar{\rho}_\delta) + W_2^2(\rho_m, \bar{\rho}_\delta) - W_2^2(\rho_m, \bar{\rho})] \\ &\leq \frac{C}{2\tau c_\delta} W_2(\bar{\rho}, \bar{\rho}_\delta) \xrightarrow{\delta \rightarrow 0} 0. \end{aligned}$$

Therefore, $\rho_{m,\delta}$ converges to the fixed minimizer ρ_m . We can then pass to the limit $\delta \rightarrow 0$, and obtain the same inequality for ρ_m .

- It is also useful to notice another feature of the problem with an exit Γ_{out} : once some mass arrives to the exit, it does not move anymore. This precisely means the following: if γ_τ^k is an optimal transport plan from the selected measure ρ_τ^k to the previous one, ρ_τ^{k-1} , and $(x, y) \in \text{supp}(\gamma_\tau^k)$ with $y \in \Gamma_{out}$, then $y = x$. This means that all the mass which was already on the door for ρ_τ^{k-1} will not move. To prove it, it is sufficient to consider the map $F : \bar{\Omega} \times \bar{\Omega} \rightarrow \bar{\Omega} \times \bar{\Omega}$ defined by $F(x, y) = (y, y)$ if $y \in \Gamma_{out}$ and $F(x, y) = (x, y)$ if $y \notin \Gamma_{out}$. The measure $F_\# \gamma_\tau^k$ is a transport plan between a new measure ρ and ρ_τ^{k-1} , which reduces the transport cost and the functional J (since D is minimal on the exit). Moreover, since ρ is obtained from ρ_τ^k by moving some mass onto the door, we have $\rho \in K$ as well. This would contradict the optimality of ρ_τ^k unless $F_\# \gamma_\tau^k = \gamma_\tau^k$, which is the thesis.

This also proves uniqueness of the optimal transport plan between ρ_τ^k and ρ_τ^{k-1} since, if we look it the other way around (from ρ_τ^{k-1} to ρ_τ^k), we can decompose the problem in one part which will not move (corresponding to $\rho_\tau^{k-1} \mathbf{1}_{\Gamma_{out}}$) and one part which is the transport of an absolutely continuous density ($\rho_\tau^{k-1} \mathbf{1}_{\bar{\Omega}}$).

We will also denote by E_τ^k the excess mass of ρ_τ^k with respect to ρ_τ^{k-1} on the exit, i.e. $E_\tau^k := \rho_\tau^k(\Gamma_{out}) - \rho_\tau^{k-1}(\Gamma_{out}) \geq 0$.

- In lemma 3.3, the solution of

$$\rho_\tau^k \in \operatorname{argmin}_{\rho \in K} \left\{ \int_{\Omega} D(x)\rho(x) dx + \frac{1}{\tau} \int_{\Omega} \bar{\varphi}(x)\rho(x) dx \right\}$$

is not necessarily the same in the general case, as there exists no limit density on Γ_{out} . Let us define: $l := \inf_{x \in \Gamma_{out}} F(x)$, and $\Gamma_{min} = \{x \in \Gamma_{out} : F(x) = l\}$. If $|[F < l]| \geq 1$, then the solution is the same as in the previous proof. However, if $|[F < l]| < 1$, it costs less to put a part of the density onto Γ_{out} . The solution is therefore given by:

$$\begin{cases} \rho_\tau^k = 1 & \text{on } [F < l], \\ \rho_\tau^k > 0 & \text{on } \Gamma_{min}, \text{ with } \rho_\tau^k(\Gamma_{min}) = 1 - |[F < l]|, \\ \rho_\tau^k \leq 1 & \text{on } [F = l] \setminus \Gamma_{min}, \\ \rho_\tau^k = 0 & \text{on } [F > l]. \end{cases}$$

The pressure p_τ^k defined by

$$p_\tau^k(x) := (l - F(x))_+ = \left(l - D(x) - \frac{\bar{\varphi}(x)}{\tau} \right)_+$$

then belongs to $H_{\rho_\tau^k}^1$, and we prove the decomposition $\mathbf{U} = \mathbf{v}_\tau^k + \nabla p_\tau^k$ as before.

In order to prove the a priori estimates of lemma 3.4, we have to take into account the singularity part of the densities on Γ_{out} . Notice that, to avoid any ambiguity where the transport does not exist, we only defined a discrete velocity vector field inside $\mathring{\Omega}$. To be clearer, we want to spend some disambiguation words on what $\tilde{\mathbf{E}}_\tau$ and \mathbf{E}_τ are in this case.

- The measure $\tilde{\mathbf{E}}_\tau$ is as usual defined as the vector measure satisfying the continuity equation with the curve $\tilde{\rho}_\tau$. We also have an explicit formula in terms of the optimal transport plans γ_τ^k from ρ_τ^k to ρ_τ^{k-1} : for any $t \in [k - 1\tau, k\tau[$ take

$$\tilde{\mathbf{E}}_\tau(t) := (\pi_{(k\tau-t)/\tau})_{\#} \left(\frac{x-y}{\tau} \cdot \gamma_\tau^k \right),$$

where $\pi_s(x, y) = (1-s)x + sy$.

- The measure \mathbf{E}_τ is simply defined as the product of $\rho_\tau \mathbf{1}_{\mathring{\Omega}}$ times the velocity vector field defined in Section 2.3, on the non-singular part only (again we use $\mathring{\Omega}$ instead of Ω to stress that the boundary is excluded). As before, the idea is that this vector measure satisfies good properties from optimality conditions, while the previous one satisfies the continuity equation. We need to compare them.
- There is also in this case a third vector measure, that we can call $\hat{\mathbf{E}}_\tau$, which is defined exactly as $\tilde{\mathbf{E}}_\tau$ but ignoring the part on Γ_{out} :

$$\hat{\mathbf{E}}_\tau(t) := (\pi_{(k\tau-t)/\tau})_{\#} \left(\frac{x-y}{\tau} \cdot \mathbf{1}_{x \in \mathring{\Omega}} \gamma_\tau^k \right).$$

The utility of $\hat{\mathbf{E}}_\tau$ is that it is more easily comparable to \mathbf{E}_τ .

We come back to the proof of lemma 3.4: as a matter of fact, we now have:

$$W_2^2(\rho_\tau^{k-1}, \rho_\tau^k) = \tau^2 \int_{\Omega} \rho_\tau^k |\mathbf{v}_\tau^k|^2 + \int_{\Gamma_{out} \times \Omega} |x-y|^2 d\gamma_\tau^k(x, y),$$

where γ_τ^k is the optimal transport plan between ρ_τ^k and ρ_τ^{k-1} . Therefore, we have

$$\int_{\Omega} \rho_\tau |\mathbf{v}_\tau|^2 \leq \tau^{-2} W_2^2(\rho_\tau^{k-1}, \rho_\tau^k),$$

and the a priori estimates (i) and (ii) are still satisfied. The proof of (iii) is unchanged, but let us remark that we have no longer the equality:

$$\int_{\Omega} \tilde{\rho}_\tau^k |\tilde{\mathbf{v}}_\tau^k|^2 = \int_{\Omega} \rho_\tau^k |\mathbf{v}_\tau^k|^2,$$

and that the geodesic $\tilde{\rho}_\tau$ does not belong to K .

Lemma 3.5 is no longer true for densities that are not absolutely continuous with respect to the Lebesgue measure. Indeed, as we have seen before, the geodesic between two densities of K does not belong to K . We prove instead the following lemma:

Lemma 4.6. *Let $\mu, \nu \in K$. Then, for all function $f \in H^1$ with $f = 0$ on Γ_{out} , we have the following inequality:*

$$\int_{\Omega} f d(\mu - \nu) \leq \|\nabla f\|_{L^2(\Omega)} L(\mu, \nu)$$

where $L(\mu, \nu)$ is the length of the shortest path in K joining μ and ν (see (24)).

Proof. The proof is an adaptation of the one of Lemma 3.5 : let σ_t be a minimal length curve in K joining μ and ν , and let \mathbf{w}_t such that (σ, \mathbf{w}) satisfies the continuity equation and $\|\mathbf{w}_t\|_{L^2(\sigma_t)} = L(\mu, \nu)$. Since $f \in H^1$ with $f = 0$ on Γ_{out} , then ∇f does not see the part of $\sigma \mathbf{w}$ on the boundary, so that we have:

$$\begin{aligned} \int_{\Omega} f d(\mu - \nu) &= \int_0^1 \frac{d}{dt} \left(\int_{\Omega} f d\sigma_t \right) = \int_0^1 \int_{\dot{\Omega}} \nabla f \cdot \mathbf{w}_t d\sigma_t dt \\ &\leq \left(\int_0^1 \int_{\dot{\Omega}} |\nabla f|^2 d\sigma_t dt \right)^{1/2} \left(\int_0^1 \int_{\dot{\Omega}} |\mathbf{w}_t|^2 d\sigma_t dt \right)^{1/2} \\ &\leq \|\nabla f\|_{L^2(\Omega)} L(\mu, \nu) \end{aligned}$$

since $\sigma_t \leq 1$ in $\dot{\Omega}$. \square

4.3. Generalization of the proof. At step 1, we need again to prove that the limits of $\tilde{\mathbf{E}}_{\tau}$ and \mathbf{E}_{τ} are the same. As far as the limits of $\tilde{\rho}_{\tau}$ et ρ_{τ} are concerned, everything works as in Section 3.2: this also proves that the limit curve ρ belongs to K , since this is the case for ρ_{τ} (but not for $\tilde{\rho}_{\tau}$).

It is easy to check that the comparison we did in Step 1 of Section 3.2 may be performed again so as to obtain that the limit of $\tilde{\mathbf{E}}_{\tau}$ and \mathbf{E}_{τ} are the same. What we need to do now is proving that the limit of $\hat{\mathbf{E}}_{\tau}$ and $\tilde{\mathbf{E}}_{\tau}$ are the same. We will prove that the mass of $\tilde{\mathbf{E}}_{\tau} - \hat{\mathbf{E}}_{\tau}$ is negligible, i.e. that

$$\int_0^T dt \int_{\bar{\Omega}} d|\tilde{\mathbf{E}}_{\tau}(t) - \hat{\mathbf{E}}_{\tau}(t)| \xrightarrow{\tau \rightarrow 0} 0.$$

To do this, it is sufficient to estimate

$$\sum_{k=0}^{T/\tau} \int_{(k-1)\tau}^{k\tau} dt \int_{\bar{\Omega} \times \bar{\Omega}} \frac{|x-y|}{\tau} \mathbf{1}_{\Gamma_{out} \times \bar{\Omega}} d\gamma_{\tau}^k = \sum_{k=0}^{T/\tau} \int_{\bar{\Omega} \times \bar{\Omega}} |x-y| \mathbf{1}_{\Gamma_{out} \times \bar{\Omega}} d\gamma_{\tau}^k.$$

Thanks to what we underlined before, namely that the mass which is on Γ_{out} does not move any more, we know that $|x-y| \mathbf{1}_{\Gamma_{out} \times \bar{\Omega}} d\gamma_{\tau}^k = |x-y| \mathbf{1}_{\Gamma_{out} \times \dot{\Omega}} d\gamma_{\tau}^k$ and the mass of $\mathbf{1}_{\Gamma_{out} \times \dot{\Omega}} d\gamma_{\tau}^k$ is exactly the excess mass E_{τ}^k . Thanks to the Lemma 4.7 below, we can go on and obtain

$$\begin{aligned} \sum_{k=0}^{T/\tau} \int_{\bar{\Omega} \times \bar{\Omega}} |x-y| \mathbf{1}_{\Gamma_{out} \times \bar{\Omega}} d\gamma_{\tau}^k &= \sum_{k=0}^{T/\tau} \int_{\bar{\Omega} \times \bar{\Omega}} |x-y| \mathbf{1}_{\Gamma_{out} \times \dot{\Omega}} d\gamma_{\tau}^k \\ &\leq \sum_{k=0}^{T/\tau} \left(\int_{\Gamma_{out} \times \dot{\Omega}} |x-y|^2 d\gamma_{\tau}^k \right)^{\frac{1}{2}} \left(\int_{\Gamma_{out} \times \dot{\Omega}} d\gamma_{\tau}^k \right)^{\frac{1}{2}} \leq \sum_{k=0}^{T/\tau} W_2(\rho_{\tau}^{k-1}, \rho_{\tau}^k)^{4/3} \\ &\leq \left(\sum_{k=1}^{E(T/\tau)} W_2(\rho_{\tau}^{k-1}, \rho_{\tau}^k)^2 \right)^{\frac{2}{3}} \left(\sum_{k=1}^{E(T/\tau)} 1 \right)^{\frac{1}{3}} \leq (C\tau)^{\frac{2}{3}} \left(\frac{T}{\tau} \right)^{\frac{1}{3}} = C \tau^{\frac{1}{3}} \xrightarrow{\tau \rightarrow 0} 0. \end{aligned}$$

Lemma 4.7. *Suppose $\mu, \nu \in K$ and set $E := |\mu(\Gamma_{out}) - \nu(\Gamma_{out})|$. Then we have $E \leq CW_2^{2/3}(\mu, \nu)$, where the constant C depends on the geometry of Ω and Γ_{out} .*

Proof. Suppose for simplicity $\nu(\Gamma_{out}) \geq \mu(\Gamma_{out})$. Take an optimal transport plan γ from μ to ν . Consider $\gamma' = \mathbf{1}_{\dot{\Omega} \times \Gamma_{out}} \gamma$. The mass of γ' is a number E' , larger than E . Let μ' be the projection of γ' on the first variable (x): it is a measure with mass E' , dominated by $\mathbf{1}_{\dot{\Omega}} \mu$ (and hence it is absolutely continuous with density smaller than 1). We have

$$W_2^2(\mu, \nu) = \int |x-y|^2 d\gamma \geq \int |x-y|^2 d\gamma' \geq \int d(x, \Gamma_{out})^2 d\gamma' = \int d(x, \Gamma_{out})^2 d\mu'.$$

It is sufficient to prove that this last integral is larger than $c(E')^3$. Set $d(x) := d(x, \Gamma_{out})$: we will use the fact that $||d \leq t|| \leq ct$. We have

$$\begin{aligned} \int d(x)^2 d\mu' &= \int_0^\infty \mu'([d^2 > t]) dt = \int_0^\infty (E' - \mu'([d \leq \sqrt{t}])) dt \\ &\geq \int_0^\infty (E' - |[d \leq \sqrt{t}]|) dt \geq \int_0^{(E'/c)^2} (E' - c\sqrt{t}) dt = c(E')^3. \end{aligned}$$

□

At step 2, we prove with the same method that \mathbf{E} is absolutely continuous with respect to the density $\bar{\rho} := \lim_{\tau \rightarrow 0} (\rho_\tau)_\Omega = \rho_\Omega$ (the decomposition of the measures into a part on Γ_{out} and a part on $\mathring{\Omega}$ passes to the limit, because of the density bound on $\mathring{\Omega}$): there exists \mathbf{u} such that $\mathbf{E} = \rho_\Omega \mathbf{u}$. Moreover (ρ, \mathbf{E}) satisfies the continuity equation, and we can prove again the equality

$$\int_\Omega \nabla q \cdot \mathbf{u} dx = 0 \quad \forall q \in H_\rho^1.$$

At step 3, the first estimates are still true, since we integrate over $\mathring{\Omega}$ ($p_\tau = 0$ on Γ_{out}). However, we can't use lemma 3.5 anymore. Instead, we apply lemma 4.6 and get the inequality

$$\int_a^b \int_\Omega p_\tau(t, x) (\rho_\tau(a, x) - \rho_\tau(t, x)) dx dt \leq \int_a^b \|\nabla p_\tau(t, \cdot)\|_{L^2(\Omega)} L(\rho_\tau(a, \cdot), \rho_\tau(t, \cdot)) dt.$$

Using proposition (4.1) and the same notation as in Section 3.2, step 3, the limit $\tau \rightarrow 0$ reads:

$$\begin{aligned} \lim_{\tau \rightarrow 0} \frac{1}{b-a} \int_a^b \int_\Omega p_\tau(t, x) (\rho_\tau(a, x) - \rho_\tau(t, x)) dx dt \\ \leq \frac{1}{b-a} \sqrt{\mu([a, b])} \left(\int_a^b L(\rho(a), \rho(t))^2 dt \right)^{\frac{1}{2}}. \end{aligned}$$

Since at the limit, the curve $\rho(t)$ belongs to K for every t , we have also:

$$L(\rho(a), \rho(t)) dt \leq \int_a^t |\rho'|_{W_2}(s) ds \leq \left(\int_a^t |\rho'|_{W_2}^2(s) ds \right)^{1/2} (t-a)^{1/2} \leq C(b-a)^{1/2}.$$

Therefore, we have the following inequality:

$$\begin{aligned} \lim_{\tau \rightarrow 0} \frac{1}{b-a} \int_a^b \int_\Omega p_\tau(t, x) (\rho_\tau(a, x) - \rho_\tau(t, x)) dx dt &\leq \frac{1}{b-a} \sqrt{\mu([a, b])} \left(\int_a^b C(b-a) dt \right)^{\frac{1}{2}} \\ &= C \sqrt{\mu([a, b])} \xrightarrow{b \rightarrow a} 0 \text{ for a.e. } a, \end{aligned}$$

and we conclude the proof as in the particular case $\Gamma_{out} = \emptyset$.

5. ILLUSTRATION: A CONVERGENT CORRIDOR

We present here an example where both the transport equation and discrete process of the gradient-flow problem can be solved quasi-explicitly. We also give numerical estimations on the convergence of the discrete scheme to the solution of the continuity equation.

We want to model the displacement of a crowd through a convergent corridor. We thus take for Ω a portion of a cone, expressed in polar coordinates as $[r \in [a, R], \theta \in [-\alpha, \alpha]]$ (see fig 2), with a possible ‘‘exit’’ $\Gamma_{out} = \{a\} \times [-\alpha, \alpha]$, and we take for D the distance to the exit (or to the apex, which is equivalent): $D(r) = r$. We assume that the initial density is uniform: $\rho^0(r) = \rho^0 < 1$. We will consider in this section two examples: the case $a = 0$ with no exit (so that people will in the end concentrate on the neighborhood of the vertex) and the case $a > 0$ with exit.




FIGURE 2. Modeling of the displacement of a crowd through a convergent corridor.

Thanks to the symmetry of this problem, the minimizing movement scheme can be written as a minimization problem on the transport function:

$$\left\{ \begin{array}{l} \rho_\tau^0 = \rho^0 \\ \rho_\tau^k = \mathbf{s}_{\mathbf{k} \# \rho_\tau^{k-1}}, \mathbf{s}_{\mathbf{k}} \in \operatorname{argmin}_{\mathbf{t} \# \rho_\tau^{k-1} \in K} \left\{ \int_a^R \left(D(\mathbf{t}(r)) + \frac{1}{2\tau} |r - \mathbf{t}(r)|^2 \right) \rho_\tau^{k-1}(r) r \, dr \right\} \end{array} \right\}.$$

Let us first consider the case where $a = 0$ (and $\Gamma_{out} = \emptyset$), where this problem can be explicitly solved: ρ_τ^k is given by

$$(25) \quad \rho_\tau^k(r) = \begin{cases} 1 & \text{on } [a, b_\tau^k[\\ \rho^0 \left(1 + \frac{k\tau}{r} \right) & \text{on } [b_\tau^k, R - k\tau[\\ 0 & \text{on } [R - k\tau, R], \end{cases}$$

where b_τ^k satisfies the recurrence relation

$$(26) \quad \begin{cases} b_\tau^0 = 0 \\ (b_\tau^k)^2 - \rho^0 (b_\tau^k + k\tau)^2 = (b_\tau^{k-1})^2 - \rho^0 (b_\tau^{k-1} + (k-1)\tau)^2, \end{cases}$$

and the solution of the continuity equation can be easily calculated:

$$\rho(t, r) = \begin{cases} 1 & \text{if } r \in [a, b(t)[\\ \rho^0 \left(1 + \frac{t}{r} \right) & \text{if } r \in [b(t), R - t] \\ 0 & \text{if } r \in [R - t, R], \end{cases} \quad \text{where } \begin{cases} b(0) = 0 \\ b'(t) = \rho^0 \frac{b(t) + t}{b(t) - \rho^0 (b(t) + t)}. \end{cases}$$

In figure 3, we represent the discrete densities ρ_τ^k at different times for the numerical values $\tau = 0.01$, $a = 0$, $R = 10$, and $\rho^0 = 0.4$. Let us remark that the recurrence relation that satisfies b_τ^k is a numerical scheme for the ODE on $b(t)$. Indeed, it writes

$$\frac{b_\tau^k - b_\tau^{k-1}}{\tau} = F \left(\frac{b_\tau^k + b_\tau^{k-1}}{2}, \frac{2k-1}{2}\tau \right), \quad \text{where } F(r, t) = \rho^0 \frac{b(t) + t}{b(t) - \rho^0 (b(t) + t)}.$$

Using the conservation of the total amount of people, it is easy to prove that this scheme is exact at every time step $k\tau$, and so is the discrete solution ρ_τ^k .




FIGURE 3. Evolution of the solution of the minimizing movement scheme in the case where $\Gamma_{out} = \emptyset$.

We now consider the case $a > 0$ with exit. The densities have then the same form, except that the evolution of the interface in the continuous case is now given by

$$\begin{cases} b'(t) = \Phi(t, b(t)) \\ b(t_0) = a, \end{cases} \quad \text{with : } \Phi(t, r) = \begin{cases} \frac{\rho_0 \left(1 + \frac{t}{r}\right) - \frac{r-a}{r \ln(r/a)}}{1 - \rho_0 \left(1 + \frac{t}{r}\right)} & \text{if } r \leq R - t \\ -\frac{r-a}{r \ln(r/a)} & \text{if } r > R - t, \end{cases}$$

whereas in the discrete case, b_τ^k satisfies now the recurrence relation

$$(b_\tau^k)^2 - a^2 - \rho_0(b_\tau^k + k\tau)^2 = (b_\tau^{k-1})^2 - r_e^2 - \rho_0(b_\tau^{k-1} + (k-1)\tau)^2$$

if $b_\tau^{k-1} < R - (k-1)\tau$, and

$$(b_\tau^k)^2 - a^2 = (b_\tau^{k-1})^2 - r_e^2$$

if $b_\tau^{k-1} \geq R - (k-1)\tau$, where r_e is the (unknown) radius such that people who were between a and r_e at step $k-1$ will exit the corridor (i.e. arrive at a) at step k . This radius is given as the minimum of an integral expression that we will not develop here. In figure 4, we represent the discrete densities for $a = 1$ and for the same numerical values as before.

It is also interesting to estimate numerically the error between the solution of the continuity equation and the solution of the minimizing movement scheme. In this purpose, we consider the case where the density is initially saturated ($\rho^0 = 1$), and we compute b and b_τ with high accuracy (high order method for the ODE on b , and precise quadrature and optimization methods to estimate r_e and b_τ), so that space discretization does not affect error estimation. We obtain numerically that b_τ converges to b when τ tends to 0 with an error of order 1 (interpolation polynomial of $\tau \mapsto |b(T) - b_\tau(T)|$ gives order 0.989 for $T = 1$), which gives also an order 1 error for the Wasserstein distance between ρ and ρ_τ .

6. MODELLING ISSUES, EXTENSIONS

We would like to conclude this paper by some remarks on the limitations of the overall approach in terms of modelling, and on possible extensions to other domains.

With its very macroscopic and eulerian nature, the model is designed to handle large populations as a whole, and does not allow to localize and follow in their path individual pedestrians. As a direct consequence, the spontaneous (or desired) velocity of an individual may depend on its location only, so that differentiated individual strategies (e.g. avoidance of crowded zone,




FIGURE 4. Evolution of the solution of the minimizing movement scheme in the case with exit.

skirting of obstacles) cannot be included straightforwardly. Besides, the macroscopic expression of the non-overlapping constraint is less restrictive than its microscopic counterpart. In the microscopic setting (people are identified with rigid discs), for highly packed situations, the non-overlapping constraints induce some kind of non-negative divergence constraint in the directions of contacts. Consider the example of a cartesian distribution of monodisperse discs (see Fig. 5, left), with a uniform spontaneous velocity directed toward a wall. The actual velocity will be 0, whereas in the macroscopic version it is not (see Fig. 5, center). Note that, if the microscopic distribution is modified, while mean density is preserved, the situation is no longer static (see Fig. 5, right). This example illustrates the deep difference between the microscopic approach (for which local structure and orientation of contacts lines play an essential role), and the macroscopic one, which only considers local density.




FIGURE 5. Differences between micro and macro approaches

As a consequence the model is unlikely to reproduce the formation of blocked arches near an exit, which are observed in practice in highly critical emergency situations, and which can be recovered by the microscopic model, even without friction. Note also that, together with the structural anisotropy we just mentioned, individual anisotropy of pedestrians is not handled (whereas it may be in microscopic models by replacing discs by ellipses, for example).

Yet, despite these limitations in the modelling of crowd motion, we believe that this new type of evolution problem may be fruitful to model phenomena, in particular in the domain

of cell dynamics. In this context, the spontaneous velocity would be replaced by some kind of chemotaxis velocity. As an illustration, let us express what could be called a unilateral version of Keller-Segel equation, in the spirit of what has been presented here:

$$\begin{aligned} \frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) &= 0 \\ \mathbf{u} &= P_{C_\rho} \mathbf{U}, \quad \mathbf{U} = \nabla c, \quad -\Delta c = \rho, \end{aligned}$$

where c denotes the concentration of some attracting agent, generated by the cells themselves. Notice that the congestion constraint prevents concentration of mass. As a matter of fact, the characteristic function of a single ball is a static solution to this system, in the whole space \mathbb{R}^3 , and it can be expected that any solution converges to such a configuration. Note also that bacterial growth could be handled by adding an appropriate term in the right-hand side of the transport equation.

We also believe that it can be fruitful in the modelling of granular media. Bouchut et al.[8] propose a model of pressureless gas for which the density is subject to remain less than 1. This model is essentially mono-dimensionnal (the construction of explicit solutions proposed in [6] uses extensively the one-dimensionality). As our model can be seen as a first order (in time) version of this second order pressureless gas model, we believe that the handling of the congestion constraint we propose here, which applies in any dimension, might be used in the future for a macroscopic description of granular flows in higher dimension. The corresponding model, which is a second order version of the model we considered here, could be written as follows

$$\begin{aligned} \frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) &= 0, \\ \frac{\partial(\rho \mathbf{u})}{\partial t} + \nabla \cdot (\rho \mathbf{u} \otimes \mathbf{u}) + \nabla p &= 0, \\ 0 \leq \rho \leq 1, \quad p(1 - \rho) &= 0, \quad \mathbf{u}^+ = P_{C_\rho} \mathbf{u}^-, \end{aligned}$$

where C_ρ is the cone of feasible velocities which we introduced, and \mathbf{u}^- (resp. \mathbf{u}^+) is the velocity before (resp. after) the collision (both are equal in case there is no collision).

Those extensions are not straightforward. In particular, the ‘‘spontaneous’’ velocity (i.e. chemotactic velocity for cell models, velocity before collision for granular flow models) may not be a gradient, and may furthermore vary in time. It rules out some arguments we used here. The discrete gradient flow construction, which is based on a simultaneous handling of advection and congestion constraint, could be replaced by a prediction-correction strategy: a first step of free advection followed by a projection (for the Wasserstein distance) onto the set of feasible densities. This approach would reproduce in the Wasserstein setting the techniques which proved successful in the context of sweeping processes in Hilbert spaces, which we mentioned in the introduction (see Refs. [36], [20], [21]).

ACKNOWLEDGMENT

We would like to thank Y. Brenier, A. Figalli and N. Gigli for fruitful discussions.

REFERENCES

- [1] L. Ambrosio, Movimenti minimizzanti, *Rend. Accad. Naz. Sci. XL Mem. Mat. Sci. Fis. Natur.* **113** (1995) 191–246.
- [2] L. Ambrosio, N. Gigli, G. Savaré, Gradient flows in metric spaces and in the space of probability measures, *Lectures in Mathematics* (ETH Zürich, 2005).
- [3] L. Ambrosio, G. Savaré, Gradient flows of probability measures, *Handbook of differential equations, Evolutionary equations* **3**, ed. by C.M. Dafermos and E. Feireisl (Elsevier, 2007).
- [4] N. Bellomo, C. Dogbe, On the modelling crowd dynamics from scalling to hyperbolic macroscopic models, *Math. Mod. Meth. Appl. Sci.* **18** Suppl. (2008) 1317–1345.

- [5] J.-D. Benamou, Y. Brenier, A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem, *Numer. Math.* **84**(3) (2001) 375–393.
- [6] F. Berthelin, Existence and weak stability for a pressureless model with unilateral constraint, *Math. Mod. Meth. Appl. Sci.* **12**(2) (2002) 249–272.
- [7] A. Blanchet, V. Calvez, J.A. Carrillo, Convergence of the mass-transport steepest descent scheme for the sub-critical Patlak-Keller-Segel model, *SIAM J. Numer. Anal.* to appear.
- [8] F. Bouchut, Y. Brenier, J. Cortes, J.-F. Ripoll, A hierarchy of models for two-phase flows, *J. nonlinear sci.* **10**(6) (2000) 639–660.
- [9] G. Buttazzo, C. Jimenez, E. Oudet, An optimization problem for mass transportation with congested dynamics, *SIAM J. Control Optim.* (2007).
- [10] G. Buttazzo, F. Santambrogio, A model for the optimal planning of an urban area, *SIAM J. Math. Anal.* **37**(2) (2005) 514–530.
- [11] J.A. Carrillo, M.P. Gualdani, G. Toscani, Finite speed of propagation in porous media by mass transportation methods, *R. Acad. Sci. Paris Ser. I* **338** (2004) 815–818.
- [12] C. Chalons, Numerical approximation of a macroscopic model of pedestrian flows, *SIAM J. Sci. Comput.* **29**(2) (2007) 539–555.
- [13] C. Chalons, Transport-equilibrium schemes for pedestrian flows with nonclassical shocks, *Traffic and Granular Flows'05*, 347–356 (Springer, 2007).
- [14] R.M. Colombo, M.D. Rosini, Pedestrian flows and non-classical shocks, *Math. Methods Appl. Sci.* **28** (2005) 1553–1567.
- [15] V. Coscia, C. Canavesio, First-order macroscopic modelling of human crowd dynamics, *Math. Mod. Meth. Appl. Sci.* **18** (2008) 1217–1247.
- [16] B. Dacorogna, J. Moser, On a partial differential equation involving the Jacobian determinant, *Annales de l'Institut Henry Poincaré Analyse Non Linéaire* **7**(1) (1990) 1–26.
- [17] E. De Giorgi, New problems on minimizing movements, *Boundary Value Problems for PDE and Applications*, C. Baiocchi and J. L. Lions eds. (Masson, 1993) 81–98.
- [18] P. Degond, L. Navoret, R. Bon, D. Sanchez, Congestion in a macroscopic model of self-driven particles modeling gregariousness, to appear.
- [19] C. Dogbe, On the numerical solutions of second order macroscopic models of pedestrian flows, *Comput. Appl. Math.* **567** (2008) 1884–1898.
- [20] J.F. Edmond, L. Thibault, Relaxation of an optimal control problem involving a perturbed sweeping process, *Math. Program Ser B* **104**(2-3) (2005) 347–373.
- [21] J.F. Edmond, L. Thibault, BV solutions of nonconvex sweeping process differential inclusion with perturbation, *J. Differential Equations* **226**(1) (2006) 135–179.
- [22] D. Helbing, A fluid dynamic model for the movement of pedestrians, *Complex Systems* **6** (1992) 391–415.
- [23] D. Helbing, P. Molnar, F. Schweitzer, Computer simulations of pedestrian dynamics and trail formation, *Evolution of Natural Structures, Sonderforschungsbereich* **230** (Stuttgart, 1994) 229–234.
- [24] D. Helbing, P. Molnár, Social force model for pedestrian dynamics, *Phys. Rev E* **51** (1995) 4282–4286.
- [25] D. Helbing, T. Vicsek, Optimal self-organization, *New J. Phys.* **1** (1999) 13.1–13.17.
- [26] D. Helbing, I. Farkas, T. Vicsek, Simulating dynamical features of escape panic, *Nature* **407** (2000) 487–490.
- [27] L.F. Henderson, The statistics of crowd fluids, *Nature* **229** (1971) 381–383.
- [28] S.P. Hoogendoorn, P.H.L. Bovy, Dynamic user-optimal assignment in continuous time and space, *Transport. Res. Part B* **38** (2004) 571–592.
- [29] S.P. Hoogendoorn, P.H.L. Bovy, Pedestrian route-choice and activity scheduling theory and models, *Transport. Res. Part B* **38** (2004) 169–190.
- [30] R. L. Hughes, A continuum theory for the flow of pedestrian, *Transport. Res. Part B* **36** (2002) 507–535.
- [31] R. L. Hughes, The flow of human crowds, *Ann. Rev. Fluid Mech.* **35** (2003) 169–183.
- [32] R. Jordan, D. Kinderlehrer, F. Otto, The variational formulation of the Fokker-Planck equation, *SIAM J. Math. Anal.* **29**(1) (1998) 1–17.
- [33] L. V. Kantorovich, On the transfer of masses, *Dokl. Akad. Nauk. SSSR* **37** (1942) 227–229.
- [34] B. Maury, J. Venel, Handling of contacts in crowd motion simulations, *Traffic and Granular Flow* (Springer, 2007).
- [35] B. Maury, J. Venel, A mathematical framework for a crowd motion model, *C.R. Acad. Sci. Paris, Ser I* **346** (2008) 1245–1250.
- [36] J.J. Moreau, Evolution problem associated with a moving convex set in a Hilbert space, *J. Differential Equations* **26**(3) (1977) 346–374.
- [37] F. Otto, The geometry of dissipative evolution equations: the porous medium equation, *Comm. Partial Differential Equations* **26**(1–2) (2001) 101–174.
- [38] B. Piccoli, A. Tosin, Time-evolving measures and macroscopic modeling of pedestrian flow (2008) to appear.

- [39] B. Piccoli, A. Tosin, Pedestrian flows in bounded domains with obstacles, *Contin. Mech. Thermodyn.* (2009) to appear.
- [40] M. Roesch, Contribution à la description géométrique des fluides parfaits incompressibles, *PhD Thesis, Université de Paris 6*.
- [41] J. Venel, Integrating strategies in numerical modelling of crowd motion, *Pedestrian and Evacuation Dynamics* (Springer, 2008).
- [42] C. Villani, Topics in optimal transportation, *Grad. Stud. Math.* **58** (AMS, Providence 2003).
- [43] C. Villani, Optimal transport, old and new, *Grundlehren der mathematischen Wissenschaften* **338** (2009).

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ PARIS-SUD,, BAT. 425, 91405 ORSAY CEDEX, FRANCE,
BERTRAND.MAURY@MATH.U-PSUD.FR

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ PARIS-SUD,, BAT. 425, 91405 ORSAY CEDEX, FRANCE,
AUDE.ROUDNEFF@MATH.U-PSUD.FR

LABORATOIRE CEREMADE, UNIVERSITÉ PARIS-DAUPHINE,, PL. DE LATTRE DE TASSIGNY, 75775 PARIS
CEDEX 16, FRANCE, FILIPPO@CEREMADE.DAUPHINE.FR