



HAL
open science

Physiological recovery from episodic acid stress does not mean population recovery of *Gammarus fossarum*

V. Felten, J.M. Baudoin, F. Guerold

► To cite this version:

V. Felten, J.M. Baudoin, F. Guerold. Physiological recovery from episodic acid stress does not mean population recovery of *Gammarus fossarum*. *Chemosphere*, 2006, 65, p. 988 - p. 998. 10.1016/j.chemosphere.2006.03.059 . hal-00452245

HAL Id: hal-00452245

<https://hal.science/hal-00452245>

Submitted on 1 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author-produced version of the article published in Chemosphère,
2006, 65, 988-998.
The original publication is available at <http://www.sciencedirect.com/>
doi : 10.1016/j.chemosphere.2006.03.059

1 **Physiological recovery from episodic acid stress does not mean**
2 **population recovery of *Gammarus fossarum***

3 **V. Felten^a, J.M. Baudoin^b & F. Guérol^c.**

4 ^a Laboratoire d'écotoxicologie, CEMAGREF, 3 bis quai Chauveau, 69336 Lyon, CP 220,
5 Cedex 09, France.

6 ^b Laboratoire Biodiversité et Fonctionnement des Ecosystèmes, Université de Metz, Campus
7 Bridoux, Rue du Général Delestraint, 57 070 Metz, France.

8 ^c Laboratoire Ecotoxicité et Santé Environnementale, CNRS UMR 7146, Université Paul
9 Verlaine – Metz Rue du Général Délestraint, 57070 Metz, France.

10

11

12 **Abstract**

13 The physiological responses of the acid-sensitive amphipod *Gammarus fossarum* exposed *in*
14 *situ* to acid stress (pH 4.5 and 5.5) and then transferred back to neutral water were
15 investigated. Survival rate and haemolymph [Cl⁻] and [Na⁺] were assessed after 24, 48 and
16 72h of exposure in acidic streams and after a recovery period of 12, 24, 36, 48 and 60h. After
17 24h, exposure to slightly acidic (pH 5.5) and strongly acidic water (pH 4.5) led to a severe
18 and significant depletion in haemolymph [Na⁺] and [Cl⁻] compared to organisms exposed in
19 circumneutral water (pH 7.3). However, after only a 12h- period of transfer back in neutral
20 water and whatever the previous exposure time (24, 48 and 72h) in both slightly and strongly
21 acidic water, haemolymph [Na⁺] and [Cl⁻] were equal or superior to the control level without
22 associated mortality. In spite of this fast physiological recovery capacity, populations of *G.*
23 *fossarum* living in streams undergoing episodic acid stresses were drastically affected thus,
24 demonstrating the high acid-sensitivity of this species. We discuss the possible reasons of
25 population regression and the absence of population recovery.

26 **Keywords:** *Gammarus fossarum*, haemolymph, ion loss, acid stress, recovery, *in situ*
27 exposure.

28

29

30 INTRODUCTION

31 Acidification of freshwater ecosystems related to anthropogenic emissions of SO₂ and NO_x
32 has been one of the most striking ecological problems throughout the northern hemisphere
33 during the 20th century. National and international legislation in the 1980s and 1990s aimed
34 to reduce the emissions of acidifying pollutants (e.g. Clean Air Act in the USA and the
35 Convention on Long-Range Transboundary Air Pollution in Europe) have led to the decline in
36 acidic depositions across wide areas of Europe and North America (Stoddard et al., 1999;
37 Lawrence et al., 2000; Likens et al., 2001). Several recent studies have shown that recovery of
38 alkalinity has occurred in many areas of Europe and North America (Stoddard et al., 1999;
39 Skjelkvale et al., 2001), but acidification of freshwater ecosystems still occurs in many areas
40 (Guérol et al., 2000; Driscoll et al., 2001; Evans et al., 2001). In addition, acidification of
41 aquatic ecosystems is now reported across other large areas of the world where high economic
42 and demographic growth rates occur, such as in China (Thorjörn et al., 1999; Tang et al.,
43 2001) and India (Aggarwal et al., 2001).

44 Episodic acidification following snowmelt or heavy rainfalls has been well documented
45 (Ormerod & Jenkins, 1994; O'Brien & Eshleman, 1995; Wigington *et al.*, 1996). These
46 hydrometeorological events induce a decrease of pH, Acid Neutralizing Capacity (ANC), and
47 base cations concentrations as well as an increase of aluminum concentrations (O'Brien et al.,
48 1993; Soulsby, 1995). The intensity of acid-stress tends to be greater in more acidic
49 environments because low ANC streams can be subject to episodic acidic stress throughout
50 the year (Colin *et al.*, 1989). Change in Al speciation accompanying large pH depressions has

51 been shown to cause stress and mortality in many aquatic species (Weatherley & Ormerod,
52 1991; Carline et al., 1992; Van Sickle et al., 1996).

53 One of the most striking consequences of freshwater acidification is the erosion of
54 biodiversity (Muniz, 1991). Numerous studies have clearly demonstrated a failure to regulate
55 blood or haemolymph Na^+ and Cl^- levels in acid-stressed fish, clams (Unionidae) and
56 decapods (Massabuau, 1985; McMahon & Stuart, 1989; Pynnönen, 1991; Masson et al.,
57 2002). However, most of the studies have focused on large species, and relatively little is
58 known about physiological responses in smaller acid-sensitive species of macroinvertebrates.

59 Crustaceans contain many of the most acid-sensitive macroinvertebrate species (Sutcliffe &
60 Carrick, 1973; Guérol et al., 2000). In previous studies, Felten & Guérol (2001, 2004)
61 showed that *Gammarus fossarum* (Crustacea: Amphipoda) also suffered a severe depletion of
62 haemolymph Na^+ and Cl^- ions when exposed to acidic conditions. Conjointly, we determined
63 relationship between acidification level and haemolymph ion losses. Thus, we proposed that
64 haemolymph ion concentrations in the acid-sensitive species *G. fossarum* could represent
65 effective biomarkers for monitoring acidification of running waters.

66 The present study aims to investigate the recovery of haemolymph $[\text{Na}^+]$ and $[\text{Cl}^-]$ in *G.*
67 *fossarum* previously exposed to different magnitudes of acid stresses. In this context we
68 assessed *in situ* the short-term response of haemolymph $[\text{Na}^+]$ and $[\text{Cl}^-]$ in *G. fossarum*
69 transferred to 3 headwater streams providing 3 different acidification levels (defined by: pH,
70 ANC, $[\text{Mg}^{2+}]$, $[\text{Ca}^{2+}]$ and $[\text{Al}_{\text{tot}}]$). For each stream and each exposure time, organisms were
71 transferred back to their native circumneutral stream not only to test for *G. fossarum* recovery
72 capacity but also to evaluate the effect of an episodic acid stress.

73

74

75

MATERIALS AND METHODS

76
77 **Study organisms.** Experiments were performed on *Gammarus fossarum* because this species
78 presents several interesting characteristics for ecotoxicological investigations : 1) it is an
79 acid-sensitive species (Guéroid *et al.*, 2000), 2) widespread and common in West Palearctica
80 (Barnard & Barnard, 1983), 3) often occurring in high density, 4) easy to identify to species
81 level, 5) characterised by a sexual dimorphism and (6) playing a major role in the leaf litter
82 breakdown process and consequently in the entire food web (Pöckl, 1995).

83 **Experimental design.** The study was conducted in the Vosges Mountains (north-eastern
84 France) in 3 headwater streams draining sandstone bedrock and providing exposure to a range
85 of acidification levels (**Fig.1.**, neutral : La Maix; Ravine: slightly acidified; Gentil Sapin:
86 strongly acidified). The 3 sites were located in the same area, the highest distance between 2
87 sites being less than 10 km representing a 20 mn transport of the organisms.

88 Males *G. fossarum* with 8-10mm body size were collected from the neutral stream, La Maix.
89 The experimental design we used is shown in **Fig. 2.** Each Plexiglas flow-through enclosure
90 (experimental unit) contained 144 *G. fossarum*. Enclosures were divided in 18 compartments
91 with 8 individual each. Two enclosures like this were placed in each river. Enclosures were
92 then transferred to each of the 3 streams including the neutral streams. The total number of *G.*
93 *fossarum* collected for this experiment was 2592.

94 For each stream, one enclosure was used for haemolymph analyses and the other for the
95 assessment of survival. For each stream, there were 3 pairs of enclosure each corresponding to
96 one exposure time (24, 48 and 72 h).

97 In order to evaluate the initial concentrations of haemolymph Cl^- and Na^+ in *G. fossarum*, 12
98 organisms were sampled in the neutral “native” stream just before the onset of the
99 experiments (T_0 , control). At 24, 48 and 72 h of exposure, survival was assessed and samples
100 of haemolymph from 8 organisms were randomly collected in each stream for analysis (in

101 enclosure corresponding to exposure time). For more details concerning the enclosure
102 protocol see Felten & Guérolde (2004).

103 After each exposure time, the sampling enclosures were transferred to the circumneutral
104 stream (La Maix) in order to initiate the recovery experiment (**Fig 2.b.**); the latest values of
105 haemolymph $[Cl^-]$ and $[Na^+]$ recorded before the transfer to circumneutral water were referred
106 as t_0 for the recovery experiment. After 12, 24, 36, 48 and 60 h of recovery, the survival was
107 assessed and samples of haemolymph from 8 organisms were randomly collected for analysis
108 in each enclosure corresponding to each stream exposure time (**Fig 2.b.**).

109 Survival rates calculated for the recovery period are based on the number of organisms
110 remaining (in the enclosure) at the time of transfer to the circumneutral stream. Thus, there
111 was a 60 h-recovery kinetic for each exposure type (circumneutral, slightly acid and strongly
112 acid).

113 **Survival, haemolymph sampling and analysis.** For each acid exposure time and recovery
114 time, the survival was assessed in each stream (3 replicates of 48 organisms). Samples of
115 haemolymph (0.8 to 1.2 μ l) were taken from the telson of each individual ($n = 8$) using a
116 microsyringe, transferred to a gauged 5- μ l microcapillary tube and centrifuged for 10 min at
117 6596 g. After centrifugation the liquid phase was diluted in 2 ml of Nanopur water to
118 determine chloride and sodium concentrations in haemolymph by ionic chromatography
119 (Dionex 4500i with Ion Pac AS4A column) and atomic absorption spectrophotometry (AAS)
120 (Perkin Elmer Analyst 100), respectively.

121 **Water analysis.** Water samples were collected at the initiation of the experiment (T_0) and at
122 each time of acid exposure (24, 48 and 72 h) and recovery (12, 24, 36, 48 and 60 h). Cations
123 were analysed by flame AAS and anions by ionic chromatography as described previously.
124 Total aluminium was determined by graphite furnace AAS (Varian Spectraa 300) after
125 acidification with 0.25% HNO_3 . Acid neutralising capacity (ANC) was measured by Gran's

126 titration and pH (glass electrode), and conductivity with multi-parametric equipment (WTW).
127 Chemical characteristics of water from each stream are given in **Table 1**.

128 **Statistical analysis.** All data are reported as mean \pm SD. Statistical comparisons of
129 experimental data were performed by two-way analysis of variance (ANOVA) and Fisher's
130 Least Significant Difference test (LSD). The analyses were carried out using STATISTICA
131 (Microsoft), with a probability limit of $p \leq 0.05$ considered as significant.

132

133 RESULTS

134 Acid exposure

135 Acidified streams were characterized by low pH, low ANC, low $[\text{Mg}^{2+}]$ and low $[\text{Ca}^{2+}]$ and
136 high $[\text{Al}_{\text{tot}}]$ (**Table 1**). The baseline levels of haemolymph $[\text{Cl}^-]$ and $[\text{Na}^+]$ prior to the
137 exposure were 54.1 ± 8.1 and 90.4 ± 15.9 mmol l^{-1} , respectively, in control organisms (**Fig.**
138 **2.a.-b.**).

139 The 2-way analysis of variance (ANOVA) indicated that stream acidity, exposure time and
140 the interaction between them (Stream \times Exposure Time) exerted a significant effect on
141 haemolymph parameters ($[\text{Na}^+]$, $[\text{Cl}^-]$) and survival (**Table 2.a.**).

142 Haemolymph $[\text{Na}^+]$ and $[\text{Cl}^-]$ in *G. fossarum* exposed to circumneutral stream remained
143 constant over a 72-h exposure period, but decreased significantly in organisms exposed to
144 slightly acidic (Ravine, mean pH = 5.5) and strongly acidic streams (Gentil Sapin, mean pH =
145 4.5) during the first 24 h (**Fig. 3.a.-b.**). Indeed, after 24 h of exposure, the loss of
146 haemolymph Cl^- ranged from 22.5% in the slightly acidic stream (mean haemolymph $[\text{Cl}^-]$ =
147 41.9 ± 6.7 mmol l^{-1}) to 48.8% in the strongly acidic one (mean haemolymph Cl^- = 27.7 ± 5.5
148 mmol l^{-1}) compared with the control (mean haemolymph Cl^- = 54.1 ± 8.1 mmol l^{-1}) (**Fig.**
149 **3.a.**). The same trend was observed for haemolymph $[\text{Na}^+]$. After 24 h of exposure, the loss of
150 haemolymph Na^+ ranged from 19.9% in Ravine (mean haemolymph Na^+ = 72.4 ± 13.6 mmol

151 l^{-1}) to 53.9% in Gentil Sapin (mean haemolymph $\text{Na}^+ = 41.7 \pm 9.6 \text{ mmol l}^{-1}$) compared with
152 the control (mean haemolymph $\text{Na}^+ = 90.4 \pm 15.9 \text{ mmol l}^{-1}$) (**Fig. 3.b.**).

153 After 48 h of exposure, the haemolymph $[\text{Na}^+]$ and $[\text{Cl}^-]$ of the organisms exposed to slightly
154 and strongly acidic streams were close to 24h-exposure values and remained constant until the
155 end of the experiment. Individuals transferred for 72 h to Ravine had a mean haemolymph
156 $[\text{Cl}^-]$ of $42 \pm 11.5 \text{ mmol l}^{-1}$ and $[\text{Na}^+]$ of $70.8 \pm 20 \text{ mmol l}^{-1}$, representing a significant 22.3%
157 loss of chloride and a significant 21.7% loss of sodium ($p < 0.05$). On the other hand,
158 individuals transferred for 72 h to Gentil Sapin had a haemolymph $[\text{Cl}^-]$ of $25.6 \pm 4.6 \text{ mmol l}^{-1}$
159 and $[\text{Na}^+]$ of $44.7 \pm 6.9 \text{ mmol l}^{-1}$, representing a chloride loss of 45.3% and a sodium loss of
160 50.5 ($p < 0.001$).

161 For each exposure time, the survival rate in *G. fossarum* transferred to the native
162 circumneutral stream (La Maix) remained very high and above 99%. After 72h of exposure,
163 the survival rates in organisms exposed to slightly and strongly acidic streams were
164 significantly different from mean control values for the same exposure time ($p < 0.05$),
165 reaching $89.6 \pm 7.5\%$ and $42.4 \pm 11.9\%$ respectively. On the contrary the survival rates after
166 24 and 48h of exposure in slightly acidified streams (Ravine) were not significantly different
167 from those observed in the neutral stream whereas significant differences were measured in
168 organisms transferred to strongly acidified stream (Gentil sapin). Thus, survival rates of
169 gammarids transferred to Gentil sapin reached $79.9 \pm 11.8\%$ and $51.7 \pm 10.8\%$ after a 24h and
170 a 48h-exposure time respectively (**Fig. 3.c.**).

171

172

Recovery

173 The 2-way analysis of variance (ANOVA) indicated that Exposure Time and/or Recovery
174 Time as well as the interaction term had a significant influence on haemolymph $[\text{Na}^+]$,
175 haemolymph $[\text{Cl}^-]$ and survival (**Table 2.b.**).

176 We observed a rapid and total recovery of haemolymph $[Cl^-]$ and $[Na^+]$ in organisms
177 previously exposed to slightly and strongly acidic stream. After only 12h transferred back in
178 the native circumneutral stream, the mean values of haemolymph parameters were similar or
179 significantly higher than those measured at To (**Fig. 4.a.-b.** and **Fig. 5. a.-b.**).

180 After a 12h-recovery in La Maix following a 72h-exposure in Gentil sapin (the longest and
181 most intense acid stress tested), the mean haemolymph $[Cl^-]$ increased from 29.6 ± 4.3 mmol
182 l^{-1} to 63.8 ± 13.9 mmol l^{-1} , representing an increase of 115.6%. Similarly the mean
183 haemolymph $[Na^+]$ increased from 44.7 ± 6.9 mmol l^{-1} to 85.2 ± 8.5 mmol l^{-1} , representing an
184 increase of 90.6%.

185 In all enclosures, the survival rates remained high throughout the recovery experiment (>93%,
186 **Fig. 4.c.** and **Fig. 5 .c.**).

187

188 DISCUSSION

189 Acid-exposure

190 Several studies have shown that crustaceans exposed to water-borne pollutants, environmental
191 stressors and pathological agents usually exhibit disruption of ionic regulation (Lignot et al.,
192 2000). Different causes include alterations in the structure and ultrastructure of the branchial
193 and excretory organs, and changes in Na^+ , K^+ -ATPase activity, ion fluxes and surface
194 permeability (Lignot et al., 2000). Ion-regulation failure leading to a severe deficiency of
195 extracellular ions (i.e. Na^+ and Cl^-) has been recognised to be the major response in fish to
196 acid stress (McDonald et al., 1989; Potts & McWilliams, 1989; Wood, 1989). Similar results
197 have been reported in crayfish (Appelberg, 1985; Fjeld et al., 1988; McMahon & Stuart,
198 1989; Jensen & Malte, 1990), gammarids (Rupprecht, 1992; Felten & Guérol, 2001, 2004)
199 and molluscs (Pynnönen, 1991). Surprisingly, and despite the numerous papers reporting
200 detrimental effects of acidification on invertebrate communities, few studies have been

201 performed on the ecophysiology of smaller acid-sensitive macroinvertebrate species
202 (Herrmann, 1987; Herrmann & Andersson, 1986).

203 In the present study we showed that exposure of *G. fossarum* to strongly acidic or slightly
204 acidic water induced early significant losses of haemolymph $[Na^+]$ and $[Cl^-]$. Moreover,
205 failure in ion-regulation was accompanied by a significant mortality. These results were in
206 agreement with those obtained in previous studies (Felten & Guérol, 2001, 2004) which
207 permitted us to demonstrate that ion losses were significantly correlated to pH.

208

209 **Recovery from acid stress**

210 Despite drastic ion losses (Cl^- and Na^+) following exposures to slightly and strongly acidic
211 stresses, organisms transferred back to the native circumneutral stream were surprisingly able
212 to recover rapidly from acid stress (as soon as 12h). Moreover, survival rates associated with
213 the recovery period were very high (>93.4%). Consequently, the ion-regulation failure
214 evidenced during the exposure under acidic conditions appears reversible even when the loss
215 of ions was severe (>50%).

216 These results are in accordance with those of several studies which have highlighted the
217 recovery capacity of several acid-sensitive species following their transfer from an acidified
218 medium to a circumneutral medium. Favrel (1998) observed a total recovery of haemolymph
219 $[Cl^-]$ and $[Na^+]$ in *Dinocras cephalotes* (Plecoptera) after 16 days of recovery in a
220 circumneutral water (pH = 6.6, $[Al] = 10 \mu g l^{-1}$) following a 8 days exposure in an acidified
221 water (pH = 4.6, $[Al] = 540 \mu g l^{-1}$). Comparable results had been also reported for *Anodonta*
222 *anatina* (Mollusc) for a 8 day exposure to pH 4.0-5.0 followed by a 8 days transfer in a water
223 with a pH 7.2-7.4 (Pynnönen, 1994). Similarly, Kroglund & Staurnes (1999) conducted an
224 experiment showing that acid-exposed (pH 6.0 and 5.6) *Salmo salar* smolts were able to re-
225 establish their plasma chloride levels within 3 days in pH 6.28 water, but the authors did not

226 sample organisms until the end of the 3 days. In our study, the recovery observed for *G.*
227 *fossarum* was definitely faster than those observed in the species mentioned above.

228 Moreover, these results are in accordance with our previous observation (Felten & Guérol, 2004) showing the physiological adaptation of *G. fossarum* exposed to slightly acidic stress
229 (pH 5.7, $[Al_{tot}] = 2.5 \mu\text{mol l}^{-1}$, $[Ca^{2+}] = 26.8 \mu\text{mol l}^{-1}$). After 24h of exposure, *G. fossarum* had
230 undergone significant ion losses but following 96h, no differences were observed in
231 haemolymph $[Cl^-]$ and $[Na^+]$ (compared to the control) without any additional mortality.

232 Several assumptions may be advanced to explain the fast recovery of haemolymph $[Cl^-]$ and
233 $[Na^+]$. On the one hand Chamier et al. (1989) reported that Na^+ turnover is very fast in *G.*
234 *pulex* (65% of total body in Na^+ per day at 9°C). In addition, Wood & Ronago (1986)
235 observed an increase in Cl^- and Na^+ uptake during the recovery following an exposure to pH 4.
236 This can be due to a reduction of the $[H^+]/[Na^+]$ ratio in water, consequently decreasing the
237 competition between these two ions at ionic transport sites, leading to an increase in Na^+
238 uptake. In addition organisms transferred back to circumneutral waters were then facing lower
239 passive ion losses and water uptake because of the increase in $[Ca^{2+}]/[Al]$ and $[Ca^{2+}]/[H^+]$
240 ratios. Thus, the easier access to Ca^{2+} allowed the reinforcement of cellular junctions allowing
241 reduced permeability. Finally, different studies have shown an increased number of chloride
242 cells in gills as well as morphological changes in the apical surface of these chloride cells, in
243 various species of fish exposed to acid water (Chevalier et al., 1985; Karlsson-Norrgrén et al.,
244 1986; Jagoe & Haines, 1990, 1997). For example, chloride cells can present apical wells
245 (Leino & McCormick, 1984; Leino et al., 1987a, 1987b) or apical evaginations (Chevalier et
246 al., 1985; Leino et al., 1987a, 1987b) increasing exchange surface. This kind of change which
247 can limit and/or offset ion losses under acidic conditions, could explain the fast recovery of
248 haemolymph Na^+ and Cl^- concentrations after a transfer back to circumneutral water.
249

250 However, to our knowledge, no study has highlighted such modifications in small
251 macroinvertebrate species.

252

253 **Extrapolation for populations**

254 In a previous study (Felten, 2003), populations of *G. fossarum* were sampled using a surber
255 sampler, on 10 occasions and in 3 sites of the same circumneutral stream (La Maix). This
256 stream was characterized by a decreasing gradient of acidification from upstream to
257 downstream (the most distant sites were 2.7 km apart). The most downstream site, studied in
258 this paper, was circumneutral throughout the year (mean pH : 7; from 6.7 to 7.4; n = 25). The
259 most upstream site was affected by episodic strong acidification events (mean pH: 6.0; from
260 4.3 to 6.80; n = 25) whereas slightly episodic stresses occurred in the intermediate site (mean
261 pH : 6.6; from 5.7 to 7; n = 25). As a result, *G. fossarum* was totally absent from the upper
262 site and, at the intermediate site, the density of the species was 10 times lower (443 ind./m²)
263 that those observed downstream (4317 ind./m²). Consequently, although *G. fossarum* was
264 able to rapidly compensate high ion losses caused by acid stress, gammarid population was
265 drastically affected by repeated episodic acid stress meaning that the rapid compensation of
266 high ion loss seems not to help the *G. fossarum* population in surviving acid episodes if these
267 are repeated.

268 Episodic acidification has been recognized to exert a drastic impact on macroinvertebrate and
269 fish populations and communities (Weatherley & Osmerod, 1991 ; Baker *et al.*, 1996 ; Van
270 Sickle *et al.*, 1996; Lepori *et al.*, 2003). McCahon & Poulton (1991) and Merrett *et al.* (1991)
271 showed the cumulative effect of multiple acid stresses on macroinvertebrate mortality.
272 Finally, several studies conclude that acid episodes can restrict or offset the recovery process
273 of acid sensitive invertebrates (Kowalik *et al.*, 2006) and fish (Kroglund *et al.*, 2001).

274 According to Davies *et al.* (1992), episodic acidifications take place over hours (rainfall) to
275 months (snowmelt). In a previous experiment, Felten & Guérolde (2005) showed that a 168h-
276 exposure of *G. fossarum* to Gentil sapin streamwater led to 80 % of mortality which can
277 explain the total absence of the specie in the upper site.

278 Several non exclusive hypotheses can conjointly explain the decrease of density or the loss of
279 population in streams subjected to episodic acid stresses, such as *i*) an increase of drift-
280 response and mortality of sensitive stages (juveniles) (McCahon & Poulton, 1991; Taylor *et*
281 *al.*, 1994), *ii*) a lower food quality (Willoughby, 1988; Willoughby & Mappin, 1988; Sutcliffe
282 & Hildrew, 1989) or/and a lower conversion efficiency of food to growth (Lee *et al.*, 1983;
283 Hargeby & Petersen, 1988), *iii*) a lower food intake (Lemly & Smith, 1985; Tierney &
284 Atema, 1986), *iv*) an enhanced energetic cost associated with osmoregulation, ion retention
285 and respiration (Økland & Økland, 1986) leading to a decrease of growth and reproduction
286 (Maltby, 1994; Seiler & Turner, 2004).

287 Moulting is known to be a critical phase for crustaceans (Wright & Frain, 1981; McCahon &
288 Pascoe, 1988; Wheatly & Gannon, 1995) requiring a lot of energy (Maltby, 1994; Wheatly &
289 Gannon, 1995). According to Pöckl (1992), in younger gammarid stages, moulting occurred
290 close together. Consequently, a higher acid-sensitivity of younger stages (Naylor *et al.*, 1990)
291 explained by the moulting frequency could partly account for population regression in acidified
292 streams. Moreover, small-sized gammarids affected by an acid stress may be unable to resist
293 to the associated hydrological stress.

294

295 **Conclusion**

296 *G. fossarum* facing acid stress fail to regulate ion losses, leading thus to an important
297 mortality when the environmental stress is intense. This failure is rapidly reversible when the
298 stress ceases. However, despite the recovery capacity of *G. fossarum* from important ion

299 losses following acid stress, populations have been severely reduced or have totally
300 disappeared from numerous headwater streams draining catchments subject to acidification.
301 Thus, it is suggested that both the frequency and the intensity of stresses clearly structure
302 gammarid population. In this context, we recommend to conduct further studies dealing with
303 the effects of episodic acidification on population structure, in relation to physiological
304 parameters (e.g. energetic cost, growth) to better understand population regression and
305 recovery. This kind of study is particularly important since the decline of base cations (mainly
306 Ca^{2+} and Mg^{2+}) in soils and surface waters has been reported in most areas where high rates of
307 sulphur depositions occurred previously. These trends should indeed lead to an increase of
308 episodic acid stress frequency and intensity.

309

310 **Acknowledgements.** We would like to thank Crane Rogers for linguistic corrections of this
311 manuscript.

312

313

314

LITERATURE CITED

315 Aggarwal, S.G., Chandrawanshi, K., Patel, R.M., Agarwal, S., Kamavishdar, A., Mundhara,
316 G., 2001. Acidification of surface water in central India. *Water Air Soil Poll.* 130, 855-862.

317 Appelberg, M., 1985. Changes in haemolymph ion concentrations of *Astacus astacus* L. and
318 *Pacifastacus leniusculus* (Dana) after exposure to low pH and aluminum. *Hydrobiologia*
319 121, 19-25.

320 Barnard, J.L., Barnard, C.M., 1983. *Freshwater Amphipoda of the World (Parts I & II)*.
321 Hayfield Associates, Mt. Vernon.

322 Baker J.P., Van Sickle J., Gagen C.J., DE Walle D.R., Sharpe W.E., Carline R.F., Baldigo
323 B.P., Murdoch P.S., Bath D.W., Kretser W.A., Simonin H.A. & Wigington P.J.Jr., 1996.

- 324 Episodic acidification of small streams in the northeastern United States: Effet on fish
325 populations. *Ecol. Appl.* 6 (2), 422-437.
- 326 Carline, R.F., DeWalle, D.R., Sharpe, W.E., Dempsey, B.A., Gagen, C.J., Swistock, B., 1992.
327 Water chemistry and fish community responses to episodic stream acidification in
328 Pennsylvania, U.S.A. *Environ. Pollut.* 78, 45–48.
- 329 Chamier, A.C., Sutcliffe, D.W., Lishman, J.P., 1989. Changes in Na, K, Ca, Mg, and Al
330 content of submersed leaf litter, related to ingestion by the amphipod *Gammarus pulex*
331 (L.). *Freshwater Biol.* 21, 181-189.
- 332 Chevalier, G., Gauthier, L., Moreau, G., 1985. Histopathological and electron microscopic
333 studies of gills of brook trout, *Salvelinus fontinalis*, from the acidified lakes. *Can. J. Zool.*
334 63, 2062-2070.
- 335 Colin, J.L., Renard, D., Lescoat, V., Jaffrezo, J.L., Gros, J.M., Strauss, B., 1989. Relationship
336 between rain and snow acidity and air mass trajectory in Eastern France. *Atmos. Environ.*
337 23 (7), 1487-1498.
- 338 Davies, T.D., Tranter, M., Wigington, P.J., & Eshleman K.N., 1992. ‘Acidic episodes’ in
339 surface waters in Europe. *Journal of Hydrology*, 132, 25–69.
- 340 Driscoll, C.T., Lawrence, G.B., Bulger, A.J., Butler, T.J., Cronan, C.S., Eagar, C., Lambert,
341 K.F., Likens, G.E., Stoddard, J.L., and Weathers, K.C., 2001. Acidic deposition in the
342 northeastern United States: Sources and inputs, ecosystem effects, and management
343 strategies. *BioScience* 51, 180-198.
- 344 Evans, C.D., Harriman, R., Monteih, D.T., Jenkins, A., 2001. Assessing the suitability of acid
345 neutralizing capacity as a measure of long-term trends in acidic waters based on two
346 parallel datasets. *Water Air Soil Poll.* 130, 1541-1546.
- 347 Favrel, C., 1998. Ecophysiologie de la larve de *Dinocras cephalotes* (Plecoptère, Perlidé)
348 exposée à un milieu acidifié, Thesis, Metz university, 279 p.

- 349 Felten, V., Guerold, F., 2001. Hyperventilation and loss of hemolymph Na^+ and Cl^- in the
350 freshwater amphipod *Gammarus fossarum* exposed to acid stress: a preliminary study. Dis.
351 Aquat. Org. 45, 77-80.
- 352 Felten, V., 2003. Effets de l'acidification des ruisseaux vosgiens sur la biologie, l'écologie et
353 l'écophysiologie de *Gammarus fossarum* Koch, 1835 (Crustacea Amphipoda): Approche
354 intégrée à différents niveaux d'organisation. Thesis, Metz university, 279 p.
- 355 Felten, V., Guerold, F., 2004. Haemolymph $[\text{Na}^+]$ and $[\text{Cl}^-]$ loss in *Gammarus fossarum*
356 exposed *in situ* to a wide range of acidic streams. Dis. Aquat. Org. 61, 113-121.
- 357 Fjeld, E., Hessen, D.O., Roos, N., Taubolt, T., 1988. Changes in gill ultrastructure and
358 haemolymph chloride concentrations in the crayfish, *Astacus astacus*, exposed to de-
359 acidified aluminum rich water. Aquaculture 72, 139-150.
- 360 Guerold, F., Boudot, J.P., Jacquemin, G., Vein, D., Merlet, D., Rouiller, J., 2000.
361 Macroinvertebrate community loss as a result of headwater stream acidification in the
362 Vosges Mountains (N-E France). Biodivers. Conserv. 9, 767-783.
- 363 Hargeby, A., Petersen, R.C. Jn., 1988. Effects of low pH and humus on the survivorship,
364 growth and feeding of *Gammarus pulex* (L.) (Amphipoda). Freshwater Biol. 19, 235-247.
- 365 Herrmann, J., Andersson K.G., 1986. Aluminium impact on respiration of lotic mayflies at
366 low pH. Water Air Soil Poll. 30, 703-709.
- 367 Herrmann, J., 1987. Sodium levels of lotic mayfly nymphs being exposed to aluminium at
368 low pH a preliminary report. Ann. Soc. R. Zool. de Belg. 117 (1), 181-188.
- 369 Jagoe, C.H., Haines, T.A., 1990. Morphometric effects of low pH and limed water on the gills
370 of atlantic salmo (*Salmo salar*). Can. J. Fish. Aquat. Sci. 47, 2451-2460.
- 371 Jagoe, C.H., Haines, T.A., 1997. Changes in gill morphology of atlantic salmon (*Salmo salar*)
372 smolts due to addition of acid and aluminum to stream water. Environ. Pollut. 97 (1-2),
373 137-146.

- 374 Jensen, F.B., Malte, H., 1990. Acid-base and electrolyte regulation, and hemolymph gas
375 transport in crayfish *Astacus astacus*, exposed to soft, acid water with and without
376 aluminum. J. Comp. Physiol. B 160, 483-490.
- 377 Karlsson-Norrgren L., Björklund I., Ljungberg, O., Runn, P., 1986. Acid water and
378 aluminium exposure : experimentally induced gill lesions in brown trout, *Salmo trutta* L. J.
379 Fish Dis. 9, 11-25.
- 380 Kroglund F, Kaste O, Rosseland B.O., Poppe T. 2001. The Return of the Salmon. Water, Air
381 and Soil Pollution. 130 : 1349-1354
- 382 Kowalik, R.A. & Ormerod, S. J., 2006. Intensive sampling and transplantation experiments
383 reveal continued effects of episodic acidification on sensitive stream invertebrates.
384 Freshwater Biol. 51, 180-191.
- 385 Kroglund, F., Staurnes, M., 1999. Water quality requirements of smolting Atlantic salmon
386 (*Salmo salar*) in limed acid rivers. Can. J. Fish. Aquat. Sci. 56, 2078–2086.
- 387 Lawrence, G.B., Lovett, G.M., Baeovsky, Y.H., 2000. Atmospheric deposition and watershed
388 nitrogen export along an elevational gradient in the Catskill Mountains, New York.
389 Biogeochemistry 50, 21-43.
- 390 Leino, R.L., McCormick, J.H., 1984. Morphological and morphometrical changes in chloride
391 cells of the gills of *Pimephales promelas* after chronic exposure to acid water. Cell. Tissue
392 Res. 236, 121-128.
- 393 Leino, R.L., McCormick, J.H., Jensen, K.M., 1987a. Changes in gill histology of fathead
394 minnows and yellow perch transferred to soft water or acidified soft water with particular
395 reference to chloride cells. Cell. Tissue Res. 250 : 389-399.
- 396 Leino, R.L., McCormick, J.H., Anderson, J.G., 1987b. Histopathological changes in the gills
397 of pearl dace, *Semotilus margarita*, and fathead minnows, *Pimephales promelas*, from
398 experimentally acidified Canadian lakes. Can. J. Fish. Aquat. Sci. 44, 126-134.

- 399 Lepori, F., Barbieri, A., Ormerod, S. J., 2003. Effects of episodic acidification on
400 macroinvertebrate assemblages in Swiss Alpine streams. *Freshwater Biol.* 48, 1873–1885.
- 401 Lignot, J.H., Spanings-Pierrot, C., Charmentier, G., 2000. Osmoregulatory capacity as a tool
402 in monitoring the physiological condition and the effect of stress in crustaceans.
403 *Aquaculture* 191, 209-245.
- 404 Lee, R.M., Gerking, S.D., Jezierska, B., 1983. Electrolyte balance and energy mobilization in
405 acid-stressed rainbow trout *Salmo gairdneri* and their relation to reproductive stress.
406 *Environmental Biology of Fishes* 8, 115–123.
- 407 Lemly, A.D., Smith, R.J.F., 1985. Effects of acute exposure to acidified water on the
408 behavioural response of Fathead Minnows, *Pimephales promelas*, to chemical feeding
409 stimuli. *Aquat. Toxicol.* 6, 625-36.
- 410 Likens, G.E., Butker, T.J., Buso, D.C., 2001. Long- and short-term changes in sulphate
411 deposition : effects of the 1990 Clean Air Act Amendments. *Biogeochemistry* 52, 1-11.
- 412 Massabuau, J.C., 1985. Pluies acides et physiologie des animaux aquatiques. *Bull. Soc.*
413 *Ecophysiol.* 10, 59-74.
- 414 McCahon, C.P., Pascoe, D., 1988. Cadmium toxicity to the freshwater amphipod *Gammarus*
415 *pulex* (L.) during the moult cycle. *Freshwater Biol.* 19, 197-203.
- 416 McCahon, C.P., Poulton, M.J., 1991. Lethal and sub-lethal effects of acid, aluminium and
417 lime on *Gammarus pulex* during repeated simulated episodes in a welsh stream.
418 *Freshwater Biol.* 25, 169-178.
- 419 McDonald, D.G., Reader, J.P., Dalziel T.R.K., 1989. The combined effects of pH and trace
420 metals on fish ionoregulation. In: Morris, R., Taylor, E.W., Brown, D.J.A., Brown, J.A.
421 (Eds.), *Acid toxicity and aquatic animals*. Cambridge University Press, Cambridge, pp
422 221-242.

- 423 McMahon, B.R., Stuart, S.A., 1989. The physiological problem of crayfish in acid water. In:
424 Morris R, Taylor EW, Brown DJA & Brown JA (Eds.), Acid toxicity and aquatic animals.
425 Cambridge University Press, Cambridge, pp 171-200.
- 426 Maltby, L., 1994. Stress, shredders and streams: using *Gammarus* energetics to assess water
427 quality. In: Sutcliffe, D.W., (Ed.), Water quality and stress indicators in marine and
428 freshwater ecosystems : linking levels of organisation (individual, population,
429 communities). Freshwater Biological association, Ambleside, pp 98-110.
- 430 Masson, N., Guerold, F., Dangles, O., 2002. Use of blood parameters in fish to assess acidic
431 stress and chloride pollution in French running waters. *Chemosphere* 47, 467-473.
- 432 Merrett, W.J., Rutt, G.P., Weatherley, N.S., Thomas, S.P., Ormerod, S.J., 1991. The response
433 of macroinvertebrates to low pH and increased aluminium concentrations in Welsh
434 streams: multiple episodes and chronic exposure. *Arch. Hydrobiol.* 121, 115-125.
- 435 Muniz, I.P., 1991. Freshwater acidification: Its effects on species and communities of
436 freshwater microbes, plants and animals. *Proc. R. Soc. Edinburgh* 97b, 227-254.
- 437 Naylor, C., Pindar, L., Calow, P., 1990. Inter- and intraspecific variation in sensitivity to
438 toxins; the effects of acidity and zinc on the freshwater crustaceans *Asellus aquaticus* (L.)
439 and *Gammarus pulex* (L.). *Water Res.* 24 (6), 757-762.
- 440 O'Brien, A.K., Rice, K.C., Kennedy, M.M., Bricker, O., 1993. Comparison of episodic
441 acidification of mid-atlantic upland and coastal plain stream. *Water Resour. Res.* 29 (9),
442 3029-3039.
- 443 O'Brien, A.K., Eshleman, K.N., 1995. Episodic acidification of a costal plain stream in
444 Virginia. *Water Air Soil Poll.* 89, 291-316.
- 445 Økland, J., Økland, K.A., 1986. The effects of acid deposition on benthic animals in lakes and
446 stream. *Experimentia* 42, 471-486.

- 447 Ormerod, S.J., Jenkins, A., 1994. The biological effect of acid episodes. In: Wright, R.,
448 Steinberg, C. (Eds.), *Acidification; Past, Present and Future*, Dahlem Workshop, Wiley,
449 Chichester, pp 259-272.
- 450 Pöckl, M., 1992. Effects of temperature, age and body size on moulting and growth in the
451 freshwater amphipods *Gammarus fossarum* and *G. roeseli*. *Freshwater Biol.* 27, 211-225.
- 452 Pöckl, M., 1995. Laboratory studies on growth, feeding, moulting and mortality in the
453 freshwater amphipods *Gammarus fossarum* and *G. roeseli*. *Arch. Hydrobiol.* 134 (2), 223-
454 253.
- 455 Potts, W.T.W., McWilliams, P.G., 1989. The effect of hydrogen and aluminum ions on fish
456 gills. In: Morris, R., Taylor, E.W., Brown, D.J.A., Brown, J.A. (Eds.), *Acid toxicity and*
457 *aquatic animals*. Cambridge University Press, Cambridge, pp 201-220
- 458 Pynnönen, K., 1991. Influence of aluminum and protons on the electrolyte homeostasis in the
459 Unionidae *Anodonta anatina* and *Unio pictorum*. *Arch. Environ. Contam. Toxicol.* 20,
460 218-225.
- 461 Pynnönen, K., 1994. Hemolymph gases, acid-base status, and electrolyte concentration in the
462 freshwater clams *Anodonta anatina* and *Unio pictorium* during exposure to and recovery
463 from acidic conditions. *Physiol. Zool.* 67 (6), 1544-1559.
- 464 Rupprecht, R., 1992. Zu den Ursachen der unterschiedlichen pH-Empfindlichkeit von
465 *Gammarus pulex* L., *G. fossarum* Koch (Amphipoda) und *Asellus aquaticus* L. (Isopoda).
466 *Int. Revue ges. Hydrobiologia* 77 (2), 255-264.
- 467 Steven, M., Seiler, S.M., Andrew, M., Turner, A.M., 2004. Growth and population size of
468 crayfish in headwater streams: individual- and higher-level consequences of acidification.
469 *Freshwater Biol.* 49, 870–881.
- 470 Skjelkvale, B.L., Torseth, K., Aas, W., Andersen, T., 2001. Decrease in acid deposition-
471 recovery in Norwegian waters. *Water Air Soil Poll.* 130, 1433-1438.

- 472 Soulsby, C., 1995. Contrasts in storm event hydrochemistry in an acidic afforested catchment
473 in upland Wales. *J. Hydrol.* 170, 159-179.
- 474 Stoddart, J.L., Jeffries, D.S., Lukewille, A., Clair, T.A., Dillon, P.J., Driscoll, C.T., Forsius,
475 M., Johannessen, M., Kahl, J.S., Kellogg, J.H., Kemp, A., Mannio, J., Monteith, D.T.,
476 Murdoch, P.S., Patrick, S., Rebsdorf, A., Skjelkvale, B.L., Stainton, M.P., Traaen, T., van
477 Dam, H., Webster, K.E., Wieting, J., Wilander, A., 1999. Regional trends in aquatic
478 recovery from acidification in North America and Europe. *Nature*, 401 (6753), 575-578.
- 479 Sutcliffe, D.W., Carrick, T.R., 1973. Studies on mountain streams in the English Lake
480 District. I. PH, calcium and the distribution of invertebrates in the River Duddon.
481 *Freshwater Biol.* 3, 437-462.
- 482 Sutcliffe, D.W., Hildrew, A.G., 1989. Invertebrate communities in acid streams. In: Morris,
483 R., Taylor, E.W., Brown, D.J.A., Brown, J.A. (Eds), *Acid toxicity and aquatic animals*.
484 Cambridge University Press, Cambridge, pp 13-29.
- 485 Tang, D., Lydersen, E., Seip, H.M., Angell, V., Eilertsen, O., Larssen, T., Liu, G., Kong, G.,
486 Mulder, J., Semb, A., Solberg, S., Torseth, K., Vogt, R.D., Xiao, J., Zhao, D., 2001.
487 Integrated monitoring program on acidification of Chinese terrestrial systems (impacts) –
488 A Chinese- Norwegian collaboration project. *Water Air Soil Poll.* 130, 1073-1078.
- 489 Taylor, E.J., Rees, E.M., Pascoe, D., 1994. Mortality and drift-related response of the
490 freshwater amphipod *Gammarus pulex* (L.) exposed to natural sediments, acidification and
491 copper. *Aquat. Toxicol.* 29, 83-101.
- 492 Thorjörn, L., Seip, H.M., Semb, A., Mulder, J., Muniz, I., Vogt, R.D., Lydersen, E., Angell,
493 V., Tang, D., Eilertsen, O., 1999. Acid deposition and its effects in China : An overview.
494 *Environmental Science & Policy* 2, 9-24.

- 495 Tierney, A.J., Atema, J., 1986. Effect of acidification on the behavioural response of
496 Crayfishes (*Orconectes virilis* and *Procambarus acutus*) to chemical stimuli. *Aquat.*
497 *Toxicol.* 9, 1-11.
- 498 Van Sickle, J., Baker, J.P., Simonin, H.A., Baldigo, B.P., Kretser, W.A., Sharpe, W.E., 1996.
499 Episodic acidification of small streams in the northeastern United States : Fish mortality in
500 field bioassays. *Ecol. Appl.* 6 (2), 408-421.
- 501 Weatherley, N.S., Ormerod, S.J., 1991. The importance of acid episodes in determining
502 faunal distributions in Wales. *Freshwater Biol.* 25, 71-84.
- 503 Wheatly, M.G., Gannon, A.T., 1995. Ion regulation in crayfish : freshwater adaptation and the
504 problem of molting. *Amer. Zool.* 35, 49-59.
- 505 Wigington, P.J.Jr., Baker, J.P., Dewalle, D.R., Kretser, W.A., Murdoch, P.S., Simonin, H.A.,
506 Van Sickle, J., McDowell, M.K., Peck, D.V., Barchet, W.R., 1996. Episodic acidification
507 of small streams in the northeastern United States : Episodic Response Project. *Ecol. Appl.*
508 6 (2), 374-388.
- 509 Willoughby, L.G., 1988. The ecology of *Baetis muticus* and *Baetis rhodani* (Insecta,
510 Ephemeroptera), with special emphasis on acid water backgrounds. *Int. Rev. Ges.*
511 *Hydrobiol.* 73, 259-273.
- 512 Willoughby, L.G., Mappin, R.G., 1988. The distribution of *Ephemerella ignita*
513 (Ephemeroptera) in streams : the role of pH and food resources. *Freshwater Biol.* 19, 145-
514 155.
- 515 Wood, C.M., & Rogano, M.S., 1986. Physiological responses to acid stress in crayfish
516 (*Orconectes*) : Hemolymph ions, acid-base status, and exchanges with the environment.
517 *Can. J. Fish. Aquat. Sci.* 43, 1017-1026.

518 Wood, C.M., 1989. The physiological problems of fish in acid water. In: Morris, R., Taylor,
519 E.W., Brown, D.J.A., Brown, J.A. (Eds.), Acid toxicity and aquatic animals. Cambridge
520 University Press, Cambridge, pp 125-152.

521 Wright, D.A., Frain, J.W., 1981. The effect of calcium on cadmium toxicity in the freshwater
522 amphipod, *Gammarus pulex* (L.). Arch. Environm. Contam. Toxicol. 10, 321-328.

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541 **Table 1.** Mean and SD values of chemical parameters of each exposure stream (n = 4). ANC:
 542 Acid Neutralizing Capacity.

	La Maix		Ravine		Gentil Sapin	
	Mean	SD	Mean	SD	Mean	SD
pH	7,3	0,0	5,5	0,1	4,5	0,1
ANC ($\mu\text{eq l}^{-1}$)	599,3	10,4	5,5	3,5	-19,4	6,5
conductivity ($\mu\text{S cm}^{-1}$)	78,6	2,2	30,5	2,0	29,9	1,0
Temperature ($^{\circ}\text{C}$)	10,3	0,3	11,9	0,5	12,9	0,5
Ca^{2+} ($\mu\text{mol l}^{-1}$)	281,7	5,2	55,9	5,6	38,8	0,8
Mg^{2+} ($\mu\text{mol l}^{-1}$)	187,1	4,7	35,7	2,4	25,9	1,3
Na^{+} ($\mu\text{mol l}^{-1}$)	47,6	0,0	52,0	4,8	37,0	2,0
K^{+} ($\mu\text{mol l}^{-1}$)	45,7	0,6	42,5	2,7	25,7	0,8
SO_4^{2-} ($\mu\text{mol l}^{-1}$)	63,4	0,4	72,5	5,1	50,7	2,6
NO_3^{-} ($\mu\text{mol l}^{-1}$)	62,8	1,0	49,6	5,5	80,3	3,4
Cl^{-} ($\mu\text{mol l}^{-1}$)	38,7	1,2	42,2	2,5	32,1	1,4
Total Al ($\mu\text{mol l}^{-1}$)	0,4	0,2	4,4	4,8	15,9	1,3

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558 **Table 2.** Summary of 2-way analysis of variance. F, F ratio; p, probability; d.f., degrees of
 559 freedom; n.s., not significant.

560

	[Cl ⁻] hemolymph			[Na ⁺] hemolymph			Survival		
	d.f.	F	p	d.f.	F	p	d.f.	F	p
a) Exposure effect									
Factor a (Stream)	2	34,1	< 10 ⁻³	2	46,2	< 10 ⁻³	2	110,4	< 10 ⁻³
Factor b (Exposure time)	3	18,0	< 10 ⁻³	3	23,0	< 10 ⁻³	3	29,8	< 10 ⁻³
Interaction	6	4,6	< 10 ⁻³	6	6,7	< 10 ⁻³	6	20,0	< 10 ⁻³
b) Recovery effect									
<i>Slightly acid exposure (Ravine)</i>									
Factor a (Exposure time)	3	3,2	< 5*10 ⁻²	3	1,7	n.s.	2	0,3	n.s.
Factor b (Recovery time)	5	14,1	< 10 ⁻³	5	17,5	< 10 ⁻³	5	2,9	< 5*10 ⁻²
Interaction	15	2,0	< 5*10 ⁻²	15	3,2	< 10 ⁻³	10	0,3	n.s.
<i>Strongly acid exposure (Gentil sapin)</i>									
Factor a (Exposure time)	3	3,9	< 5*10 ⁻²	3	0,5	n.s.	2	11,7	< 10 ⁻³
Factor b (Recovery time)	5	37,3	< 10 ⁻³	5	53,6	< 10 ⁻³	5	0,7	n.s.
Interaction	15	5,2	< 10 ⁻³	15	7,9	< 10 ⁻³	10	0,7	n.s.

561

562

563

564

565

566

567


568

569

570

571

572



573

574 **Figure 1.** Location of the 3 study sites in north-eastern France.

575

576

577

578

579

580

581

582

583

584


585

586

587

588

589



590

591 **Figure 2.** Experimental design of the exposure and recovery experiment

592

593

594

595

596


597

598


599

600


601



625 **Figure 3.** Mean (\pm SD) [Cl⁻] haemolymph (a), [Na⁺] haemolymph (b), and survival (c) of
626 *Gammarus fossarum* exposed to circumneutral, slightly and strongly acidic streams waters.
627 Significant differences against T₀ are indicated by asterisks (Fischer's Least Significant
628 Difference test; *: p<0.05; **: p<0.01; ***: p<0.001).



647 **Figure 4.** Mean (\pm SD) [Cl⁻] haemolymph (a) and [Na⁺] haemolymph, (b) and survival (c) of
 648 *Gammarus fossarum* exposed to slightly acidic waters (Ravine) and transferred in the
 649 circumneutral stream (La Maix) to test recovery capacity. Significant differences against T₀
 650 are indicated by asterisks (Fischer's Least Significant Difference test; *: p<0.05; **: p<0.01;
 651 ***: p<0.001). t₀ represents the last exposure time before organism transfer in circumneutral
 652 stream.



675 **Figure 5.** Mean (\pm SD) [Cl⁻] haemolymph (a) and [Na⁺] haemolymph, (b) and survival (c) of
 676 *Gammarus fossarum* exposed to strongly acidic waters (Gentil sapin) and transferred in the
 677 circumneutral stream (La Maix) to test recovery capacity. Significant differences against T₀
 678 are indicated by asterisks (Fischer's Least Significant Difference test; *: p<0.05; **: p<0.01;
 679 ***: p<0.001). t₀ represents the last exposure time before organism transfer in circumneutral
 680 stream.