



Double Dirichelet Averages and Complex B-Splines

Peter Massopust

► To cite this version:

Peter Massopust. Double Dirichelet Averages and Complex B-Splines. SAMPTA'09, May 2009, Marseille, France. pp.Poster session. hal-00453557

HAL Id: hal-00453557

<https://hal.science/hal-00453557>

Submitted on 5 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Double Dirichlet Averages and Complex B-Splines

Peter Massopust ^(1,2)

(1) Institute for Biomathematics and Biometry, Helmholtz Zentrum München, Neuberberg, Germany.

(2) Center of Mathematics, Technische Universität München, Garching, Germany.

massopust@ma.tum.de

Abstract:

A relation between double Dirichlet averages and multivariate complex B-splines is presented. Based on this relationship, a formula for the computation of certain moments of multivariate complex B-splines is derived.

1. Introduction

Recently, a new class of B-splines with complex order z , $\operatorname{Re} z > 1$, was introduced in [4]. It was shown that complex B-splines generate a multiresolution analysis of $L^2(\mathbb{R})$. Unlike the classical cardinal B-splines, complex B-splines B_z possess an additional modulation and phase factor in the frequency domain:

$$\widehat{B}_z(\omega) = \widehat{B}_{\operatorname{Re} z}(\omega) e^{i \operatorname{Im} z \ln |\Omega(\omega)|} e^{-\operatorname{Im} z \arg \Omega(\omega)},$$

where $\Omega(\omega) := (1 - e^{-i\omega})/(i\omega)$. The existence of these two factors allows the extraction of additional information from sampled data and the manipulation of images.

In [6] and [9], some further properties of complex B-splines were investigated. In particular, connections between complex derivatives of Riemann-Liouville or Weyl type and Dirichlet averages were exhibited. Whereas in [6] the emphasis was on univariate complex B-splines and their applications to statistical processes, multivariate complex B-splines were defined in [9] using a well-known geometric formula for classical multivariate B-splines [7, 10]. It was also shown that Dirichlet averages are especially well-suited to explore the properties of multivariate complex B-splines. Using Dirichlet averages, several classical multivariate B-spline identities were generalized to the complex setting. There also exist interesting relationships between complex B-splines, Dirichlet averages and difference operators, several of which are highlighted in [5].

This short paper presents a generalization of some results found in [3, 12] to complex B-splines. For this purpose, the concept of double Dirichlet average [1] was introduced and its definition extended via projective limits to an infinite-dimensional setting suitable for complex B-splines. Moments of complex B-splines are defined and a formula for their computation in terms of a special double Dirichlet average presented.

2. Complex B-Splines

Let $n \in \mathbb{N}$ and let Δ^n denote the standard n -simplex in \mathbb{R}^{n+1} :

$$\Delta^n := \left\{ u := (u_0, \dots, u_n) \in \mathbb{R}^{n+1} \mid \begin{array}{l} u_j \geq 0; \\ j = 0, 1, \dots, n; \sum_{j=0}^n u_j = 1 \end{array} \right\}.$$

The extension of Δ^n to infinite dimensions is done via projective limits. The resulting infinite-dimensional standard simplex is given by

$$\Delta^\infty := \left\{ u := (u_j)_j \in (\mathbb{R}_0^+)^{\mathbb{N}_0} \mid \sum_{j=0}^\infty u_j = 1 \right\},$$

and endowed with the topology of pointwise convergence, i.e., the weak*-topology. We denote by $\mu_b = \varprojlim \mu_b^n$ the projective limit of *Dirichlet measures* μ_b^n on the n -dimensional standard simplex Δ^n with density

$$\frac{\Gamma(b_0) \cdots \Gamma(b_n)}{\Gamma(b_0 + \cdots + b_n)} u_0^{b_0-1} u_1^{b_1-1} \cdots u_n^{b_n-1}. \quad (1)$$

Here, $\Gamma : \mathbb{C} \setminus \mathbb{Z}_0^- \rightarrow \mathbb{C}$ denotes the Euler Gamma function. Let $\mathbb{R}^+ := \{x \in \mathbb{R} \mid x > 0\}$ and let $\mathbb{C}_+ := \{z \in \mathbb{C} \mid \operatorname{Re} z > 0\}$.

Definition 1 ([6]). Given a weight vector $b \in \mathbb{C}_+^{\mathbb{N}_0}$ and an increasing knot sequence $\tau := \{\tau_k\}_k \in \mathbb{R}^{\mathbb{N}_0}$ with the property that $\lim_{k \rightarrow \infty} \sqrt[k]{\tau_k} \leq \varrho$, for some $\varrho \in [0, e)$, a complex B-spline $B_z(\bullet \mid b; \tau)$ of order z , $\operatorname{Re} z > 1$, with weight vector b and knot sequence τ is a function satisfying

$$\int_{\mathbb{R}} B_z(t \mid b; \tau) g^{(z)}(t) dt = \int_{\Delta^\infty} g^{(z)}(\tau \cdot u) d\mu_b(u) \quad (2)$$

for all $g \in \mathcal{S}(\mathbb{R})$.

Here, $\mathcal{S}(\mathbb{R})$ denotes the space of Schwartz functions on \mathbb{R} , and $\tau \cdot u = \sum_{k \in \mathbb{N}_0} \tau_k u_k$ for $u = \{u_k\}_{k \in \mathbb{N}_0} \in \Delta^\infty$. In addition, we used the Weyl or Riemann-Liouville fractional derivative [8, 11, 13] of complex order z , $\operatorname{Re} z > 0$, $W^z : \mathcal{S}(\mathbb{R}) \rightarrow \mathcal{S}(\mathbb{R})$, defined by

$$(W^z f)(x) := \frac{(-1)^n}{\Gamma(\nu)} \frac{d^n}{dx^n} \int_x^\infty (t-x)^{\nu-1} f(t) dt,$$

with $n = \lceil \operatorname{Re} z \rceil$, and $\nu = n - z$. Here $\lceil \cdot \rceil : \mathbb{R} \rightarrow \mathbb{Z}$, $x \mapsto \min\{n \in \mathbb{Z} \mid n \geq x\}$, denotes the *ceiling function*. To simplify notation, we write $f^{(z)}$ for $W^z f$.

It is easy to show that the univariate complex B-spline $B_z(t \mid b; \tau)$ is an element of $L^2(\mathbb{R})$ [5].

Remark 2. For finite $\tau = \tau(n)$ and $b = b(n)$ and $z := n \in \mathbb{N}$, (2) defines also *Dirichlet splines* if g is chosen in $C^n(\mathbb{R})$. For, Dirichlet splines $D_n(\cdot \mid b; \tau)$ of order n are defined as those functions for which

$$\int_{\mathbb{R}} g^{(n)}(t) D_n(t \mid b; \tau) dt = \int_{\Delta^n} g^{(n)}(\tau \cdot u) d\mu_b(u),$$

holds true for $\tau \in \mathbb{R}^{n+1}$ and for all $g \in C^n(\mathbb{R})$, and thus for $g \in \mathcal{S}(\mathbb{R})$.

To define a multivariate analogue of the univariate complex B-splines, we proceed as follows. Let $\lambda \in \mathbb{R}^s \setminus \{0\}$ be a direction, and let $g : \mathbb{R} \rightarrow \mathbb{C}$ be a function. The *ridge function* corresponding to g is defined as $g_\lambda : \mathbb{R}^s \rightarrow \mathbb{C}$,

$$g_\lambda(x) = g(\langle \lambda, x \rangle) \quad \text{for all } x \in \mathbb{R}^s.$$

We denote the canonical inner product in \mathbb{R}^s by $\langle \bullet, \bullet \rangle$ and the norm induced by it by $\| \bullet \|$.

Definition 3 ([9]). Let $\tau = \{\tau^n\}_{n \in \mathbb{N}_0} \in (\mathbb{R}^s)^{\mathbb{N}_0}$ be a sequence of knots in \mathbb{R}^s with the property that

$$\exists \varrho \in [0, e) : \limsup_{n \rightarrow \infty} \sqrt[n]{\|\tau^n\|} \leq \varrho. \quad (3)$$

The multivariate complex B-spline $B_z(\bullet \mid b, \tau) : \mathbb{R}^s \rightarrow \mathbb{C}$ of order z , $\operatorname{Re} z > 1$, with weight vector $b \in \mathbb{C}_+^{\mathbb{N}_0}$ and knot sequence τ is defined by means of the identity

$$\int_{\mathbb{R}^s} g(\langle \lambda, x \rangle) B_z(x \mid b, \tau) dx = \int_{\mathbb{R}} g(t) B_z(t \mid b, \lambda \tau) dt, \quad (4)$$

where $g \in \mathcal{S}(\mathbb{R})$, and where $\lambda \in \mathbb{R}^s \setminus \{0\}$ such that $\lambda \tau := \{\langle \lambda, \tau^n \rangle\}_{n \in \mathbb{N}_0}$ is separated.

As consequence of the fact that $B_z(\bullet \mid b; \tau) \in L^2(\mathbb{R})$, one obtains from the above definition that $B_z(\bullet \mid b, \tau) \in L^2(\mathbb{R}^s)$ [5]. Moreover, it follows from the Hermite-Genocchi formula for the univariate complex B-splines $B_z(\bullet \mid b, \lambda \tau)$ and (4), that $B_z(x \mid b, \tau) = 0$, when $x \notin [\tau]$, the convex hull of τ .

3. Dirichlet Averages

Let Ω to be a nonempty open convex set in \mathbb{C}^s , $s \in \mathbb{N}$, and let $b \in \mathbb{C}_+^{\mathbb{N}_0}$. Let $f \in \mathcal{S}(\Omega) := \mathcal{S}(\Omega, \mathbb{C})$ be a measurable function. For $\tau \in \Omega^{\mathbb{N}_0} \subset (\mathbb{C}^s)^{\mathbb{N}_0}$ and $u \in \Delta^\infty$, define $\tau \cdot u$ to be the bilinear mapping $(\tau, u) \mapsto \sum_{i=1}^\infty u_i \tau^i$. The infinite sum exists if there exists a $\varrho \in [0, e)$ so that

$$\limsup_{n \rightarrow \infty} \sqrt[n]{\|\tau^n\|} \leq \varrho. \quad (5)$$

Here, $\| \cdot \|$ now denotes the canonical Euclidean norm on \mathbb{C}^s . (See also [6].)

Definition 4. Let $f : \Omega \subset \mathbb{C}^s \rightarrow \mathbb{C}$ be a measurable function. The Dirichlet average $F : \mathbb{C}_+^{\mathbb{N}_0} \times \Omega^{\mathbb{N}_0} \rightarrow \mathbb{C}$ over Δ^∞ is defined by

$$F(b; \tau) := \int_{\Delta^\infty} f(\tau \cdot u) d\mu_b(u),$$

where $\mu_b = \varprojlim \mu_b^n$ is the projective limit of Dirichlet measures on the n -dimensional standard simplex Δ^n .

We remark that the Dirichlet average is holomorphic in $b \in (\mathbb{C}_+)^{\mathbb{N}_0}$ when $f \in C(\Omega, \mathbb{C})$ for every fixed $\tau \in \Omega^{\mathbb{N}_0}$. (See [2] for the finite-dimensional case and [9] for the infinite-dimensional setting.)

Definition 5. [1] Let $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$ be continuous. Let $b \in \mathbb{C}_+^{k+1}$ and $\beta \in \mathbb{C}_+^{\varkappa+1}$. Suppose that for fixed $k, \varkappa \in \mathbb{N}$, $X \in \mathbb{C}^{(k+1) \times (\varkappa+1)}$ and that the convex hull $[X]$ of X is contained in Ω . Then the double Dirichlet average of f is defined by

$$\mathcal{F}(b; X; \beta) := \int_{\Delta^k} \int_{\Delta^\varkappa} f(u \cdot Xv) d\mu_b^k(u) d\nu_\beta^\varkappa(v),$$

where $u \cdot Xv := \sum_{i=0}^k \sum_{j=0}^\varkappa u_i X_{ij} v_j$.

Note that $\mathcal{F}(b; X; \beta)$ is holomorphic on Ω in the elements of b, β , and X .

We again use projective limits to extend the notion of double Dirichlet average to an infinite-dimensional setting. To this end, let $u, v \in \Delta^\infty$ and let $\mu_b = \varprojlim \mu_b^n$ and $\nu_\beta = \varprojlim \nu_\beta^n$ be the projective limits of Dirichlet measures μ_b^n and ν_β^n of the form (1) on the n -dimensional standard simplex, where $b, \beta \in \mathbb{C}_+^{\mathbb{N}_0}$. Now suppose that $X \in \mathbb{C}^{\mathbb{N}_0 \times \mathbb{N}_0}$ is a infinite matrix with the property that $\sum_{i=0}^\infty \sum_{j=0}^\infty |X_{ij}|$ converges. Let

$$u \cdot Xv := \sum_{i=0}^\infty \sum_{j=0}^\infty u_i X_{ij} v_j.$$

Suppose that $\Omega \subset \mathbb{C}$ contains the convex hull $[X]$ of X and that $f : \Omega \rightarrow \mathbb{C}$ is continuous. The double Dirichlet average of f over Δ^∞ is then given by

$$\mathcal{F}(b; X; \beta) := \int_{\Delta^\infty} \int_{\Delta^\infty} f(u \cdot Xv) d\mu_b(u) d\nu_\beta(v). \quad (6)$$

(We use the same symbol for the (double) Dirichlet average over Δ^∞ and its finite-dimensional projections Δ^n .) It is easy to show that

$$\mathcal{F}(b; X; \beta) = \int_{\Delta^\infty} F(\beta; uX) d\mu_b(u), \quad (7)$$

where $uX := \{\langle u, X_j \rangle\}_{j \in \mathbb{N}_0}$, with X_j denoting the j -column of X .

We note that $\mathcal{F}(b; X; \beta)$ is holomorphic in the elements of b, β , and X over Δ^∞ .

For $z \in \mathbb{C}_+$, we define

$$\mathcal{F}^{(z)}(b; X; \beta) := \int_{\Delta^\infty} \int_{\Delta^\infty} f^{(z)}(u \cdot Xv) d\mu_b(u) d\nu_\beta(v).$$

(See also [9] for the case of a single Dirichlet average.)

4. Double Dirichlet Averages and Complex B-Splines

Assume now that the matrix X is real-valued and of the form $X_{ij} = 0$, for $i \geq s$ and all $j \in \mathbb{N}_0$, some $s \in \mathbb{N}$. In other words, $X \in \mathbb{R}^{s \times \mathbb{N}_0}$.

Theorem 6. *Suppose that $\beta \in \mathbb{R}_+^\infty$ and that $\operatorname{Re} z > 1$. Let $b := (b_0, b_1, \dots, b_{s-1}) \in \mathbb{R}^s$ be such that $\sum_{i=0}^{s-1} b_i \notin -\mathbb{N}_0$. Assume that $f \in \mathcal{S}(\mathbb{R}^+)$. Further assume that uX is separated for all $u \in \Delta^{s-1}$. Then*

$$\mathcal{F}^{(z)}(b; X; \beta) = \int_{\mathbb{R}^s} B_z(x | \beta, X) F^{(z)}(b; x) dx.$$

Proof. We prove the formula first for $b \in \mathbb{R}_+^s$. To this end, we identify $u = (u_0, u_1, \dots, u_{s-1}, 0, 0, \dots) \in \Delta^\infty$ with $(u_0, u_1, \dots, u_{s-1}) \in \Delta^{s-1}$. By the Hermite-Genocchi formula for complex B-splines (see [6] and to some extend [9]), we have that

$$\begin{aligned} F^{(z)}(\beta; uX) &= \int_{\Delta^\infty} f^{(z)}(u' \cdot uX) d\mu_\beta(u') \\ &= \int_{\mathbb{R}} f^{(z)}(t) B_z(t | \beta, uX) dt \end{aligned}$$

Substituting this expression into (7) and using (4) gives

$$\begin{aligned} \mathcal{F}^{(z)}(b; X; \beta) &= \\ &= \int_{\Delta^\infty} \int_{\mathbb{R}^s} f^{(z)}(\langle u, x \rangle) B_z(x | \beta, uX) dx d\mu_b(u). \end{aligned}$$

Interchanging the order of integration yields the statement for $b \in \mathbb{R}_+^s$. To obtain the general case $b \in \mathbb{R}^s$, we note that by Theorem 6.3-7 in [2], the Dirichlet average F can be holomorphically continued in the b -parameters provided $\sum_{i=0}^{s-1} b_i \notin -\mathbb{N}_0$. \square

Remark 7. Theorem 6 extends Theorem 6.1 in [12] to complex B-splines and the Δ^∞ -setting.

5. Moments of Complex B-Splines

Following [2], we define the R -hypergeometric function $R_a(b; \tau) : \mathbb{R}_+^s \times \Omega^s \rightarrow \mathbb{C}$ by

$$R_a(b; \tau) := \int_{\Delta^{s-1}} (\tau \cdot u)^a d\mu_b^{s-1}(u), \quad (8)$$

where $\Omega := H$, H a half-plane in $\mathbb{C} \setminus \{0\}$, if $a \in \mathbb{C} \setminus \mathbb{N}$, and $\Omega := \mathbb{C}$, if $a \in \mathbb{N}$. It can be shown (see [2]) that R_{-a} , $a \in \mathbb{C}_+$, has a holomorphic continuation in τ to \mathbb{C}_0 , where $\mathbb{C}_0 := \{\zeta \in \mathbb{C} \mid -\pi < \arg \zeta < \pi\}$.

Taking in the definition of the double Dirichlet average (6) for f the real-valued function $t \mapsto t^{-c}$, where $c := \sum_{i=0}^{s-1} b_i$, the resulting double Dirichlet average is denoted by $\mathcal{R}_{-c}(b; X; \beta)$ and generalizes power functions. The corresponding single Dirichlet average $R_{-c}(b; x)$, where $x = (x_0, \dots, x_{s-1})$, is given by

$$R_{-c}(b; x) = \prod_{i=0}^{s-1} x_i^{-b_i}, \quad x \notin [X]. \quad (9)$$

(See, [2], (6.6-5).)

Now, let $p = (p_0, p_1, \dots, p_{s-1}) \in \mathbb{R}^s$, $s \in \mathbb{N}$, be a multi-index all of whose components satisfy $p_i < -\frac{1}{2}$. The moment $M_{|p|}^{(z)}(b; X)$ of order $|p| := \sum_{i=1}^s p_i$ of the complex B-spline $B_z(\bullet | \beta, X)$ is defined by

$$M_{|p|}^{(z)}(b; X) := \int_{\mathbb{R}^s} x^p B_z(x | \beta, X) dx.$$

Note that since $B_z(\bullet | \beta, X) \in L^2(\mathbb{R}^s)$ and $B_z(\bullet | \beta, X) = 0$, for $x \notin [X]$, an easy application of the Cauchy-Schwartz inequality shows that the above integral exists provided the multi-index p satisfies the aforementioned condition on its components.

Using a result from [8], namely Property 2.5 (b), and requiring that $\operatorname{Re} z < \operatorname{Re} c$, we substitute the function $f := \frac{\Gamma(c-z)}{\Gamma(c)} (\bullet)^{-(c-z)}$ into (8) to obtain

$$R_{-(c-z)}^{(z)}(b; x) = R_{-c}(b; x) = \prod_{i=0}^{s-1} x_i^{b_i}.$$

The above considerations together with Theorem 6 immediately yield the next result.

Corollary 8. *Suppose that $\beta \in \mathbb{R}_+^\infty$ and that $\operatorname{Re} z > 1$. Let $b := (b_0, b_1, \dots, b_{s-1}) \in (-\infty, -\frac{1}{2})^s$ be such that $c := \sum_{i=0}^{s-1} b_i \notin -\mathbb{N}_0$. Moreover, suppose that $\operatorname{Re} z < \operatorname{Re} c$. Then*

$$M_{-c}^{(z)}(b; X) = \mathcal{R}_{-(c-z)}^{(z)}(b; X; \beta). \quad (10)$$

6. Acknowledgements

This work was partially supported by the grant MEXT-CT-2004-013477, Acronym MAMEBIA, of the European Commission.

References:

- [1] B. C. Carlson. Appell functions and multiple averages. *SIAM J. Math. Anal.*, 2(3):420–430, August 1971.
- [2] B. C. Carlson. *Special Functions of Applied Mathematics*. Academic Press, New York, 1977.
- [3] B. C. Carlson. B-splines, hypergeometric functions, and Dirichlet averages. *J. Approx. Th.*, 67:311–325, 1991.
- [4] B. Forster, T. Blu, and M. Unser. Complex B-splines. *Appl. Comp. Harmon. Anal.*, 20:261–282, 2006.
- [5] B. Forster and P. Massopust. Multivariate complex B-splines, Dirichlet averages and difference operators. *accepted SAMPTA 2009*, 2009.
- [6] B. Forster and P. Massopust. Statistical encounters with complex B-splines. *Constr. Approx.*, 29(3):325–344, 2009.
- [7] S. Karlin, C. A. Micchelli, and Y. Rinott. Multivariate splines: A probabilistic perspective. *Journal of Multivariate Analysis*, 20:69–90, 1986.

- [8] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo. *Theory and Applications of Fractional Differential Equations*. Elsevier B. V., Amsterdam, The Netherlands, 2006.
- [9] P. Massopust and B. Forster. Multivariate complex B-splines and Dirichlet averages. *to appear in J. Approx. Th.*, 2009.
- [10] C. A. Micchelli. A constructive approach to Kergin interpolation in \mathbb{R}^k : Multivariate B-splines and Lagrange interpolation. *Rocky Mt. J. Math.*, 10(3):485–497, 1980.
- [11] K. S. Miller and B. Ross. *An Introduction to the Fractional Calculus and Fractional Differential Equations*. Wiley, New York, 1993.
- [12] E. Neuman and P. J. Van Fleet. Moments of Dirichlet splines and their applications to hypergeometric functions. *J. Comput. and Appl. Math.*, 53:225–241, 1994.
- [13] S. G. Samko, A. A. Kilbas, and O. I. Marichev. *Fractional Integrals and Derivatives*. Gordon and Breach Science Publishers, Minsk, Belarus, 1987.