The *Leishmania infantum* cytosolic SIR2 related protein 1 (LiSIR2RP1) is an NAD⁺-dependent deacetylase and ADP-ribosyltransferase

Joana Tavares, Ali Ouaiissi, Nuno Santarém, Denis Sereno, Baptiste Vergnes,
Paula Sampaio, Anabela Cordeiro-Da-Silva

► To cite this version:

Joana Tavares, Ali Ouaiissi, Nuno Santarém, Denis Sereno, Baptiste Vergnes, et al.. The *Leishmania infantum* cytosolic SIR2 related protein 1 (LiSIR2RP1) is an NAD⁺-dependent deacetylase and ADP-ribosyltransferase. *Biochemical Journal*, 2008, 415 (3), pp.377-386. 10.1042/BJ20080666 . hal-00479004

HAL Id: hal-00479004

<https://hal.science/hal-00479004>

Submitted on 30 Apr 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The *Leishmania infantum* cytosolic SIR2 related protein 1 (LiSIR2RP1) is an NAD⁺-dependent deacetylase and ADP-ribosyltransferase

Joana TAVARES ^{*†§}, Ali OUAISSI ^{*‡§¹}, Nuno SANTARÉM ^{*†}, Denis SERENO [§], Baptiste VERGNES [§], Paula SAMPAIO ^{*} and Anabela CORDEIRO-DA-SILVA ^{*†¹²}

^{*} IBMC – Instituto de Biologia Molecular e Celular, Universidade do Porto, Rua do Campo Alegre, 823, 4150-180 Porto, Portugal, [†] Laboratório de Bioquímica, Faculdade de Farmácia da Universidade do Porto, R. Aníbal Cunha n.º 164, 4050-047 Porto, Portugal, [‡] INSERM, CNRS, UMR 5235, Université de Montpellier 2, Bât 24 - CC 107, Pl. Eugène Bataillon, 34095 Montpellier Cx 5, France and [§]Centre IRD de Montpellier, 911 avenue Agropolis, 34394 Montpellier, France.

¹ Both authors have supervised the implementation of the work.

² To whom correspondence should be addressed (e-mail: cordeiro@ibmc.up.pt).

Short (page heading) title: LiSIR2RP1 deacetylates and ADP-ribosylates α -tubulin

Keywords: Parasites, LiSIR2RP1, α -tubulin, ADP-ribosyltransferase, NAD⁺-dependent deacetylase, cytoskeleton.

Abbreviations used:

FBS, Foetal bovine serum; IPTG, Isopropyl- β -D-thiogalactopyranoside; MAA, Medium for axenic amastigotes; OAADPr, O-acetyl-ADP-ribose; RT, Room temperature; SIR2, Silent Information Regulator 2; TCA, trichloroacetic acid; TSA, trichostatin A; DTT, dithiothreitol; PI, propidium iodide; wt, wild type;

Synopsis

Proteins of the SIR2 family are characterized by a conserved catalytic domain that exerts unique NAD⁺-dependent deacetylase activity on histone and various other cellular substrates. Previous reports from our team have identified a *Leishmania infantum* gene encoding a cytosolic protein termed LiSIR2RP1 that belongs to the SIR2 family. Targeted disruption of one LiSIR2RP1 gene allele led to decreased amastigote virulence, *in vitro* as well as *in vivo*. In the present study, attempts were made for the first time to explore and characterize the LiSIR2RP1 enzymatic functions. The LiSIR2RP1 exhibited robust NAD⁺-dependent deacetylase and ADP-ribosyltransferase activities. Moreover, LiSIR2RP1 is capable of deacetylating tubulin, either in dimers or, when present, in taxol-stabilized microtubules or in promastigote and amastigote extracts. Furthermore, the immunostaining of parasites revealed a partial co-localization of α -tubulin and LiSIR2RP1 with punctuate labelling, seen on the periphery of both promastigote and amastigote stages. Isolated parasite cytoskeleton reacted with antibodies showed that part of LiSIR2RP1 is associated to the cytoskeleton network of both promastigote and amastigote forms. Moreover, the Western blot analysis of the detergent's soluble and insoluble fractions of promastigote and amastigote forms revealed the presence of α -tubulin in the insoluble fraction, and the LiSIR2RP1 distributed in both soluble and insoluble fractions of promastigotes as well as amastigotes. Collectively, our results demonstrate that LiSIR2RP1 is an NAD⁺-dependent deacetylase that also exerts an ADP-ribosyltransferase activity. The fact that tubulin could be among the targets of LiSIR2RP1 may have significant implications during the remodelling of the parasite's morphology and its interaction with the host cell.

INTRODUCTION

The yeast Silent Information Regulator 2 (SIR2) protein is the founding member of a large conserved family of proteins that are present in many organisms ranging from bacteria to humans [1]. This protein is an NAD⁺-dependent histone deacetylase that modulates the chromatin structure, and silences the transcription at the level of the silent mating type loci, telomeres and ribosomal DNA [2-4]. In addition to epigenetic silencing, the yeast SIR2 also has a role in DNA repair, recombination and DNA replication [5]. The first evidence that the SIR2 proteins could be involved in functions other than transcriptional silencing came from the presence of the *Leishmania major* SIR2 related protein in cytoplasmic granules [6]. Seven SIR2 genes were identified in humans, and the proteins classified as Sirtuins (SIRT1-7) present several subcellular localizations [7, 8]. However, not all of the sirtuin family members seem to be active deacetylases [9, 10]. SIRT1 is localized in the nucleus and deacetylates the histones H3 and H4, p53, TAF168 and PCAF/MyoD [4, 11-14]. Despite the SIRT2's cytosolic localization and its ability to deacetylate α -tubulin, this protein shuttles the nucleus during G2/M transition and thus plays a role in the regulation of microtubule dynamics and cell cycle progression [15-17]. The SIRT3 was found in the mitochondria controlling the activity of the acetyl-CoA synthetase 2 [18]. Histone and non-histone substrates are deacetylated by sirtuins in the presence of NAD⁺ and lead to the production of the deacetylated protein, nicotinamide and O-acetyl-ADP-ribose (OAADPr) [19, 20]. Several questions have been raised on understanding the sirtuin's NAD⁺ dependency, since there is no energetic reason that favours the coupling. Among the possibilities is the connection between the proteins deacetylation and the cell's metabolic status or the formation of a second messenger, OAADPr [20].

Some members of the SIR2 family, like the yeast SIR2, the human SIRT2, the mouse SIRT6, the *T. brucei* SIR2RP1 and the *P. falciparum* SIR2, exhibit ADP-ribosyltransferase activity [7, 21-24]. This kind of enzymatic activity was for the first time attributed to this family of proteins due to the ability of the *Salmonella typhimurium* SIR2 homologue, Cob B, to compensate for the loss of Cob T in the cobalamine biosynthesis [25].

Additionally, sirtuins, and particularly the one associated with caloric restriction regimens in several organisms like *C. elegans*, yeast and fly might also be involved in longevity [26-28].

The agent of leishmaniasis, the protozoan parasite *Leishmania*, is responsible for serious diseases characterized by diverse clinical manifestations varying from localized ulcerative skin lesions and destructive mucosal inflammation to the disseminated visceral infection also known as Kala azar. The visceral leishmaniasis caused by *L. infantum* is a serious problem throughout the Mediterranean basin, where dogs are the most important reservoirs of the parasite. A vertebrate host becomes infected by *Leishmania* after the bite of a sandfly (*Phlebotomus* and *Lutzomyia spp*) during a blood meal, through the inoculation of infective flagellated promastigotes that invade or are phagocytosed by local or recruited host cells. In the phagolysosomes, the promastigotes will differentiate into non-flagellated amastigotes that multiply and are able to infect other adjacent or distant macrophages. The classical treatment is currently unsatisfactory due to side effects, the appearance of resistances and the need for increased efficacy in immunosuppressed patients, especially due to HIV co-infection. During the last few years, our group has been interested in the *Leishmania* SIR2 proteins due to the role of *LiSIR2RP1* in the parasite's survival and virulence [29, 30].

Although few reports have disclosed the SIR2 protein function in the protozoan parasites, such as *Plasmodium falciparum*, whose PfSIR2, localized in the nucleus, is

involved in *var* gene silencing through heterochromatin formation [31], and *Trypanosoma brucei*, which has a chromosome-associated NAD⁺-dependent deacetylase and ADP ribosyltransferase that confers resistance to DNA damage [22], little is known about the *Leishmania* cytosolic LiSIR2RP1 biological properties. Here, we provide molecular and biochemical evidences supporting the notion that LiSIR2RP1 is an NAD⁺-dependent deacetylase enzyme that may also express ADP-ribosyltransferase activity. In addition, we show that the protein deacetylates α -tubulin and is partially associated with the microtubule network. Further studies on LiSIR2RP1 may provide new insights into the relationships among cytoskeleton dynamic, stage transition and parasite-host cell interaction.

EXPERIMENTAL

Parasites

The *L. infantum* (clone MHOM/MA/67/ITMAP-263) wild type (wt) promastigotes were grown at 27°C in RPMI 1640 medium (Cambrex) supplemented with 10% heat inactivated foetal bovine serum (FBS-Cambrex), 2mM L-glutamine (Cambrex), 20mM Hepes (Cambrex), 100U/ml penicillin (Cambrex) and 100µg/ml streptomycin (Cambrex). *L. infantum* axenic amastigote forms were grown at 37°C with 5% CO₂ in a cell free medium called MAA/20 (medium for axenic amastigotes). MAA/20 consisted of modified medium 199 (GIBCO) with Hank's balanced salt solution supplemented with 0.5% trypto-casein (Oxoid), 15mM D-glucose (SIGMA), 5mM glutamine (Cambrex), 4mM NaHCO₃ (Sigma), 0.023mM bovine hemin (FLUKA), and 25mM HEPES to a pH of 5.8 and supplemented with 20% heat inactivated FBS (Cambrex). The heterozygote LiSIR2^{+/−} derived from the wt clone was generated as previously described, and maintained in their respective selective medium [30]. The levels of LiSIR2 synthesis in LiSIR2^{+/−} and wt parasites were evaluated by western blotting, using an anti-LiSIR2 monoclonal antibody as described in previous reports [30].

Heterologous Expression and Purification of LiSIR2RP1

The LiSIR2RP1 encoding sequence (GenBank accession number AF487351) from *L. infantum* genomic DNA was amplified by PCR using two primers, a forward primer (5'-CAATTTGCATATGACAGCGTCTCCGAGAGCGCC A-3') and a reverse primer (5'-CCCAAGCGAATTCTCACGTCTCAT TCGGCGC-3'). The amplified product of about 1200 bp was cloned into the *Nde*I and *Eco*RI restriction sites of the prokaryotic expression vector pET28a (Novagen, Madison, WI, USA) and the fusion protein LiSIR2RP1 with an N-terminal tail of six histidines was produced in *E. coli* BL21 (DE3). The transformants were grown in a Luria Broth medium containing 30µg/ml of kanamycin, and the recombinant protein expression was induced in mid-logarithmic phase with 1mM of isopropyl-β-D-thiogalactopyranoside (IPTG) for 3 hours at 37°C. The bacterial pellet was re-suspended in a buffer containing 500mM NaCl, 20mM Tris-HCl pH 7.6 and disrupted by sonication. The clarified supernatant was applied to a Ni-NTA agarose column (Quiagen) and the recombinant LiSIR2RP1 protein eluted with an imidazole gradient from 25 to 1000mM. The eluted fractions were controlled for the LiSIR2RP1 presence using an SDS-PAGE. The fractions containing the recombinant protein were pooled, applied to PD-10 columns (Amersham Pharmacia Biotech) and eluted with PBS pH 7.4.

Deacetylase assay

The NAD⁺-dependent deacetylase activity was evaluated by using a commercially available CycLex SIRT1/Sir2 deacetylase fluorometric assay kit (Cyclex Co., Ltd., Nagano, Japan). The enzymatic reaction containing 0.25µg of rLiSIR2RP1 was carried out in accordance with the manufacturer's instructions and monitored every 30 seconds for 1 hour. The results are expressed as the rate of reaction for the first 20 minutes, due to the linear correlation between the fluorescence and this period of time.

Tubulin deacetylation assay

The tubulin deacetylation reactions were performed using either purified tubulin (Pure, Cytoskeleton Inc, Denver, USA) or *L. infantum* promastigote and axenic amastigote extracts, prepared in PEM buffer (80mM PIPES pH 6.8, 1mM MgCl₂, 1mM EGTA) by sonication. In both cases, the tubulin was either left in heterodimers, or polymerized in PEM buffer in the presence of 20µM taxol (Cytoskeleton Inc, Denver, USA) and 1mM

GTP (Cytoskeleton Inc, Denver, USA) for 45 minutes at 37°C. The reactions were carried out in deacetylase buffer (10mM Tris-HCl, pH 8.0 and 10mM NaCl), in the presence or absence of 1mM NAD⁺ with constant agitation at room temperature (RT) for 3 hours. The reactions were stopped by adding 5x Laemmli sample buffer and the proteins separated by 10% SDS-PAGE and western blotted.

ADP-ribosylation assays

The ADP-ribosylation assays were performed according to Garcia-Salcedo et al. [22]. Briefly, the assays were carried out with 1µg of either of the following proteins: rLiSIR2RP1, recombinant human SIRT1 (rSIRT1, Cyclex kit), or with the recombinant *L. infantum* cytosolic peroxiredoxin (LicTXNPx), in the presence of 5µCi of [³²P] NAD and 5µg of the following acceptor proteins: calf thymus histones (Sigma) or BSA (Sigma). Some assays were conducted with 0.5 µg of rLiSIR2RP1 or rLicTXNPx in the presence of 2.5µCi of [³²P] NAD and 1µg of pure tubulin (Cytoskeleton Inc, Denver, USA) or calf thymus histones (Sigma). The reactions were performed at RT for 2 hours in a buffer containing 150mM NaCl, 10mM dithiothreitol (DTT), 50mM Tris-HCl pH 8.8, stopped with Laemmli loading buffer and dried in Whatman paper after electrophoresis, and exposed to a film 2 for 12 hours at -80°C. As for quantitative experiments, the reaction products were precipitated by 20% trichloroacetic acid (TCA), washed and counted after the addition of 200µl scintillation fluid.

Antibodies

Antibodies to LiSIR2RP1 were obtained from BALB/c mouse immunization. The mouse was injected intraperitoneally with 50µg of rLiSIR2RP1 in 200µl of PBS three times at a 7 day interval. Two weeks after the final immunization, the serum was collected. The mouse monoclonal antibody, IIIG4, against the LmSIR2, was obtained according to Vergnes et al., 2002. Mouse monoclonal anti-α-tubulin antibody (clone DM1A) and rabbit polyclonal anti-α-tubulin antibody were purchased from NeoMarkers, USA. Mouse monoclonal anti-acetylated tubulin antibody (clone 6-11B-1) came from Sigma Chemical Co, USA. Rabbit anti-*Trypanosoma brucei* RAB11 antibodies were a gift from Prof. Field, MC.

Cytoskeleton Preparation

The cytoskeleton preparation was performed according to Schneider et al. [32]. Briefly, the *L. infantum* promastigotes and axenic amastigotes were collected and washed three times in ice-cold PBS. The parasites were resuspended at a concentration of 4x10⁷ cells/ml in a buffer containing 10mM MOPS, pH 6.9, 0.1mM EGTA, 1mM MgSO₄ and 0.1% Triton X-100 supplemented with protease inhibitor cocktail (Roche), and incubated on ice for 10 minutes followed by a 5 minute 3000g centrifugation. The supernatant was collected and concentrated to 1/10 of the initial volume, and a 0.25 volume of 4x Laemmli sample buffer was added. The pellet was washed once with PBS and resuspended in 1/20 of the initial volume with a 1:1 mixture of PBS/Laemmli sample buffer. For immunofluorescence, the pellets were prepared as described and after being washed with PBS, they were attached to poly-L-lysine-coated slides.

Immunofluorescence assays

L. infantum promastigotes and axenic amastigotes were fixed with 4% paraformaldehyde in PBS for 1 hour at RT. After several washes, the parasites were made permeable with 0.1% (v/v) Triton X-100 in PBS and incubated for 30 minutes in PBS containing 1% bovine serum albumin (PBS-1% BSA). Parasites were then

incubated with primary antibodies diluted in PBS-1% BSA for 1 hour at RT, washed several times with PBS, and then incubated for 1 hour with the Alexa Fluor labelled secondary antibodies (Molecular Probes) diluted in PBS-1% BSA at RT. The immunofluorescence assays using *Leishmania* cytoskeletons were performed as described above, with the exception of the fixation process, which was done using 100% methanol at 20°C for 15 minutes, and the permeabilization step, which was not performed. Washed parasites or cytoskeletons were mounted in Vectashield with DAPI (Vector Laboratories, Burlingame, CA). Z-series optical sections were collected through the Zeiss Axio Imager Z1 microscope (Carl Zeiss, Oberkochen, Germany), using an AxioCam. Data stacks were deconvolved using either the Axiovision AxioVs40 V 4.2.0.0 (Carl Zeiss) or the Huygens Essential version 3.0.2p1 (Scientific Volume Imaging, Hilversum, The Netherlands). Images were treated using Adobe Photoshop CS (Adobe Microsystems, CA).

RESULTS

LiSIR2RP1 is an NAD⁺ dependent deacetylase and ADP-ribosyltransferase

The *LiSIR2RP1* gene encoding sequence was cloned in the bacterial expression vector pET28a. The fusion protein with an N-terminal tail of six histidines was purified under non-denaturing conditions, with the objective of studying its enzymatic activity. After purification and dialysis, the protein showed a good level of purity as evaluated by SDS-PAGE Coomassie blue staining (Fig. 1, panel A, lane 1). As expected, the recombinant protein was recognized by the monoclonal antibodies anti-*L. major* SIR2 protein (III G4), previously generated by our group (Fig. 1, panel A, lane 2) [29]. Moreover, the mouse polyclonal antibodies produced against the rLiSIR2RP1 recognized the parasite protein in total extracts from different parasite developmental stages (Fig 1, panel B: Pl, logarithmic phase promastigotes; Ps, stationary phase promastigotes; A, axenic amastigotes). It is well established that the proteins belonging to the SIR2 family exhibit NAD⁺ dependent deacetylase activity due to the presence of a well conserved enzymatic core domain of ~250 amino acids [7, 8], and this seems also to be the case for LiSIR2RP1. Indeed, as shown in Fig 2A, LiSIR2RP1 expresses an NAD⁺-dependent deacetylase activity that is not inhibited by 1 μ M of the class I and II NAD⁺ independent deacetylase inhibitor, trichostatin A (TSA). A dose-dependent increase of the deacetylase activity catalyzed by rLiSIR2RP1 was observed with increasing concentrations of NAD⁺ up to $\approx$ 0.5 mM. However, concentrations of NAD⁺ higher than 1 mM led to a reduction in the deacetylase activity (Fig 2B). This phenomenon is likely to be due to the release of nicotinamide, which in turn could inhibit the LiSIR2RP1 enzymatic activity (see below). This enzyme exhibits co-factor specificity for the oxidative form of NAD that was confirmed by the absence of deacetylase activity in the presence of the reduced form of NAD (data not shown). Nicotinamide is one of the several products originated in the deacetylation reaction catalyzed by the SIR2 family members. As shown in Fig 2C, the rLiSIR2RP1 is inhibited by nicotinamide in a dose-dependent manner, and the calculated nicotinamide concentration that inhibits 50% of the enzymatic activity (IC₅₀) is 39.35 $\pm$ 5.01 μ M. Several enzymatic reactions in the presence or absence of 50 μ M of nicotinamide and different NAD⁺ concentrations were performed to determine whether NAD⁺ and nicotinamide are competing for the same binding site. The Lineweaver-Burk plot of the data demonstrates that nicotinamide is an NAD⁺ non-competitive inhibitor (Fig 2D). Some members of the SIR2 family can mediate the transference of the ribose 5'-phosphate from nicotinic acid mononucleotide to the amino acid residues of BSA, histones or SIR2 proteins themselves [7, 21-24]. To assess whether LiSIR2RP1 is an ADP-ribosyltransferase, [³²P]NAD was used as a donor and BSA or calf thymus histones as acceptor proteins. A non-related *Leishmania* recombinant protein, rLicTXNPx1, which is produced and purified as the rLiSIR2RP1, was included as a control. The analysis of the reaction products by autoradiography revealed that LiSIR2RP1 was capable of inducing ADP-ribosylation of the histones and BSA (Fig. 3A). However, the label intensity was stronger when histones rather than BSA were used as acceptor proteins. Additionally, the parasite protein was subject to auto ADP-ribosylation when incubated with the histones. No signal was detected when the proteins were incubated with [³²P]NAD without an acceptor protein. A quantitative ADP-ribosyltransferase reaction catalyzed by rLiSIR2RP1 using histones as the acceptor protein was carried out to characterize the factors affecting the extent of the enzymatic reaction. Increasing the concentration of rLiSIR2RP1 (2 times) led to a decrease in the extent of the enzymatic transfer of the label, whereas a 2-time increase of histone concentration led to an enhancement of ADP-ribosyl transfer to the acceptor

target protein (Fig. 3B). This unexpected reduction of the ribosylation reaction when adding more enzyme, might be due to some level of protein aggregation. In fact, we have experienced the purification of recombinant LiSIR2RP1 enzyme and found that after dialysis, increasing the concentration of LiSIR2RP1 may lead to the formation of some aggregates.

Given that LiSIR2RP1 is a cytosolic protein, we decided to verify whether it was capable of ADP-ribosylating tubulin. Thus, purified tubulin or calf thymus histones as a control were incubated with LiSIR2RP1 or LicTXNPx in the presence of [32 P]NAD. The autoradiography evidenced a positive but weak labelling of tubulin and LiSIR2RP1 itself, when compared with the reaction where histones were used as substrates. Even though tubulin ADP-ribosylation is an already reported modification, this is the first evidence of such ADP-ribosylation transfer mediated by a protein belonging to the SIR2 family [33]. This modification seems to either inhibit the *in vitro* assembly of microtubules or induce microtubule depolymerization of assembled microtubules [33].

The LiSIR2RP1 is a microtubule associated protein

We have previously reported that both *L. major* (LmSIR2RP1) and *L. infantum* (LiSIR2RP1) are cytosolic proteins [6, 29]. A similar cytosolic localization was reported in the case of human SIRT2 (yeast SIR2 ortholog) and a human NAD $^{+}$ -independent histone deacetylase, HDAC6, both proteins have been shown to express α -tubulin deacetylase activity [15, 34]. Based on these observations and on the fact that tubulin is one of the most notable cytoplasmic proteins subjected to modification by acetylation [35], we first investigated the cell distribution of LiSIR2RP1 and α -tubulin. The immunostaining of both promastigote and amastigote forms revealed partial co-localization of α -tubulin with LiSIR2RP1 (Fig 4A). In addition to the positive signals observed in the parasite cytoplasm, punctuate labelling could be seen on the periphery of both promastigote and amastigote stages. This observation suggests the possible association of LiSIR2RP1 with the cytoskeleton. In order to examine this possibility, attempts were made to isolate the parasite cytoskeleton according to the method described by Schneider et al. [32], that allows subpellicular microtubules and the flagellar axoneme microtubules to be separated from the membrane and cytosolic components by a non-ionic detergent extraction using Triton X-100, followed by centrifugation. The cytoskeleton preparations were then reacted with the anti-LiSIR2RP1 antibodies. The results shown in (Fig 4B) suggest that part of LiSIR2RP1 is associated to the cytoskeleton network of both promastigote and amastigote forms. Interestingly, punctuate labelling could also be seen over the flagellar axoneme microtubules when the cytoskeleton preparation was reacted with anti-LiSIR2RP1 antibodies. In agreement with this, the Western blot analysis of the detergent-soluble and insoluble fractions of promastigote and amastigote forms revealed, as expected, the presence of α -tubulin in the insoluble fraction, and of LiSIR2RP1 distributed in both soluble and insoluble fractions of promastigotes and amastigotes (Fig 4C). Although in the case of the promastigotes the detergent-insoluble fraction contains a band of ~64.6kD recognized by the anti-LiSIR2RP1 antibodies, its nature is presently unknown. It might be due to non-complete reduction of LiSIR2RP1 present in a molecular complex with other parasite polypeptides. As a control, we have used an antibody to *T. brucei* RAB11 that cross-react with *Leishmania* homologous protein already used by others to control the cytoskeletons purity preparations of *L. major* [37]. Indeed, the RAB11 is a protein found on the membrane of endosomes not associated with the cytoskeleton [36]. As shown in Fig 4A, the RAB11 has different cell distribution pattern when compared to tubulin and LiSIR2RP1. Interestingly, no positive signal could be

seen in immunofluorescence preparations of parasite cytoskeletons reacted with anti-RAB11 antibodies (Fig 4B). Moreover, the WB membranes dehybridized and re-probed with anti-RAB11 antibodies showed that the RAB11 protein is present exclusively in the detergent soluble fraction of parasite extracts (Fig 4C), supporting therefore the specificity of the LiSIR2RP1 and α -tubulin interaction.

LiSIR2RP1 accumulates in the posterior pole of promastigotes after taxol treatment

Leishmania is a protozoan parasite characterized by a digenetic life cycle exhibiting a particular range of cell shapes mostly defined by their internal cytoskeleton. The latter is composed mainly of microtubules that are polymers of repeating α/β tubulin heterodimers, and a variety of minor components known as microtubule-associated proteins. The microtubules play an important role in many cellular processes such as mitosis, cell shape, motility, intracellular vesicle transport and organelle position. Several compounds have been described as microtubule-interacting agents, based on their ability to induce or inhibit the tubulin assembly in microtubules. Even though a high degree of tubulin amino acid conservation is found throughout the evolution, there are some reports suggesting that colchicine and benzimidazoles, two classical mammalian tubulin inhibitors, had only a slight effect on the trypanosomatids tubulin polymerization [38, 39]. On the other hand, taxol, a microtubule network stabilizing agent, affects the parasite cytoskeleton in a dose-dependent manner [40]. The *Leishmania* parasites were treated for 16 hours with 1 μ M of the microtubule stabilizing agent, taxol, or with 5 μ g/ml of the microtubule network disrupting agent, nocodazole, to analyze if the SIR2RP1 cell distribution is affected by changes in the parasite cytoskeleton. Treatment of promastigotes with taxol led to a weak but distinguishable accumulation of SIR2RP1 in the posterior pole of the cell (Fig. 5A). The treatment of parasites with nocodazole modifies the tubulin distribution, being mainly localized near the nucleus in the parasite body. However, LiSIR2RP1 showed no particular modification in terms of cell distribution, compared to non-treated cells. Although the morphology of treated parasites differed from non-treated ones, their viability measured by propidium iodide (PI) incorporation and flow cytometry analysis revealed no differences. Indeed, after 16 hr of treatment with 1 μ M taxol or 5 μ g/ml nocodazole, the percentage of PI negative parasites $\pm$ standard deviation was 98.2 $\pm$ 0.1 and 97.6 $\pm$ 0.2 % respectively, compared to 97.4 $\pm$ 0.1 % for untreated ones. Interestingly, after taxol treatment, acetylated α -tubulin accumulated in the posterior pole of the cell (Fig. 5B), coinciding with the high LiSIR2RP1 reactive zone seen in panel A. Moreover, punctuate labelling could be seen on the whole periphery of the parasites as well as the flagellum. These observations are in agreement with the notion of LiSIR2RP1 and tubulin being interacting proteins.

LiSIR2RP1 deacetylates α -tubulin

In most eukaryotes, tubulin is subject to several post-translational modifications, including detyrosination, acetylation, phosphorylation, palmitoylation, polyglutamylation and polyglycylation [41]. Tubulin acetylation occurs on lysine-40 of the α -tubulin subunit [42]. To test LiSIR2RP1's ability to deacetylate α -tubulin *in vitro*, several amounts of rLiSIR2RP1 were incubated with purified tubulin in the presence or absence of NAD⁺. The deacetylation of α -tubulin was analyzed by Western blot with specific antibodies to either total α -tubulin or its acetylated form. As shown in Fig.6, panel A, incubation of a defined amount of tubulin (0.5 μ g) with increasing quantities of rLiSIR2RP1 (0.125 to 0.5 μ g) led to a dose-dependent deacetylation of α -tubulin.

Interestingly, the addition of increasing concentrations of nicotinamide, a known physiological inhibitor of SIR2 deacetylases, led to significant inhibition of tubulin deacetylation by rLiSIR2RP1.

We further explored whether LiSIR2RP1 is able to deacetylate tubulin present within polymerized microtubules. Thus, addition of taxol and GTP to tubulin heterodimers led to microtubule polymerization. Under these conditions, LiSIR2RP1 still had the capacity to efficiently deacetylate tubulin. Moreover, the addition of increasing concentrations of nicotinamide in the assay samples led to the inhibition of LiSIR2RP1-mediated tubulin deacetylation (Fig. 6, panel B).

We then examined whether this activity also occurred when using taxol and GTP-treated parasite extracts as a source of polymerized tubulin. As shown in Fig. 6, panels C-D, when total extracts from promastigotes and amastigotes treated with taxol were used as a source of polymerized microtubules, LiSIR2RP1 was also able to deacetylate tubulin. Taken together, these observations demonstrate that tubulin is a substrate of LiSIR2RP1 deacetylase.

We further examined the LiSIR2RP1 NAD^+ -dependent deacetylase substrate's specificity by using acetylated BSA or cytochrome c and Western blot technique. In contrast to tubulin, the level of BSA and cytochrome c acetylation presented no changes after incubation with LiSIR2RP1 in the presence or absence of NAD^+ . This allows us to conclude that, at least from the approach we have used, LiSIR2RP1 does not deacetylate either BSA or cytochrome c, suggesting some substrate specificity of LiSIR2RP1 as a deacetylase (data not shown).

In order to evaluate whether LiSIR2RP1 is deacetylating tubulin *in vivo*, we have analyzed the ratio of acetylated tubulin over total α -tubulin in the wild type and single LiSIR2RP1 knockout promastigotes and axenic amastigotes. Only a slight difference in the level of tubulin acetylation between the wild type and single knockout axenic amastigotes was detected (Fig. 6, panel E). However, the disruption of a single allele from the LiSIR2RP1 gene may not be enough to affect the tubulin acetylation level, even though it is enough to reduce amastigotes ability to multiply, be it *in vitro* as free forms or inside macrophages, or *in vivo* in Balb/c mice [30].

DISCUSSION

In this report, we describe and characterize the enzymatic activities of the LiSIR2RP1 for the first time. Assuming that this protein is involved in the parasite's virulence and survival, it could constitute a new and potential drug target against leishmaniasis that needs to be exploited. We have demonstrated that, although LiSIR2RP1 differs from its *T. brucei* ortholog, TbSIR2RP1 [22] in terms of subcellular localization (cytosolic versus nuclear), it expresses similar enzymatic activities, being able to act as an NAD⁺-dependent deacetylase and to transfer [³²P]NAD⁺ to histones and, to a lesser extent, to BSA or tubulin. Initial reports on the yeast SIR2 have shown that the protein has a weak mono-ADP-ribosyltransferase activity and a more robust NAD⁺-dependent deacetylase activity [21]. This finding led to the general idea that the ADP-ribosyltransferase activity might be the result of a non-enzymatic reaction. However, our data and the investigations using TbSIR2RP1 [22], the human SIRT6 [23] or the *P. falciparum* Sir2 [24], provided evidence supporting the notion that these SIR2 family members could mediate the enzymatic transfer of the ADP-ribosyl group. Studies from Kowieski et al. [43] have recently shed light on the ADP-ribosylation mechanism catalyzed by the TbSIR2RP1. The authors demonstrated that ADP-ribosylation of histone by TbSIR2RP1 required an acetylated substrate. Additionally, two distinct ADP-ribosylation pathways that involve an acetylated substrate, NAD⁺ and TbSIR2RP1, were described. One is a non-catalytic reaction between the deacetylation product o-acetyl-ADP-ribose (or its hydrolysis product, ADPr) and histones, and the other a more efficient mechanism involving interception of an ADP-ribose-acetylpeptide-enzyme intermediate by a side-chain nucleophile from bound histone [43]. This might also explain the low level of tubulin ADP-ribosylation catalyzed by the *Leishmania* SIR2 protein. If the enzyme preferentially deacetylates tubulin and the ADP-ribosylation requires an acetylated protein, the occurrence of the latter will be dependent on the first one [43].

The LiSIR2RP1 possesses the expected NAD⁺-dependent deacetylase activity that characterizes the SIR2 family members, and one of their reaction products, nicotinamide, is an NAD⁺ non-competitive inhibitor. Several biological activities have been attributed to the proteins belonging to the SIR2 family, that, even though sharing a conserved enzymatic core domain, are flanked at their N- and C-terminal parts by divergent sequences between family members, which may promote the binding to specific substrates [44]. Two main Sirtuin substrates that have been identified are large fibrous macromolecular complexes, chromatin and tubulin [15]. The partial co-localization of *L. infantum* SIR2RP1 with α -tubulin and its association with the parasite cytoskeleton, suggested that tubulin could be a substrate of this enzyme as it is for the human enzymes SIRT2 and HDAC6 [15, 34]. Control tests using antibodies to a parasite cytosolic protein, RAB11, which confirmed the absence of RAB11-tubulin relationship, strongly support the notion of LiSIR2RP1-tubulin specific interaction. In fact, LiSIR2RP1 was either able to deacetylate tubulin in dimers or, when present, in taxol stabilized microtubules, or in total promastigote or amastigote extracts. Although the α -tubulin acetylation was already reported a few years ago, its importance in cell biological processes is not yet completely understood. The presence of acetylated tubulin in stable microtubules and the absence of highly acetylated microtubules in the fibroblast leading edge, that is a dynamic structure involved in cell motility, suggested that this post-translational modification could play a role in microtubule stability [45]. However, Palazzo et al., [46] demonstrated that enhanced tubulin acetylation does not increase the level of stable microtubules. Moreover, ciliated or flagellated protists can use a nonacetylatable α -tubulin with no observable ill effects on the microtubules or the

organisms [47]. By contrast, the level of tubulin acetylation seems to affect a number of cellular events that includes the infection of CD4⁺ T cells by HIV1 [48], the binding and transport of a kinesin 1 [49], the correct organization of the immune synapse [50], and the dynamics of cellular adhesions [51]. Interestingly, the promastigotes treatment with the microtubule stabilizing agent, taxol, increased the number of parasites showing an accumulation of LiSIR2RP1 in the posterior pole of the cell that is enriched in acetylated α -tubulin. It is likely that LiSIR2RP1 could bind to α -tubulin and just deacetylate it in a specific step during the parasite's life cycle. Additionally, the promastigotes infect macrophages through a receptor-mediated mechanism initiated at the posterior pole of the parasite [52]. One can hypothesize that the accumulation of LiSIR2RP1 in the posterior pole could be important to the contact site with the APC during the infection. Indeed, translocation and accumulation of HDAC 6, a human cytosolic deacetylase, in the immune synapse at the contact site of T cells with APC has recently been reported [50]. Furthermore, it is noteworthy that the influence of the deacetylase activity in the cell motility versus adhesion reported by Tran et al. [51] could also be relevant to the parasites when they are infecting the host cell.

The *Leishmania* parasites are subject to several modifications in their morphology during their complex life cycle. Although some environmental factors, such as pH, temperature and nutrients, can trigger morphological changes in the parasite, relatively little is known about the molecular processes that mediate the cellular remodelling. Further functional studies of LiSIR2RP1 could help disclose the role of the tubulin acetylation/deacetylation and/or ADP-ribosylation in the remodelling of the parasite's morphology and the infection of host cells.

ACKNOWLEDGEMENTS

This work has been funded by Fundação para a Ciência e Tecnologia (FCT) POCI 2010, co-funded by FEDER grant number POCI/SAU- FCF/59837/2004, Centre Franco Indien pour la promotion de la recherche avancée; Indo-French Centre for the promotion of advanced research CEFIPRA/IFCPAR grant number 3603, INSERM and IRD UR 008. JT is supported by a fellowship from FCT number SFRH/BD/18137/2004; NS has a fellowship in the above-mentioned FCT/FEDER project. The authors thank I. Silveira for her support in the radioactive experiments, A. Tomás for providing the LicTXNPx bacterial clone, Professor Field M.C. for the gift of the antibodies to TbRAB11, and A. Ferreira for the English revision of the manuscript.

REFERENCES

- 1 Brachmann, C.B., Sherman, J.M., Devine, S.E., Cameron, E.E., Pillus, L. and Boeke, J.D. (1995) The SIR2 gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. *Genes Dev.* **9**, 2888-2902
- 2 Bryk, M., Banerjee, M., Murphy, M., Knudsen, K.E., Garfinkel, D.J. and Curcio, M.J. (1997) Transcriptional silencing of Ty1 elements in the RDN1 locus of yeast. *Genes Dev.* **11**, 255-269
- 3 Smith, J.S. and Boeke, J.D. (1997) An unusual form of transcriptional silencing in yeast ribosomal DNA. *Genes Dev.* **11**, 241-254
- 4 Imai, S., Armstrong, C.M., Kaerberlein, M. and Guarente, L. (2000) Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* **403**, 795-800
- 5 Blander, G. and Guarente, L. (2004). The Sir2 family of protein deacetylases. *Annu. Rev. Biochem.* **73**, 417-435
- 6 Zemzoumi, K., Sereno, D., Francois, C., Guilvard, E., Lemesre, J.L., Ouassii, A. (1998) Leishmania major: cell type dependent distribution of a 43kDa antigen related to silent information regulatory-2 protein family. *Biol. Cell* **90**, 239-245
- 7 Frye, R.A. (1999) Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity. *Biochem. Biophys. Res. Commun.* **260**, 273-279
- 8 Frye R.A. (2000) Phylogenetic classification of prokaryotic and eukaryotic Sir2-like proteins. *Biochem. Biophys. Res. Commun.* **273**, 793-798
- 9 Mostoslavsky, R., Chua, K.F., Lombard, D.B., Pang, W.W., Fischer, M.R., Gellon, L. et al. (2006) Genomic instability and aging-like phenotype in the absence of the of mammalian SIRT6. *Cell* **124**, 315-329
- 10 Haigis, M.C., Mostoslavsky, R., Haigis, K.M., Fahie, K., Christodoulou, D.C., Murphy et al. (2006) SIRT4 inhibits glutamate dehydrogenase and opposes the effects of calorie restriction in pancreatic beta cells. *Cell* **126**, 941-954
- 11 Vaziri, H., Dessain, S.K., Ng Eaton, E., Imai, S.I., Frye, R.A., Pandita, T.K., Guarente, L. and Weinberg, R.A. (2001) hSIR2 (SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell* **107**, 149-59
- 12 Luo, J., Nikolaev, A.Y., Imai, S., Chen, D., Su, F., Shiloh, A., Guarente, L. and Gu, W. (2001) Negative control of p53 by Sir2alpha promotes cell survival under stress. *Cell* **107**, 137-148
- 13 Muth, V., Nadaud, S., Grummt, I., Voit, R. (2001) Acetylation of TAF(I)68, a subunit of TIF-IB/SL1, activates RNA polymerase I transcription. *EMBO J.* **20**, 1353-1362
- 14 Fulco, M., Schiltz, R.L., Iezzi, S., King, M.T., Zhao, P., Kashiwaya, Y., Hoffman, E., Veech, R.L. and Sartorelli, V. (2003) Sir2 regulates skeletal muscle differentiation as a potential sensor of the redox state. *Mol. Cell* **12**, 51-62
- 15 North, B.J., Marshall, B.L., Borra, M.T., Denu, J.M. and Verdin, E. (2003) The human Sir2 ortholog, SIRT2, is an NAD⁺-dependent tubulin deacetylase. *Mol. Cell.* **11**, 437-444
- 16 Dryden, S.C., Nahhas, F.A., Nowak, J.E., Goustin, A.S. and Tainsky MA. (2003) Role for human SIRT2 NAD-dependent deacetylase activity in control of mitotic exit in the cell cycle. *Mol. Cell Biol.* **23**, 3173-3185
- 17 Vaquero, A., Scher, M.B., Lee, D.H., Sutton, A., Cheng, H.L., Serrano, L., Sternglanz, R. and Reinberg D. (2006) SirT2 is a histone deacetylase with preference for histone H4 Lys 16 during mitosis. *Genes Dev.* **20**, 1256-1261

- 18 Schwer, B., Bunkenborg, J., Verdin, R.O., Andersen, J.S. and Verdin, E. (2006) Reversible lysine acetylation controls the activity of the mitochondrial enzyme acetyl-CoA synthetase 2. *Proc. Natl. Acad. Sci. U S A* **103**, 10224-10229
- 19 Tanny, J.C. and Moazed, D. (2001) Coupling of histone deacetylation to NAD breakdown by the yeast silencing protein Sir2: Evidence for acetyl transfer from substrate to an NAD breakdown product. *Proc. Natl. Acad. Sci. U S A* **98**, 415-420
- 20 Sauve, A.A., Celic, I., Avalos, J., Deng, H., Boeke, J.D. and Schramm, V.L. (2001) Chemistry of gene silencing: the mechanism of NAD⁺-dependent deacetylation reaction. *Biochemistry* **40**, 15456-15463
- 21 Tanny, J.C., Dowd, G.J., Huang, J., Hilz, H. and Moazed, D. (1999) An enzymatic activity in the yeast Sir2 protein that is essential for gene silencing. *Cell* **99**, 735-745
- 22 Garcia-Salcedo, J.A., Gijon, P., Nolan, D.P., Tebabi, P. and Pays, E. (2003) A chromosomal SIR2 homologue with both histone NAD-dependent ADP-ribosyltransferase and deacetylase activities is involved in DNA repair in *Trypanosoma brucei*. *EMBO J.* **22**, 5851-5862
- 23 Liszt, G., Ford, E., Kurtev, M. and Guarente, L. (2005) Mouse Sir2 homolog SIRT6 is a nuclear ADP-ribosyltransferase. *J. Biol. Chem.* **280**, 21313-21320
- 24 Merrick, C.J. and Duraisingh, M.T. (2007) *Plasmodium falciparum* Sir2: an unusual sirtuin with dual histone deacetylase and ADP-ribosyltransferase activity. *Eukaryot. Cell.* **6**, 2081-2091
- 25 Tsang, A.W. and Escalante-Semerena, J.C. (1998) CobB, a new member of the SIR2 family of eucaryotic regulatory proteins, is required to compensate for the lack of nicotinate mononucleotide:5,6-dimethylbenzimidazole phosphoribosyltransferase activity in cobT mutants during cobalamin biosynthesis in *Salmonella typhimurium* LT2. *J. Biol. Chem.* **273**, 31788-31794
- 26 Tissenbaum, H.A. and Guarente, L. (2001) Increased dosage of a sir-2 gene extends lifespan in *Caenorhabditis elegans*. *Nature* **410**, 227-230
- 27 Kaerberlein, M., McVey, M. and Guarente, L. (1999) The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes Dev.* **13**, 2570-2580
- 28 Rogina, B. and Helfand, S.L. (2004) Sir2 mediates longevity in the fly through a pathway related to calorie restriction. *Proc. Natl. Acad. Sci. U S A* **101**, 15998-156003
- 29 Vergnes, B., Sereno, D., Madjidian-Sereno, N., Lemesre, J.L. and Ouaisi, A. (2002) Cytoplasmic SIR2 homologue overexpression promotes survival of *Leishmania* parasites by preventing programmed cell death. *Gene* **296**, 139-50
- 30 Vergnes, B., Sereno, D., Tavares, J., Cordeiro-da-Silva, A., Vanhille, L., Madjidian-Sereno, N., Depoix, D., Monte-Alegre, A. and Ouaisi, A. (2005) Targeted disruption of cytosolic SIR2 deacetylase discloses its essential role in *Leishmania* survival and proliferation. *Gene* **363**, 85-96
- 31 Freitas-Junior, L.H., Hernandez-Rivas, R., Ralph, S.A., Montiel-Condado, D., Ruvalcaba-Salazar, O.K., Rojas-Meza A.P., Mancio-Silva, L., Leal-Silvestre R.J., Gontijo, A.M., Shorte, S. and Scherf, A. (2005) Telomeric heterochromatin propagation and histone acetylation control mutually exclusive expression of antigenic variation genes in malaria parasites. *Cell* **121**, 25-36
- 32 Schneider, A., Sherwin, T., Russell, D.G., Gull, K. and Seebeck, T. (1987) Subpellicular and flagellar microtubules of *Trypanosoma brucei* contain the same alpha tubulin isoforms. *J. Cell Biol.* **104**, 431-438

- 33 Scaife RM, Wilson L, Purich DL. (1992) Microtubule protein ADP-ribosylation in vitro leads to assembly inhibition and rapid depolymerization. *Biochemistry* **31**, 310-316
- 34 Hubbert, C., Guardiola, A., Shao, R., Kawaguchi, Y., Ito, A., Nixon, A., Yoshida, M., Wang, X.F. and Yao, T.P. (2002) HDAC6 is a microtubule-associated deacetylase. *Nature* **417**, 455-458
- 35 Kouzarides, T. (2000) Acetylation: a regulatory modification to rival phosphorylation? *EMBO J.* **19**, 1176-1179
- 36 Ambit, A., Fasel, N., Coombs, G.H., Mottram, J.C. (2008) An essential role for the *Leishmania* major metacaspase in cell cycle progression. *Cell Death. Differ.* **15**, 113-22
- 37 Jeffries, T.R., Morgan, G.W., Field, M.C. (2001) A developmentally regulated rab11 homologue in *Trypanosoma brucei* is involved in recycling processes. *J. Cell Sci.* **114**, 2617-26
- 38 MacRae, T.H. and Gull, K. (1990) Purification and assembly in vitro of tubulin from *Trypanosoma brucei brucei*. *Biochem. J.* **265**, 87-93
- 39 Vickerman, K. and Preston, T.M. (1970) Spindle microtubules in the dividing nuclei of trypanosomes. *J. Cell Sci.* **6**, 365-383
- 40 Baum, S.G., Wittner, M., Nadler, J.P., Horwitz, S.B., Dennis, J.E., Schiff, P.B. and Tanowitz, H.B. (1981) Taxol, a microtubule stabilizing agent, blocks the replication of *Trypanosoma cruzi*. *Proc. Natl. Acad. Sci. U S A* **78**, 4571-4575
- 41 Rosenbaum, J. (2000) Cytoskeleton: Functions for tubulin modifications at last. *Current Biology* **10**, 801-803
- 42 Nogales, E., Whittaker, M., Milligan, R.A. and Downing, K.H. (1999) High-resolution model of the microtubule. *Cell* **96**, 79-88
- 43 Kowieski, T.M., Lee, S., Denu, J.M. (2008) Acetylation-dependent ADP-ribosylation by *Trypanosoma brucei* Sir2. *J. Biol. Chem.* **283**, 5317-5326
- 44 Cuperus, G., Shafaatian, R. and Shore, D. (2000) Locus specificity determinants in the multifunctional yeast silencing protein Sir2. *EMBO J.* **19**, 2641-2651
- 45 Wadsworth, P. (1999) Regional regulation of microtubule dynamics in polarized, motile cells. *Cell Motil. Cytoskeleton* **42**, 48-59
- 46 Palazzo, A., Ackerman, B. and Gundersen, G.G. (2003) Cell biology: Tubulin acetylation and cell motility. *Nature* **16**, 230
- 47 Gaertig, J., Cruz, M.A., Bowen, J., Gu, L., Pennock, D.G. and Gorovsky MA. (1995) Acetylation of lysine 40 in alpha-tubulin is not essential in *Tetrahymena thermophila*. *J. Cell Biol.* **129**, 1301-1310
- 48 Valenzuela-Fernandez, A., Alvarez, S., Gordon-Alonso, M., Barrero, M., Ursa, A., Cabrero, J.R., Fernandez, G., Naranjo-Suarez, S., Yanez-Mo, M., Serrador, J.M., Munoz-Fernandez, M.A. and Sanchez-Madrid, F. (2005) Histone deacetylase 6 regulates human immunodeficiency virus type 1 infection. *Mol. Biol. Cell* **11**, 5445-5454.
- 49 Reed, N.A., Cai, D., Blasius, T.L., Jih, G.T., Meyhofer, E., Gaertig, J. and Verhey, K.J. (2006) Microtubule acetylation promotes kinesin-1 binding and transport. *Curr. Biol.* **16**, 2166-2172
- 50 Serrador, J.M., Cabrero, J.R., Sancho, D., Mittelbrunn, M., Urzainqui, A. and Sanchez-Madrid, F. (2004) HDAC6 deacetylase activity links the tubulin cytoskeleton with immune synapse organization. *Immunity* **20**, 417-428
- 51 Tran, A.D., Marmo, T.P., Salam, A.A., Che, S., Finkelstein, E., Kabarriti, R., Xenias, H.S., Mazitschek, R., Hubbert, C., Kawaguchi, Y., Sheetz, M.P., Yao, T.P.

- and Bulinski, J.C. (2007) HDAC6 deacetylation of tubulin modulates dynamics of cellular adhesions. *J. Cell. Sci.* **120**, 1469-1479
- 52 Courret, N., Frehel, C., Gouhier, N., Pouchelet, M., Prina, E., Roux, P. and Antoine, J.C. (2002) Biogenesis of *Leishmania*-harbouring parasitophorous vacuoles following phagocytosis of the metacyclic promastigote or amastigote stages of the parasites. *J. Cell Sci.* **111**, 2303-2316

FIGURE LEGENDS

Figure 1 LiSIR2RP1 characterization. The purity of the rLiSIR2RP1 was confirmed by SDS-PAGE Coomassie blue staining (A, lane 1). A monoclonal antibody IIIG4 to *L. major* recombinant SIR2RP1 showed strong positive signals when reacted in Western blot against rLiSIR2RP1 (A, lane 2). Western blot analysis of LiSIR2RP1 expression in total parasite extracts (B, PI-logarithmic promastigotes; Ps-stationary promastigotes and A-amastigotes) evidenced by using a mouse polyclonal antibody raised against the rLiSIR2RP1. Samples of protein extracts from each parasite developmental stage also reacted with anti- α -tubulin monoclonal antibody, DM1A.

Figure 2 LiSIR2RP1 is a NAD⁺-dependent deacetylase. The NAD⁺-dependent deacetylase activity was measured using a fluorometric deacetylase kit (CycLex SIRT1/Sir2 fluorometric assay kit) in the presence of 0.25 μ g of rLiSIR2RP1. LiSIR2RP1 is an NAD⁺-dependent deacetylase that is not inhibited by 1 μ M of TSA (A). The rate of the deacetylase reaction was measured in the presence of several NAD⁺ concentrations (B). Nicotinamide, a deacetylase reaction product, inhibited the LiSIR2RP1 NAD⁺-dependent deacetylase activity in a dose-dependent manner (C) and according to the Lineweaver-Burk plot, 50 μ M of nicotinamide promotes an NAD⁺ non-competitive inhibition (D). The results are expressed as the mean rate of reaction $\pm$ standard deviation for the first 20 minutes.

Figure 3 LiSIR2RP1 has ADP-ribosyltransferase activity. (A) The rLiSIR2RP1 ($\blacktriangleright$) ADP-ribosyltransferase activity was assessed using [³²P]NAD as donor and BSA (*) or calf thymus histones (#) as acceptor proteins. A non-related *Leishmania* recombinant protein, rLicTXNPx1 ($\rightarrow$), was included as controls. The reaction products were analyzed by SDS-PAGE Coomassie blue stained and by autoradiography. (B) Quantitative analysis of the effect of NAD⁺, acceptor protein or rLiSIR2RP1 amount on the ribosylation reaction. The reaction products were precipitated with TCA, collected, washed and then counted after the addition of scintillation liquid. (C) Assessment of tubulin ($\blacksquare$) ADP-ribosylation mediated by rLiSIR2RP1. Tubulin and histones were included as acceptor proteins, ³²P[NAD] as donor and rLicTXNPx as control. The reaction products were analyzed by SDS-PAGE Coomassie blue staining and by autoradiography.

Figure 4 Cellular distributions of LiSIR2RP1 and α -tubulin. The *L. infantum* promastigotes and axenic amastigotes (A) and their respective cytoskeletons (B) were double stained by immunofluorescence with mouse anti-LiSIR2RP1 antibodies and rabbit anti- α -tubulin antibodies (left panels) or with rabbit anti-TbRAB11 antibodies and mouse anti- α -tubulin antibodies followed by their respective Alexa Fluor 488 (green) or Alexa Fluor 568 (red) conjugated IgG and counterstained with DAPI (blue). The parasites were visualized under a 1000x magnification using a Zeiss Axio Imager Z1 microscope, and Z-series optical sections were collected using an AxioCam. Data stacks were deconvolved using either the Axiovision AxioVs40 V 4.2.0.0 (Carl Zeiss) or the Huygens Essential version 3.0.2p1. Western blot analysis of Triton X-100 extracted *L. infantum* promastigote (P) and amastigote (A) fractions. S designates the supernatant and I the cytoskeleton pellet. The fractions were separated in a 10% SDS-PAGE, transferred to a nitrocellulose membrane and probed with either anti- α -tubulin, anti-acetylated tubulin, anti-LiSIR2RP1 or anti-TbRAB11 antibodies (C).

Figure 5 Cellular distribution of LiSIR2RP1, α -tubulin and acetylated α -tubulin after parasite treatment with taxol or nocodazole.

The cellular distribution of the *LiSIR2RP1*, α -tubulin and acetylated α -tubulin were analyzed after 16 hours of *L. infantum* promastigote treatment with 1 μ M of taxol or 5 μ g/ml of nocodazole at 27°C. The parasites were fixed with paraformaldehyde, made permeable with triton X-100 and double labelled with mouse anti-*LiSIR2RP1* antibodies and rabbit anti- α -tubulin antibodies (A) or with mouse anti-acetylated α -tubulin antibodies (6-11B-1) and rabbit anti- α -tubulin antibodies (B) followed by their respective Alexa Fluor 488 (green) or Alexa Fluor 568 (red) conjugated IgG and counterstained with DAPI (blue). The parasites were visualized under a 1000x magnification using a Zeiss Axio Imager Z1 microscope, and Z-series optical sections were collected using an AxioCam. Data stacks were deconvolved using either the Axiovision AxioVs40 V 4.2.0.0 (Carl Zeiss) or the Huygens Essential version 3.0.2p1.

Figure 6 α -Tubulin is deacetylated by LiSIR2RP1. Purified tubulin dimers (A), taxol stabilized microtubules (B), taxol stabilized microtubules from whole promastigote lysate (C) and taxol stabilized microtubules from whole amastigote lysate (D) were incubated with several amounts of purified rLiSIR2RP1 in the presence or absence of 1mM of NAD⁺ and several nicotinamide concentrations for 3 hours at room temperature. The reaction products were analyzed by western blotting with specific antibodies to acetylated α -tubulin, α -tubulin and *LiSIR2RP1*. Analysis by Western blot of the acetylated α -tubulin and total α -tubulin levels in the wt and *LiSIR2*[±] mutant (clone N2) promastigote (P) and axenic amastigote (A) total extracts (E).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 6