Identification of 5-lipoxygenase and microsomal prostaglandin E synthase-1 as functional targets of the anti-inflammatory and anti-carcinogenic garcinol

Andreas Koeberle, Hinnak Northoff, Oliver Werz

► To cite this version:

Andreas Koeberle, Hinnak Northoff, Oliver Werz. Identification of 5-lipoxygenase and microsomal prostaglandin E synthase-1 as functional targets of the anti-inflammatory and anti-carcinogenic garcinol. *Biochemical Pharmacology*, 2009, 77 (9), pp.1513. 10.1016/j.bcp.2009.02.005 . hal-00493492

HAL Id: hal-00493492

<https://hal.science/hal-00493492>

Submitted on 19 Jun 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Identification of 5-lipoxygenase and microsomal prostaglandin E₂ synthase-1 as functional targets of the anti-inflammatory and anti-carcinogenic garcinol

Authors: Andreas Koeberle, Hinnak Northoff, Oliver Werz

PII: S0006-2952(09)00105-1
DOI: doi:10.1016/j.bcp.2009.02.005
Reference: BCP 10084

To appear in: *BCP*

Received date: 9-12-2008
Revised date: 27-1-2009
Accepted date: 11-2-2009

Please cite this article as: Koeberle A, Northoff H, Werz O, Identification of 5-lipoxygenase and microsomal prostaglandin E₂ synthase-1 as functional targets of the anti-inflammatory and anti-carcinogenic garcinol, *Biochemical Pharmacology* (2008), doi:10.1016/j.bcp.2009.02.005

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Identification of 5-lipoxygenase and microsomal prostaglandin E₂ synthase-1 as functional targets of the anti-inflammatory and anti-carcinogenic garcinol

Running title: Garcinol inhibits 5-lipoxygenase and mPGES-1

Andreas Koeberle^a, Hinnak Northoff^b, and Oliver Werz^{c*}

^a Department for Pharmaceutical Analytics, Pharmaceutical Institute, University of Tuebingen, Auf der Morgenstelle 8, D-72076 Tuebingen, Germany

^b Institute for Clinical and Experimental Transfusion Medicine, University Medical Center Tuebingen, Hoppe-Seyler-Straße 3, 72076 Tuebingen, Germany

* Corresponding author: Dr. Oliver Werz, Department for Pharmaceutical Analytics, Pharmaceutical Institute, University of Tuebingen, Auf der Morgenstelle 8, D-72076 Tuebingen, Germany. Phone: +4970712978793; Fax: +497071294565; e-mail: oliver.werz@uni-tuebingen.de

Abstract

Garcinol (camboginol) from the fruit rind of Guttiferae species shows anti-carcinogenic and anti-inflammatory properties, but the underlying molecular mechanisms are unclear. Here we show that garcinol potently interferes with 5-lipoxygenase (EC 7.13.11.34) and microsomal prostaglandin (PG)E₂ synthase (mPGES)-1 (EC 5.3.99.3), enzymes that play pivotal roles in inflammation and tumorigenesis. In cell-free assays, garcinol inhibited the activity of purified 5-lipoxygenase and blocked the mPGES-1-mediated conversion of PGH₂ to PGE₂ with IC₅₀ values of 0.1 and 0.3 μ M, respectively. Garcinol suppressed 5-lipoxygenase product formation also in intact human neutrophils and reduced PGE₂ formation in interleukin-1 β -stimulated A549 human lung carcinoma cells as well as in human whole blood stimulated by lipopolysaccharide. Moreover, garcinol interfered with isolated cyclooxygenase (COX)-1 (EC 1.14.99.1, IC₅₀ = 12 μ M) and with the formation of COX-1-derived 12(S)-hydroxy-5-cis-8,10-trans-heptadecatrienoic acid and thromboxane B₂ in human platelets. In contrast, neither Ca²⁺-ionophore (A23187)-induced arachidonic acid release in neutrophils nor COX-2 activity in A549 cells or whole blood, measured as formation of 6-keto PGF_{1 α} , or isolated human recombinant COX-2 were significantly affected by garcinol ($\leq$ 30 μ M). Together, the high potency of garcinol to selectively suppress PGE₂ synthesis and 5-lipoxygenase product formation provides a molecular basis for the anti-inflammatory and anti-carcinogenic effects of garcinol and rationalizes its therapeutic use.

Abbreviations used: AA, arachidonic acid; COX, cyclooxygenase; cPL, cytosolic phospholipase; ERK, extracellular signal-regulated kinase; FLAP, 5-lipoxygenase-activating protein; 12-HHT, 12(*S*)-hydroxy-5-cis-8,10-trans-heptadecatrienoic acid; iNOS, inducible nitric oxide synthase; 5-LO, 5-lipoxygenase; LT, leukotriene; mPGES, microsomal prostaglandin E₂ synthase; NSAIDs, non steroidal anti-inflammatory drugs; PG, prostaglandin.

1. Introduction

Garcinol (**Fig. 1A**) is a polyprenylated acylphloroglucinole contained in the rind of Guttiferae (*Garcinia indica*, *G. huillkensis* and *G. cambogia*), named Kokum, which is used as culinary spice and garnish for curry. In India, Kokum is used in ayurvedic medicine for the treatment of inflammatory and infectious diseases [1]. In rats, garcinol protects against 4-nitroquinoline-1-oxide-induced tongue carcinogenesis [2] and azoxymethane-induced colon carcinogenesis [3]. Functional investigations have revealed anti-oxidative [4], anti-proliferative [5], pro-apoptotic [6], anti-invasive [7], and anti-inflammatory properties of garcinol [6, 8, 9]. Garcinol reduced the expression or activation of kinases involved in cell proliferation (i.e., focal adhesion kinase, extracellular signal-regulated kinase (ERK)-1/2, p38 mitogen-activated protein kinase [8, 10]), blocked the expression of enzymes involved in the development and progression of cancer and inflammation (i.e., cyclooxygenase (COX)-2 (EC 1.14.99.1) and inducible nitric oxide synthase (iNOS)), and suppressed nuclear factor (NF)κB signalling [8, 10]). However, molecular targets of garcinol are less defined, and thus far, only histone acetyltransferase p300 ($IC_{50} = 7 \mu M$) [11], P300/CBP-associated factor ($IC_{50} = 5 \mu M$) [11], and acetylcholine esterase [12] ($IC_{50} = 0.7 \mu M$) were identified.

Certain COX-1/2-derived prostanoids and 5-lipoxygenase (5-LO)-derived leukotrienes (LTs) initiate and maintain inflammatory and allergic reactions (for review see [13]), and pharmacological intervention with their biosynthesis (i.e., utilizing COX- and 5-LO (EC 7.13.11.34) inhibitors) is applied in therapy of respective diseases. COX-2 and/or 5-LO are also overexpressed in various cancer cells (e.g., prostate, breast, lung, and colon) [14-16], and pre-clinical studies indicate tumor-preventive effects of COX inhibition by non steroidal anti-inflammatory drugs (NSAIDs) [17]. Moreover, pharmacological suppression of the 5-LO pathway, using 5-LO inhibitors or inhibitors of the 5-LO-activating protein (FLAP), causes

apoptosis of various tumor cells [16, 18]. Prostaglandin (PG)E₂ is the most prominent COX product in inflammation and tumorigenesis and is formed from PGH₂ by PGE₂ synthases. The microsomal PGE₂ synthase (mPGES)-1 (EC 5.3.99.3) is an inducible isoform functionally coupled to COX-2. Accumulating evidence suggests that mPGES-1 is responsible for massive PGE₂ formation associated with pathologies [19]. Therefore, mPGES-1 may represent an alternative drug target for treatment of inflammation and cancer, which is supported by data from mPGES-1 deficient mice and animal studies with selective mPGES-1 inhibitors [19-21].

Despite the anti-inflammatory and anti-tumorigenic potential of garcinol, its effects on the formation of eicosanoids were rarely analyzed. Hong et al. recently showed that garcinol modulates arachidonic acid (AA) metabolism in lipopolysaccharide-stimulated macrophages by blocking cytosolic phospholipase (cPL)A₂ (EC 3.1.1.4) activation due to inhibition of ERK1/2-mediated phosphorylation [10]. However, modulation of PG and/or LT biosynthesis by garcinol or effects on the catalytic activity of the key enzymes involved (e.g., COX-1/2, PG synthases, 5-LO) have not been addressed yet. Here, we demonstrate that garcinol inhibits prostanoid and LT formation in cellular and cell-free assays by preferential and potent interference with mPGES-1 and 5-LO.

2. Materials and Methods

2.1. Reagents

Garcinol (Biomol, Hamburg, Germany) was dissolved in dimethyl sulfoxide (DMSO) and kept in the dark at -20 °C, and freezing/thawing cycles were kept to a minimum. The thromboxane synthase inhibitor CV4151 [22] and anti-6-keto PGF_{1α} antibody were generous gifts by Dr. S. Laufer (University of Tuebingen, Germany) and Dr. T. Dingermaier (University of Frankfurt, Germany), respectively. Materials used: DMEM/High Glucose (4.5 g/l) medium, penicillin, streptomycin, trypsin/EDTA solution, PAA (Coelbe, Germany); PGH₂, Larodan (Malmö, Sweden); 11β-PGE₂, PGB₁, MK-886 (3-[1-(4-chlorobenzyl)-3-*t*-butyl-thio-5-isopropylindol-2-yl]-2,2-dimethylpropanoic acid), 6-keto PGF_{1α}, human recombinant COX-2, ovine COX-1, Cayman Chemical (Ann Arbor, MI); [5, 6, 8, 9, 11, 12, 14, 15-³H]arachidonic acid ([³H]AA), BioTrend Chemicals GmbH (Cologne, Germany); Ultima Gold^{TM} XR, Perkin Elmer (Boston, MA). All other chemicals were obtained from Sigma-Aldrich (Deisenhofen, Germany) unless stated otherwise.

2.2. Cells and cell viability assay

A549 cells were cultured in DMEM/High glucose (4.5 g/l) medium supplemented with heat-inactivated fetal calf serum (10%, v/v), penicillin (100 U/ml), and streptomycin (100 µg/ml) at 37 °C in a 5% CO₂ incubator. After 3 days, confluent cells were detached using 1 × trypsin/EDTA solution and reseeded at 2 × 10⁶ cells in 20 ml medium in 175 cm² flasks. Cell viability was assessed by trypan blue staining and light microscopy. A549 cells (4 × 10⁵) were plated in a 75 cm³ cell culture flask and incubated at 37 °C and 5% CO₂ for 72 h. Then, garcinol was added and the incubation was continued for 5 or 24 h before cell viability was

determined. Treatment with garcinol (10 or 30 μ M) for 5 h did not significantly reduced cell viability, excluding acute cytotoxic effects in the cellular assays used in this study. However, prolonged incubation (24 h) with 10 or 30 μ M garcinol led to a strong increase in the ratio of dead cells to total number of cells (50% and 71% dead cells, respectively).

Blood cells were freshly isolated from leukocyte concentrates obtained at the Blood Center of the University Hospital Tuebingen (Germany) as described [23]. In brief, venous blood was taken from healthy adult donors that did not take any medication for at least 7 days, and leukocyte concentrates were prepared by centrifugation (4000 $\times$ g, 20 min, 20 $^{\circ}$ C). Cells were immediately isolated by dextran sedimentation and centrifugation on Nycoprep cushions (PAA, Coelbe, Germany). Platelet-rich-plasma was obtained from the supernatants, mixed with phosphate-buffered saline (PBS) pH 5.9 (3:2 v/v), centrifuged (2100 $\times$ g, 15 min, room temperature), and the pelleted platelets were resuspended in PBS pH 5.9/0.9% NaCl (1:1, v/v). Washed platelets were finally resuspended in PBS pH 7.4 and 1 mM CaCl_2 . Neutrophils were immediately isolated from the pellet after centrifugation on Nycoprep cushions, and hypotonic lysis of erythrocytes was performed as described [24]. Cells were finally resuspended in PBS pH 7.4 (PBS) containing 1 mg/ml glucose and 1 mM CaCl_2 (PGC buffer) (purity > 96-97%).

2.3. Release of arachidonic acid in neutrophils

Freshly isolated human neutrophils (3×10^7 /ml PGC buffer) were pre-incubated with the test compounds for 10 min prior to stimulation with Ca^{2+} -ionophore A23187 (2.5 μ M). After 5 min at 37 $^{\circ}$ C, the reaction was stopped by addition of 1 ml of methanol. After adjusting pH 3 and addition of magarinic acid (11.1 nmol) as internal standard, released AA was separated by solid phase extraction (RP-18 material and elution with methanol) and coupled to 2,4-

dimethoxyaniline hydrochloride (0.14 mg, 0.75 μ mol) using *N*-Ethyl-*N'*-(3-dimethylaminopropyl)carbodiimide (1.46 mg, 9.4 μ mol) as coupling reagent for 1 h at 37 °C. Derivatized AA was analyzed by RP-HPLC (gradient: 71% methanol / 29% water to 100% methanol in 20 min, detection at 272 nm).

2.4. Determination of COX-1 product formation in washed platelets

Freshly isolated platelets (10^8 /ml PBS containing 1 mM CaCl_2) were pre-incubated with the indicated agents for 5 min at room temperature. After addition of 5 μ M AA and further incubation for 5 min at 37 °C, the COX-1-derived TxB_2 was quantified using a TxB_2 High Sensitivity EIA Kit (Assay Designs, Ann Arbor, MI). 12(*S*)-Hydroxy-5-*cis*-8,10-*trans*-heptadecatrienoic acid (12-HHT, non-enzymatically formed from PGH_2) was extracted and analyzed by HPLC as described [23].

2.5. Determination of PGE_2 and 6-keto $\text{PGF}_{1\alpha}$ formation in intact A549 cells

Cellular PGE_2 and 6-keto $\text{PGF}_{1\alpha}$ formation was determined according to [25]. A549 cells (2×10^6 cells) were incubated for 16 h at 37 °C and 5% CO_2 , the medium was changed, and the cells were stimulated with interleukin-1 β (1 ng/ml) for 48 h. After trypsination, cells were washed with PBS twice. For determination of PGE_2 , 4×10^6 cells per ml PBS containing CaCl_2 (1 mM) were pre-incubated with the indicated compounds at 37 °C for 10 min, and PGE_2 formation was started by addition of ionophore A23187 (2.5 μ M), AA (1 μ M), and [^3H]AA (18.4 kBq). The reaction was stopped after 15 min at 37 °C, and the samples were put on ice. After centrifugation (800 $\times$ g, 5 min, 4 °C), the supernatant was acidified (pH 3) by addition of citric acid (20 μ l, 2 M), and the internal standard 11 β - PGE_2 (2 nmol) was added. Radiolabeled PGE_2 was separated and analyzed by RP-18 solid phase extraction and HPLC

analysis. The amount of 11β -PGE₂ was quantified by integration of the area under the eluted peaks. For quantification of radiolabeled PGE₂, fractions (0.5 ml) were collected and mixed with Ultima Gold^{TM} XR (2 ml) for liquid scintillation counting in a LKB Wallac 1209 Rackbeta Liquid Scintillation Counter.

For determination of 6-keto PGF_{1 α} , 1×10^6 cells, resuspended in 1 ml PBS containing CaCl₂ (1 mM), were pre-incubated with the indicated compounds for 15 min at 37 °C, and 6-keto PGF_{1 α} formation was initiated by addition of AA (30 μ M). After 15 min at 37 °C, the reaction was stopped by cooling on ice. Cells were centrifuged (300 $\times$ g, 5 min, 4 °C), and the amount of released 6-keto PGF_{1 α} was assessed by ELISA as described [25].

2.6. Activity assays of isolated COX-1 and -2

Inhibition of the activities of isolated ovine COX-1 and human COX-2 was performed as described [25]. Briefly, purified COX-1 (ovine, 50 units) or COX-2 (human recombinant, 20 units) were diluted in 1 ml reaction mixture containing 100 mM Tris buffer pH 8, 5 mM glutathione, 5 μ M haemoglobin, and 100 μ M EDTA at 4 °C and pre-incubated with the test compounds for 5 min. Samples were pre-warmed for 60 sec at 37 °C, and AA (5 μ M for COX-1, 2 μ M for COX-2) was added to start the reaction. After 5 min at 37 °C, the COX-derived 12-HHT was extracted and then analyzed by HPLC as described [23].

2.7. Preparation of crude mPGES-1 in microsomes of A549 cells and determination of PGE₂ synthase activity

Preparation of A549 cells and determination of mPGES-1 activity was performed as described [25]. In brief, cells were incubated for 16 h at 37 °C and 5% CO₂, the medium was replaced,

interleukin-1 β (1 ng/ml) was added and cells were incubated for another 48 h. Cells were harvested, frozen in liquid nitrogen, and ice-cold homogenisation buffer (0.1 M potassium phosphate buffer pH 7.4, 1 mM phenylmethanesulphonyl fluoride, 60 μ g/ml soybean trypsin inhibitor, 1 μ g/ml leupeptin, 2.5 mM glutathione, and 250 mM sucrose) was added. After 15 min, cells were sonicated on ice (3 $\times$ 20 sec) and the homogenate was subjected to differential centrifugation at 10,000 $\times$ g for 10 min and 174,000 $\times$ g for 1 h at 4 $^{\circ}$ C. The pellet (microsomal fraction) was resuspended in 1 ml homogenization buffer and the total protein concentration was determined by Coomassie protein assay [26]. Microsomal membranes were diluted in potassium phosphate buffer (0.1 M, pH 7.4) containing 2.5 mM glutathione. Test compounds or vehicle were added and after 15 min at 4 $^{\circ}$ C, the reaction (100 μ l total volume) was initiated by addition of PGH₂ (20 μ M, final concentration). After 1 min at 4 $^{\circ}$ C, the reaction was terminated using stop solution (100 μ l; 40 mM FeCl₂, 80 mM citric acid, and 10 μ M of 11 β -PGE₂). PGE₂ was separated by solid phase extraction on reversed phase (RP)-C18 material using acetonitrile (200 μ l) as eluent and analyzed by RP-HPLC (30% acetonitrile / 70% water + 0.007% TFA (v/v)) with UV detection at 195 nm. 11 β -PGE₂ was used as internal standard to quantify PGE₂ product formation by integration of the area under the peaks.

2.8. Determination of PGE₂, 6-keto PGF_{1 α} , and 12-HHT formation in human whole blood

Peripheral blood from healthy adult volunteers, who had not received any medication for at least two weeks under informed consent, was obtained by venipuncture and collected in syringes containing heparin (20 U/ml). For determination of PGE₂ and 6-keto PGF_{1 α} , aliquots of whole blood (0.8 ml) were mixed with the thromboxane synthase inhibitor CV4151 (1 μ M) and with aspirin (50 μ M). A total volume of 1 ml was adjusted with sample buffer (10 mM

potassium phosphate buffer pH 7.4, 3 mM KCl, 140 mM NaCl, and 6 mM D-glucose). After pre-incubation with the indicated compounds for 5 min at room temperature, the samples were stimulated with lipopolysaccharide (10 μ g/ml) for 5 h at 37 °C. Prostanoid formation was stopped on ice, the samples were centrifuged (2300 $\times$ g, 10 min, 4 °C), and 6-keto PGF_{1 α} was quantified in the supernatant using a 6-keto PGF_{1 α} High Sensitivity EIA Kit (Assay Designs, Ann Arbor, MI) according to the manufacturer's protocol. PGE₂ was determined as described [25]. In brief, the supernatant was acidified with citric acid (30 μ l, 2 M), and after centrifugation (2300 $\times$ g, 10 min, 4 °C), solid phase extraction, and HPLC, analysis of PGE₂ was performed to isolate PGE₂. The PGE₂ peak (3 ml), identified by co-elution with the authentic standard, was collected, and acetonitrile was removed under a nitrogen stream. The pH was adjusted to 7.2 by addition of 10 $\times$ PBS buffer pH 7.2 (230 μ l) before PGE₂ contents were quantified using a PGE₂ High Sensitivity EIA Kit (Assay Designs, Ann Arbor, MI) according to the manufacturer's protocol.

For determination of 12-HHT, human whole blood (2 ml) was pre-incubated with the indicated compounds at 37 °C for 5 min, and formation of 12-HHT was initiated by addition of 30 μ M Ca²⁺-ionophore A23187 and 100 μ M AA. After 10 min at 37 °C, the reaction was stopped on ice, and the samples were centrifuged (600 $\times$ g/10 min/4 °C). Aliquots of the resulting plasma (500 μ l) were then mixed with 2 ml of methanol, and 200 ng of PGB₁ was added as internal standard. The samples were placed at -20 °C for 2 h and centrifuged again (600 $\times$ g/15 min/4 °C). The supernatants were collected and diluted with 2.5 ml PBS and 75 μ l HCl 1 N. Formed 12-HHT was extracted and analyzed by HPLC as described [23].

2.9. Expression of human recombinant 5-lipoxygenase in *E.coli*, preparation of homogenates and 100,000 $\times$ g supernatants, and semi-purification of 5-lipoxygenase

protein

E.coli MV1190 was transformed with pT3-5-LO plasmid, and recombinant 5-LO protein was expressed at 27 °C as described [27]. Cells were lysed in 50 mM triethanolamine/HCl pH 8.0, 5 mM EDTA, soybean trypsin inhibitor (60 µg/ml), 1 mM phenylmethanesulphonyl fluoride, and lysozyme (500 µg/ml), homogenized by sonication (3 × 15 sec), and centrifuged at 100,000×g for 70 min at 4 °C. The 100,000×g supernatant (S100) was applied to an ATP-agarose column to partially purify 5-LO as described previously [28]. S100 or semi-purified 5-LO were immediately used for activity assays.

2.10. Determination of 5-lipoxygenase product formation in cell-free systems

Aliquots of S100 or semi-purified 5-LO were diluted with ice-cold PBS containing 1 mM EDTA, and 1 mM ATP was added. Samples were pre-incubated with the test compounds as indicated. After 10 min at 4 °C, samples were pre-warmed for 30 sec at 37 °C, and 2 mM CaCl₂ plus 20 µM AA was added to start 5-LO product formation. The reaction was stopped after 10 min at 37 °C by addition of 1 ml ice-cold methanol, and the formed metabolites were analyzed by RP-HPLC as described [24]. 5-LO products include the all-trans isomers of LTB₄ and 5(*S*)-hydro(pero)xy-6-*trans*-8,11,14-*cis*-eicosatetraenoic acid.

2.11. Determination of 5-lipoxygenase product formation in intact cells

Freshly isolated neutrophils (5 × 10⁶/ml PGC buffer) were pre-incubated with the test compounds for 15 min at room temperature and 5-LO product formation was started by addition of 2.5 µM ionophore A23187 plus 20 µM AA. After 10 min at 37 °C, the reaction was stopped with 1 ml of methanol and 30 µl of 1 N HCl, and then, 200 ng PGB₁ and 500 µl PBS were added. Formed 5-LO metabolites were extracted and analyzed by HPLC as described [24]. 5-LO products include LTB₄ and its all-trans isomers, and 5(*S*)-

hydro(pero)xy-6-*trans*-8,11,14-*cis*-eicosatetraenoic acid. Cysteinyl-LTs C₄, D₄, and E₄ were not detected, and oxidation products of LTB₄ were not determined.

2.12. Statistics

Data are expressed as mean $\pm$ S.E. IC₅₀ values are approximations determined by graphical analysis (linear interpolation between the points between 50% activity). The program Graphpad InStat (Graphpad Software Inc., San Diego, CA) was used for statistical comparisons. Statistical evaluation of the data was performed by one-way ANOVAs for independent or correlated samples followed by Tukey HSD post-hoc tests. A P value of <0.05 (*) was considered significant.

3. Results

3.1. Effects of garcinol on arachidonic acid release from human neutrophils

Previously, it was suggested that garcinol interferes with eicosanoid formation at the level of gene expression (COX-2) [8], or ERK1/2 phosphorylation resulting in reduced AA release in lipopolysaccharide-stimulated RAW264.7 murine macrophages [10]. We investigated the effects of garcinol on cellular cPLA₂-mediated AA release in human neutrophils using the Ca²⁺-ionophore A23187 as stimulus that acts by substantial elevation of [Ca²⁺]_i, independent of cellular signalling pathways. Neutrophils, pre-incubated with garcinol (3 to 33 μM), were stimulated with Ca²⁺-ionophore (2.5 μM), and the released AA was derivatized with 2,4-dimethoxyaniline and analyzed by RP-HPLC. Garcinol failed to significantly suppress the release of AA up to 33 μM (**Fig. 1B**), whereas the cPLA₂α-inhibitor (control) potently inhibited AA liberation.

3.2. Effects of garcinol on cellular prostanoid formation

Next, we investigated the effects of garcinol on the transformation of AA to prostanoids in intact cells. Prostanoid formation was initiated by addition of exogenous AA in order to circumvent endogenous AA supply, and thus, to exclude inhibition of prostanoid formation at the level of PLA₂-mediated substrate release. For determination of COX-1 inhibition, human platelets were pre-incubated with garcinol (1 to 33 μM), and formation of COX-1-derived TxB₂ and 12-HHT (non-enzymatically formed from PGH₂ in platelets [29]) was initiated by AA (5 μM). Garcinol significantly suppressed TxB₂ (**Fig. 1C**) and 12-HHT (**Fig. 1D**) formation at 10 μM with an IC₅₀ of 16 and 11 μM, respectively. To investigate inhibition of cellular COX-2 activity, we assessed the formation of PGE₂ and the stable PGI₂ degradation

product 6-keto PGF_{1α} (as biomarker for PGI₂ formation) in A549 cells. For measuring 6-keto PGF_{1α}, higher concentrations of AA (30 instead of 1 μM) were required than for determination of PGE₂ to get detectable amounts of 6-keto PGF_{1α} (detection limit: 14 ng/ml at a signal to noise ratio of 3). A contribution of COX-1 can be excluded in interleukin-1β-stimulated A549 cells because COX-1 is neither detectable on the protein nor on the RNA level [30, 31]. COX-2-mediated synthesis of PGE₂ was significantly inhibited at 10 μM with an apparent IC₅₀ of approx. 10 μM (**Fig. 2A**), whereas the synthesis of 6-keto PGF_{1α} was not significantly reduced by garcinol up to 33 μM (**Fig. 2B**). PGE₂ formation is inhibited by garcinol within a narrow concentration range (3 to 10 μM), and higher concentrations of garcinol failed to completely block PGE₂ synthesis. Also MK-886 (33 μM), a mPGES-1 inhibitor, did not interfere with 6-keto PGF_{1α} synthesis and failed to entirely suppress PGE₂ formation [25]. In contrast, the COX-2 inhibitor celecoxib (5 μM) efficiently suppressed both formation of PGE₂ (**Fig. 2A**) and 6-keto PGF_{1α} (**Fig. 2B**), respectively.

3.3. Effects of garcinol on the activity of COX-1/2 in cell-free assays

We determined whether garcinol may directly interfere with the catalytic activity of isolated COX-1 and/or COX-2. The formation of 12-HHT, the major COX-1/2-derived product under the experimental conditions chosen [32], was initiated by addition of AA (5 μM for COX-1, 2 μM for COX-2). Garcinol concentration-dependently and significantly inhibited COX-1-mediated 12-HHT formation at 10 μM with an IC₅₀ = 12 μM, which is in good agreement with the efficacy in intact platelets (IC₅₀ = 11 μM, see above). In contrast, COX-2 activity was not significantly reduced up to 33 μM (**Fig. 3A**). Indomethacin (10 μM, for COX-1) and celecoxib (5 μM, for COX-2), used as reference compounds, potently inhibited 12-HHT formation, as expected (**Fig. 3B**).

3.4. Effects of garcinol on the activity of mPGES-1

Because garcinol failed to inhibit COX-2 in cell-free assays and did not suppress 6-keto PGF_{1α} synthesis in A549 cells but reduced PGE₂ formation, it appeared reasonable that garcinol might selectively suppress PGE₂ formation via inhibition of PGE₂ synthases downstream of COX-2. Microsomal preparations of interleukin-1β-stimulated A549 cells, used as source of mPGES-1 [33], were pre-incubated with garcinol or vehicle (DMSO) for 15 min, and then, PGE₂ formation was initiated by addition of 20 μM PGH₂. In agreement with the literature [25, 34], the mPGES-1 inhibitor MK-886 (used as control) inhibited PGE₂ formation with an IC₅₀ of 2.1 μM (data not shown). Garcinol was slightly superior over MK-886 and concentration-dependently inhibited PGE₂ formation with an IC₅₀ of 1.2 μM (**Fig. 4A**). Decreasing the PGH₂ concentration to 1 μM even enhanced the potency of garcinol to suppress mPGES-1 activity (IC₅₀ = 0.3 μM, **Fig. 4B**) suggesting a substrate competitive inhibitory mechanism.

To investigate whether garcinol inhibits PGE₂ synthesis in a reversible manner, microsomal preparations of A549 cells were pre-incubated with garcinol and MK-886, and wash-out experiments were carried out. MK-886 and garcinol failed to efficiently block PGE₂ synthesis at 0.3 μM, whereas PGE₂ formation was efficiently inhibited at 3 μM (**Fig. 4C**). However, 10-fold dilution of the samples containing 3 μM garcinol and MK-886 restored mPGES-1 activity (**Fig. 4C**) implying a reversible mode of inhibition.

3.5. Effects of garcinol on prostanoid formation in human whole blood

Many potent inhibitors of prostanoid biosynthesis in cell-free assays or in intact isolated cells lose their efficacy under physiologically relevant conditions such as in whole blood.

Therefore, we investigated whether garcinol could affect prostanoid biosynthesis also in a human whole blood assay. To assess inhibition of COX-1, freshly drawn human venous blood was pre-incubated with the test compounds or vehicle (DMSO) for 5 min and then treated with Ca^{2+} -ionophore plus AA (100 μM). Formation of 12-HHT, which mainly derives from constitutively expressed COX-1 under these experimental conditions, was not significantly affected by garcinol up to 33 μM (**Fig. 5A**). In control experiments, indomethacin (20 μM) almost completely blocked 12-HHT synthesis.

For determination of COX-2 inhibition, heparinized blood was pre-incubated with garcinol or vehicle (5 min) and stimulated with lipopolysaccharide for 5 h. Then, the levels of PGE_2 and 6-keto $\text{PGF}_{1\alpha}$ were determined in the corresponding plasma by ELISA. For measuring PGE_2 levels, PGE_2 was first separated from other AA metabolites by RP-HPLC [25]. In agreement with the results obtained with A549 cells, garcinol suppressed PGE_2 formation in whole blood in a concentration-dependent manner starting at 3 μM with an apparent IC_{50} value of 30 μM (**Fig. 5B**). Significant inhibition of PGE_2 formation was only reached at 30 μM for both garcinol and the mPGES-1 reference inhibitor MK-886, and also prolonged stimulation of whole blood with LPS up to 24 h did not increase the potency of garcinol. Neither garcinol nor MK-886 reduced 6-keto $\text{PGF}_{1\alpha}$ formation up to 30 μM under the same conditions, suggesting a selective interference of garcinol with PGE_2 synthesis (**Fig. 5C**). Note that garcinol as well as MK-886 were less efficient in inhibiting PGE_2 formation as compared to indomethacin or celecoxib.

3.6. Inhibition of 5-lipoxygenase product formation by garcinol

To determine the effects of garcinol on 5-LO product formation in intact cells, freshly isolated human neutrophils were pre-incubated with garcinol for 15 min and 5-LO product synthesis

1 was initiated by Ca^{2+} -ionophore A23187 (2.5 μM) plus exogenous AA (20 μM). 5-LO
2 product formation was concentration-dependently inhibited by garcinol with an $\text{IC}_{50} = 1.9 \mu\text{M}$
3
4 (**Fig. 6A**). Differences in the potency of garcinol to inhibit the formation of LTB_4 (formed via
5 LTA_4 hydrolase) and 5(*S*)-hydro(pero)xy-6-*trans*-8,11,14-*cis*-eicosatetraenoic acid ($\text{IC}_{50} = 1.5$
6 and 1.8 μM , respectively) were not observed, excluding an interference of garcinol with LTA_4
7 hydrolase at these concentrations. The efficiency of garcinol was somewhat improved in a
8 cell-free assay using 100,000 $\times$ g supernatant of homogenized neutrophils as source of enzyme
9 ($\text{IC}_{50} = 0.5 \mu\text{M}$, **Fig. 6B**) and was even higher for semi-purified recombinant 5-LO ($\text{IC}_{50} = 0.1$
10 μM , **Fig. 6C**). In contrast, the iron-ligand type 5-LO inhibitor BWA4C (1 μM , used as
11 control) was equally effective independent of the assay conditions (data not shown).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

4. Discussion

Animal studies have revealed an anti-carcinogenic potential of garcinol [2, 3], and *in vitro* studies have demonstrated potent growth-inhibitory and apoptotic [5, 6] as well as anti-oxidant [4] and anti-inflammatory effects [8, 10]. Interference with lipopolysaccharide-induced NF κ B signalling, lowering COX-2 and iNOS expression, may explain the long-term anti-inflammatory and anti-carcinogenic effects of garcinol [8], whereas the short term effects were ascribed to impaired cPLA₂ activation through inhibition of ERK-1/2 phosphorylation restricting the supply of AA for eicosanoid formation [10]. Although suppression of eicosanoid formation may be a crucial mode of action of garcinol, direct inhibitory effects on enzymes involved in eicosanoid biosynthesis have not been investigated yet.

Our data demonstrate that garcinol is a direct and potent inhibitor of purified 5-LO and cell-free mPGES-1 with IC₅₀ = 0.1 and 0.3 μ M, respectively, and also a direct inhibition of COX-1 by garcinol is evident. Inhibition of other molecular targets (i.e., histone acetyltransferase p300 [11], P300/CBP-associated factor [11], and acetylcholine esterase [12]) required higher concentrations of garcinol (IC₅₀ = 0.7 to 7 μ M). However, because of the different experimental settings applied for analysis of the interference of garcinol with these targets, direct comparison of garcinol's potencies is difficult. 5-LO and mPGES-1 are crucial enzymes in the biosynthesis of key mediators in inflammatory processes [16, 19], providing a rationale for the anti-inflammatory effects of garcinol. Moreover, 5-LO and mPGES-1 are overexpressed in many tumors, and PGE₂ as well as 5-LO products play important roles in tumorigenesis [16, 18, 19]. Together, we conclude that the interference of garcinol with 5-LO and mPGES-1 provides a molecular basis for its documented anti-carcinogenic and anti-inflammatory properties.

1 Inhibition of 5-LO and mPGES-1 is evident also in intact cell assays and in human whole
2 blood. On the other hand, inhibition of COX-1-derived 12-HHT formation in human whole
3 blood was minute and not significant up to 33 μ M, although COX-1 activity is obviously
4 inhibited in intact platelets and in the cell-free assay (IC_{50} = 11 and 12 μ M, respectively). This
5 loss of potency in whole blood might be related to albumin-binding of garcinol but also to
6 competition with high AA concentrations (that are required in whole blood for analysis of 12-
7 HHT formation) for binding to the active site of COX-1.
8
9
10
11
12
13
14
15
16
17
18

19 The experimental settings (stimuli used, availability of respective substrates, and products
20 determined) of these cell-based assays allowed us to attribute the effects of garcinol to the
21 interference with the putative targets and to exclude other points of attack. In particular,
22 garcinol may simply block the supply of AA by interference with cPLA₂ activity, an enzyme,
23 whose activation and cellular functionality has been previously proposed to be affected by
24 garcinol [10]. Hong et al. showed that in lipopolysaccharide-stimulated RAW264.7 murine
25 macrophages garcinol blocked AA release by impairing the activation of cPLA₂. This was
26 ascribed to the interference with defined signalling steps (i.e., ERK signalling,
27 phosphorylation of cPLA₂) [10]. To circumvent such signalling pathways for cPLA₂
28 activation, we challenged human neutrophils with the Ca²⁺-ionophore A23187, where the
29 massive increase in intracellular Ca²⁺ essentially stimulates cPLA₂. Garcinol failed to block
30 AA liberation under these experimental conditions, indicating that PLA₂-mediated AA release
31 is not generally inhibited by garcinol. Note that in our cell-based assays analyzing AA
32 transformations by COX-1/2 or by 5-LO, AA was exogenously supplemented to avoid effects
33 on eicosanoid formation due to inhibition of PLA₂. On the other hand, the effects of garcinol
34 on PGE₂ and 6-keto PGF_{1 α} formation in whole blood were analyzed under conditions
35 (lipopolysaccharide challenge) where AA was supplied by endogenous sources. In contrast to
36 PGE₂ formation, the synthesis of 6-keto PGF_{1 α} was unaffected by garcinol, excluding general
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 suppression of AA and suggesting PGE₂ synthase (i.e., mPGES-1) as molecular target of
2 garcinol. However, our data cannot exclude that garcinol also inhibits other PGE₂ synthases
3
4 (such as cytosolic PGE₂ synthase or mPGES-2) in addition to mPGES-1.
5
6
7
8

9 PGE₂ and 6-keto PGF_{1α} are synthesized via the COX-2 pathway in lipopolysaccharide-
10 stimulated whole blood [35]. Because 6-keto PGF_{1α} formation in whole blood was unaffected
11
12 by garcinol, our data imply that garcinol-mediated suppression of PGE₂ synthesis in the same
13
14 assay is independent of an interference with COX-2. Also, garcinol failed to inhibit isolated
15
16 COX-2. Co-expression studies have revealed a preferred functional coupling of COX-2 and
17
18 mPGES-1 [36] suggesting that mPGES-1 is the major contributing PGE₂ synthase for PGE₂
19
20 synthesis in lipopolysaccharide-stimulated blood. Nevertheless, in analogy to other mPGES-1
21
22 inhibitors [25], garcinol failed to completely suppress PGE₂ synthesis in whole blood and the
23
24 remaining PGE₂ synthesis of approx. 40% might be attributable to other PGE₂ synthases (i.e.,
25
26 cytosolic PGE₂ synthase or mPGES-2) [25].
27
28
29
30
31
32
33
34
35

36 For inhibition of purified 5-LO (IC₅₀ = 0.1 μM) and cellular 5-LO product formation (IC₅₀ =
37
38 1.9 μM), a discrepancy in the potencies of garcinol was obvious. Interestingly, a similar
39
40 pattern was found for the structurally related hyperforin [23] that possesses an unique
41
42 inhibitory mode of action by interfering with the membrane-binding region of 5-LO's C2-like
43
44 domain (Feißt et al., 2009, manuscript submitted). However, garcinol is an effective
45
46 antioxidant [4] and may act as redox-type inhibitor of 5-LO reducing the active site iron. Such
47
48 reduction might be more vigorous for cell-free 5-LO as compared to intracellular enzyme, a
49
50 common phenomenon observed for many redox-active 5-LO inhibitors [37].
51
52
53
54
55
56
57

58 Despite the obvious inhibition of PGE₂ and LT formation by garcinol *in vitro*, the question
59
60 whether the interference of garcinol with mPGES-1 and 5-LO accounts for its anti-
61
62
63
64
65

inflammatory and anti-tumorigenic properties *in vivo*, can not be clearly answered yet. Unpublished studies, mentioned by Hong et al. [10], showed that garcinol inhibited phorbol myristate acetate-induced mouse ear edema accompanied by decreased PGE₂ and LTB₄ production. Of interest, in these experiments the peak plasma levels of garcinol (12 µM) obtained after oral application of garcinol were in a concentration range, where cellular inhibition 5-LO and mPGES-1 is evident in our study. Therefore, one may speculate that garcinol, after oral administration, could achieve plasma levels sufficient to markedly suppress PGE₂ and LT biosynthesis under inflammatory conditions. PGE₂ is one of the most-established key players in the initiation and progression of inflammation [38], and NSAIDs exert their beneficial effects *in vivo* essentially by repressing PGE₂ formation [13]. Animals deficient in COX-2 or mPGES-1 clearly show reduced inflammatory symptoms [19, 39]. Moreover, LTB₄ is a potent chemotactic and chemokinetic agent that recruits pro-inflammatory cells towards sites of inflammation [40]. But also tumorigenesis, particularly of colon, breast, prostate, and lung carcinoma has been associated with excessive PGE₂ and LT formation [16,18,19]. COX-2 selective inhibitors have shown a chemopreventive potential [41], and mPGES-1 is overexpressed in various cancers such as non-small cell lung cancer, invasive breast cancer, colorectal cancer, and gastric cancer [19]. Also 5-LO and its helper-protein FLAP are overexpressed in cancer cells, and 5-LO products promote cell proliferation and/or inhibit apoptosis, whereas 5-LO or FLAP inhibitors caused cell death [18].

In conclusion, we have identified human mPGES-1 and 5-LO as molecular targets of garcinol, and we showed that garcinol potently interferes with the catalytic activity of these enzymes in cell-free and cell-based assays. Inhibition of mPGES-1 and 5-LO may contribute to the anti-carcinogenic and anti-inflammatory effectiveness of garcinol observed in previous studies. Our data indicate that garcinol exerts a promising pharmacological profile combining potent inhibition of PGE₂ and LT formation without marked inhibition of COX isoenzymes.

References

- [1] Krishnamurthy N, Lewis YS, Ravindranath B. On the structures of garcinol, isogarcinol and camboginol. *Tetrahedron Letters* 1981;22:793-6.
- [2] Yoshida K, Tanaka T, Hirose Y, Yamaguchi F, Kohno H, Toida M, et al. Dietary garcinol inhibits 4-nitroquinoline 1-oxide-induced tongue carcinogenesis in rats. *Cancer Lett* 2005;221:29-39.
- [3] Tanaka T, Kohno H, Shimada R, Kagami S, Yamaguchi F, Kataoka S, et al. Prevention of colonic aberrant crypt foci by dietary feeding of garcinol in male F344 rats. *Carcinogenesis* 2000;21:1183-9.
- [4] Yamaguchi F, Ariga T, Yoshimura Y, Nakazawa H. Antioxidative and anti-glycation activity of garcinol from *Garcinia indica* fruit rind. *J Agric Food Chem* 2000;48:180-5.
- [5] Hong J, Kwon SJ, Sang S, Ju J, Zhou JN, Ho CT, et al. Effects of garcinol and its derivatives on intestinal cell growth: Inhibitory effects and autoxidation-dependent growth-stimulatory effects. *Free Radic Biol Med* 2007;42:1211-21.
- [6] Pan MH, Chang WL, Lin-Shiau SY, Ho CT, Lin JK. Induction of apoptosis by garcinol and curcumin through cytochrome c release and activation of caspases in human leukemia HL-60 cells. *J Agric Food Chem* 2001;49:1464-74.
- [7] Liao CH, Sang S, Ho CT, Lin JK. Garcinol modulates tyrosine phosphorylation of FAK and subsequently induces apoptosis through down-regulation of Src, ERK, and Akt survival signaling in human colon cancer cells. *J Cell Biochem* 2005;96:155-69.
- [8] Liao CH, Sang S, Liang YC, Ho CT, Lin JK. Suppression of inducible nitric oxide synthase and cyclooxygenase-2 in downregulating nuclear factor-kappa B pathway by Garcinol. *Mol Carcinog* 2004;41:140-9.

- [9] Matsumoto K, Akao Y, Kobayashi E, Ito T, Ohguchi K, Tanaka T, et al. Cytotoxic benzophenone derivatives from *Garcinia* species display a strong apoptosis-inducing effect against human leukemia cell lines. *Biol Pharm Bull* 2003;26:569-71.
- [10] Hong J, Sang S, Park HJ, Kwon SJ, Suh N, Huang MT, et al. Modulation of arachidonic acid metabolism and nitric oxide synthesis by garcinol and its derivatives. *Carcinogenesis* 2006;27:278-86.
- [11] Balasubramanyam K, Altaf M, Varier RA, Swaminathan V, Ravindran A, Sadhale PP, et al. Polyisoprenylated benzophenone, garcinol, a natural histone acetyltransferase inhibitor, represses chromatin transcription and alters global gene expression. *J Biol Chem* 2004;279:33716-26.
- [12] Lenta BN, Vonthron-Senecheau C, Weniger B, Devkota KP, Ngoupayo J, Kaiser M, et al. Leishmanicidal and cholinesterase inhibiting activities of phenolic compounds from *Allanblackia monticola* and *Symphonia globulifera*. *Molecules* 2007;12:1548-57.
- [13] Funk CD. Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science* 2001;294:1871-5.
- [14] Zha S, Yegnasubramanian V, Nelson WG, Isaacs WB, De Marzo AM. Cyclooxygenases in cancer: progress and perspective. *Cancer Lett* 2004;215:1-20.
- [15] Gupta S, Srivastava M, Ahmad N, Sakamoto K, Bostwick DG, Mukhtar H. Lipoxygenase-5 is overexpressed in prostate adenocarcinoma. *Cancer* 2001;91:737-43.
- [16] Werz O, Steinhilber D. Therapeutic options for 5-lipoxygenase inhibitors. *Pharmacol Ther* 2006;112:701-18.
- [17] Gasparini G, Longo R, Sarmiento R, Morabito A. Inhibitors of cyclo-oxygenase 2: a new class of anticancer agents? *Lancet Oncol* 2003;4:605-15.

- [18] Chen X, Sood S, Yang CS, Li N, Sun Z. Five-lipoxygenase pathway of arachidonic acid metabolism in carcino-genesis and cancer chemoprevention. *Curr Cancer Drug Targets* 2006;6:613-22.
- [19] Samuelsson B, Morgenstern R, Jakobsson PJ. Membrane prostaglandin E synthase-1: a novel therapeutic target. *Pharmacol Rev* 2007;59:207-24.
- [20] Cote B, Boulet L, Brideau C, Claveau D, Ethier D, Frenette R, et al. Substituted phenanthrene imidazoles as potent, selective, and orally active mPGES-1 inhibitors. *Bioorg Med Chem Lett* 2007;17:6816-20.
- [21] Xu D, Rowland SE, Clark P, Giroux A, Cote B, Guiral S, et al. MF63 {2-(6-chloro-1H-phenanthro[9,10-d]imidazol-2-yl)isophthalonitrile}, a selective microsomal prostaglandin E synthase 1 inhibitor, relieves pyresis and pain in preclinical models of inflammation. *J Pharmacol Exp Ther* 2008;326:754-63.
- [22] Kato K, Ohkawa S, Terao S, Terashita Z, Nishikawa K. Thromboxane synthetase inhibitors (TXSI). Design, synthesis, and evaluation of a novel series of omega-pyridylalkenoic acids. *J Med Chem* 1985;28:287-94.
- [23] Albert D, Zundorf I, Dingermann T, Muller WE, Steinhilber D, Werz O. Hyperforin is a dual inhibitor of cyclooxygenase-1 and 5-lipoxygenase. *Biochem Pharmacol* 2002;64:1767-75.
- [24] Werz O, Burkert E, Samuelsson B, Rådmark O, Steinhilber D. Activation of 5-lipoxygenase by cell stress is calcium independent in human polymorphonuclear leukocytes. *Blood* 2002;99:1044-52.
- [25] Koeberle A, Siemoneit U, Buehring U, Northoff H, Laufer S, Albrecht W, et al. Licofelone suppresses prostaglandin E2 formation by interference with the inducible microsomal prostaglandin E2 synthase-1. *J Pharmacol Exp Ther* 2008;326:975-82.

- [26] Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248-54.
- [27] Fischer L, Szellas D, Rådmark O, Steinhilber D, Werz O. Phosphorylation- and stimulus-dependent inhibition of cellular 5-lipoxygenase activity by nonredox-type inhibitors. *FASEB J* 2003;17:949-51.
- [28] Burkert E, Arnold C, Hammarberg T, Rådmark O, Steinhilber D, Werz O. The C2-like beta-barrel domain mediates the Ca²⁺-dependent resistance of 5-lipoxygenase activity against inhibition by glutathione peroxidase-1. *J Biol Chem* 2003;278:42846-53.
- [29] Hamberg M, Samuelsson B. Prostaglandin endoperoxides. Novel transformations of arachidonic acid in human platelets. *Proc Natl Acad Sci U S A* 1974;71:3400-4.
- [30] Asano K, Lilly CM, Drazen JM. Prostaglandin G/H synthase-2 is the constitutive and dominant isoform in cultured human lung epithelial cells. *Am J Physiol* 1996;271:L126-31.
- [31] Thoren S, Jakobsson PJ. Coordinate up- and down-regulation of glutathione-dependent prostaglandin E synthase and cyclooxygenase-2 in A549 cells. Inhibition by NS-398 and leukotriene C4. *Eur J Biochem* 2000;267:6428-34.
- [32] Capdevila JH, Morrow JD, Belosludtsev YY, Beauchamp DR, DuBois RN, Falck JR. The catalytic outcomes of the constitutive and the mitogen inducible isoforms of prostaglandin H2 synthase are markedly affected by glutathione and glutathione peroxidase(s). *Biochemistry* 1995;34:3325-37.
- [33] Jakobsson PJ, Thoren S, Morgenstern R, Samuelsson B. Identification of human prostaglandin E synthase: a microsomal, glutathione-dependent, inducible enzyme, constituting a potential novel drug target. *Proc Natl Acad Sci U S A* 1999;96:7220-5.
- [34] Claveau D, Sirinyan M, Guay J, Gordon R, Chan CC, Bureau Y, et al. Microsomal prostaglandin E synthase-1 is a major terminal synthase that is selectively up-

- regulated during cyclooxygenase-2-dependent prostaglandin E2 production in the rat adjuvant-induced arthritis model. *J Immunol* 2003;170:4738-44.
- [35] Patrignani P, Panara MR, Greco A, Fusco O, Natoli C, Iacobelli S, et al. Biochemical and pharmacological characterization of the cyclooxygenase activity of human blood prostaglandin endoperoxide synthases. *J Pharmacol Exp Ther* 1994;271:1705-12.
- [36] Murakami M, Naraba H, Tanioka T, Semmyo N, Nakatani Y, Kojima F, et al. Regulation of prostaglandin E2 biosynthesis by inducible membrane-associated prostaglandin E2 synthase that acts in concert with cyclooxygenase-2. *J Biol Chem* 2000;275:32783-92.
- [37] Werz O. Inhibition of 5-lipoxygenase product synthesis by natural compounds of plant origin. *Planta Med* 2007;73:1331-57.
- [38] Smith WL. The eicosanoids and their biochemical mechanisms of action. *Biochem J* 1989;259:315-24.
- [39] Langenbach R, Loftin CD, Lee C, Tiano H. Cyclooxygenase-deficient mice. A summary of their characteristics and susceptibilities to inflammation and carcinogenesis. *Ann N Y Acad Sci* 1999;889:52-61.
- [40] Claesson HE, Dahlen SE. Asthma and leukotrienes: antileukotrienes as novel anti-asthmatic drugs. *J Intern Med* 1999;245:205-27.
- [41] Rao CV, Reddy BS. NSAIDs and chemoprevention. *Curr Cancer Drug Targets* 2004;4:29-42.

Acknowledgments

The authors thank Bianca Jazzar and Gertrud Kleefeld for expert technical assistance. The financial support by the Deutsche Forschungsgemeinschaft is acknowledged.

Conflict of Interest Statement: None declared.

Figure Legends

Fig. 1. Effects of garcinol on cellular arachidonic acid release and prostanoid formation.

(A) Chemical structure of garcinol. (B) AA release from neutrophils. Neutrophils ($3 \times 10^7/\text{ml}$ PGC buffer) were pre-incubated with the indicated concentrations of garcinol or vehicle (DMSO, w/o) for 10 min. Then, AA release was initiated by Ca^{2+} -ionophore ($2.5 \mu\text{M}$). After 5 min at 37°C , AA was coupled to 2,4-dimethoxyaniline and analyzed by HPLC as described in the experimental section. In the absence of test compounds (100%, control), 180 ng/ml AA were released. cPLA $_2\alpha$ -inhibitor ($10 \mu\text{M}$) was used as control. (C) TxB $_2$ and (D) 12-HHT formation in intact platelets. Platelets ($10^8/\text{ml}$ PBS containing 1 mM CaCl_2) were pre-incubated with the indicated concentrations of garcinol or vehicle (DMSO, w/o) for 5 min prior to stimulation with AA ($5 \mu\text{M}$). After another 5 min at 37°C , the formation of TxB $_2$ was assessed by ELISA. TxB $_2$ formed in the absence of test compounds (100%, control) was 360 ng/ml. The formation of 12-HHT was determined by RP-HPLC as described. 12-HHT formed in the absence of test compounds (100%, control) was 290 ng/ml. Indomethacin (Indo, $10 \mu\text{M}$) was used as control. Data are given as mean $\pm$ S.E., $n = 3$, $**p < 0.01$ or $***p < 0.001$ vs. vehicle (0.1% DMSO) control, ANOVA + Tukey HSD *post-hoc* tests.

Fig. 2. Effects of garcinol on cellular arachidonic acid release and prostanoid formation.

(A) PGE $_2$ formation in A549 cells. Interleukin-1 β -stimulated A549 cells ($4 \times 10^6/\text{ml}$) were pre-incubated with garcinol or vehicle (DMSO, w/o) for 10 min, and then, $2.5 \mu\text{M}$ A23187 plus $1 \mu\text{M}$ AA and [^3H]AA (18.4 kBq) were added. After 15 min at 37°C , formed [^3H]PGE $_2$ was analyzed by RP-HPLC and liquid scintillation counting as described. The 100% value corresponds to 30 ng/ml PGE $_2$. MK-886 ($33 \mu\text{M}$) and celecoxib (Cele, $5 \mu\text{M}$) were used as

controls. **(B)** 6-Keto PGF_{1α} formation in A549 cells. Interleukin-1β-stimulated A549 cells (1×10^6 /ml) were pre-incubated with the indicated concentrations of garcinol or vehicle (DMSO) for 15 min prior to addition of 30 μM AA. After 15 min at 37 °C the amount of 6-keto PGF_{1α} was assessed by ELISA as described. The 100% value corresponds to 70 ng/ml 6-keto PGF_{1α}. Indomethacin (Indo, 10 μM) and celecoxib (Cele, 5 μM) were used as control. Data are given as mean $\pm$ S.E., $n = 3$, $*p < 0.05$, $**p < 0.01$ or $***p < 0.001$ vs. vehicle (0.1% DMSO) control, ANOVA + Tukey HSD *post-hoc* tests.

Fig. 3. Effects of garcinol on the activity of isolated COX-1 and -2.

(A) Purified ovine COX-1 (50 units) or **(B)** human recombinant COX-2 (20 units) were added to a COX reaction mix, containing 5 mM glutathione. After pre-incubation with the test compounds or vehicle (DMSO, w/o) for 5 min, the reaction was started with 5 μM (COX-1) or 2 μM (COX-2) AA. After 5 min at 37 °C, the formation of 12-HHT was determined by RP-HPLC as described. The 100% values correspond to 80 (COX-1) or 70 (COX-2) ng/ml 12-HHT. Indomethacin (Indo, 10 μM) and celecoxib (Cele, 5 μM) were used as controls. Data are given as mean $\pm$ S.E., $n = 3$, $**p < 0.01$, $***p < 0.001$ vs. vehicle (0.1% DMSO) control, ANOVA + Tukey HSD *post-hoc* tests.

Fig. 4. Effects of garcinol on the activity of mPGES-1.

(A) Concentration-response curves for garcinol. Microsomal preparations of interleukin-1β-stimulated A549 cells were pre-incubated with vehicle (DMSO) or the test compounds at the indicated concentrations for 15 min at 4 °C, and the reaction was started with 20 μM PGH₂. After 1 min at 4 °C, the reaction was terminated using a stop solution containing FeCl₂ and 11β-PGE₂ (1 nmol) as internal standard. The 100% value corresponds to 0.8 nmol PGE₂. **(B)**

The potency of garcinol for mPGES-1 inhibition was compared at 1 and 20 μM PGH_2 as substrate. The amount of PGE_2 was quantified for 1 μM PGH_2 by use of a PGE_2 High Sensitivity EIA Kit according to the manufacturer's protocol. The 100% value corresponds to 34 pmol PGE_2 . (C) Reversibility of mPGES-1 inhibition by garcinol and MK-886. Microsomal preparations of interleukin-1 β -stimulated A549 cells were pre-incubated with 3 μM inhibitor for 15 min at 4 $^\circ\text{C}$. An aliquot was diluted 10-fold to obtain an inhibitor concentration of 0.3 μM . For comparison, microsomal preparations were pre-incubated for 15 min with vehicle (DMSO, w/o) or 0.3 μM MK-886 or garcinol, and then, 20 μM PGH_2 was added (no dilution). All samples were incubated for 1 min on ice, and PGE_2 formation was analyzed as described by RP-HPLC. Data are given as mean + S.E., $n = 2-4$, $*p < 0.05$ or $***p < 0.001$ vs. vehicle (0.1% DMSO) control, ANOVA + Tukey HSD *post-hoc* tests.

Fig. 5. Effects of garcinol on prostanoid formation in human whole blood.

(A) 12-HHT formation. Heparinized whole blood was pre-incubated with garcinol or vehicle (DMSO, w/o) at the indicated concentrations for 5 min, and AA (100 μM) and Ca^{2+} -ionophore (30 μM) were added to induce COX-product formation. After 10 min at 37 $^\circ\text{C}$, 12-HHT was extracted and analyzed by RP-HPLC as described. The 100% value corresponds to 1.0 $\mu\text{g/ml}$ 12-HHT. Indomethacin (Indo, 20 μM) was used as control. (B, C) Heparinized human whole blood, treated with 1 μM thromboxane synthase inhibitor and 50 μM aspirin, was pre-incubated with the indicated concentrations of garcinol or vehicle (DMSO, w/o) for 5 min at room temperature, and then, prostanoid formation was induced by addition of 10 $\mu\text{g/ml}$ lipopolysaccharide. After 5 h at 37 $^\circ\text{C}$, (B) PGE_2 was extracted from plasma by RP-18 solid phase extraction, separated by RP-HPLC, and quantified by ELISA as described. The 100% value corresponds to 9 ng/ml PGE_2 . (C) 6-keto $\text{PGF}_{1\alpha}$ was directly determined in blood plasma by ELISA. The 100% value corresponds to 4 ng/ml 6-keto $\text{PGF}_{1\alpha}$. MK-886 (30 μM),

indomethacin (Indo, 50 μ M), celecoxib (Cele, 20 μ M), or vehicle (DMSO) were used as controls. Data are given as mean $\pm$ S.E., $n = 3-4$, $**p < 0.01$ or $***p < 0.001$ vs. vehicle (0.1% DMSO) control, ANOVA + Tukey HSD *post-hoc* tests.

Fig. 6. Inhibition of 5-lipoxygenase activity by garcinol.

(A) Neutrophils (5×10^6 cells/ml) were pre-incubated with the indicated concentrations of garcinol or vehicle (DMSO) for 15 min. 5-LO product formation was initiated by addition of 2.5 μ M Ca^{2+} -ionophore plus 20 μ M AA. (B) S100 of *E.coli* lysates expressing human recombinant 5-LO or (C) semi-purified human recombinant 5-LO were supplemented with garcinol and 1 mM ATP, and product formation was started by addition of 2 mM CaCl_2 and 20 μ M AA. The 100% values correspond to 1.8, 1.3, and 1.4 μ g/ml 5-LO products for intact cells, S100, and purified enzyme, respectively. Data are given as mean $\pm$ S.E., $n = 3-6$, $*p < 0.05$, $**p < 0.01$ or $***p < 0.001$ vs. vehicle (0.1% DMSO) control, ANOVA + Tukey HSD *post-hoc* tests.

Fig. 1

A

B

C

D

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

graphical abstract