An extracellular transglutaminase is required for apple pollen tube growth

Alessia Di Sandro, Stefano del Duca, Elisabetta Am Verderio, Alan Jeffrey Hargreaves, Alessandra Scarpellini, Giampiero Cai, Mauro Cresti, Claudia Faleri, Rosa Anna Iorio, Shigehisa Hirose, et al.

► To cite this version:

Alessia Di Sandro, Stefano del Duca, Elisabetta Am Verderio, Alan Jeffrey Hargreaves, Alessandra Scarpellini, et al.. An extracellular transglutaminase is required for apple pollen tube growth. Biochemical Journal, 2010, 429 (2), pp.261-271. 10.1042/BJ20100291 . hal-00495492

HAL Id: hal-00495492

<https://hal.science/hal-00495492>

Submitted on 28 Jun 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2

1
2
3
4
5
6
7
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

An extracellular transglutaminase is required for apple pollen tube growth

Alessia Di Sandro¹, Stefano Del Duca¹, Elisabetta Verderio², Alan J. Hargreaves², Alessandra Scarpellini^{1,2}, Giampiero Cai³, Mauro Cresti³, Claudia Faleri³, Rosa Anna Iorio¹, Shigehisa Hirose⁴, Yutaka Furutani⁴, Ian G.C. Coutts², Martin Griffin⁵, Philip L.R. Bonner², Donatella Serafini-Fracassini¹

¹Dipartimento di Biologia Evoluzionistica Sperimentale, Università degli Studi di Bologna, Via Irnerio 42, 40126, Bologna, Italy

²School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK

³Dipartimento di Scienze Ambientali, Università di Siena, 53100 Siena, Italy

⁴Department of Biological Sciences, Tokyo Institute of Technology, Yokohama, Japan

⁵School of Life and Health Sciences, Aston University, Birmingham B47ET, UK

Correspondence should be addressed to Donatella Serafini-Fracassini, Dipartimento di Biologia Evoluzionistica Sperimentale, Sede Botanica, Università di Bologna Via Irnerio 42, 40126 Bologna, Italy.

Phone: +39-0512091292; Fax: +39-051242576. E-mail: donatella.serafini@unibo.it or to Philip L.R. Bonner School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK.

Phone: +44-1158483528; Fax: +44-1158483384. E-mail: philip.bonner@ntu.ac.uk

Running title: *extracellular pollen transglutaminase*

Abbreviations used: BSA, bovine serum albumin; DMC, N',N'-dimethyl casein; DTT, dithiothreitol; ECM, extra-cellular matrix; FA, focal adhesion; FITC, fluorescein isothiocyanate; GFP, green fluorescent protein; gpl TGase, guinea pig liver transglutaminase; HIC, hydrophobic interaction chromatography; PA, polyamine; PBS, phosphate-buffered saline; RFU, relative fluorescence units; SD, spermidine; SDS-PAGE, sodium dodecyl sulfate polyacrylamide gel electrophoresis; TGase, transglutaminase; tTGase, tissue transglutaminase; TRITC, tetramethyl rhodamine isothiocyanate.

5
651 **SYNOPSIS**

52
53 An extracellular form of the Ca^{2+} -dependent protein-cross-linking enzyme
54 transglutaminase (TGase) was demonstrated to be involved in the apical growth of
55 *Malus domestica* pollen tube. Apple pollen transglutaminase and its substrates were co-
56 localized within aggregates on the pollen tube surface, as determined by indirect
57 immuno-fluorescence staining and the *in situ* cross-linking of fluorescently labeled
58 substrates. Transglutaminase-specific inhibitors and an anti-TGase monoclonal
59 antibody blocked pollen tube growth, whereas incorporation of a recombinant
60 fluorescent mammalian TGase substrate (histidine-tagged green fluorescent protein:
61 His₆-Xpr-GFP) into the growing tube wall enhanced tube length and germination,
62 consistent with a role of TGase as modulator of cell wall building and strengthening.
63 The secreted pollen TGase catalyzed the cross-linking of both polyamines (PAs) into
64 proteins (released by the pollen tube) and His₆-Xpr-GFP into endogenous or
65 exogenously added substrates. A similar distribution of TGase activity was observed *in*
66 *planta* on pollen tubes germinating inside the style, consistent with a possible additional
67 role for TGase in the interaction between the pollen tube and the style during
68 fertilization.

69
70
71 **Key words:** extracellular localization, *Malus domestica*, pollen tube growth,
72 polyamines, protein cross-link, transglutaminase.
73

7
8

2

9

10

74 INTRODUCTION

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

In angiosperms, sexual reproduction requires the apical growth of pollen tubes through the style, towards the female gametes located into the gynoecium. In the apical zone of the pollen tube, the growth of plasma membrane and cell wall is maintained by the continuous supply of precursors that are provided by the cytoplasm. The dynamics of vesicle and organelle movement along the pollen tube is sustained by the cytoskeletal network controlled by the influx of calcium at the apical tip [1]. The cell wall strength and focal anchorage are essential requisites to push the extending pollen tip through the style, especially for solid stylar transmitting tracts, as in *Rosaceae* [2]. A number of yet to be completely characterized pollen proteins, either newly synthesised or located in the pollen wall intine, are released into the style [3]. During pollen growth, some of these pollen proteins interact with stylar glycoproteins to anchor the pollen tube to the pollen/stylar extracellular matrix (psECM), via molecular mechanisms that are not fully understood [4, 5]. Polyamines (PAs) are among other secreted molecules necessary for the apical growth of germinating pollen [6, 7], helping to sustain pollen germination as demonstrated with inhibitors of PAs-biosynthetic enzymes [7-11]. The role of PAs maybe related to the structure and assembly of vegetative cell walls [12, 13] or the organization of the pollen cell wall [14] and these functions may be relevant to pollen from a variety of plant species [15]. The precise role of PAs secreted from the germinating pollen tube and their interaction with the psECM are not completely understood.

The interactions between the pollen tube and the psECM are reminiscent of the cell-ECM interactions in animal cells [5]. One key modulator of the ECM in mammals is tissue transglutaminase (tTGase also known as TGase 2), a member of a large family of enzymes [16, 17] that catalyzes the calcium dependent posttranslational modification of proteins, by forming covalent cross-links between glutamyl- and either lysyl-residues (N^ε(γ-glutamyl) lysine isodipeptide) or PAs (*mono* and *bis*-(γ-glutamyl) polyamines) resulting in a bridge between two or more proteins [18]. In animal cells, extracellular tTGase is concentrated at focal adhesions (FA), which are sites of communication between the ECM and intracellular signaling molecules [19-21]. After secretion, tTGase becomes a non-enzymically active structural component of the ECM, forming a close association with the glycoprotein fibronectin and heparan sulphate proteoglycans important in cellular adhesion and migration [20, 22, 23]. Transglutaminases have been shown to be present in plant tissue [24], including apple pollen, where it is an intracellular Ca²⁺-dependent enzyme catalysing the incorporation of PAs into α-tubulin, actin monomers and uncharacterised large protein complexes [25]. Recently monoclonal antibodies raised against mammalian tTGase (e.g. ID10) have been shown to cross-react with apple pollen TGase and that the enzyme's *in vitro* activity can be blocked by an animal TGase inhibitor. In addition, this partially purified apple pollen TGase has been shown to be capable of catalysing the *in vitro* cross-linking of a number of purified cytoskeletal components. This suggests that pollen TGase may be capable of modulating the properties of the pollen cytoskeleton and possibly the subsequent development of pollen tubes [26]. The presence of TGase substrates in germinating apple pollen has also been demonstrated by their *in situ* visualization in the pollen grain, and the elongated pollen tubes as well as the pollen surface [27]. Other studies have reported the presence of a TGase like activity in the cell wall of higher plant parenchyma or of lower plant cells whose cell wall composition is different from that of pollen tubes [24, 28, 29]. The above data suggests that an extracellular TGase maybe present in the cell wall of apple pollen tubes, and may contribute to the

11

12

3

interaction of pollen proteins with psECM proteins, by catalyzing protein-protein linkages.

The current study aims to demonstrate the presence of an extracellular TGase linked to the cell wall of pollen tube or released into the extracellular medium. Apple pollen germination was studied *in vitro* (by incubating the pollen in the germination medium) confirming the presence of TGase, and this was also verified *in planta* using pollen germinated inside the style. The role of TGase in the modulation of pollen tube apical growth was investigated by the application of specific TGase inhibitors, antibodies towards both mammalian and plant TGases and by assaying the activity of the enzyme in the presence of known TGase protein substrates and PAs.

MATERIAL AND METHODS

Chemicals and Antibodies

All chemicals (unless otherwise indicated) were obtained from Sigma-Aldrich (Milan, Italy). The tTGase inhibitors 283 (1,3-dimethyl-2[(oxopropyl)thio]imidazolium) [30] (now available also at Zedira, Darmstadt, Germany) and 281 (N-Benzoyloxycarbonyl-Lphenylalanyl-6-dimethylsulfonium-5-oxo-L-norleucine), two specific site-directed inhibitors of TGase, developed to covalently bind the Cys-based active site of the calcium activated form of mammalian TGase [31- 34] were synthesised at Nottingham Trent University, UK. The mouse monoclonal antibody ID10 raised against purified guinea pig liver (gpl) TGase (tTGase) was produced at Nottingham Trent University and selected by its reactivity to gpl TGase and ion exchange-purified extracts of 14 day old *Pisum sativum* leaf tissue and germinated *Vicia faba* cotyledons [35]. The TGase-reactive clone antibody was partially purified from hybridoma culture supernatant by cation exchange chromatography on Fast Flow S-sepharose (Amersham Biosciences, Little Chalfont, UK). ELISA-positive fractions were pooled, concentrated in a dialysis tube against PEG 8,000 and stored in aliquots at -20°C. The immuno-reactivity of ID10 for pollen TGase, purified by hydrophobic interaction chromatography, was compared to TGase antibody, AbIII (Neomarker, Fremont, CA). Reactivity of ID10 was also tested against different cell fractions of *Agaricus* and compared to the cross-reactivity obtained by CUB7402 on the same cells (personal communication of M. Della Mea, University of Bologna).

Guinea pig liver TGase was obtained from Sigma-Aldrich (Milan, Italy) and further purified in our laboratory. Recombinant His₆-Xpr-GFP (a TGase substrate) was produced at the Tokyo Institute of Technology (Yokohama, Japan) [36]. Primary antibodies include: monoclonal anti-TGase 2 Ab-I [clone Cub7402] (Neomarker, Fremont, CA); polyclonal antibody AtPng1P (produced in chicken at the CSIC of qBarcelona, Spain, as already reported [37]; 81D4 monoclonal anti-N^ε(γ-glutamyl)lysine (Covalab, Lyon, France); 2c10 monoclonal anti-horse albumin (a gift from Prof. E.E. Billett, Nottingham Trent University). The following antibodies were obtained from Sigma-Aldrich (Milan, Italy) unless otherwise indicated. T1A2 (anti-tyrosinated α-tubulin subunit); B512 (anti-total α-tubulin); non-immune mouse IgG (used as negative control); polyclonal anti α-tubulin antibody. The list of secondary antibodies includes: HRP-conjugated rabbit anti-mouse IgG; FITC-conjugated rabbit anti-mouse IgG (DakoCytomation, Glostrup, Denmark); FITC-conjugated goat anti-

mouse IgG secondary antibody (Cappell, Malvern, PA); FITC-conjugated goat anti-rabbit IgGs; goat anti-mouse IgG 15nm gold conjugated (Biocell).

Plant material and growth

Mature pollen of *Malus domestica* Borkh. cv Golden Delicious was collected from plants grown in experimental plots (Dipartimento di Colture Arboree, University of Bologna). Handling and storage were performed as previously reported [25].

In vitro germination

Pollen was re-hydrated and allowed to germinate as previously described [25] into glass Petri dishes or in N',N'-dimethyl casein (DMC) coated 96-well microtitre plates (Nunc Life Technologies, Paisley, UK) or 96-well microtitre fluo black plates (Nunc Life Technologies, Paisley, UK) for up to 120 min, in the presence and absence of His₆-Xpr-GFP, TGase inhibitors (281, 283, iodoacetamide, cystamine) and anti-TGase antibodies added as specified in the figure legends at zero time of germination or for the last 30 min to allow pollen tube to partially grow. As the monoclonal antibody ID10 gave the strongest labeling it was used for further experiments. Light microscopy digital images of at least three non-overlapping fields covering the central part of each chamber were captured using a video digital camera (Olympus DP10) and quantified in terms of germinated versus ungerminated pollen, when the pollen tube was either absent or shorter than the pollen grain diameter. Pollen viability was assessed by staining with fluorescein diacetate (10 µg ml⁻¹).

Pollen protein extraction

Pollen was collected after germination and separated from the culture medium by filtration with vacuum pump on Millipore disks (5 µm pore size). Pollen proteins and those secreted in the germination medium were extracted following the instruction described in the literature [38] and quantified [26]. Proteins were also isolated from pollen cell walls according to a previously published method [39].

Partial purification of pollen TGase

The partial purification was performed by a hydrophobic interaction chromatography (HIC) with Phenyl-Sepharose 6 Fast Flow (Sigma-Aldrich, Milan, Italy) at pH 8 [26]. Partially purified TGase was immuno-detected after SDS-PAGE and western blotting with two tTGase antibodies (ID10 and AbIII).

Immuno-precipitation of pollen TGase

Aliquots (200 µl) of HIC purified pollen extract (0.2 mg ml⁻¹) were incubated with increasing amount of ID 10 monoclonal antibody for 2 h on ice, and the samples immuno-precipitated as described [23]. The supernatant was assayed for TGase activity [19] and compared to that of the pre-cleaned sample obtained after incubation of HIC purified pollen extract (see above) directly with protein G-agarose beads (control) (Sigma-Aldrich, Poole, UK). The difference between the activities in the supernatants obtained after ID10 treatment and the control gave a quantitative indication of the

TGase activity lost due to the binding of the pollen TGase to the protein G-coupled ID10.

SDS-PAGE and Western blot analysis

A total of 50 µg of germinated pollen proteins obtained after (i) extraction after low speed centrifugation or from pollen cell walls, or (ii) following purification via HIC or (iii) after the concentration of the germination medium, were electrophoresed on 10% (w/v) polyacrylamide SDS-PAGE slab gels [40], using 0.2 µg gpl TGase as a standard. Western blotting was performed as described previously and protein bands that cross-reacted with ID10, AbIII, CUB 7402 or AtPng1P revealed accordingly to the literature [37].

Transglutaminase activity assays

Extracellular pollen TGase activity was measured by the conjugation of biotinylated cadaverine to DMC as described [19], with minor modifications. Native hydrated pollen grains (1 mg ml⁻¹ in germination medium) were allowed to germinate in the wells for 120 min at 30°C in germination medium (see above) buffered with 5 mM MES pH 6.5, containing 0.1 mM biotinylated cadaverine and 50 µM DTT. Germination was monitored by light microscopy throughout. After removal of germinated pollen by extensive washings with phosphate-buffered saline (PBS), the level of biotinylated cadaverine incorporated into immobilised DMC was revealed according to an established protocol [41] with EGTA replacing EDTA in negative controls. Data are expressed as Ca²⁺-dependent increase in A₄₅₀, after subtraction of the value of the 5 mM EGTA-treated control. Specific activity was determined as a change in A₄₅₀ of 0.1 per hour per mg of non-hydrated pollen.

To further verify extracellular pollen TGase protein cross-linking activity, His₆-Xpr-GFP was substituted for the biotin cadaverine. Pollen was germinated for 90 min into 96-microwell fluo black plate (Nunc Life Technologies, Paisley, UK) pre-coated with DMC in buffered germination medium. His₆-Xpr-GFP (25 µg ml⁻¹) was added for the last 30 min of germination in the absence or in the presence of increasing concentrations of anti-TGase antibody ID10 (3.5 - 21 µg ml⁻¹), or 283 and 281 (0.05 - 5 mM) inhibitors. After removal of the pollen (utilised for the confocal analysis of the His₆-Xpr-GFP cross-link to pollen tube wall proteins) and of the germination medium, the microwells were extensively washed with PBS. The fluorescence of His₆-Xpr-GFP cross-linked to immobilised DMC was measured in an ELISA plate reader (Titertek Multiscan ELISA Spectrophotometer, Flow Laboratories, Irvine, UK) at 485 nm excitation and 535 nm emission. The cross-linking activity was quantified as relative fluorescence units (RFU). Data were expressed as RFU with 1 mM Ca²⁺ in the reaction buffer minus background values with 0.125 mM EDTA and autofluorescence. Positive control values were adjusted to 100%, with the samples incubated with antibody or inhibitors reported as percentage (mean ± SD) of the control.

Analysis of mono- and bis-γ-glutamyl-polyamine derivatives

Proteins (0.3 mg) secreted into the culture medium by pollen germinated for 2 h were assayed for extracellular TGase activity and for pollen natural substrates [42] in the presence of 10 µl of [1,4(n)-³H] spermidine (specific activity 1.5 TBq mmol⁻¹; NEN,

25
26

271 Milan, Italy), 40 mM MES pH 6.5, 10 mM DTT, and 5 mM CaCl₂ in a final volume of
272 300 µl. In the negative control, CaCl₂ was substituted with 10 mM EDTA.

273
274
275

Histochemical staining of germinating pollen *in vitro*

276 The distribution of immuno-reactive TGase and its cross-linked reaction products was
277 determined by immuno-histochemical and immuno-fluorescence staining. Intracellular
278 immuno-staining of TGase was carried out on pollen germinated for up to 120 min,
279 after fixation, by digesting pollen for 7 min with 0.75% (w/v) pectinase (from
280 *Rhizopus*) and 0.75% (w/v) cellulase (from *Penicillium*) [43]. Cell wall digestion and
281 membrane permeabilization was omitted for extracellular staining of TGase. Non-
282 specific protein binding was blocked by incubating fixed pollen with 3% (w/v) BSA in
283 PBS pH 7.4, for 2 hours at 37°C. For immuno-fluorescence staining, further blocking
284 with 20% (v/v) non-immune serum was carried out. Once transferred on poly-lysine
285 coated-slides, pollen was incubated with the following mouse monoclonal antibodies:
286 T1A2 (anti-tyrosinated α -tubulin subunit), B512 (antitotal α -tubulin), ID10 (anti-
287 TGase) or 81D4 (anti-cross-link) using non-immune mouse IgG as a negative control.
288 Antibody-antigen binding was revealed using either HRP-conjugated rabbit anti-mouse
289 IgG followed by incubation with 3-amino-9-ethylcarbazole (AEC) as a peroxidase
290 substrate, or FITC-conjugated rabbit anti-mouse IgG. Staining was observed by a video
291 digital camera (Olympus DP10) producing digital images of at least 6 non-overlapping
292 fields covering the central portion of each sample. In some experiments, germinated
293 pollen was also treated with 2 M NaCl, 1% (w/v) SDS or 1% (w/v) DTT either prior to
294 or after fixation. Polyclonal anti α -tubulin antibody was used as the positive control and
295 non-immune mouse IgGs raised in rabbit as the negative control, which were revealed
296 by FITC-conjugated anti-rabbit IgGs. Immuno-gold detection of TGase was performed
297 exactly as previously described [44]; the primary anti-TGase antibody was used at the
298 dilution of 1:5 while the secondary goat anti-mouse IgG 15-nm gold conjugated was
299 used at 1:20. *In situ* TGase activity was visualized by incubation of germinated pollen
300 with His₆-Xpr-GFP, a specific glutamine and lysine rich fluorescent substrate for ECM
301 mammalian TGase modified as reported [36] at different concentrations (from 0.25 to
302 50 µg ml⁻¹). Fluorescence staining, was observed using a Leica TCSNT confocal laser
303 scanning microscope system (Leica Lasertechnik, Heidelberg, Germany), with the
304 photo-multiplier tube (PMT) adjusted to minimize autofluorescence emission in the
305 negative control. Images from at least 10 random fields, were scored for the distribution
306 of immuno-reactive TGase and N^ε(γ -glutamyl)-PA or lysine cross-link, present in
307 pollen proteins with the aid of the Leica TCSNT (version 1.5-451) image processing
308 menu. For double staining, germinated pollen was incubated with recombinant His₆-
309 Xpr-GFP as described before, and after extensive washes, further incubated with either
310 81D4 or ID10, which were revealed by TRITC-conjugated anti-mouse antibody.
311 Double staining was performed by adjusting the argon/krypton laser at 488 and 560 nm
312 for fluorescein and rhodamine excitation, respectively. As a further control, individual
313 antibody staining was also performed in parallel to double staining.

314
315
316

Histochemical staining of germinating pollen *in planta*

317 Fresh pollen was applied to the styles in apple flowers and after 48 h the styles were
318 collected and solubilized in buffer (100 mM Pipes-KOH pH 6.8, containing 10 mM
319 MgCl₂, 10 mM EGTA, 0.1%w/v NaN₃, and 0.05% v/v Triton X-100) supplemented
320 with fixatives (1.5% w/v paraformaldehyde, 0.05% v/v glutaraldehyde). After 60 min at

27
28

7

29
30

4°C the styles were cut longitudinally into four pieces using razor blades with the aid of a stereo microscope. The cell walls were digested with 0.75% (w/v) cellulysine and 0.75% (w/v) pectinase in buffer for 7 min in the dark. After washing with buffer, the primary antibody (ID10, diluted 1:20 in buffer) was added to samples and incubated for 1 h at 37°C. Samples were then washed with buffer and incubated for 1 h at 37°C with the FITC-conjugated goat anti-mouse IgG secondary antibody (Cappel, Malvern, PA) diluted 1:50 in buffer. Styles were finally mounted in anti fade mountant (Citifluor, London, UK) and observed with a Zeiss Axiophot fluorescence microscopy equipped with an AxioCam MRc5 and a 63x oil-immersion objective (Zeiss Inc., Oberkochen, Germany).

Statistics

The values reported are expressed as mean $\pm$ SD and represent one of at least 3 or 4 different experiments undertaken in triplicate. Differences between sample sets were determined by the Student's t-test with 95% confidence limits. When indicated statistical analysis was performed using GraphPad Prism (version 5.0a McIntosh GraphPad software, San Diego, CA, USA). Percent data from pollen germination were analyzed after arcsin $\rightarrow$ p transformation. Pearson correlation analysis was performed when necessary.

RESULTS

Localization of transglutaminase in pollen tubes by immunological techniques

To localize TGase in pollen tubes, pollen was allowed to germinate for 120 min *in vitro*, after which the pollen tube wall was incubated with or without cellulase and pectinase, and its membrane permeabilized [43] prior to probing with the ID10 anti-TGase monoclonal antibody. This monoclonal antibody has been shown to immuno-react with partially purified TGase extracted from apple pollen [26]. Two well-characterized anti-tubulin antibodies were utilised as positive controls and anti-mouse IgGs as negative controls. Cellulase-digested and membrane permeabilized pollen tubes resulted in intracellular staining for TGase in the neck region and along the shank of the pollen tube, especially at the apical tip (Figure 1A a-c). Anti α -tubulin staining was distributed along the tube, whereas tyrosinated α -tubulin was enriched at the growing tip (Figure 1A d-f), in agreement with previous findings [45-46]. The distribution of TGase along the pollen tube was confirmed by confocal immuno-staining (Figure 1B). To specifically visualize extracellular TGase, the staining with ID10 was replicated in non-digested and non-permeabilised germinating pollen (Figure 1D a-c). In these conditions, extracellular TGase was detected along the pollen tube surface, in aggregates outside the tube and also on the pollen grain. Treatment with 2 M NaCl, 1% (w/v) SDS or 1% (w/v) DTT prior to fixation and immuno-labeling did not dissociate the TGase from the tube wall, suggesting either a covalent or a stable interaction between the enzyme and the pollen tube cell wall had occurred (Figure 1D a-b). Fixation did not affect the immuno-labeling, as incubation with the antibody prior to fixation led to a similar pattern of TGase staining (Figure 1D c). Digestion of the cell wall with cellulase and pectinase following the immuno-staining with ID10 resulted in an almost complete loss of the detection of extracellular TGase, except from around the pollen grain wall (which is protected by the cellulase-resistant sporopollenin) (Figure

31
32

8

33
34

1D d). No sign of osmotic stress was observed. The localization of TGase was also investigated by immuno-gold labeling, which showed the presence of TGase in the pollen tube wall and surface (Figure 1E, arrows). Western blotting utilizing anti-TGase antibody revealed two immuno-reactive bands of 70 and 75 kDa, the latter more marked, in crude extracts of germinating pollen (Figure 1C, lane CE), in concentrated germination medium in which germinated pollen had been removed by filtration (Figure 1C, lane GM), and in cell walls (Figure 1C lane CW). A weakly reactive band of lower molecular weight (around 55 kDa) was observed in the cell wall extracts (Figure 1C lane CW). As a positive control, western blotting was conducted on partially purified 70 kDa TGase isolated from pollen supernatant by hydrophobic ion chromatography [26], (Figure 1C, lane C, WB). This EGTA-eluted fraction was immuno-detected in a dose dependent manner by ID10 and by AbIII, another monoclonal anti-TGase antibody that recognizes plant TGases (Supplementary Figure S1) as reported by Della Mea *et al.* [29]. The partially purified 70 kDa TGase catalyses transamidating and cross-linking reactions, as described by Del Duca *et al.* [26].

Inhibition of extracellular TGase activity leads to inhibition of pollen germination and tube apical growth

Evidence for the activity of the extracellular TGase immuno-detected in viable pollen tubes was obtained by allowing pollen to germinate in the wells of microtitre plates (to 78±5% germination) followed by *in situ* determination of Ca²⁺-dependent TGase-mediated incorporation of biotin cadaverine into DMC immobilized in the plate wells (EGTA-treated samples were used as the control). Typical values of extracellular TGase activity are shown in Figure 2A,B (control bars, bottom histograms). The pH optimum for the extracellular pollen TGase activity was determined to be 6.5, which is also the optimum pH of the medium to allow pollen germination. Extracellular pollen TGase activity (Figure 2, bars), percentage germination (Figure 2, lines) and tube growth (Figure 2, top images) were then measured in the presence of TGase inhibitors and a TGase specific antibody (Figure 2A,B, respectively). All three events were inhibited in a dose dependent manner by the monoclonal antibody ID10 (Figure 2A) and by the two site-directed irreversible inhibitors of TGase activity, 283 and 281 (Figure 2B). The presence of the inhibitors resulted in shorter and thicker pollen tubes with a decreased growth rate within the first 30 min of germination, followed by an inability to further extend the tip, between 30 and 60 min after germination (Figure 2A,B). Eventually, tube-burst occurred in a dose dependent manner for antibody concentrations higher than 7 µg ml⁻¹ and at inhibitor concentrations above 100 µM. Inhibition of TGase activity and pollen tube growth were not observed with the control antibody (Figure 2A). The significant correlation between pollen germination and TGase activity when inhibited by ID10, 281 and 283 respectively, is supported by high value Pearson coefficients. In a similar fashion, other less specific inhibitors of TGase activity, cystamine and iodoacetamide (50 –1000 µM) inhibited enzyme activity and pollen tube growth in a dose dependent manner; at 1 mM, cystamine and iodoacetamide inhibited the TGase activity by 57±3% and 44±5% respectively; the *in vivo* supply of 1 mM cystamine inhibited the tube growth by 40±4%. A competitive substrate of TGase activity, putrescine (assayed in the range 1 µM-2.5 mM) led to an inhibition of tube growth between 30±4% and 53±2%. Further confirmation that ID10 inhibited pollen TGase was obtained by immuno-precipitation of partially purified pollen TGase [26], which led to a dose-dependent inhibition of TGase activity, whereas immuno-precipitation using the negative control antibody (monoclonal anti-horse albumin)

35
36

9

37
38

produced no effect (Supplementary Figure S2). Western blot analysis to determine the specificity of ID10 showed that it reacted only to a single 75 kDa band corresponding to TGase 2 in gpl homogenates and did not recognize TGase 1 or TGase 3 confirming an anti-TGase 2 antibody (Supplementary Figure S3).

Effect of Gln and Lys-rich TGase substrate on pollen germination and tube elongation

Experiments were performed to evaluate the effect on pollen of His₆-Xpr-GFP, a specific glutamine- and lysine-rich substrate of mammalian ECM TGases [36]. Incubation of His₆-Xpr-GFP in the germination medium during pollen germination and subsequent *in situ* localization by confocal microscopy, in non-digested and non-permeabilised pollen, revealed that His₆-Xpr-GFP was associated extracellularly with the pollen tube wall (Figure 3A). The addition of His₆-Xpr-GFP at 0.25 µg ml⁻¹ gave rise to a staining located within 20 µm from the tip; increasing the concentrations (1-50 µg ml⁻¹) resulted in a proportional increase in the incorporation along the tube and into aggregates (Figure 3A). At the highest concentration (50 µg ml⁻¹), equivalent to that used for mammalian tissue [36], longer segments of tube were fluorescently labelled, with the additional formation of larger extracellular structures similar to those recognized by the anti-TGase monoclonal antibody ID10 (Figure 1D). Extracellular pollen TGase-mediated crosslinking of His₆-Xpr-GFP to pollen tube proteins was confirmed by inhibition with 281, 283 and by the blocking effect of ID10 (Figure 3B). Unexpectedly, at concentrations between 1 and 25 µg ml⁻¹, His₆-Xpr-GFP resulted in a significant increase in the percentage of germinated pollen, when compared to the pollen incubated with control mouse IgG (*p* < 0.05; Table 1). Incubation with 5 µg ml⁻¹ of this fluorescent TGase substrate resulted in longer pollen tubes when compared to incubation with the control mouse IgG (Figure 3C). These effects were not produced by unmodified GFP, rGFPuv (data not shown). A modified microtiter "fluor black" plate assay was set up for the detection of cross-linking activity of extracellular TGase of germinating pollen, which involved incubating pollen in germination medium containing His₆-Xpr-GFP in wells coated with immobilised DMC. Fluorescence measurement showed cross-linking of His₆-Xpr-GFP to DMC (Figure 3D, control). This activity was inhibited using both the anti-tTGase ID10 and the inhibitors 281 and 283 (Figure 3D). Both sets of data suggest that during pollen germination the detected cross-linking is mediated by extracellular TGase activity. Evidence that endogenous pollen proteins are substrates for pollen TGase was obtained by immuno-histochemical staining with 81D4 (Figure 3E). This antibody specifically binds TGase-mediated protein-protein and amine-protein cross-links, reacting with bis-PA derivatives > gln-lys isodipeptide linkage > mono-PA derivatives but not with their free counterparts [47]. The distribution of anti cross-link staining was observed along the pollen tube surface (Figure 3E). The ability of pollen extracellular TGase to incorporate the exogenously supplied His₆-Xpr-GFP in the same sites of pollen protein-protein and protein-amine cross-links revealed by 81D4 was demonstrated by their co-localisation in confocal micrographs (Figure 4A). In addition, the extracellular pollen TGase co-localized within the same accumulation sites of His₆-Xpr-GFP cross-link (Figure 4B).

Presence of pollen extracellular TGase in the germination medium

The ability of extracellular pollen TGase to conjugate PAs to pollen proteins secreted during germination *in vitro* was analyzed by incubation with the TGase amine substrate

39
40

10

41
42

[³H]spermidine as a tracer. The proteins were extracted from the concentrated germination medium after 120 min of germination and the removal of pollen. The germination medium contained 1560 DPM and 1816 DPM of *mono*-(γ -glutamyl) spermidine and *bis*-(γ -glutamyl) spermidine derivatives per g of germinated pollen, respectively. These data confirm the presence of TGase catalyzed reaction products in the pool of *in vitro* secreted pollen proteins.

***In planta* localization of extracellular pollen transglutaminase**

To confirm the presence of extracellular TGase *in vivo*, pollen tubes germinated (*in planta*) in the stylar tissue of carpels were fixed and stained by indirect immunofluorescence using ID10. As shown in Figure 5, anti-TGase reactivity was localized in extracellular aggregates along the pollen tube surface in a similar distribution to that observed *in vitro* (Figures 1D and 3A). By contrast, no ID10 staining was observed on the surface of stylar tissue, as shown by the black background (Figure 5, arrows).

DISCUSSION

The data presented here provides evidence that in germinating apple pollen there is the a catalytically-active TGase, which is localized extracellularly and is involved in pollen tube growth.

Localization

The model presented in Figure 6 summarizes the main results discussed below. In the non-permeabilised *Malus* pollen tube TGase was found in discrete aggregates outside the tube both *in vitro* (Figure 1D,E) and *in planta* (Figure 5). In permeabilised germinated pollen, TGase antibody labelling was observed at the apical tip of the growing pollen tube (Figure 1A), suggesting that accumulated TGase is present in and/or released from the pollen tube apex. As the pollen tube elongates during its growth, the released TGase could be deposited along the tube wall, as shown by extracellular TGase labeling on the pollen tube surface both *in vitro* and *in planta* (Figures 1D,E and 5). However, the mechanism by which pollen TGase and tTGase of mammalian cells [22] is actually secreted in the psECM or ECM respectively remains unknown. The intracellular distribution of TGase (Figure 1B) in the pollen tubes appears to be within vesicle-like structures. These may be the same or similar to those containing cell wall precursors and enzymes that are released from the apical tube tip [5]. Sub-cellular fractionation of *Nicotiana* petal cells led to the detection of TGase antigen and activity in the cell wall and microsomal fraction [29], suggesting the compartmentalisation of plant TGase in membrane structures.

The cross-reactivity of antibodies raised against animal TGases to react with plant TGases has previously been reported in studies on different plant tissues [24, 29]. In germinated apple pollen, two immuno-reactive polypeptides of 75 and 70 kDa (Figure 1C) cross-reacted with the same anti-TGase antibody (ID10) used to visualize the 70 kDa TGase partially purified from ungerminated apple pollen [26]. It is not known if the two TGase forms have a specific role or different characteristics; they may differ in their calcium binding properties which allowed only the 70 kDa band to bind to the hydrophobic interaction column used for purification. Supplementary Figure S1 shows that 70 kDa band matches with both ID10 and AbIII. The 70 kDa TGase showed *in*

43
44

11

45
46

vitro Ca²⁺-dependent transamidating activity on cytoskeletal substrates and was inhibited by TGase inhibitors [26] well characterised by Balklava *et al.* [22] and by Baumgartner *et al.* [33]. The detection of the 70 and the 75 kDa forms in both the cell wall and in the germination medium (Figure 1C), suggests that they are both secreted. We cannot exclude the possibility that the 70kDa form has been derived from the 75kDa form from a post-translational modification and that this may have occurred prior to or post secretion. The reduction in size has also altered the ability of the 70kDa forms to interact with an HIC column in the presence of Ca²⁺. Further work will be needed to characterise the roles of these TGase isoforms in pollen germination.

The additional 55 kDa immuno-reactive polypeptide observed on Western blots (Figure 1C) of the cell wall fraction has the same molecular mass as an active TGase extracted from *Nicotiana* petal cell wall and may be involved in cell wall strengthening [29]. A similar role in cell wall organization has also been suggested for a TGase from the unicellular green alga, *Chlamydomonas reinhardtii* [28]. Only one TGase gene in apple genome seems to occurs similarly to AtPNglp in Arabidopsis genome (personal communication of Dr. R.Velasco and Dr. A. Cestaro, IASMA).

Substrates

The presence of PAs in apple pollen (ungerminated or germinated), as well as their release from the pollen tube has previously been established [6, 7]. In kiwi pollen, free, TCA-soluble and -insoluble conjugated PAs were also detected, with spermidine (SD) conjugates being the most abundant conjugated PA [9]. Inhibition of PAs biosynthesis caused a severe inhibition of pollen tube growth, possibly due to reduced binding to cell wall polysaccharides [11]. The PA presence in the cell wall of higher plants and their deficiency, which induced loosening of the fibrillar component, are pertinent to the structure of the cell wall. In vegetative tobacco cells microfibrillar cellulose structures and pectic substances of the cell wall were altered under treatments with PA biosynthesis inhibitors [12]. PAs are also suggested to contribute to vegetative cell wall assembly by forming amide bonds with pectins [13]. Hydroxycinnamic acid (HCA)-PAs have been found in the pollen of various plant species, thus suggesting a general role in modulating the rigidity of the cell wall [15]. The triferuloyl-SD conjugate has a possible role in the organization of the *Arabidopsis* pollen cell wall, and as a component of the tryphine (constituents of the pollen coat), which is involved in pollination and in pollen-stigma interaction [14].

PAs are known to be substrates for TGase enzymes [18]. The intracellular PAs of apple pollen were reported to be conjugated by TGase to cytoskeletal proteins regulating their functional properties [26]. In the present study the occurrence of pollen protein cross-links in the germination medium was demonstrated by the detection of labeled glutamyl-PAs derivatives. The detection of a TGase in the apple pollen cell wall (Figure 3 D, E), could support a role for protein-conjugated PAs in pollen tube growth and stability. TGase activity in apple pollen extracts could be detected using a PA incorporation assay and inhibited by an anti-TGase antibody and specific TGase inhibitors (Figure 2A,B).

The secreted apple pollen TGase catalyzed the incorporation of two exogenous substrates (i.e. H₆-Xpr-GFP and DMC) into the pollen cell wall proteins (Figure 3A,D), and this incorporation was inhibited by specific TGase inhibitors (Figure 3B). The presence of TGase mediated protein cross-links was demonstrated by *in situ* immuno-detection using the anti cross-link monoclonal antibody 81D4 (Figures 3E and 4A). The identity of these protein substrates is not known.

47
48

12

49
50

In pollen, we have shown that TGase activity could be stimulated by high concentration of substrates (Figure 3A for higher His₆-Xpr-GFP concentration), which possibly act as a nucleation centre for further cross-linking. The concomitant enhancement of pollen tube growth associated with His₆-Xpr-GFP protein cross-linking at the pollen tube surface suggests that His₆-Xpr-GFP facilitates pollen tube apical growth. The finding that both anti-TGase antibodies and inhibitors (mainly acting extracellularly) block His₆-Xpr-GFP binding to cell wall and inhibit pollen tube growth is a clear indication that the catalytic activity of extracellular pollen TGase is critical for correct pollen tube growth in *Malus*.

The co-localisation between extracellular TGase antigen and cross-linked products suggests that apple pollen TGase could become embedded, under the effect of local micro-environmental conditions, into the same accumulation sites of the protein cross-links catalyzed by its enzyme activity. The localization of TGase-catalyzed cross-linked aggregates along the shank of pollen tubes is similar to that observed in the cell wall during the pulsatory growth of pollen tubes in flowering plants with a solid style. However, the distribution of pectins and AGPs reported in the literature does not fit exactly with the irregular distribution of the TGase-catalyzed aggregates [48]. These discontinuous cross-linked structures may reflect oscillations of other factors influencing TGase activity and pollen growth, e.g. changes in Ca²⁺ and pH [1]. These collective results indicate that when the apple pollen TGase is externalised it remains biologically active.

Common aspects with other TGases

The evidence that extracellular TGase activity from germinating pollen tubes is inhibited by compounds that inhibit animal tTGase (281 and 283, iodoacetamide and cystamine), [16, 30-33] and by anti-TGase antibodies (Figure 2A,B) strongly suggests that the plant enzymes are catalytically similar to animal TGases. In mammalian cells, imidazolium derivatives such as 283 and 281 have been described as potent inhibitors of mammalian TGases [22, 30, 32]. The inhibitor 281 is reported to act primarily outside the cell [33] and to inhibit pollen TGase activity by reducing the fluorescein-cadaverine cross-linking to pollen proteins during germination [27].

When the germination occurs *in planta* inside the style, the secreted TGase could have a potential role in the adhesion of pollen tube to stylar cells, similar to the role of mammalian tTGase in cell-ECM interactions [19-23], thus allowing style anchorage and subsequent tube migration. Interestingly, in mammalian cells extracellular tTGase remains embedded in the ECM thus acting as a structural adhesive protein as well as a cross-linking enzyme involved in cell adhesion [20]. tTGase has been shown to co-localise with fibronectin and integrins-β1 at focal adhesion structures [49,50], where it acts as an integrin co-receptor; once secreted, tTGase modifies the ECM by cross-linking fibronectin and collagen thus leading to the modulation of cell adhesion's [20].

The above reported similarity between animal tTGase and pollen TGase is confirmed by the evidence that partially purified 70 kDa TGase extracted from pollen shares similar properties to gpl TGase in affecting the functional properties of the cytoskeleton [26]. The inhibition of pollen TGase by 281, which covalently binds the cysteine at the active site of mammalian TGase, also suggests the presence of a similar cysteine at the active site in *Malus* pollen TGase. This is in agreement with the observation that the typical catalytic triad of the TGase family (His-Cys-Asp) [16, 17] is conserved in *Arabidopsis* TGase (AM745095) [37] and in an EST sequence in the *Malus* database (Tree Fruit Technology, Michigan State University), which shows 62%

51
52

13

53

54

621 homology with *Arabidopsis* TGase (increasing to 77% homology with TGase active-
622 site containing regions).

623 In conclusion, the data presented in this work suggest that pollen TGase plays an
624 essential role in successful apple pollen tube growth. The enzyme could, therefore,
625 function by *in vitro* protein cross-linking and amine protein conjugation, strengthening
626 the cell wall scaffold of the extending pollen tube. *In planta*, we hypothesize that the
627 extracellular TGase might be involved in connecting the pollen tube to surrounding
628 stylar cells as it progresses through the psECM. Work in progress will help to further
629 characterize the molecular mechanisms by which pollen TGase facilitates successful
630 pollen tube growth.

631

632

633 ACKNOWLEDGEMENTS

634

635 Financial sources. This work was supported by the Italian Ministero dell'Università
636 (PRIN 2007 and - Progetto Strategico d'Ateneo Unibo 2006 Crossallergenicity) to
637 DSF. The ESF Transglutaminase Program, the COST Action 844 and Progetto Giovani
638 Ricercatori (University of Bologna) supported the travel grants to Dr. A. Di Sandro and
639 Prof S. Del Duca. We are also grateful to the British-Italian partnership project 2005-
640 2006 to Dr. A. Di Sandro and Dr E. Verderio for partially supporting the work.

641 We thank Dr. R. Velasco and Dr. A. Cestaro (IASMA, Trento) for the TGase sequence
642 investigation on the not yet published apple genome.

643 We thank Mr. N. Mele and Dr. C. Pagnucco (University of Bologna) respectively for
644 the image elaboration and glutamyl-derivatives analysis and Dr R. Saints (Nottingham
645 Trent University) for the production of inhibitors.

646

55

56

14

57

58

647References

648

6491 Holdaway-Clarke, T.L. and Hepler, P.K. (2003) Control of pollen tube growth: role of
650 ion gradients and fluxes. *New Phytol.* **159**, 539–563.

651

6522 Heslop-Harrison, J. (1979) Pollen-stigma interaction in grasses: a brief review. *New*
653 *Zealand J. Bot.* **17**, 537–546.

654

6553 de Graaf, B.H.J., Knuiman, B.A., Derksen, J. and Mariani, C. (2003) Characterization
656 and localization of the transmitting tissue-specific PELP III proteins of *Nicotiana*
657 *tabacum*. *J. Exp. Bot.* **54**, 55–63.

658

6594 Cheung, A.Y., Wang, H. and Wu, H. (1995) A floral transmitting tissue-specific
660 glycoprotein attracts pollen tubes and stimulates their growth. *Cell* **82**, 383–393.

661

6625 Lord, E.M. and Russell, S.D. (2002) The mechanisms of pollination and fertilization in
663 plants. *Ann. Rev. Cell Develop. Biol.* **18**, 81–105.

664

6656 Speranza, A. and Calzoni, G.L. (1980) Compounds released from incompatible apple
666 pollen during in vitro germination. *Z. Pflanzenphysiol.* **97**, 95–102.

667

6687 Bagni, N., Adamo, P. and Serafini-Fracassini, D. (1981) RNA, proteins and polyamines
669 during tube growth in germinating apple pollen. *Plant Physiol.* **68**, 727–730.

670

6718 Biasi, R., Altamura, M.M. and Bagni, N. (1999) Identification of markers for sexual
672 determination in *Actinidia deliciosa*. *Acta Hort.* **498**, 93–97.

673

6749 Falasca, G., Franceschetti, M., Bagni, N., Altamura, M.M. and Biasi, R. (2010)
675 Polyamine biosynthesis and control of the development of functional pollen in kiwifruit.
676 *Plant Physiol. Biochem.* In press.

677

67810 Chibi, F., Matilla, A.J., Angosto, T. and Garrido, D. (1994) Changes in polyamine
679 synthesis during anther development and pollen germination in tobacco (*Nicotiana*
680 *tabacum*). *Physiol. Plant.* **92**, 61–68.

681

68211 Antognoni, F. and Bagni, N. (2008) Bis(guanylhya zones) negatively affected in vitro
683 germination of kiwifruit pollen and alter the endogenous polyamine pool. *Plant Biol.* **10**,
684 334–341.

685

68612 Berta, G., Altamura, M.M., Fusconi, A., Cerruti, F., Capitani, F. and Bagni, N. (1997)
687 The plant cell wall is altered by inhibition of polyamine biosynthesis. *New Phytol.* **137**,
688 569–577.

689

69013 Lenucci, M., Piro, G., Miller, J.G., Dalessandro, G. and Fry, S.C. (2005) Do polyamines
691 contribute to plant cell wall assembly by forming amide bonds with pectins?
692 *Phytochemistry* **66**, 2581–2594.

693

69414 Grienberger, E., Besseau, S., GeoVroy, P., Debayle, D., Heintz, D., Lapierre, C.,
695 Pollet, B., Heitz, T. and Legrand, M. (2009) A BAHD acyltransferase is expressed in the

59

60

15

61
62

696 tapetum of *Arabidopsis* anthers and is involved in the synthesis of hydroxycinnamoyl
697 spermidines. *Plant J.* **58**, 246-259.

698

69915 Bokern, M., Witte, L., Wray, V., Nimtz, M. and Meurer-Grimes, B. (1995) Trisubstituted
700 hydroxycinnamic acid spermidines from *Quercus dentata* pollen. *Phytochemistry* **39**,
701 1371-1375.

702

70316 Lorand, L. and Graham, R.M. (2003) Transglutaminases: crosslinking enzymes with
704 pleiotropic functions. *Nature Rev. Mol. Cell Biol.* **4**, 140-156.

705

70617 Griffin, M., Casadio, R. and Bergamini, C. (2002). Transglutaminases: nature's
707 biological glues. *Biochem. J.* **368**, 377-396.

708

70918 Folk, J.E., Park, M.H., Chung, S.I., Schrode, J., Lester, E.P. and Cooper, H. (1980)
710 Polyamines as physiological substrates for transglutaminases. *J. Biol. Chem.* **255**, 3695-
711 3700.

712

71319 Jones, R.A., Nicholas, B., Mian, S., Davies, P.J. and Griffin, M. (1997) Reduced
714 expression of tissue transglutaminase in a human endothelial cell line leads to changes in
715 cell spreading, cell adhesion and reduced polymerisation of fibronectin. *J. Cell Science*
716 **110**, 2461-2472.

717

71820 Verderio, E., Telci, D., Okoye, A., Melino, G. and Griffin, M. (2003) A novel RGD
719 independent cell adhesion pathway mediated by fibronectin-bound tissue
720 transglutaminase rescues cells from anoikis. *J. Biol. Chem.* **278**, 42604-42614.

721

72221 Belkin, A.M., Tsurupa, G., Zemskov, E., Veklich, Y., Weisel, J.W. and Medved, L.
723 (2005) Transglutaminase-mediated oligomerization of the fibrin(ogen) α C-domains
724 promotes integrin-dependent cell adhesion and signaling. *Blood* **105**, 3561-3568.

725

72622 Balklava, Z., Verderio, E., Collighan, R., Gross, S., Adams, J. and Griffin, M. (2002)
727 Analysis of tissue transglutaminase function in the migration of Swiss 3T3 fibroblasts:
728 the active-state conformation of the enzyme does not affect cell motility but is important
729 for its secretion. *J. Biol. Chem.* **277**, 16567-16575.

730

73123 Scarpellini, A., Germack, R., Lortat-Jacob, H., Muramatsu, T., Billett, E., Johnson, T.
732 and Verderio, E. (2009) Heparan sulfate proteoglycans are receptors for the cell-surface
733 trafficking and biological activity of transglutaminase-2. *J. Biol. Chem.* **284**, 18411-
734 18423.

735

73624 Serafini-Fracassini, D. and Del Duca, S. (2008) Transglutaminases: widespread cross-
737 linking enzymes in plants. *Ann. Bot.* **102**, 145-152.

738

73925 Del Duca, S., Bregoli, A.M., Bergamini, C. and Serafini-Fracassini, D. (1997)
740 Transglutaminase-catalyzed modification of cytoskeletal proteins by polyamines during
741 the germination of *Malus domestica* pollen. *Sex. Plant Reprod.* **10**, 89-95.

742

74326 Del Duca, S., Serafini-Fracassini, D., Bonner, P., Cresti, M. and Cai, G. (2009) Effects of
744 post-translational modifications catalyzed by pollen transglutaminase on the functional
745 properties of microtubules and actin filaments. *Biochem. J.* **418**, 651-664.

63
64

16

65

66

- 74627 Iorio, R.A., Di Sandro, A., Scarpellini, A., Del Duca, S., Serafini-Fracassini, D. and
 747 Verderio, E. (2008) Visualisation of transglutaminase-mediated cross-linking activity in
 748 germinating pollen by laser confocal microscopy. *Plant Biosystems* **142**, 360-365.
 749
- 75028 Waffenschmidt, S., Kusch, T. and Woessner, J.P. (1999) A transglutaminase
 751 immunologically related to tissue transglutaminase catalyzes cross-linking of cell wall
 752 proteins in *Chlamydomonas reinhardtii*. *Plant Physiol.* **121**, 1003-1015.
 753
- 75429 Della Mea, M., De Filippis, F., Genovesi, V., Serafini-Fracassini, D. and Del Duca, S.
 755 (2007) The acropetal wave of developmental cell death (DCD) of *Nicotiana tabacum*
 756 corolla is preceded by activation of transglutaminase in different cell compartments.
 757 *Plant Physiol.* **144**, 1211-1222.
 758
- 75930 Freund, K.F., Doshi, K.P., Gaul, S., Claremon, D.A., Remy, D.C., Baldwin, J.J.,
 760 Pitzenberger, S.M. and Stern, A.M. (1994) Transglutaminase inhibition by 2-[(2-
 761 Oxopropyl)thio]imidazolium derivatives: mechanism of Factor XIIIa inactivation.
 762 *Biochemistry* **33**, 10109-10119.
 763
- 76431 Griffin, M., Coutts, I.G. and Saint, R.E. (2004) Novel compounds and methods of using
 765 them. International Publication No WO 2004/113363 A2.
 766
- 76732 Griffin, M., Mongeot, A., Collighan, R., Saint, R.E., Jones, R.A., Coutts, I.G. and
 768 Rathbone, D.L. (2008) Synthesis of potent water-soluble tissue transglutaminase
 769 inhibitors. *Bioorg. Med. Chem. Letters* **18**, 5559-5562.
 770
- 77133 Baumgartner, W., Golenhofen, N., Weth, A., Hiiragi, T., Saint, R., Griffin, M. and
 772 Drenckhahn, D. (2004) Role of transglutaminase 1 in stabilisation of intercellular
 773 junctions of the vascular endothelium. *Histochem. Cell Biol.* **122**, 17-25.
 774
- 77534 Beck, K.E., De Girolamo, L.A., Griffin, M. and Billet, E.E. (2006) The role of tissue
 776 transglutaminase in 1-methyl-4-phenylpyridinium(MPP+)-induced toxicity in different
 777 human SH-SY5Y neuroblastoma cells. *Neurosci. Lett.* **405**, 46-51.
 778
- 77935 Durose, L.J. (2001) Partial purification and characterisation of transglutaminases from
 780 dicotyledonous plant tissues. PhD Thesis, Nottingham Trent University, UK.
 781
- 78236 Furutani, Y., Kato, A., Notoya, M., Ghoneim, M.A. and Hirose, S. (2001) A Simple
 783 assay and histochemical localization of transglutaminase activity using a derivative of
 784 green fluorescent protein as a substrate. *J. Histochem. Cytochem.* **9**, 247-258.
 785
- 78637 Della Mea, M., Caparròs-Ruiz, D., Claparols, I., Serafini-Fracassini, D. and Rigau, J.
 787 (2004) AtPng1p. The first plant transglutaminase. *Plant Physiol.* **135**, 2046-2054.
 788
- 78938 Cai, G., Romagnoli, S., Moscatelli, A., Ovidi, E., Gambellini, G., Tiezzi, A. and Cresti,
 790 M. (2000) Identification and characterization of a novel microtubule-based motor
 791 associated with membranous organelles in tobacco pollen tubes. *Plant Cell* **12**, 1719-
 792 1736.
 793
- 79439 Yi-Qin, L., Croes, A.F. and Linskens, H.F. (1983) Cell-wall proteins in pollen and roots
 795 of *Lilium longiflorum*: extraction and partial characterization. *Planta* **158**, 422-427.

67

68

17

69

70

- 79640 Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of
797 bacteriophage T4. *Nature* **227**, 680-685.
798
- 79941 Lilley, G.R., Skill, J., Griffin, M. and Bonner, P.L. (1998) Detection of Ca^{2+} -dependent
800 transglutaminase activity in root and leaf tissue of monocotyledonous and
801 dicotyledonous plants. *Plant Physiol.* **117**, 1115-1123.
802
- 80342 Del Duca, S., Beninati, S. and Serafini-Fracassini, D. (1995). Polyamines in chloroplasts:
804 identification of their glutamyl- and acetyl-derivatives. *Biochemistry J.* **305**, 233-237.
805
- 80643 Miller, D.D., Scordilis, S.P. and Hepler, P.K. (1995) Identification and localization of
807 three classes of myosins in pollen tubes of *Lilium longiflorum* and *Nicotiana glauca*. *J.*
808 *Cell Sci.* **108**, 2549-2653.
809
- 81044 Persia, D., Cai, G., Del Casino, C., Faleri, C., Willemse, M.T.M. and Cresti, M. (2008)
811 Sucrose synthase is associated with the cell wall of tobacco pollen tubes. *Plant Physiol.*
812 **147**, 1603-1618.
813
- 81445 Del Casino, C., Li, Y., Moscatelli, A., Scali, M., Tiezzi, A. and Cresti, M. (1993)
815 Distribution of microtubules during the growth of tobacco pollen tubes. *Biol. Cell* **79**,
816 125-132.
817
- 81846 Westermann, S. and Weber, K. (2003) Post-translational modifications regulate
819 microtubule function. *Nature* **4**, 938-947.
820
- 82147 Thomas, V., Fournet, G., Simonet, F., Roch, A.M., Ceylan, I., El Alaouia, S. and Quash,
822 G. (2004) Definition of the fine specificity of the monoclonal antibody 81D4: its
823 reactivity with lysine and polyamine isopeptide cross-links. *J. Immunol. Met.* **292**, 83-95.
824
- 82548 Li, Y.Q., Fang, C., Faleri, C., Ciampolini, F., Linskens, H.F. and Cresti, M. (1995)
826 Presumed phylogenetic basis of the correlation of pectin deposition pattern in pollen tube
827 walls and the stylar structure of angiosperms. *Proc.Kon.Ned.Akad.v.Wetensch.* **98**, 39-
828 44.
829
- 83049 Akimov, S.S., Krylov, D., Fleischman, L.F. and Belkin, A.M. (2000) Tissue
831 transglutaminase is an integrin-binding adhesion coreceptor for fibronectin. *J. Cell Biol.*
832 **148**, 825-838.
833
- 83450 Gaudry, C.A., Verderio, E., Aeschlimann, D., Cox, A., Smith, C. and Griffin, M. (1999)
835 Cell surface localisation of tissue transglutaminase is dependent on a fibronectin-binding
836 site in its N-terminal beta-sandwich domain. *J. Biol. Chem.* **274**, 30707-30714.
837
838
839
840
841

71

72

18

73
74

Table 1. Evaluation of the effect of His₆-Xpr-GFP on pollen germination.

His₆-Xpr-GFP and control mouse IgG were added at different concentrations into the medium containing hydrated pollen at time zero and left up to 120 min germination. The data are expressed as percentage of germination (mean $\pm$ SD). P values were obtained by paired *t*-Student test between control and His₆-Xpr-GFP-treated pollen of 10 samples of three independent experiments. Significantly different at $p < 0.05$ ^(a).

848
849
850

Table 1. The effect of His₆-Xpr-GFP and IgG on pollen germination		
$\mu\text{g ml}^{-1}$	% Germination	
	His ₆ -Xpr-GFP	IgG
0.10	68 $\pm$ 6	67 $\pm$ 1
0.25	69 $\pm$ 4	72 $\pm$ 3
0.50	82 $\pm$ 5	77 $\pm$ 1
1.00	88 $\pm$ 3 ^a	78 $\pm$ 5
5.00	88 $\pm$ 2 ^a	76 $\pm$ 2
25.00	80 $\pm$ 2 ^a	70 $\pm$ 2
50.00	82 $\pm$ 2	76 $\pm$ 2

THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20100291

75
76

77
78

LEGENDS

Figure 1. Immuno-localisation of pollen TGase during germination.

(A) Pollen was germinated for 2 h and fixed, permeabilised and probed with monoclonal antibodies directed to TGase 2 (ID10) (a-c), α -tubulin (e) and α -tyrosinated tubulin (f). Negative control: non-specific mouse IgG (d). Magnifications of the distal part (b), the neck region and the proximal parts (c) of germinated pollen stained for TGase. Scale bar: 8 μ m.

(B) Laser scanning confocal microscopy of germinated pollen tubes treated as in (A), and probed with secondary FITC-conjugated antibody. TGase staining along the tube and at the apex (a) and visualization of the same sample in phase contrast (b). Staining with non-specific mouse IgG followed by secondary antibody as 2 μ m-single section (ct) or overlaid sections (d); arrows indicate the pollen tubes, whose autofluorescence was minimized. Scale bar: 8 μ m.

(C) TGase-active fractions (50 μ g protein) of germinated pollen extract (lane CE) germination medium (lane GM) and cell wall extract (lane CW) were analysed by western blotting with the monoclonal anti-TGase antibody ID10. The partially-purified TGase from pollen supernatant (isolated by hydrophobic ion chromatography and eluted with EGTA) was evaluated both by coomassie gel staining (lane C) and by western blotting with anti-TGase antibody ID10 (lane WB). "tTGase" in the first lane indicates the commercial gpl TGase used as a positive control.

(D) The cell-wall of non digested and non membrane-permeabilised 120 min-germinated pollen was treated with 2 M NaCl, 1% (w/v) SDS or 1% (w/v) DTT, fixed and then stained with anti-TGase antibody ID10 (a-b). In order to prove that the staining was not affected by fixation, ID10 was incubated for the last 30 min of germination in the medium prior to fixation (c). The arrows point the ID10 immuno-labelling. In some cases, the cell wall was further digested and the residual labeling was localized around the grain (arrows) (d). Scale bar: 8 μ m.

(E) Immuno-gold labeling of apple pollen tubes conducted with anti-TGase ID10. Labeling was mainly found in association with the amorphous layer of cell wall (arrows) in the flanks of pollen tubes. Scale bar: 200 nm.

Figure 2. Inhibition of TGase activity affects pollen tube growth.

(A) Effect of anti-TGase ID10 on pollen tube morphology, germination and extracellular TGase activity. Pollen was incubated at time zero with ID10 or control antibodies against horse albumin, in microtiter plates, and allowed to germinate up to 120 min. Pollen tube morphology was observed by light microscopy (upper panels) in the presence of control anti-horse albumin (left) and increasing concentrations of ID10 (centre and right images). Pollen tube morphological malformations and diffuse bursting are indicated by arrows. Scale bar: 24 μ m. Bottom plot: lines show the percentage (mean $\pm$ SD; $n = 10$) of germination with ID10 (black squares) or control antibody (white squares); microplate assay of TGase activity of germinating pollen (bars) treated with ID10 (crossed lines) or anti-horse albumin (vertical lines) and expressed as units (U) mg^{-1} pollen. The correlation analysis between pollen germination and TGase activity, when inhibited by ID10, gave a Pearson coefficient of $R = 0.92$ ($p = 0.03$).

(B) Effect of two irreversible inhibitors of TGase on pollen tube morphology, germination and extracellular TGase activity. Pollen was germinated, as in (A), and, at time zero, incubated with 50 μ M 283 and 281 or control MES buffer (0). Pollen tube morphological malformations are indicated by arrows (upper panels). Scale bar: 24 μ m.

79
80

20

81
82

Bottom plot: lines show the percentage of germination (mean $\pm$ SD; $n = 10$) in the presence of inhibitors 283 (white squares) and 281 (grey squares). TGase activity was reported, as in (A), in the presence of 283 (bars with horizontal lines) or 281 (bars with oblique lines). The correlation analysis between pollen germination and TGase activity when inhibited by 281 and 283 respectively, gave a Pearson coefficient of $R = 0.94$ ($p < 0.0001$) and 0.92 ($p < 0.0005$).

Figure 3. Extracellular TGase cross-linking during pollen germination.

(A) Laser scanning confocal microscopy at 120 min-germination of undigested and non-permeabilised pollen, His₆-Xpr-GFP was added for the last 30 min of germination in the medium prior to fixation. Superimposition of the GFP labeling on the phase contrast image (upper panel); GFP labeling (bottom panel). Control: non-specific mouse IgG followed by FITC-antibody. Arrows indicate His₆-Xpr-GFP fluorescence. Scale bar: 8 μ m.

(B) Laser confocal microscopy of pollen, incubated and treated as in (A), but in the presence of either the anti-TGase ID10 or 281/283 inhibitors for the final 30 min of germination. Scale bar: 8 μ m.

(C) Pollen was germinated from time zero in the presence of 5 μ g ml⁻¹ of His₆-Xpr-GFP (black bars) or mouse IgG (grey bars) and the tube length was evaluated via a digital camera at 30, 60, 90 and 120 min and reported as relative frequency (mean $\pm$ SD of 100-1000 μ m tube length). Square brackets report the lowest and the highest number of measured pollen tubes for each optical field.

(D) Microplate assay of extracellular TGase cross-linking activity in the presence of increasing concentrations of the anti-TGase antibody ID10 and the inhibitors 283 and 281 supplied to pollen for the last 30 min of germination, as indicated above in (A). The fluorescence of His₆-Xpr-GFP cross-linked by the pollen extracellular TGase to immobilised DMC was measured and compared to the values obtained by standard free gpl TGase (control, oblique line bar). Relative fluorescence units (RFU) of treated samples were reported as percentage (mean $\pm$ SD) of the control normalized to 100% ($12,580 \pm 1,133$ RFU).

(E) TGase-mediated crosslinking products immuno-localized by the 81D4 (monoclonal antibody anti-N^ε(γ -glutamyl)-lysine) on 120 min-germinated unfixed pollen (neither digested nor permeabilized). Arrows indicate the cross-link immuno-detection. Scale bar: 8 μ m.

Figure 4. Co-association of protein cross-linking and extracellular TGase on the surface of 120 min-germinated pollen.

(A) Germinating pollen tubes were incubated *in vivo* for the last 30 min of germination with His₆-Xpr-GFP prior to fix and, omitting cell wall digestion and permeabilisation, incubated with 81D4 (monoclonal antibody anti-N^ε(γ -glutamyl)-lysine and -PAs), followed by TRITC-conjugated secondary antibody. In the different panels, the positive staining indicates the localisation of TGase-mediated cross-linking products (His₆-Xpr-GFP incorporation) (upper), immuno-staining with 81D4 (centre), and superimposition of the His₆-Xpr-GFP and 81D4/TRITC immuno-labeled pollen tubes (merge bottom panel). Scale bar: 8 μ m.

(B) Germinating pollen tubes were treated as in A but stained with anti-TGase antibody ID10. Localization of TGase mediated cross-linking products by His₆-Xpr-GFP (upper panel), immuno-staining with ID10 (centre panel), and superimposition of His₆-Xpr-GFP and ID10/TRITC immuno-labeling (merge bottom panel). Scale bar: 8 μ m.

83
84

21

85
86

Figure 5. Immuno-localisation of pollen TGase during germination *in planta*.

Following the flower pollination, the pollen tubes were left to grow for 48 h into the stigma/style. The latter was fixed and anti-TGase ID10-probed (pollen tubes delineated as interrupted lines). Pollen tubes growing inside the style are shown in the upper panel; magnification of the clusters around the pollen tubes is reported in lower panel. Scale bar: 8 μ m.

Figure 6. Model of secreted TGase regulating *in vitro* pollen tube growth.

In the pollen tube apical zone distally growing (grey arrows), TGase would accumulate: i) intracellularly at the tip as such or in vesicles, ii) outside the membrane (M) in the thin wall (under construction) and iii) released outside (pink arrows). In the apical tube wall, TGase cross-links tube wall (Tw) proteins without forming aggregates. Once released into the Ca^{2+} rich-germination medium, TGase may cross-link proteins or incorporate PAs into pollen proteins, also secreted. As the pollen tip is continuously elongating during pollen germination, TGase might remain behind along the Tw and be embedded, under the effect of local micro-environmental conditions, into the same accumulation sites of the protein cross-links catalysed by its enzymic activity.

LEGENDS FOR SUPPLEMENTAL FIGURES

Figure S1. Western blot of an active pollen fraction partially purified from hydrated pollen by a hydrophobic interaction chromatography (HIC), set up to purify Ca^{2+} -dependent TGases, assayed at 30, 20 and 10 μ g of protein and both not purified (NP) and purified (P) gpl TGase, comparatively probed with anti-TGase antibodies ID10 and AbIII.

Figure S2. Percentage of residual TGase activity in the pollen SN after immuno-precipitation obtained with increasing amounts of anti-TGase antibody ID10. Proteins were extracted from hydrated pollen grains.

Figure S3. (A) Western blots of 1 μ g and 0.1 μ g human recombinant TGase 1 (1), TGase 2 (2) and TGase 3 (3) probed with the anti-TGase antibody ID10; **(B)** Western blots of 0.1 μ g human recombinant TGase 1 (1), TGase 2 (2) and TGase 3 (3) probed with anti-TGase antibodies ID10, CUB7402, anti-TGase 1 and anti-TGase 3. These results show the specificity of ID10 toward TGase 2.

87
88

22

89
90

988
989

990 **Figure 1.** Immuno-localisation of pollen TGase during germination.

91
92

23

93
94

991
992
993 **Figure 2.** Inhibition of TGase activity affects pollen tube growth.

95
96

24

97
98

994
995

996 **Figure 3.** Extracellular TGase cross-linking during pollen germination.

99
100

25

101
102

997
998
999 **Figure 4.** Co-association of protein cross-linking and extracellular TGase on the surface of
1000 120 min-germinated pollen. Scale bar: 8 μ m.

103
104

105
106

1001
1002

1003 **Figure 5.** Immuno-localisation of pollen TGase during germination *in planta*.

107
108

109
110

1004
1005
1006
1007
1008

Figure 6. Model of externalised TGase regulating *in vitro* pollen tube growth.

THIS IS NOT THE VERSION OF RECORD - see doi:10.1042/BJ20100291

111
112