Accepted Manuscript

Prognostic values of α_2 -macroglobulin, fibrinogen and albumin in regards to mortality and frailty in old rats

Gilles Mayot, Karine Vidal, Jean-François Martin, Denis Breuillé, Stephanie Blum, Christiane Obled, Isabelle Papet

PII: S0531-5565(07)00021-6
DOI: [10.1016/j.exger.2007.01.003](https://doi.org/10.1016/j.exger.2007.01.003)
Reference: EXG 8299

To appear in: *Experimental Gerontology*

Received Date: 16 June 2006
Revised Date: 9 January 2007
Accepted Date: 16 January 2007

Please cite this article as: Mayot, G., Vidal, K., Martin, J-F., Breuillé, D., Blum, S., Obled, C., Papet, I., Prognostic values of α_2 -macroglobulin, fibrinogen and albumin in regards to mortality and frailty in old rats, *Experimental Gerontology* (2007), doi: [10.1016/j.exger.2007.01.003](https://doi.org/10.1016/j.exger.2007.01.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Prognostic values of α_2 -macroglobulin, fibrinogen and albumin in regards to mortality and frailty in old rats

Gilles Mayot ¹, Karine Vidal ², Jean-François Martin ¹, Denis Breuillé ², Stephanie Blum ²,
Christiane Obled ¹, Isabelle Papet ^{1*}

¹ *INRA, Centre Clermont-Ferrand – Theix, UMR 1019 Unité de Nutrition Humaine, F-63122 Saint-Genès-Champanelle, France*

² *Nutrition and Health, Nestlé Research Center, 1000 Lausanne 26, Switzerland*

* To whom correspondence should be addressed: Dr. Isabelle Papet, Unité de Nutrition Humaine, INRA Theix, 63 122 Saint-Genès-Champanelle, France.

Telephone: 33 4 73 62 42 01

Fax: 33 4 73 62 47 55

Email: papet@clermont.inra.fr

Running title: Mortality and frailty in inflamed old rat

Keywords: Frailty; Inflammation; Mortality; α_2 -macroglobulin; Fibrinogen; Albumin.

Abbreviations used: APP: acute phase protein, CI: confidence interval, CRP: C-reactive protein, IL-6: interleukin-6, LBP: Lipopolysaccharide binding protein, TNF- α : tumor necrosis factor-alpha, sTNFR-1: soluble TNF- α receptor-1.

Abstract

The study aimed to determine if acute phase proteins (APP) are markers of frailty in old rats. We evaluated in male Wistar rats at 96 weeks of age (n=72) whether single measurements of α_2 -macroglobulin, fibrinogen and albumin are predictive of mortality, body weight loss and inflammatory status during a 10-week follow-up period. Rats were clustered depending on levels of these APP at baseline. Rats with extremely high levels of α_2 -macroglobulin or fibrinogen (upper quartiles), or extremely low level of albumin (lower quartile), had an 11.6, 8.1 and 5.3-fold higher risk of mortality, respectively, than other rats. Body weight loss was negatively correlated with α_2 -macroglobulin, a trend was observed with fibrinogen ($P = 0.08$) but not with albumin. Rats with fibrinogen levels > 4.0 g/L or α_2 -macroglobulin levels > 91 mg/L (respective top halves) at 96 weeks of age had higher levels of α_2 -macroglobulin and fibrinogen and lower levels of albumin throughout the follow-up period and higher levels of sTNFR-1 and lipopolysaccharide-binding protein at 106 weeks of age. Highest levels of α_2 -macroglobulin, fibrinogen and lowest albumin were predictive of mortality, whereas moderate levels of α_2 -macroglobulin and fibrinogen were, according to body weight loss and inflammatory status, markers of frailty in old rats.

1. Introduction

Frailty syndrome is a physiological state of susceptibility that places older individuals at high risk for adverse outcomes such as falls, disability, morbidity and institutionalization (Dayhoff et al., 1998; Fried et al., 2001; Ferrucci et al., 2002a; Walston, 2004). Frailty ultimately leads to death. Frailty consists of multi-system decline and is considered to be a consequence of changes in neuromuscular, endocrine and immune functions that occur as people progress in age (Ferrucci et al., 2002a; Walston, 2004). Frailty can and does result in several “vicious loops”: neuromuscular impairment induces increased rate of falls and fractures in old age; the act of falling causes itself hospitalisation and immobilisation, which

aggravates sarcopenia; sarcopenia leads to deficit in continuous neuromuscular training which brings about neuromuscular impairment (Muhlberg and Sieber, 2004; Walston, 2004). Moreover, frailty has nutritional and metabolic implications which have themselves deleterious implications (Chevalier et al., 2003; Muhlberg and Sieber, 2004; Puts et al., 2005).

Early screening of frailty in the elderly population is essential to reduce the adverse outcomes described above. Despite the growing awareness of this problem over the past two decades, there is no clear guidelines for identifying and describing older adults as frail (Markle-Reid and Browne, 2003; Walston, 2004). Screening for frailty syndrome in humans is often based on the measurement of parameters related to mobility and motor performance. These include weakness (grip strength), weight loss, fatigue, slow walking speed and low levels of physical activity (Fried et al., 2001). Subjects meeting three or more of these criteria are categorized as frail (Fried et al., 2001). Several researchers have sought to find sensitive biomarkers of frailty (Ferrucci et al., 2002a). Among the candidate biomarkers are compounds associated with the inflammatory process (Ferrucci et al., 2002a; De Martinis et al., 2006). Frail subjects have evidence of increased inflammation (Leng et al., 2002; Puts et al., 2005). Increased levels of certain acute phase reactants, including C-reactive protein (CRP), clotting factors such as fibrinogen and factor VIII, and decreased level of albumin, were associated with increasing frailty level in elderly (Walston et al., 2002). In accordance, the levels of inflammatory markers, such as interleukin-6 (IL-6) and CRP, increase with ageing, and that elevated levels are associated with disability and mortality (Harris et al., 1999; Ferrucci et al., 2002b; Cohen et al., 2003; Bautmans et al., 2005).

In spite of the fact that rat is a common model for human nutritional and metabolic studies, to our knowledge, no investigation has assessed frailty in old rats. Frail rats are expected to exhibit the human frailty phenotype, including vulnerability to mortality, body weight loss and inflammation. Physical performance which is classically assessed in human clinical settings cannot be easily evaluated in old rats. So, there is a need to identify plasma

1 biomarkers of rat frailty. Measurements of inflammatory markers, that correlate with frailty,
2 has recently been proposed for identification of frail animals (Walston et al., 2006). In the
3 present study, we investigated whether acute phase proteins (APP) are potential biomarkers of
4 frailty in old rats.

5 The APP α_1 -acid glycoprotein, α_2 -macroglobulin and fibrinogen are known to
6 increase with inflammation in rats (Fuller, G. M., 1993; Roberts, R. C., 1993; Breuillé et al.,
7 1998; Ruot et al., 2002). CRP is commonly used as a marker of inflammation in human
8 clinical trials and research settings (Kushner, I. and Maciewicz, A., 1993). But CRP is a poor
9 marker of inflammation compared to fibrinogen in the rat (Giffen et al., 2003). By contrast,
10 α_2 -macroglobulin in rats is an APP characterized by a kinetic relatively similar to human CRP
11 kinetic since both of them peak very quickly after inflammatory stress and thus α_2 -
12 macroglobulin is a good marker of inflammation in the rat. The APP albumin is known to
13 decline with inflammation in rats (Aldred, A. R. and Schreiber, G., 1993; Breuillé et al., 1998;
14 Ruot et al., 2002). We have recently shown that α_2 -macroglobulin and fibrinogen
15 concentrations were higher and albumin concentration was lower in old rats (95 week-old)
16 compared to adults (35 week-old), whereas plasma α_1 -acid glycoprotein concentration was
17 similar in both groups (Papet et al., 2003). In addition, greater variability was observed in the
18 APP in the old rats compared to their adult rats counterparts. Taken together, α_2 -
19 macroglobulin, fibrinogen and albumin are potential candidates for assessing frailty in old
20 rats.

21 The purpose of the study was to determine if the APP, α_2 -macroglobulin, fibrinogen
22 and albumin, are biomarkers of frailty in old rats. The parameters assessed included mortality,
23 body weight loss and persistent inflammatory status evaluated in a 10-week longitudinal study
24 on a cohort of 96-week old rats. Inflammatory status was assessed by measuring the APP
25 throughout the follow-up period and IL-6, tumor necrosis factor- α (TNF- α), soluble TNF- α
26 receptor-1 (sTNFR-1) and lipopolysaccharide binding protein (LBP) at the end of the study.

1

2 **2. Materials and methods**

3 *2.1. Animals*

4 Male Wistar rats were produced and bred in our conventional (not specific-pathogen-free)
5 animal facility (Unité Expérimentale de Nutrition Comparée, INRA, Saint-Genès-
6 Champanelle, France). They were maintained in collective cages (3 to 4 rats per cage) under
7 controlled conditions (temperature 21°C, relative humidity 55 %, 12-h light/dark cycles with
8 free access to water and standard pelleted food (A04 from SAFE (Scientific Animal Food and
9 Engineering), Villemoisson-sur-orge, France). The composition of the diet was 16 % protein,
10 3% fat, 60 % carbohydrates, 12 % water, fibers, vitamins and minerals. The study was
11 performed according to the current legislation on animal experiments in France.

12 *2.2. Experimental design*

13 The experimental cohort consisted of 119 rats at the age of 13 weeks, born over a two
14 week period. Blood samples were withdrawn from the lateral vein of 32 rats at 35 week-old in
15 order to estimate the adult cohort values for plasma levels of the three APP: α_2 -
16 macroglobulin, fibrinogen and albumin. The longitudinal study was initiated when the rats
17 were 96 week-old (n = 72) and lasted for an additional 10 weeks. Body weight was measured
18 at weeks 96 and 106 and sample collection from a lateral tail vein was performed at weeks 96,
19 100, 104 and 106. Mortality was recorded weekly. Samples collected throughout the study
20 were assayed for α_2 -macroglobulin, fibrinogen and albumin. Additionally, samples collected
21 at 106 weeks of age were also assayed for IL-6, TNF- α , sTNFR-1 and LBP. All samples were
22 stored at -80°C until subjected to analyses.

23 *2.3. Acute-phase protein concentrations*

24 Plasma fibrinogen level was measured by turbidimetry (Macart et al., 1989) on a
25 Cobas Mira analyzer (ABX Diagnostics, Montpellier, France). Concentration is expressed as
26 g human equivalent/L since human fibrinogen was used as reference (Ingen, Rungis, France).

Plasma levels of α_2 -macroglobulin and albumin were measured by single radial immunodiffusion (Breuillé et al., 1998). Serum concentration of LBP was measured by Enzyme Linked ImmunoSorbent Assay (ELISA) that detects free LBP (Biometec, Greifswald, Germany); its detection limit was 5 ng/mL.

2.4. Serum levels of cytokines and cytokine receptor

Serum levels of IL-6, TNF- α and sTNFR-1 were quantified with specific ELISA kits (R6000B, RTA00 and MRT10, respectively, R&D Systems, Abingdon, UK). All quantifications were realized according to the manufacturer. Detection limits were 21, 12 and 16 pg/mL for IL-6, TNF- α and sTNFR-1, respectively. All samples and standards were measured in duplicate and the means were used in statistical analyses.

2.5. Statistical analysis

All values are given as means $\pm$ SD. Statistical analyses were performed using StatView for Windows, version 5 software (SAS Institute, Cary, NC). Values of $P < 0.05$ are considered significant for all analyses.

The effect of age was analyzed using one-way ANOVA and Fisher's PLSD test and Student's t test for unpaired data was used to compare alive and dead old rats. Linear regression was used to determine the relationship among APP themselves and between the percentage of body weight change and each APP.

Several statistical tests were used to analyse mortality data obtained for all rats ($n = 72$). Rats were divided into quartile categories of either α_2 -macroglobulin, fibrinogen or albumin plasma levels at the age of 96 weeks. Survival curves were generated by the Kaplan-Meier method and compared by the log-rank test in order to analyze the impact of APP plasma levels measured in 96 week-old rats on mortality across the 10 weeks of follow-up. The relative risk of death was calculated using the Cox proportional hazards model for categorical variables. Univariate Cox and multivariate Cox regressions for continuous variables were used to analyze how APP were predictive of mortality.

Changes in APP during the follow-up period and the values of sTNFR-1 and LBP at the age of 106 weeks were analyzed on rats which were still alive at 106 weeks ($n = 48$). For each APP, data from alive rats from quartiles 1 and 2 (bottom half) were compared to those of alive rats from quartiles 3 and 4 (top half). Two-way ANOVA for repeated measurements with group as the between-rat factor and time as the within-rat factor was performed to analyze APP changes. When significant effects were observed for group and time, Fisher's PLSD test and Student's t test for unpaired data were performed as post ANOVA tests. Student's t test for unpaired data was also used to analyze the group effect for sTNFR-1 and LBP at the age of 106 weeks.

3. Results

3.1. Rat characteristics

The three APP, α_2 -macroglobulin, fibrinogen and albumin, and body weight were significantly affected by age (Table 1). α_2 -macroglobulin and fibrinogen plasma levels were higher and albumin plasma level lower at the age of 96 and 106 weeks than at the age of 35 weeks. In 96 week-old rats, α_2 -macroglobulin and fibrinogen were highly positively correlated to each other ($r = 0.73$, $P < 0.001$, $n = 72$) and each of them was negatively correlated with albumin ($r = -0.71$ and -0.67 , respectively, $P < 0.001$, $n = 72$). Mean body weight was higher in 96 week-old rats than in 35 week-old rats. At 96 weeks of age, rat body weight was negatively correlated with α_2 -macroglobulin and fibrinogen ($r = -0.46$ and -0.29 , respectively, $P < 0.05$, $n = 72$) and was positively correlated with albumin ($r = 0.46$, $P < 0.001$, $n = 72$).

3.2. APP measurements at 96 weeks of age are predictive of the mortality

Between 96 and 106 weeks of age, 24 rats died which represented a 33 % of mortality rate. The rats that died prior to 106 weeks had higher α_2 -macroglobulin (851 %) and

fibrinogen (47 %) and lower albumin (24 %) plasma levels than alive rats (Table 2). In addition, the mean body weight of deceased rats was 12 % lower than their alive counterparts.

Kaplan-Meier analyses show that each APP measured at 96 weeks of age was associated with 10-week mortality (Figure 1). α_2 -macroglobulin plasma level in the top quartile (≥ 506 mg/L) was the strongest predictor of mortality (relative risk = 11.6; 95 % confidence interval (CI) = 4.8 to 27.7; $P < 0.0001$). Rats belonging to the top fibrinogen quartile had an 8.1-fold higher risk of mortality (95 % CI = 3.5 to 18.8; $P < 0.0001$) than those of the other quartiles (< 5.4 g/L). The mortality risk of rats in the bottom albumin quartile (≤ 10 g/L) relative to the three other quartiles was 5.3 (95 % CI = 2.3 to 11.9; $P < 0.0001$). Univariate Cox analyses show that each of the three APP measured at the age of 96 weeks was predictive of mortality. The multivariable Cox regression model using forward stepwise selection method conducted with the three APP show that α_2 -macroglobulin alone was sufficient to explain rat mortality during the 10-week follow-up.

3.3. APP levels at 96 weeks of age and body weight change

Percentage of body weight loss throughout 10-week follow-up in rats alive at 106 weeks ($n = 48$) was negatively correlated with α_2 -macroglobulin levels measured at 96 weeks old ($P < 0.01$) (Table 3). A similar correlation with fibrinogen level was observed, but did not attain statistical significance ($P = 0.08$). However, no associations between body weight changes and albumin levels were observed.

3.4. APP levels at 96 weeks of age and inflammatory status throughout 10-week follow-up

To determine if single measurements of APP were predictive of the consecutive inflammatory status, changes in the APP throughout the follow-up period and values of other inflammatory markers at 106 weeks of age were assessed in surviving rats. This analysis was performed on the top half and the bottom half groups constituted on the basis of each plasma APP measured in all rats at 96 weeks of age (Table 4). Rats with α_2 -macroglobulin levels $>$

91 mg/L at 96 weeks of age exhibited higher levels of plasma α_2 -macroglobulin (~ 450 %) and fibrinogen (~ 34 %) and lower levels of plasma albumin (~ 20 %) at all tested times during the follow-up period than the other rats, excepted for albumin at the age of 96 weeks. Interestingly serum sTNFR-1, detected in all rats at the age of 106 weeks, had a 66 % higher level in rats having the highest α_2 -macroglobulin level. No difference was observed in levels of serum LBP. No detectable level of serum IL-6 and TNF- α was observed. Rats with fibrinogen levels > 4 g/L at 96 weeks of age exhibited higher levels of plasma α_2 -macroglobulin (~ 280 %) and fibrinogen (~ 47 %) and lower levels of plasma albumin (~ 17 %) at all tested times during the follow-up than other rats, excepted for albumin at the age of 96 weeks and for α_2 -macroglobulin at the age of 100 weeks. The levels of both serum sTNFR-1 and LBP at the age of 106 weeks were higher in rats having the highest fibrinogen level (70 and 130 %, respectively). Rats with albumin levels lower than 12 g/L at 96 weeks of age exhibited similar values for the three APP throughout 10-week period (except albumin at the age of 96 weeks) and similar levels of sTNFR-1 and LBP at the age of 106 weeks than other rats.

4. Discussion

In this paper, we report that highest plasma levels of α_2 -macroglobulin, fibrinogen and lowest albumin are predictive of mortality. According to body weight loss and persistent inflammatory status, moderate levels of α_2 -macroglobulin and fibrinogen are biomarkers of frailty in old rats, with α_2 -macroglobulin being slightly more powerful than fibrinogen.

Regardless of the type of statistical analysis performed, α_2 -macroglobulin, fibrinogen and albumin measured in 96 week-old rats were associated with 10-week mortality. Rats that died during the 10-week follow-up had higher concentrations of α_2 -macroglobulin and fibrinogen, and lower concentration of albumin (Student's *t* test). Kaplan and univariate Cox

analyses were significant for each APP. The multivariate Cox regression model showed that α_2 -macroglobulin alone was sufficient, likely due to the fact that the three APP are correlated to each other. This is in agreement with the highest log rank test for α_2 -macroglobulin alone.

The mortality was significantly enhanced in rats exhibiting extremely high concentrations of α_2 -macroglobulin (506 to 3390 mg/L) and fibrinogen (5.4 to 10.6 g/L), and extremely low concentrations of albumin (10 to 4 g/L). These values correspond to those observed during acute inflammation, such as sepsis (Breuillé et al., 1998; Breuillé et al., 1999; Ruot et al., 2002). In addition, these rats had the lowest body weight. This inflammation could be due to aging *per se*; however, age-associated inflammation in humans is a much lower grade inflammation (Bruunsgaard and Pedersen, 2003). These rats may have been infected by chronic pathogen since they were bred in none specific-pathogen-free animal facility. Moreover, these rats may have had some acute or chronic illness that activated inflammatory pathways such as chronic nephropathy, cardiomyopathy, and neoplasia which are the major age-related pathologies in rats (Maeda et al., 1985). Unfortunately, no data were collected on any illnesses and death causes in the present study. Even if the etiology of this inflammation is not known, it definitively enhanced mortality rate. It has been shown in elderly that an increase in CRP level, corresponding to either the third tertile or the fourth quartile in most human studies, was associated with increased risk of mortality (Harris et al., 1999; Reuben et al., 2002; Tice et al., 2003; Kritchevsky et al., 2005). In a recent meta-analysis, the age-adjusted relative risk per 1 g/L increase in usual fibrinogen level was 2.06-2.76 for cardiovascular disease mortality and 2.03 for nonvascular mortality in healthy middle-aged adults (Danesh et al., 2005). In addition, lowest albumin levels were associated with an increased mortality risk in community-dwelling elderly subjects (Corti et al., 1994; Sahyoun et al., 1996; Reuben et al., 2002). Altogether, our results obtained in rats are in line with data known in humans and give a prognostic value of some acute phase proteins in regards to mortality.

1 Body weight loss is both a characteristic and an outcome of frailty in humans (Fried et
 2 al., 2001). In the survivors of our cohort of rats, among the three APP tested, only α_2 -
 3 macroglobulin was significantly associated with 10-week body weight loss. A similar trend
 4 was observed for fibrinogen, but not with albumin. In the elderly, the influence of
 5 inflammatory status on body weight loss has not been directly assessed, but acute
 6 inflammation plays a major role in the pathophysiological of cachexia (Morley et al., 2006).
 7 Albumin decreases in inflammatory and malnutrition states (Don and Kaysen, 2004).
 8 Considering that old rats, which died were the ones that lost the most weight, it is surprising
 9 that low level of albumin was only associated with high mortality but not with body weight
 10 loss. This questions the validity of albumin as a nutritional marker in old rats. Unfortunately,
 11 measurement of food intake or other nutritional markers were not assessed in this study.

12 Single measurements of α_2 -macroglobulin and fibrinogen, but not albumin, were
 13 predictive of consecutive inflammatory status in survivors, assessed not only by these APP
 14 but also with additional inflammatory markers, including LBP and sTNFR-1. Plasma level of
 15 LBP, which is recognized as an APP (Tobias et al., 1986), rises dramatically after
 16 inflammatory challenges (infectious or not) (Gallay et al., 1994; Myc et al., 1997; Fenton and
 17 Golenbock, 1998). It was higher in old rats characterized as inflamed from repartitions based
 18 on fibrinogen. It is well-known that TNF- α plays an important pathophysiological role in a
 19 wide range of chronic disorders, but sTNFRs are stronger markers of severity and clinical
 20 outcome than TNF- α itself (Aderka, 1996) and are essential for evaluation of the TNF system
 21 (Ferrari et al., 1995). Unfortunately, the pro-inflammatory cytokines TNF- α and IL-6 were
 22 undetectable in the sera of our old rats, probably due to the low sensitivity of the ELISA.
 23 Interestingly, sTNFR-1 levels were consistently higher in rats characterized as inflamed based
 24 on α_2 -macroglobulin or fibrinogen levels. Inflammatory status in old rats, assessed with α_2 -
 25 macroglobulin or fibrinogen was not transient but persistent for at least 10 weeks.

Accordingly, it has been shown in a human study that CRP values appear relatively tightly regulated over a 6-month period (Macy et al., 1997).

In the present study, moderate α_2 -macroglobulin and fibrinogen levels in old rats was clearly associated with worsening of body weight loss and persistent inflammation, which are strong criteria of frailty in humans. The etiology of this moderate inflammation is not known, it could be either associated to age *per se* or to subclinical diseases (Grimble, 2003). In this regard, a recent paper shows an association between chronic asymptomatic infection, inflammation and frailty in elderly (Schmaltz et al., 2005). Even if rats exhibiting moderate inflammation were not at high risk of mortality during the 10-week follow-up, it is not excluded that they would be in a longer follow-up study. Hence, moderate α_2 -macroglobulin and fibrinogen levels can be considered as markers of frailty in old rats.

Taken together, the results presented here demonstrate that the highest levels of α_2 -macroglobulin and fibrinogen and the lowest levels of albumin are predictive of mortality, whereas, moderate levels of α_2 -macroglobulin and fibrinogen are biomarkers of frailty in old rat (as evaluated by body weight loss and persistent inflammatory status). These data are consistent with a human study showing that elevated levels of CRP (> 5.5 mg/L) or fibrinogen (> 3.9 g/L) are associated with higher risks for prevalent frailty, the relation between CRP levels and frailty being higher than that between fibrinogen levels and frailty (Walston et al., 2002). Thus, in old rats α_2 -macroglobulin, as CRP in humans, is a more powerful biomarker of frailty than fibrinogen.

References

1. Aderka, D., 1996. The potential biological and clinical significance of the soluble tumor necrosis factor receptors. *Cytokine & Growth Factor Reviews* 7, 231-240.
2. Aldred, A.R., Schreiber, G., 1993. The negative acute phase proteins. In: Mackiewicz,

- 1 A., Kushner, I., Baumann, H. (Eds.), Acute Phase Proteins. Molecular Biology,
2 Biochemistry, and Clinical Applications. CRC Press, Inc., Boca Raton, pp. 21-37.
- 3 3. Bautmans, I., Njemini, R., Lambert, M., Demanet, C., Mets, T., 2005. Circulating acute
4 phase mediators and skeletal muscle performance in hospitalized geriatric patients.
5 Journals of Gerontology Series aBiological Sciences and Medical Sciences 60, 361-
6 367.
- 7 4. Breuillé, D., Arnal, M., Rambourdin, F., Bayle, G., Levieux, D., Obled, C., 1998.
8 Sustained modifications of protein metabolism in various tissues in a rat model of long-
9 lasting sepsis. Clinical Science 94, 413-423.
- 10 5. Breuillé, D., Voisin, L., Contrepolis, M., Arnal, M., Obled, C., 1999. A sustained rat
11 model for studying long-lasting catabolic state of sepsis. Infection and Immunity 67,
12 1079-1085.
- 13 6. Bruunsgaard, H., Pedersen, B.K., 2003. Age-related inflammatory cytokines and
14 disease. Immunology and Allergy Clinics of North America 23, 15-39.
- 15 7. Chevalier, S., Gougeon, R., Nayar, K., Morais, J.A., 2003. Frailty amplifies the effects
16 of aging on protein metabolism: role of protein intake. American Journal of Clinical
17 Nutrition 78, 422-429.
- 18 8. Cohen, H.J., Harris, T., Pieper, C.F., 2003. Coagulation and activation of inflammatory
19 pathways in the development of functional decline and mortality in the elderly.
20 American Journal of Medicine 114, 180-187.
- 21 9. Corti, M.C., Guralnik, J.M., Salive, M.E., Sorkin, J.D., 1994. Serum albumin level and
22 physical disability as predictors of mortality in older persons. Jama-Journal of the
23 American Medical Association 272, 1036-1042.

- 1 10. Danesh, J., Lewington, S., Thompson, S.G., Lowe, G.D.O., Collins, R., 2005. Plasma
2 fibrinogen level and the risk of major cardiovascular diseases and nonvascular mortality:
3 an individual participant meta-analysis. *Jama-Journal of the American Medical*
4 *Association* 294, 1799-1809.
- 5 11. Dayhoff, N.E., Suhrheinrich, J., Wigglesworth, J., Topp, R., Moore, S., 1998. Balance
6 and muscle strength as predictors of frailty among older adults. *Journal of*
7 *Gerontological Nursing* 24, 18-27.
- 8 12. De Martinis, M., Franceschi, C., Monti, D., Ginaldi, L., 2006. Inflammation markers
9 predicting frailty and mortality in the elderly. *Experimental and Molecular Pathology*
10 80, 219-227.
- 11 13. Don, B.R., Kaysen, G., 2004. Serum albumin: relationship to inflammation and
12 nutrition. *Seminars in Dialysis* 17, 432-437.
- 13 14. Fenton, M.J., Golenbock, D.T., 1998. LPS-binding proteins and receptors. *Journal of*
14 *Leukocyte Biology* 64, 25-32.
- 15 15. Ferrari, R., Bachetti, T., Confortini, R., Opasich, C., Febo, O., Corti, A., Cassani, G.,
16 Visioli, O., 1995. Tumor necrosis factor soluble receptors in patients with various
17 degrees of congestive heart failure. *Circulation* 92, 1479-1486.
- 18 16. Ferrucci, L., Cavazzini, C., Corsi, A., Bartali, B., Russo, C.R., Lauretani, F., Ferrucci,
19 L., Cavazzini, C., Corsi, A.M., Bartali, B., Russo, C.R., Lauretani, F., Bandinelli, S.,
20 Bandinelli, S., Guralnik, J.M., 2002a. Biomarkers of frailty in older persons. *Journal of*
21 *Endocrinological Investigation* 25, 10-15.
- 22 17. Ferrucci, L., Penninx, B.W., Volpato, S., Harris, T.B., Bandeen-Roche, K., Balfour, J.,
23 Leveille, S.G., Fried, L.P., Md, J.M., 2002b. Change in muscle strength explains

- 1 accelerated decline of physical function in older women with high interleukin-6 serum
- 2 levels. *Journal of American Geriatrics Society* 50, 1947-1954.
- 3 18. Fried, L.P., Tangen, C.M., Walston, J., Newman, A.B., Hirsch, C., Gottdiener, J.,
- 4 Seeman, T., Tracy, R., Kop, W.J., Burke, G., McBurnie, M.A., 2001. Frailty in older
- 5 adults: evidence for a phenotype. *Journals of Gerontology Series A-Biological Sciences*
- 6 & *Medical Sciences* 56, M146-M156.
- 7 19. Fuller, G.M., 1993. Fibrinogen: a multifunctional acute phase protein. In: Mackiewicz,
- 8 A., Kushner, I., Baumann, H. (Eds.), *Acute Phase Proteins. Molecular Biology,*
- 9 *Biochemistry, and Clinical Applications.* CRC Press, Inc., Boca Raton, pp. 169-183.
- 10 20. Gallay, P., Barras, C., Tobias, P.S., Calandra, T., Glauser, M.P., Heumann, D., 1994.
- 11 Lipopolysaccharide (LPS)-binding protein in human serum determines the tumor-
- 12 Necrosis-Factor Response of Monocytes to LPS. *Journal of Infectious Diseases* 170,
- 13 1319-1322.
- 14 21. Giffen, P.S., Turton, J., Andrews, C.M., Barrett, P., Clarke, C.J., Fung, K.W., Munday,
- 15 M.R., Roman, I.F., Smyth, R., Walshe, K., York, M.J., 2003. Markers of experimental
- 16 acute inflammation in the Wistar Han rat with particular reference to haptoglobin and C-
- 17 reactive protein. *Archives of Toxicology* 77, 392-402.
- 18 22. Grimble, R.F., 2003. Inflammatory response in the elderly. *Current Opinion in Clinical*
- 19 *Nutrition and Metabolic Care* 6, 21-29.
- 20 23. Harris, T.B., Ferrucci, L., Tracy, R.P., Corti, M.C., Wacholder, S., Ettinger, W.H. Jr,
- 21 Heimovitz, H., Cohen, H.J., Wallace, R., 1999. Associations of elevated interleukin-6
- 22 and C-reactive protein levels with mortality in the elderly. *American Journal of*
- 23 *Medicine* 106, 506-512.

- 1 24. Kritchevsky, S.B., Cesari, M., Pahor, M., 2005. Inflammatory markers and
2 cardiovascular health in older adults. *Cardiovascular Research* 66, 265-275.
- 3 25. Kushner, I., Maciewicz, A., 1993. The acute phase response: an overview. In:
4 Maciewicz, A., Kushner, I., Baumann, H. (Eds.), *Acute Phase Proteins. Molecular*
5 *Biology, Biochemistry, and Clinical Applications*. CRC Press, London, pp. 3-19.
- 6 26. Leng, S., Chaves, P., Koenig, K., Walston, J., 2002. Serum interleukin-6 and
7 hemoglobin as physiological correlates in the geriatric syndrome of frailty: a pilot study.
8 *Journal of American Geriatrics Society* 50, 1268-1271.
- 9 27. Macart, M., Koffi, A., Henocque, G., Mathieu, J.F., Guilbaud, J.C., 1989. Optimized
10 microturbidimetric assay for fibrinogen. *Clinical Chemistry* 35, 211-214.
- 11 28. Macy, E.M., Hayes, T.E., Tracy, R.P., 1997. Variability in the measurement of C-
12 reactive protein in healthy subjects: implications for reference intervals and
13 epidemiological applications. *Clinical Chemistry* 43, 52-8.
- 14 29. Maeda, H., Gleiser, C.A., Masoro, E.J., Murata, I., McMahan, C.A., Yu, B.P., 1985.
15 Nutritional influences on aging of fisher 344 rats: II Pathology. *Journal of Gerontology*
16 40, 671-688.
- 17 30. Markle-Reid, M., Browne, G., 2003. Conceptualizations of frailty in relation to older
18 adults. *Journal of Advanced Nursing* 44, 58-68.
- 19 31. Morley, J.E., Thomas, D.R., Wilson, M.M., 2006. Cachexia: pathophysiology and
20 clinical relevance. *American Journal of Clinical Nutrition* 83, 735-743.
- 21 32. Muhlberg, W., Sieber, C., 2004. Sarcopenia and frailty in geriatric patients: implications
22 for training and prevention. *Zeitschrift Fur Gerontologie Und Geriatrie* 37, 2-8.

- 1 33. Myc, A., Buck, J., Gonin, J., Reynolds, B., Hammerling, U., Emanuel, D., 1997. The
2 level of lipopolysaccharide-binding protein is significantly increased in plasma in
3 patients with the systemic inflammatory response syndrome. *Clinical and Diagnostic*
4 *Laboratory Immunology* 4, 113-116.
- 5 34. Papet, I., Dardevet, D., Sornet, C., Bechereau, F., Prugnaud, J., Pouyet, C., Obled, C.,
6 2003. Acute phase protein levels and thymus, spleen and plasma protein synthesis rates
7 differ in adult and old rats. *Journal of Nutrition* 133, 215-219.
- 8 35. Puts, M.T.E., Visser, M., Twisk, J.W.R., Deeg, D.J.H., Lips, P., 2005. Endocrine and
9 inflammatory markers as predictors of frailty. *Clinical Endocrinology* 63, 403-411.
- 10 36. Reuben, D.B., Cheh, A.I., Harris, T.B., Ferrucci, L., Rowe, J.W., Tracy, R.P., Seeman,
11 T.E., 2002. Peripheral blood markers of inflammation predict mortality and functional
12 decline in high-functioning community-dwelling older persons. *Journal of American*
13 *Geriatrics Society* 50, 638-644.
- 14 37. Roberts, R.C., 1993. Rat alpha₂-macroglobulin and related alpha-macroglobulins in the
15 acute phase response. In: Mackiewicz, A., Kushner, I., Baumann, H. (Eds.), *Acute Phase*
16 *Proteins. Molecular Biology, Biochemistry, and Clinical Applications*. CRC Press, Inc.,
17 Boca Raton, pp. 223-238.
- 18 38. Ruot, B., Béchereau-Rambourdin, F., Bayle, G., Breuillé, D., Obled, C., 2002. The
19 response of liver albumin synthesis to infection in rats varies with the phase of the
20 inflammatory process. *Clinical Science* 102, 107-114.
- 21 39. Sahyoun, N.R., Jacques, P.F., Dallal, G., Russell, R.M., 1996. Use of albumin as a
22 predictor of mortality in community dwelling and institutionalized elderly populations.
23 *Journal of Clinical Epidemiology* 49, 981-988.

- 1 40. Schmaltz, H.N., Fried, L.P., Xue, Q.L., Walston, J., Leng, S.X., Semba, R.D., 2005.
2 Chronic cytomegalovirus infection and inflammation are associated with prevalent
3 frailty in community-dwelling older women. *Journal of the American Geriatrics Society*
4 53, 747-754.
- 5 41. Tice, J.A., Browner, W., Tracy, R.P., Cummings, S.R., 2003. The relation of C-reactive
6 protein levels to total and cardiovascular mortality in older U.S. women. *American*
7 *Journal of Medicine* 114, 199-205.
- 8 42. Tobias, P.S., Soldau, K., Ulevitch, R.J., 1986. Isolation of a lipopolysaccharide-binding
9 acute phase reactant from rabbit serum. *Journal of Experimental Medicine* 164, 777-
10 793.
- 11 43. Walston, J., 2004. Frailty--the search for underlying causes. *Sci. Aging Knowledge*
12 *Environ.* 2004, pe4.
- 13 44. Walston, J., Hadley, E.C., Ferrucci, L., Guralnik, J.M., Newman, A.B., Studenski, S.A.,
14 Ershler, W.B., Harris, T., Fried, L.P., 2006. Research agenda for frailty in older adults:
15 Toward a better understanding of physiology and etiology: summary from the american
16 geriatrics society/national institute on aging research conference on frailty in older
17 adults. *Journal of the American Geriatrics Society* 54, 991-1001.
- 18 45. Walston, J., McBurnie, M.A., Newman, A., Tracy, R.P., Kop, W.J., Hirsch, C.H.,
19 Gottdiener, J., Fried, L.P., 2002. Frailty and activation of the Inflammation and
20 coagulation systems with and without clinical comorbidities - results from the
21 cardiovascular health study. *Archives of Internal Medicine* 162, 2333-2341.

FIGURE LEGEND

FIGURE 1 Kaplan-Meier survival curves of old rats by quartiles of levels of each acute phase protein measured at 96 weeks of age. (A) α_2 -macroglobulin (B) Fibrinogen (C) Albumin. n = 18 for each quartile.

ACCEPTED MANUSCRIPT

FIGURE 1

TABLE 1*Age-related changes in plasma levels of acute phase proteins and body weight in ageing rats*

	Age (weeks)		
	35	96	106
	(n = 32)	(n = 72)	(n = 48)
α_2 -macroglobulin (mg/L)	38 ± 22^a	469 ± 811^b	215 ± 257^a
Fibrinogen (g/L)	3.0 ± 0.3^a	4.4 ± 1.6^b	4.1 ± 1.5^b
Albumin (g/L)	18 ± 3^a	12 ± 3^b	12 ± 3^b
Body weight (g)	600 ± 24^a	648 ± 87^b	625 ± 101^{ab}

All rats of the cohort were included at 96 and 106 weeks of age, whereas 32 rats were chosen when they were 35 weeks old. Values are means $\pm$ SD. One-way ANOVA was significant for all parameters. Data not sharing a common letter within a row are significantly different (Fischer's PLSD).

TABLE 2*Plasma levels of acute phase proteins and body weight in 96 week-old rats*

	All (n = 72)	Died before 106 weeks (n = 24)	Alive at 106 weeks (n = 48)
α_2 -macroglobulin (mg/L)	469 $\pm$ 811	1162 $\pm$ 1110 **	122 $\pm$ 153
Fibrinogen (g/L)	4.4 $\pm$ 1.6	5.6 $\pm$ 2.1 **	3.8 $\pm$ 0.9
Albumin (g/L)	12 $\pm$ 3	10 $\pm$ 4 **	13 $\pm$ 2
Body weight (g)	648 $\pm$ 87	596 $\pm$ 86 *	674 $\pm$ 76

Values are means $\pm$ SD. * $P < 0.001$, ** $P < 0.0001$ versus age-matched rats alive at 106 weeks (Student's t test for unpaired data).

TABLE 3

Correlations between percentage of body weight change throughout 10-week follow-up and plasma levels of acute phase proteins measured at 96 weeks of age.

Correlate of percentage of body weight change	r	P
α_2 -macroglobulin	-0.448	0.0012
Fibrinogen	-0.251	0.0849
Albumin	0.074	0.6210

Linear regressions were performed between body weight change throughout the follow-up and plasma levels of each of the three APP measured at the age of 96 weeks in rats alive at 106 weeks (n = 48), r : Pearson correlation coefficient.

TABLE 4

Plasma acute phase protein kinetics and serum levels of soluble tumor necrosis factor- α receptor 1 and lipopolysaccharide-binding protein in rats, alive at 106 weeks, in bottom half and top half categories based on plasma levels of acute phase proteins measured at 96 weeks of age.

		APP levels at 96 weeks of age					
		α_2 -macroglobulin (mg/L)		Fibrinogen g/L		Albumin g/L	
		< 91 (n = 32)	> 91 (n = 16)	< 4.0 (n = 29)	> 4.0 (n = 19)	<12 (n=19)	> 12 (n =29)
α_2 -macroglobulin (mg/L)							
Age (weeks)	96	43 $\pm$ 21	281 $\pm$ 179 **	68 $\pm$ 99	205 $\pm$ 184 *	107 $\pm$ 114	132 $\pm$ 175
	100	107 $\pm$ 355	430 $\pm$ 479 *	141 $\pm$ 382	327 $\pm$ 470	247 $\pm$ 472	194 $\pm$ 397
	104	65 $\pm$ 55	513 $\pm$ 640 *	77 $\pm$ 94	424 $\pm$ 612 *	207 $\pm$ 348	219 $\pm$ 470
	106	115 $\pm$ 200	415 $\pm$ 245 **	91 $\pm$ 102	404 $\pm$ 306 **	212 $\pm$ 274	217 $\pm$ 250
Fibrinogen (g/L)							
Age (weeks)	96	3.4 $\pm$ 0.7 ^{ab}	4.6 $\pm$ 0.7 ^{ab} **	3.3 $\pm$ 0.5 ^a	4.7 $\pm$ 0.5 ^{ab} **	3.8 $\pm$ 0.9 ^a	3.8 $\pm$ 0.8 ^{ab}
	100	3.2 $\pm$ 0.8 ^a	4.2 $\pm$ 1.0 ^a *	3.1 $\pm$ 0.8 ^a	4.2 $\pm$ 0.9 ^a **	3.7 $\pm$ 1.2 ^a	3.4 $\pm$ 0.9 ^c
	104	3.3 $\pm$ 0.9 ^{ab}	4.5 $\pm$ 1.3 ^{ab} *	3.1 $\pm$ 0.5 ^a	4.7 $\pm$ 1.2 ^{ab} **	3.8 $\pm$ 1.2 ^a	3.7 $\pm$ 1.2 ^{ac}

	106	3.6 ± 1.3^b	$5.0 \pm 1.4^{b*}$	3.3 ± 0.8^a	$5.2 \pm 1.6^{b**}$	4.0 ± 1.5^a	4.1 ± 1.5^b
Albumin (g/L)							
	96	14 ± 3^{ac}	12 ± 2^a	14 ± 2^{ac}	13 ± 2^a	11 ± 1^{ab}	$15 \pm 2^{a**}$
Age (weeks)	100	12 ± 2^b	$9 \pm 2^{b*}$	12 ± 2^b	$10 \pm 2^{b*}$	10 ± 3^b	12 ± 2^b
	104	14 ± 3^a	$10 \pm 3^{b**}$	14 ± 3^a	$11 \pm 3^{b**}$	12 ± 3^a	13 ± 3^c
	106	13 ± 2^c	$10 \pm 2^{b*}$	13 ± 2^c	$11 \pm 2^{b*}$	12 ± 3^a	12 ± 2^b
sTNFR-1 (pg/mL)							
Age (weeks)	106	289 ± 78	$479 \pm 261^*$	275 ± 57	$468 \pm 241^*$	361 ± 162	341 ± 195
LBP (μ g/mL)							
Age (weeks)	106	7.8 ± 7.8	12.5 ± 11.8	6.2 ± 4.6	$14.2 \pm 12.7^*$	11.0 ± 12.8	8.1 ± 6.2

Values are means $\pm$ SD. Groups corresponded to rats alive at 106 weeks from quartiles 1 and 2 in one hand and quartiles 2 and 3 in another hand for each of the three APP (see figure 1). Two-way ANOVA for repeated measurements with group as the between-rat factor and time as the within-rat factor: α_2 -macroglobulin groups: significant effect of group for the three APP and time for fibrinogen and albumin and significant interaction between group and time for albumin only, fibrinogen groups: significant effect of group for the three APP and time for fibrinogen and albumin and significant interaction between group and time for albumin only, albumin groups: significant effect of group for albumin only and

time for fibrinogen and albumin and significant interaction between group and time for albumin only. * $P < 0.05$, ** $P < 0.0001$ (group effect: Student's t test). Data not sharing a common letter during the follow-up are different (Fischer's PLSD). sTNFR-1: soluble tumor necrosis factor- α receptor 1, LBP: lipopolysaccharide binding protein

ACCEPTED MANUSCRIPT