Enzymatic Activity in the Rhizosphere of: Potential Contribution for Phytoremediation of metals

Rosa Reboreda, Isabel Caçador

► To cite this version:

Rosa Reboreda, Isabel Caçador. Enzymatic Activity in the Rhizosphere of: Potential Contribution for Phytoremediation of metals. *Marine Environmental Research*, 2007, 65 (1), pp.77. 10.1016/j.marenvres.2007.09.001 . hal-00501924

HAL Id: hal-00501924

<https://hal.science/hal-00501924>

Submitted on 13 Jul 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Enzymatic Activity in the Rhizosphere of *Spartina maritima*: Potential Contribution for Phytoremediation of metals

Rosa Reboreda, Isabel Caçador

PII: S0141-1136(07)00112-2
DOI: [10.1016/j.marenvres.2007.09.001](https://doi.org/10.1016/j.marenvres.2007.09.001)
Reference: MERE 3141

To appear in: *Marine Environmental Research*

Received Date: 12 January 2007
Revised Date: 3 September 2007
Accepted Date: 4 September 2007

Please cite this article as: Reboreda, R., Caçador, I., Enzymatic Activity in the Rhizosphere of *Spartina maritima*: Potential Contribution for Phytoremediation of metals, *Marine Environmental Research* (2007), doi: [10.1016/j.marenvres.2007.09.001](https://doi.org/10.1016/j.marenvres.2007.09.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

- 1   **Type of contribution:** Regular paper
- 2   **Date of preparation:** June 2007
- 3   **Text pages:** 17
- 4   **Tables:** 4
- 5   **Figures:** 1
- 6   **Title:** Enzymatic Activity in the Rhizosphere of *Spartina maritima*: Potential
- 7   **Contribution for Phytoremediation of metals.**
- 8   **Authors:** Rosa Reboreda¹ and Isabel Caçador^{2*}
- 9   **Affiliations:** ^{1, 2} Institute of Oceanography, Faculty of Sciences, University of Lisbon,
- 10   Campo Grande, 1749-016 Lisbon, Portugal
- 11   ***Corresponding author:** E-mail: micacador@fc.ul.pt   Telephone +351 2175 00104
- 12   Fax +351 21 7500009
- 13   *** Address for proofs**

---

* Institute of Oceanography, Faculty of Sciences, University of Lisbon, Rua Ernesto de Vasconcelos s/n, Campo Grande, 1749-016 Lisbon, Portugal

1 Abstract

2 Extracellular Enzymatic Activity (EEA) of five enzymes (peroxidase, phenol
3 oxidase, β -glucosidase, β -N-acetylglucosaminidase and acid phosphatase) was analysed
4 in sediments colonised by *Spartina maritima* in two salt marshes (Rosário and Pancas)
5 of the Tagus estuary (Portugal) with different characteristics such as sediment
6 parameters and metal contaminant levels. Our aim was a better understanding of the
7 influence of the halophyte on microbial activity in the rhizosphere under different site
8 conditions, and its potential consequences for metal cycling and phytoremediation in
9 salt marshes. Acid phosphatase and β -N-acetylglucosaminidase presented significantly
10 higher EEA in Rosário than in Pancas, whereas the opposite occurred for peroxidase.
11 This was mainly attributed to differences in organic matter between the two sites. A
12 positive correlation between root biomass and EEA of hydrolases (β -glucosidase, β -N-
13 acetylglucosaminidase and acid phosphatase) was found, indicating a possible influence
14 of the halophyte in sediment microbial function. This would potentially affect metal
15 cycling in the rhizosphere through microbial reactions.

16
17 Keywords: enzyme activity, halophytes, metals, phytoremediation, salt marsh, *Spartina*
18 *maritima*.

19
20 Abbreviations: EEA (extracellular enzyme activity), β -N-acetyl-glucosaminidase
21 (chitinase)

1. Introduction

Salt marshes usually receive high inputs of contaminants coming from nearby urban areas, river transport and tidal inundation. They are considered to act as sinks for contaminants, namely heavy metals (review by Williams et al., 1994; Doyle and Otte, 1997). Metals accumulate mainly in the sediments, especially in vegetated areas of the marsh (Caçador et al., 1996).

It has been shown that halophytes have a dominant role in the metal cycling in salt marshes (Otte, 1991; Caçador et al., 2000). The roots take up metals, which are mainly accumulated in root tissues and only a small fraction is translocated to the above ground parts of the plants (Alberts, 1990). When plants die, litter decomposition leads to the mobilisation of metals and their return to sediment, at a rate that is dependent on the type of tissue but also on plant species (Windham et al., 2003). Apart from direct uptake of metals, wetland plants contribute to metal removal through other important processes such as substrate stabilization, rhizosphere oxidation and supply of organic matter for microorganisms (Kosolapov et al., 2004).

Microbial activity in sediments is involved in several processes useful for bioremediation of metals, like metal precipitation by sulphate reduction and redox transformations of metals among others (Kosolapov et al., 2004). Wetland plants interact with microorganisms of the rhizosphere by supplying organic matter that is required for such reactions and thus providing an appropriate environment for metal cycling (Jacob and Otte, 2003). For instance, stimulated sulphate reducing bacteria activity caused by *S. maritima* colonisation was thought to be one of the reasons for variations in metal sulphides in salt marshes of Ría de Ortigueira, Spain (Otero and Macías, 2002). In fact, some phytoremediation strategies involve inoculation of plants with rhizosphere microbiota and arbuscular mycorrhiza (Khan, 2005).

Enzymes activities are a useful index of microbial activity (Sinsabaugh, 1994; Ravit et al., 2003). Enzymes can exist extracellularly in sediments, attached to clay minerals or complexed with humic colloids (Burns, 1982). Extracellular enzymes excreted by fungi and bacteria effectively metabolize humic compounds reducing their molecular weight, consequently increasing the solubility and mobility of these potential carriers of heavy metals (Gramss et al., 1999). The enzyme activity assay gives the maximum potential activity of enzymes occurring in sediment rather than the actual enzyme activity (Alef and Nannipieri, 1995).

In this paper we compared the EEA (Extracellular Enzymatic Activity) of five enzymes in sediments colonised by *Spartina maritima* in two salt marshes of the Tagus estuary (Portugal) with marked differences in their characteristics such as sediment parameters and metal contaminant levels (Caçador et al., 1999). Our aim was to compare the possible influence of the plant on the microbial function of the rhizosphere under different site conditions. This would help to elucidate how halophyte-microorganism interactions may contribute for the potential use of halophytes in phytoremediation of metal contaminated environments.

2. Materials and Methods

2.1 Site description

Between February-March 2005 five sediment cores were collected in areas colonised by *Spartina maritima* in each of two salt marshes of the Tagus estuary: Rosário (38° 40'N, 9° 01'W) and Pancas (38° 48'N, 8° 55'W). Rosário is a mature salt marsh (Valiela et al., 2000) with muddy sediments located in the southern part of the estuary, in the vicinity of urbanised and industrial areas. Pancas is a young salt marsh

(Valiela et al., 2000) with muddy-sand sediments located in the northeast part of the estuary, within the Tagus Nature Reserve. Distance between these sites is about 18 km. The studied salt marshes contain extensive intertidal mudflat areas colonised by halophyte plant communities dominated by *Spartina maritima* (Curtis) Fernald, *Halimione portulacoides* (L.) Aellen, *Sarcocornia fruticosa* (L.) A. J. Scott and *Sarcocornia perennis* (Mill.) A. J. Scott. These sites experience two tidal flushing a day (for a more detailed description of the estuary see Caçador et al. (1996)). Rosário presents higher levels of metal contamination than Pancas (Table 1, after Caçador and Vale, 2001).

2.2 Sampling

Collection of samples was made at low tide. Sediment was removed using a stainless steel corer with 6.8 cm inner diameter and 50 cm depth. In the field, cores were sliced to separate the following segments: 0-5 cm, 10-15 cm and 40-45 cm. These segments were expected to be representative of surface sediment, sediment with high root biomass and deep sediment with little or inexistent root influence, respectively. Redox potential (Eh) and pH values of fresh sediment were determined *in situ* using a HANNA pH/mV meter (HI 9025). The samples were placed in plastic bags and transported to the laboratory refrigerated. Those samples for determination of enzymatic activity were frozen at -20°C until enzymatic analysis was carried out (Kuperman and Carreiro, 1997). Root biomass was determined in fresh sediment separating root material from sediment and then drying to constant weight at 70°C (Gross et al., 1991).

2.3 Sediment parameters

Sediment (~20 g) was dried 48 h at 60°C and relative moisture calculated (percentage of fresh weight). In order to determine organic matter content (LOI, loss on ignition) 2g of dry sediment was burned at 600°C for 2 hours.

2.4 Enzyme activities in sediment

Enzymatic activity of 5 enzymes (two oxidoreductases and three hydrolases) was analysed: phenol oxidase, peroxidase, β -glucosidase, acid phosphatase and β -N-acetylglucosaminidase (chitinase). These enzymes are involved in metabolic processes such as C cycling (β -glucosidase, catalyzes the final step of cellulose degradation), N and C cycling (β -N-acetylglucosaminidase, hydrolyzes the glucosamine), P cycling (acid phosphatase, hydrolyzes organic-P compounds) and decomposition of plant litter (phenol oxidase, catalyzes polyphenol oxidation; and peroxidase, catalyzes the oxidation of various compounds such as phenols and amines). We used the method described in Ravit et al. (2003) with a modification in the incubation temperature and without dilution of the supernatant. Sediment samples were prepared for analysis adding 75 ml of sodium acetate buffer (pH 5) to 5 g fresh sediment. The sample was mixed for 1 min to obtain the sediment slurry. The substrates used for β -glucosidase, acid phosphatase and β -N-acetylglucosaminidase (chitinase) were p-Nitrophenyl- β -D-glucopyranoside, p-Nitrophenyl phosphate and p-Nitrophenyl-N-acetyl- β -D-glucosaminide, respectively. All of them released p-nitrophenol (pNP) when hydrolyzed. Two millilitres of substrate (5 mM) was added to 2 ml of the slurry in a test tube and incubated at 30°C in gentle agitation for 60 min (β -glucosidase), 30 min (acid phosphatase) and 2 hours (chitinase). After incubation the samples were centrifuged at 2000 rev min⁻¹ for 15 min and 0.2 ml of 1 N NaOH was added immediately to stop the reaction and cause colour change. Determination of phenol oxidase and peroxidase activities was carried out using L-DOPA (L-3,4-dihydroxyphenylalanine) as substrate

(5 mM) following the same procedure: 2 ml slurry + 2 ml substrate (adding 0.1 ml of 0.3% H₂O₂ for peroxidase). Incubation time was 60 min for both enzymes, followed by centrifugation. Absorbance of supernatant was read on a TECAN Absorbance Microplate Reader (SPECTRA Rainbow) at 410 nm for pNP enzymes and at 460 nm for L-DOPA enzymes. Absorbance for phenol oxidase was subtracted from absorbance of peroxidase total (which assay gives the activity of peroxidase + phenol oxidase) in order to calculate absorbance for peroxidase alone. Controls of sediment slurry and substrate were made, and their absorbance subtracted from the absorbance read for each sample. Enzymatic activity was expressed as $\mu\text{moles product g dry wt}^{-1} \text{ h}^{-1}$.

2.5 Statistic analyses

Significant differences in soil characteristics between sites were tested using one-way ANOVA. Significant differences in extracellular enzymatic activities between the two sites at the different depths were compared using one-way ANOVA. The same analysis was used to test differences with depth at each site separately. This was followed by a Tukey- HSD test when appropriate. Pearson's coefficient was calculated to test correlation between activities of different enzymes and correlation between enzymes activities and soil characteristics.

3. Results and discussion

Sediment characteristics (Eh, pH, LOI, relative moisture) and root biomass of *S. maritima* in Rosário and Pancas are shown in Table 2. Sediment redox potential (Eh) was significantly higher ($p < 0.05$) in Pancas, except on the surface, where Rosário showed significantly higher Eh. Organic matter (LOI) and moisture were significantly higher ($p < 0.05$) in Rosário at every depth studied in areas colonised by *S. maritima*,

whereas pH was significantly higher in Rosário only at the root depth (10-15 cm). Root biomass of *S. maritima* was significantly higher in Rosário than in Pancas ($p < 0.05$).

The extracellular enzymatic activity (EEA) of peroxidase was much higher (up to 10 fold) than the EEA of the other four enzymes at every depth studied (Figure 1). The dominance of peroxidase over hydrolytic enzymes may indicate the presence of refractory organic matter in the salt marsh (Sinsabaugh et al., 2003). Gramss et al. (1999) demonstrated that extracellular enzymes released by fungi and bacteria, as well as commercial enzymes including peroxidase and β -glucosidase, were able to effectively metabolize soil humic extracts (to which heavy metal ions and xenobiotics are potentially attached), reducing their molecular weight and consequently increasing their solubility and motility. Therefore, high extracellular activity of peroxidase in sediments colonised by *S. maritima* might influence the mobilization of metals in the studied salt marshes. In addition to supplying plant litter for decomposition, the halophyte may contribute to high EEA of peroxidase by the release of peroxidases from plant roots (Salt et al., 1998). However, it should be taken into account that the EEA assay gives the maximum potential activity of enzymes occurring in sediment rather than the actual enzyme activity (Alef and Nannipieri, 1995).

When comparing EEA's in sediments colonised by *S. maritima* in Rosário and Pancas, significant differences were observed for peroxidase, acid phosphatase and chitinase ($p < 0.05$) (Figure 1). Peroxidase presented higher EEA in Pancas than in Rosário, whereas acid phosphatase and chitinase presented higher EEA in Rosário than in Pancas. Correlation coefficients between EEA's and sediment characteristics revealed that acid phosphatase and chitinase activities were positively correlated with organic matter content and moisture (Table 3). In addition, chitinase activity was positively

1 correlated with pH. A negative correlation occurred between EEA of peroxidase and
2 organic matter content. These results suggest that differences in the organic matter
3 content between the two sites could partly explain the observed pattern in extracellular
4 activities of the enzymes in areas colonised by *S. maritima*. Higher organic matter
5 content and moisture in sediment from Rosário seemed to be related to higher EEA's of
6 acid phosphatase and chitinase (also influenced by higher pH) in this salt marsh,
7 whereas peroxidase activity was higher in Pancas, where organic matter content of the
8 sediment was lower. Results from Caçador et al. (2004) showed that litter
9 decomposition in Pancas was faster than in Corroios, another salt marsh with
10 characteristics similar to Rosário (located about 12 Km to the southwest from this)
11 (Caçador et al., 1999). This may be related to higher activities of peroxidase in Pancas,
12 as it is involved in the degradation of complex organic compounds. There was no
13 correlation between EEA's and redox potential of the sediment. This was surprising,
14 especially for phenol oxidase, that needs O₂ for its activity. An oxygen limitation on
15 phenol oxidase is associated with inhibition of hydrolytic enzymes in peatlands through
16 accumulation of phenolic compounds (Freeman et al, 2001; Freeman et al. 2004). The
17 low EEA observed for phenol oxidase (in the same order of magnitude as the
18 hydrolases) is likely to be caused by O₂ limitation in the sediment, since it is expected to
19 be scarce in salt marsh sediments. Although no reductive conditions were found in the
20 sediments studied (positive Eh) it should be taken into account that it does not indicate
21 abundance of O₂, as other oxidant compounds could be present. Accumulation of
22 phenolic compounds due to restricted phenol oxidase activity would have important
23 consequences for the cycling of heavy metals in the salt marsh, since phenols are known
24 to bind metals (Ambrosi et al., 2003; Mota et al., 2005).

25

1   Hydrolytic enzymes activities (β -glucosidase, acid phosphatase and chitinase)
2   were positively correlated with root biomass of *S. maritima* (Table 3). Microbial
3   activity is stimulated around plant roots through exudation of organic anions and sugars
4   (Ryan et al., 2001). Karjalainen et al. (2001) found that *Lobelia dortmanna* L., a
5   submersed plant species, enhanced microbial activity and biomass in sediments through
6   organic carbon release. Thus, it would be reasonable that higher root biomass would
7   result in higher exudation of organic compounds, enhancing microbial activity around
8   plant roots. Burke et al. (2002) found a positive interaction between a group of
9   *Protobacteria* and root biomass of *Spartina patens* and *Phragmites australis* in salt
10   marsh sediments. The release of soluble sugars by roots supply the substrates needed by
11   β -glucosidase and chitinase (β -N-acetylglucosaminidase), probably inducing the
12   excretion of higher amounts of these enzymes by microorganisms (Burns, 1982). In
13   spite of this, β -glucosidase activity was not significantly higher in sediment from
14   Rosário, where *S. maritima* root biomass was significantly higher (Figure 1; Table 3).
15   This might be caused by an inhibition of this enzyme in the presence of higher metal
16   concentrations in Rosário (Table 1) (Geiger et al., 1998). Exudation of phosphatases
17   from plant roots is one of the strategies used by plants to cope with P deficiency (Ryan
18   et al., 2001). This might be a strategy of *S. maritima* as well, given the correlation found
19   between acid phosphatase activity and root biomass of the plant (Table 3). Ravit (2005)
20   concluded that, in salt marsh environments of the USA, the sediment microbial
21   community structure and function is determined by the interaction of hydrological
22   conditions and the effect of plant roots. From the observed results we deduce that *S.*
23   *maritima* roots could influence the extracellular enzymatic activity of the hydrolytic
24   enzymes analysed and, accordingly, it may affect the microbial function in the
25   rhizosphere of the plant in Tagus estuary salt marshes. Given the importance of

microbially mediated reactions in metal cycling around plant roots in wetlands (Jacob and Otte, 2003), and the proposed use of halophytes like *S. maritima* for phytostabilization of metals in metal contaminated salt marshes (Reboreda and Caçador, 2007; Reboreda et al., *In press*), the halophyte-microorganism interactions arise as an additional concern for phytoremediation studies.

Variations in EEA with depth were only significant ($p < 0.05$) for β -glucosidase and acid phosphatase from Rosário. Still, enzymatic activities were in general lower in the horizon 40-45 cm, most likely as a consequence of low organic matter content and oxygen availability (Table 2). In Pancas, the lack of significant differences in EEA with depth ($p > 0.05$) was surprising considering the 2-orders of magnitude change in root biomass and organic matter (Table 2). Other sediment parameters such as grain size (muddy-sand in Pancas; Reboreda et al., *in press*) may have a stronger effect on enzymes distribution (Kandeler et al., 2000).

In Rosário, EEA's of β -glucosidase and chitinase were positively correlated with EEA of phenol oxidase (Table 4). This might suggest a similar behaviour of these enzymes, in organic matter rich/oxygen limited salt marshes, as that found by Freeman et al. (2001) and Freeman et al. (2004) in peatlands, which was described above. EEA of β -glucosidase also showed correlation with peroxidase activity. There were no significant correlations between EEA's of the studied enzymes in Pancas.

4. Conclusions

Sediments colonised by *S. maritima* show differences in some extracellular enzymes activities between salt marshes in the Tagus estuary, which appears to be related to differences in sediment organic matter and root biomass. The results support

the idea that *S. maritima* could influence to some extent the microbial function in the rhizosphere, which in turn may potentially influence microbially mediated metal cycling in the sediment. In addition, site characteristics seem to play a relevant role in enzymatic activity as well, given the differences observed between the two sites. Consequently, we suggest that halophyte-microorganism interactions are worthy of further study for its usefulness in phytoremediation strategies.

6. Acknowledgements

This research was funded by “Fundação para a Ciência e Tecnologia, MECTIS POCI/MAR/58548/2004”. The authors are grateful to Dr. Beth Ravit for the clarifications about the experimental procedure. Thank you to the anonymous referees who contributed to improve the manuscript.

6. References

- Alberts, J.J., Price, M.T., Kania, M., 1990. Metal concentrations in tissues of *Spartina alterniflora* (Loisel.) and sediments of Georgia salt marshes. Estuarine Coastal and Shelf Science 30, 47-58.
- Alef, K., Nannipieri, P. (Eds.), 1995. Methods in Applied Soil Microbiology and Biochemistry. Academic Press, London
- Ambrosi, G., Formica, M., Fusi, V., Giorgi, L., Guerri, A., Micheloni, M., Paoli, P., Pontellini, R., Rossi, P., 2003. Heavy metal ion complexes with a simple phenolic ligand. Solid state and solution studies. Inorganica Chimica Acta-Articles 356, 203-209.
- Burke, D.J., Hamerlynck, E.P., Hahn, D., 2002. Interactions among plant species and microorganisms in salt marsh sediments. Applied and Environmental Microbiology 68(3), 1157-1164.
- Burns, R.G., 1982. Enzyme activity in soil: location and a possible role in microbial ecology. Soil Biology and Biochemistry 14, 423-427.
- Caçador, I., Costa, A.L., Vale, C., 2004. Carbon storage in Tagus salt marsh sediments. Water, Air and Soil Pollution 4, 701-714.

- 1 Caçador, I., Vale, C., Catarino, F., 1996 Accumulation of Zn, Pb, Cu and Ni in
2 sediments between roots of the Tagus estuary salt marshes, Portugal. Estuarine Coastal
3 and Shelf Science 42(3), 393-403.
4
5
6 Caçador, I., Vale, C., Catarino, F., 1999. Relationships between metal concentrations
7 and organic matter content in the Tagus estuary salt marsh sediments. In: Lieth, H.,
8 Moschenko, M., Lohmann, M., Koyro, H-W., Hamdy A. (Eds.). Halophyte uses in
9 different climates I, Backhuys Publishers, Leiden, The Netherlands: 105-112.
10
11 Caçador, I., Vale, C., Catarino, F., 2000. Seasonal variation of Zn, Pb, Cu and Cd
12 concentrations in the roots-sediment system of *Spartina maritima* and *Halimione*
13 *portulacoides* from Tagus estuary salt marshes. Marine Environmental Research 49(3),
14 279-290.
15 Caçador, I., Vale, C., 2001. Retention of Heavy Metals in Salt Marshes and Estuarine
16 Water Quality Metals in the Environment: Analysis by Biodiversity. MNV. Prasad (Ed.)
17 Marcel Dekker, Inc, New York: 95-116.
18
19 Doyle, M.O., Otte, M.L., 1997. Organism-induced accumulation of Fe, Zn and AS in
20 wetland soils. Environmental Pollution 96 (1), 1-11.
21
22 Freeman, C., Ostle, N.J., Fenner, N., Kang, H., 2004. A regulatory role for phenol
23 oxidase during decomposition in peatlands. Soil Biology and Biochemistry 36, 1663-
24 1667.
25

- 1 Freeman, C., Ostle, N.J., Kang, H., 2001. An enzymic 'latch' on a global carbon store.
2 Nature 409, 149.
3
- 4 Geiger, G., Furrer, G., Funk, F., Brandl, H., Schulín, R., 1999. Heavy metal effects on
5 beta-glucosidase activity influenced by pH and buffer systems. Journal of Enzyme
6 Inhibition 14 (5), 365-379.
7
- 8 Gramss, G., Ziegenhagen, D., Sorge, S., 1999. Degradation of soil humic extract by
9 wood- and soil-associated fungi, bacteria and commercial enzymes. Microbial Ecology
10 37, 140-151.
11
- 12 Gross, M.F., Hardisky, M.A., Wolf, P.L., Klemas, V., 1991. Relationship between
13 aboveground and belowground biomass of *Spartina alterniflora* (smooth cordgrass).
14 Estuaries 14, 180-191.
15
- 16
- 17 Jacob, D.L., Otte, M.L., 2003. Conflicting processes in the wetland plant rhizosphere:
18 metal retention or mobilization? Water, Air and Soil Pollution 3, 91-104.
19
- 20
- 21 Kandeler, E., Tschérko, D., Bruce, K.D., Stemmer, M., Hobbs, P.J., Bardgett, R.D.,
22 Amelung, W., 2000. Structure and function of the soil microbial community in
23 microhabitats of a heavy metal polluted soil. Biology and Fertility of Soils 32 (5), 390-
24 400.
25

- 1 Khan, A.G., 2005. Role of soil microbes in the rhizospheres of plants growing on trace
2 metal contaminated soils in phytoremediation. *Journal of Trace Elements in Medicine*
3 *and Biology* 18, 355-364.
- 4
- 5 Kosolapov, D.B., Kusch, P., Vainshtein, M.B., Vatsourina, A.V., Wießner, A.,
6 Kästner, M., Müller, R.A., 2004. Microbial processes of heavy metal removal from
7 carbon-deficient effluents in constructed wetlands. *Engineering of Life Science* 4 (5),
8 403-411.
- 9
- 10 Kuperman, R.G., Carreiro, M.M., 1997. Soil heavy metal concentrations, microbial
11 biomass and enzyme activities in a contaminated grassland ecosystem. *Soil Biology and*
12 *Biochemistry* 29 (2), 179-190.
- 13
- 14
- 15 Mota, A.M., Cruz, P., Vilhena, C., Gonçalves, M.L.S., 2005. Influence of the sediment
16 on lead speciation in the Tagus estuary. *Water Research* 39, 1451-1460.
- 17
- 18 Otero, X.L., Macías, F., 2002. Variation with depth and season in metal sulphides in salt
19 marsh soils. *Biogeochemistry* 61 (3), 247-268.
- 20
- 21 Otte, M.L., Besterbroer, S.J., Van der Linden, J.M., Rozema, J., Broekman, R.A., 1991.
22 A survey of Zn, Cu and Cd concentrations in salt marsh plants along the Dutch coast.
23 *Environmental Pollution* 72, 175-189.
- 24

- 1 Ravit, B., 2005. Macrophytes and microbes: *Spartina alterniflora* and *Phragmites*
2 *australis* affect brackish sediment microbial community structure and function.
3 Dissertation Abstracts International Part B: Science and Engineering 66 (5), 2488.
4
- 5 Ravit, B., Ehrenfeld, J.G., Haggblom, M.M., 2003. A comparison of sediment microbial
6 communities associated with *Phragmites australis* and *Spartina alterniflora* in two
7 brackish wetlands of New Jersey. Estuaries 26(2B), 465-474.
8
- 9 Reboreda, R., Caçador, I., 2007. Halophyte vegetation influences in salt marsh retention
10 capacity for heavy metals. Environmental pollution 146, 147-154.
11
- 12 Reboreda, R., Caçador, I., Pedro, S., Raposo, P., *In press*. Mobility of metals in salt
13 marsh sediments colonised by *Spartina maritima* (Tagus estuary, Portugal).
14 Hydrobiologia.
15
- 16 Rieling, T., Gerbersdorf, S., Stodian, I., Black, H.J., Dahlke, S., Köster, M.,
17 Meyercordt, J., Meyer-Reil, L-A., 2000. Benthic microbial decomposition of organic
18 matter and nutrient fluxes at the sediment-water interface in a shallow coastal inlet of
19 the southern Baltic Sea (Nordrügenschke Boden). In: Flemming, B.W., Delafontaine,
20 M.T., Liebezeit, G. (Eds.) Muddy Coast Dynamics and Resource Management, Elsevier
21 Science BV, Amsterdam.
22
- 23 Ryan, P.R., Delhaize, E., Jones, D.L., 2001. Function and mechanism of organic anion
24 exudation from plant roots. Annual Review of Plant Physiology and Plant Molecular
25 Biology 52, 527-560.

- 1
- 2 Salt, D.E., Smith, R.D., Raskin, I., 1998. Phytoremediation. Annual Reviews of Plant
- 3 Physiology and Plant Molecular Biology 49, 643-668.
- 4
- 5 Sinsabaugh, R.L., 1994. Enzymatic analysis of microbial pattern and process. Biology
- 6 and Fertility of Soils 17 (1), 69-74.
- 7
- 8 Sinsabaugh, R.L., Saiya-Cork, K., Long, T., Osgood, M.P., Neher, D.A., Zak, D.R.,
- 9 Norby, R.J., 2003. Soil microbial activity in a *Liquidambar* plantation unresponsive to
- 10 CO₂-driven increases in primary production. Applied Soil Ecology 24, 263-271.
- 11
- 12 Williams, T.P., Bubb, J.M., Lester, J.N., 1994. Metal accumulation within salt marsh
- 13 environments: a review. Marine Pollution Bulletin 28 (5), 277-290.
- 14
- 15 Windham, L., Weis, J.S., Weis, P., 2003. Uptake and distribution of metals in two
- 16 dominant salt marsh macrophytes, *Spartina alterniflora* (cordgrass) and *Phragmites*
- 17 *australis* (common reed). Estuarine, Coastal and Shelf Science 56, 63–72.
- 18
- 19 Valiela, I., Cole, M., McClelland, J. Cebrian, J., Joye, S.B. 2000. Role of salt marshes
- 20 as part of coastal landscapes. Concepts and Controversies in Tidal Marsh Ecology (eds)
- 21 Weinstein, M.P. & Kreeger, D.A. Kluwer Academic Publishers. London
- 22

Figure caption

Figure 1. Extracellular enzymatic activities (EEA) ($\mu\text{moles h}^{-1} \text{ g dry weight}^{-1}$) in three depths of sediment at Rosário and Pancas sites colonised by *Spartina maritima* (mean $\pm$ SD). *Significant differences ($p<0.05$) between sites.

Figure 1

Table 1. Concentrations (mean $\pm$ standard deviation) of Zn, Pb, Cu and Cd ($\mu\text{g g}^{-1}$ dry weight) in surface sediments (0-5 cm) ; intermediate rooted sediment layers (10-15 cm) and deeper layers (40-45 cm) in areas colonised by *Spartina maritima*, from Pancas and Rosário salt marshes (after Caçador & Vale, 2001)

Sites	Depth (cm)	Zn $\mu\text{g g}^{-1}$ dw	Pb $\mu\text{g g}^{-1}$ dw	Cu $\mu\text{g g}^{-1}$ dw	Cd $\mu\text{g g}^{-1}$ dw
Pancas	(0-5)	95 $\pm$ 17	42 $\pm$ 4	22 $\pm$ 3	1.1 $\pm$ 0.2
	(10-15)	177 $\pm$ 18	53 $\pm$ 4	33 $\pm$ 6	1.3 $\pm$ 0.1
	(40-45)	93 $\pm$ 5	43 $\pm$ 9	14.7 $\pm$ 3	0.4 $\pm$ 0.1
Rosário	(0-5)	402 $\pm$ 27	395 $\pm$ 16	51 $\pm$ 2	1.2 $\pm$ 0.2
	(10-15)	1169 $\pm$ 74	577 $\pm$ 21	78 $\pm$ 15	1.7 $\pm$ 0.2
	(40-45)	98 $\pm$ 10	55.3 $\pm$ 11	12.7 $\pm$ 3	0.4 $\pm$ 0.1

Table 2. Eh (mV), pH, organic matter (Loss on Ignition, %), moisture (%) and root biomass (g m^{-2}) in sediments from Rosário and Pancas colonised by *S. maritima* at three sediment horizons (mean $\pm$ SD).

	Depth	Eh (mV)	pH (14°C)	LOI	Moisture (%)	Biomass (g m^{-2})
Pancas	0-5 cm	84.0 $\pm$ 24.7*	5.9 $\pm$ 0.5	10.3 $\pm$ 0.6*	52.3 $\pm$ 0.7*	264.6 $\pm$ 26.3*
	10-15 cm	80.9 $\pm$ 23.2*	5.9 $\pm$ 0.4*	8.0 $\pm$ 0.1*	50.7 $\pm$ 1.7*	681.0 $\pm$ 9.6*
	40-45 cm	76.3 $\pm$ 11.9*	5.5 $\pm$ 1.3	0.6 $\pm$ 0.7*	16.4 $\pm$ 0.6*	9.6 $\pm$ 3.0*
Rosário	0-5 cm	107.4 $\pm$ 23.0*	5.5 $\pm$ 0.1	19.0 $\pm$ 5.9*	64.1 $\pm$ 5.0*	798.8 $\pm$ 50.2*
	10-15 cm	29.0 $\pm$ 7.0*	6.7 $\pm$ 0.2*	14.5 $\pm$ 2.0*	60.0 $\pm$ 2.8*	1305.0 $\pm$ 124.3*
	40-45 cm	20.0 $\pm$ 6.8*	5.8 $\pm$ 0.1	11.8 $\pm$ 2.1*	49.7 $\pm$ 8.5*	31.8 $\pm$ 6.0*

*Significant differences between sites (one-way ANOVA, $p < 0.05$).

Table 3. Correlation coefficients (Pearson's) between sediment characteristics and extracellular enzymatic activities of phenol oxidase, peroxidase, β -glucosidase, acid phosphatase and chitinase in salt marsh sediments colonised by *Spartina maritima*.

	Eh (mV)	pH	LOI (%)	Moisture (%)	Biomass (g m ⁻²)
Phenol ox.	0.48	-0.01	-0.38	-0.17	-0.13
Peroxidase	0.46	0.00	-0.54*	-0.31	-0.11
β -glucosidase	0.32	0.46	0.26	0.31	0.86**
Phosphatase	0.47	-0.05	0.70**	0.51*	0.58*
Chitinase	-0.17	0.57*	0.69*	0.59*	0.88**

* Correlation is significant at the 0.05 level.

** Correlation is significant at the 0.01 level.

Table 4. Correlation between the activities of phenol oxidase, peroxidase, β -glucosidase, acid phosphatase and chitinase in sediments of Rosário and Pancas colonised by *S. maritima* (Pearson's Correlation).

	Enzyme	Phenol ox.	Peroxidase	β -gluc	Phosphatase	Chitinase
Rosário	Phenol ox.					
	Peroxidase	0.454				
	β -glucosidase	0.591*	0.650*			
	Phosphatase	0.211	0.198	0.398		
	Chitinase	0.707**	0.473	0.796**	0.530	
Pancas	Phenol ox.					
	Peroxidase	0.497				
	β -glucosidase	0.089	-0.230			
	Phosphatase	0.025	0.192	-0.046		
	Chitinase	-0.079	0.328	0.204	0.033	

* Correlation is significant at the 0.05 level.

** Correlation is significant at the 0.01 level.