Nematode community dynamics over an annual production cycle in the central North Sea

M. Schratzberger, R.M. Forster, F. Goodsir, S. Jennings

► To cite this version:

M. Schratzberger, R.M. Forster, F. Goodsir, S. Jennings. Nematode community dynamics over an annual production cycle in the central North Sea. *Marine Environmental Research*, 2008, 66 (5), pp.508. 10.1016/j.marenvres.2008.08.004 . hal-00501980

HAL Id: hal-00501980

<https://hal.science/hal-00501980>

Submitted on 13 Jul 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Nematode community dynamics over an annual production cycle in the central North Sea

M. Schratzberger, R.M. Forster, F. Goodsir, S. Jennings

PII: S0141-1136(08)00196-7
DOI: [10.1016/j.marenvres.2008.08.004](https://doi.org/10.1016/j.marenvres.2008.08.004)
Reference: MERE 3278

To appear in: *Marine Environmental Research*

Received Date: 10 June 2008
Revised Date: 5 August 2008
Accepted Date: 6 August 2008

Please cite this article as: Schratzberger, M., Forster, R.M., Goodsir, F., Jennings, S., Nematode community dynamics over an annual production cycle in the central North Sea, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.08.004](https://doi.org/10.1016/j.marenvres.2008.08.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Nematode community dynamics over an annual production cycle
in the central North Sea**

M. Schratzberger*, R. M. Forster, F. Goodsir, S. Jennings

*The Centre for Environment, Fisheries and Aquaculture Science, Lowestoft Laboratory,
Pakefield Road, Lowestoft, Suffolk NR33 0HT, UK*

* Corresponding author: Tel. +44(0)1502-527743, Fax. +44(0)1502-513865

E-mail address: michaela.schratzberger@cefas.co.uk (M. Schratzberger)

Abstract

Nematode species composition, trophic structure and body size distributions were followed over an annual production cycle in the central North Sea; to test responses to temporally changing food quality and quantity in the sediment. Changes in the phytoplankton concentration in the water column were quantitatively reflected in the concentration of chlorophyll *a* and breakdown products in the sediment, with higher concentrations in spring and autumn following blooms, and lower concentrations in summer and winter. The taxonomic and trophic structure of nematode communities differed significantly among stations over relatively short distances, potentially masking some of the temporal dynamics. Spatio-temporal differences in nematode species composition were linked to changes in the quality and quantity of organic material reaching the seabed, reflecting a species-specific response to the nutritional quality of sedimenting organic material and the biochemical changes in the sediment associated with its decomposition. The size distributions of selected nematode species indicated that most species bred continuously throughout the sampling period, although one species, the epigrowth feeder *Spilophorella paradoxa*, had periods of increased growth following the deposition of the spring phytoplankton bloom. There was no consistent temporal relationship between the trophic composition of nematode communities and spring chlorophyll *a* or carbon sedimentation, most likely a result of the trophic plasticity of most feeding types and the capacity of the community to use both freshly sedimented material as well as the subsequent breakdown products and refractory organic matter. Community metrics implied that there were small responses to the seasonal production cycle, but these belied strong responses of a few species with life histories that allowed them to track the availability of suitable food resources.

Key words: Nematodes; Taxonomic and functional community composition; Benthic-pelagic coupling; Food quality; Food quantity; North Sea

39 1. Introduction

40

41 The ecology of marine food webs at higher latitudes is strongly influenced by broadly
 42 predictable production cycles, driven by changes in nutrient supply, solar radiation and wind
 43 stress (Colebrook and Robinson, 1961). The production cycle in stratified temperate seas is
 44 characterised by one or more distinct phytoplankton blooms. The spring bloom is often
 45 dominated by larger-celled groups such as diatoms, followed by proliferation of nano- and
 46 picoplankton populations throughout the summer and autumn. The fate of carbon generated
 47 by phytoplankton depends on whether energy and nutrients fixed during blooms are recycled
 48 in the water column, exported to higher trophic levels or transferred to the seabed (Jago et al.,
 49 2007). After the bloom, fresh organic material sinking to the bottom is partly incorporated
 50 into the sediment as detrital particles or becomes adsorbed to sediment grains.

51 Sedimentation of organic matter represents the major input of production to the benthic
 52 community on an annual basis and is a major factor influencing benthic community structure,
 53 biomass and metabolism (Graf et al., 1982). Energy produced during annual production
 54 cycles propagates through the food web, with increasing lag and attenuation as it passes to
 55 larger animals with slower turnover times (Pope et al., 1994). Smaller benthic animals are
 56 likely to track the production cycle most closely, as their high relative rates of consumption
 57 and fast turnover times allow them to rapidly assimilate available production. The ecology
 58 and life histories of meiofauna (invertebrates that pass through a 500 – 1000 μm mesh sieve
 59 but are retained on a 32 – 63 μm mesh sieve; Giere, 1993) suggest that their population and
 60 community dynamics will respond to the seasonality of production.

61 Meiofauna play an important role as intermediaries between the microbial/detrital
 62 compartments (by influencing bacterial density and biomass; Mikola, 1998), and larger
 63 organisms (by serving as major food sources for higher trophic levels; Danovaro et al., 2007)
 64 in food webs (Chardy and Dauvin, 1992; Schmid-Araya et al., 2002). However, little is
 65 known about how food quality and quantity affect natural populations (Findlay, 1982;
 66 Danovaro and Gambi, 2002). Seasonal data on meiofauna abundance, reproduction and

67 species composition are essential for conceptualising and modelling trophic dynamics
68 (Fleeger, 1985) but to date, relationships between faunal characteristics and phytoplankton
69 sedimentation have proven elusive. This is partly a result of the rarity of surveys which
70 monitor phytoplankton sedimentation and faunal responses adequately. Thus most ecological
71 studies of meiofauna have focused on intertidal or shallow subtidal habitats. When deeper
72 water environments were studied, sampling designs rarely permitted repeated sampling at
73 multiple stations or over more than one season (e.g. Vanaverbeke et al., 2004a and references
74 therein).

75 Phytoplankton sedimentation is likely to affect meiofauna communities as many species live
76 close to the sediment surface, feeding on diatoms and epigrowth (i.e. algal cells attached to
77 particles). Fresh organic material is expected to reach deeper dwelling, deposit and bacteria
78 feeding species during later stages of phytoplankton decomposition (Graf et al., 1982;
79 Rudnick, 1989). Indeed, temperature and food availability are the factors most often invoked
80 to explain seasonal changes in meiofaunal abundance (Heip et al., 1985; Ólafsson and
81 Elmgren, 1997).

82 Taxon-based approaches are commonly used to study temporal dynamics of meiofauna.
83 Aggregating taxonomically similar species or different developmental stages of the same
84 species, however, does not take account of the diverse biology and autecological requirements
85 of the organisms (Schratzberger et al., 2007). These issues have been taken into account by
86 adopting functional and size-based approaches. The functional group approach classifies
87 organisms by morphological characteristics that are thought or known to represent an
88 important ecological function (Chalcraft and Resetarits, 2003). However, links between
89 morphology and function can be weak and inconsistent, in some cases leading to an uncertain
90 separation of taxonomic and functional approaches. The size-based approach focuses on the
91 aggregation of organisms based on individual body weight, often independent of their
92 taxonomy. Size is linked to processes such as metabolism and productivity, but size-based
93 approaches may not reflect complex differences in the feeding ecology of different sized
94 organisms.

Since taxonomic, functional and size-based approaches focus on different aspects of assemblage structure, a combination of all three methods is likely to provide better understanding of seasonal changes in assemblage structure, dynamics and responses to changing food supply. Here, we study meiofaunal nematode community dynamics in the central North Sea to assess how taxonomic, trophic and biomass distribution patterns varied during a seasonal production cycle. We relate community changes to changing food quality and quantity in the sediment to investigate two questions:

- (1) Does a local increase in food promote consistent temporal changes in the taxonomic, trophic and size structure of nematode communities?
- (2) Do selected nematode species, representing the main taxonomic and trophic categories available to the phylum, respond in a differential way to a pulsed food supply?

2. Material and methods

2.1. Study site

We studied nematode communities in the Silver Pit, central North Sea (Fig. 1). The Silver Pit is predominantly 60 – 80 m deep with a muddy-sand substratum and the water column stratifies in summer months. Parts of the Silver Pit are fished by beam trawlers but there is limited evidence that trawling disturbance has significant effects on nematode community structure. Thus a previous experimental study in the Southern North Sea, investigating acute effects of trawling disturbance on nematodes, showed that seasonal changes in assemblage structure were greater than those attributed to trawling (Schratzberger et al., 2002). Sub-surface temperature at 35 m depth at the sampling stations was low during spring and summer (6 – 8 °C February to June, 9 – 10 °C in July), peaked in late summer (11 – 16 °C August to October) and declined again during autumn and winter months (8 – 10 °C November to January) (Paul McCloghrie, unpublished data).

2.2. Sample collection

Sediment samples were collected with a NIOZ corer on six cruises in 2005 and 2006 (21 – 26 May, 12 – 16 July, 29 September – 4 October, 5 – 10 November 2005; 13 – 17 February, 26 April – 5 May 2006) from seven randomly chosen locations within 50 m of each of three stations (Fig. 1). The NIOZ corer takes a circular core of 0.1 m² to a depth of > 30 cm. The surface of the sediment core is relatively undisturbed, and the corer retains a few cm of the overlying water.

From each deployment, two sediment sub-samples were collected with a perspex corer (2 cm diameter, 3.14 cm² surface area) to a depth of 5 cm. On each sampling date, sub-samples were combined by station and frozen at -20 °C prior to the analysis of particle size and organic carbon content. From each of the first three deployments, two further sub-samples were collected, one for the analysis of sediment chlorophyll *a*, the other for the analysis of sediment porosity. These sub-samples were frozen separately at -20 °C pending analysis. As sediments reflect a time-averaged input of phytodetritus to the benthic system, sediment chlorophyll *a* concentrations are likely to show a more reliable pattern of annual algal biomass and sedimentation than near-bottom water samples. Boon et al. (1998) and Franco et al. (2007) showed that the typical algal spring bloom is quantitatively reflected in the chlorophyll *a* concentration of the majority of North Sea sediments.

From each of the last four deployments at each location, a meiofauna sub-sample was collected with a 5.5 cm internal diameter perspex tube (surface area 23.76 cm²) to a depth of 5 cm. All faunal samples were fixed and preserved in 10 % formalin in filtered seawater.

2.3. Sample processing

148 After thawing, sediment samples were wet sieved through a 500 μm sieve, and the fraction >
 149 500 μm was oven dried at 90 $^{\circ}\text{C}$ for 24 h. This fraction was then dry sieved at 0.5 phi
 150 intervals, down to 1 phi (500 μm). The fraction < 500 μm was freeze dried and analysed on a
 151 Coulter LS 130 Laser sizer. The laser sizer results were combined with the dry sieve results to
 152 give the full particle size distribution. The mean particle size diameter (hereafter referred to as
 153 grain size) was calculated from these results (Dyer, 1986). The total organic carbon content of
 154 the sediment fraction < 63 μm was determined with a Leeman CE 440 elemental analyser.
 155 Total organic carbon in the sediment consists of a variety of compounds with different
 156 decomposition rates based on their structures and susceptibility to biological attack (Henrichs
 157 and Doyle, 1986). Organic matter in the majority of marine sediments is complex and
 158 generally of less degradable (i.e. refractory) nature (Fabiano and Danovaro, 1994).
 159 Defrosted sediment chlorophyll *a* samples were extracted in neutralised 90 % acetone for 24 h
 160 at 4 $^{\circ}\text{C}$ in the dark. Measurements of extracted chlorophyll fluorescence were made (before
 161 and after acidification with 8 % hydrochloric acid to distinguish phaeopigments) using a
 162 Turner Design (Model 10 AU) filter fluorometer (Tett, 1987). Sediment chlorophyll *a* and
 163 phaeopigment concentrations in the interstitial water were corrected for porosity, determined
 164 from the dry weights and wet weights of known volumes of sediment. Sediment chlorophyll *a*
 165 concentrations were used as an index of phytodetrital concentration (Boon and Duineveld,
 166 1998). Temporal changes in chlorophyll *a* breakdown products (phaeopigments) and
 167 chlorophyll *a*:phaeopigment ratios were tracked as indices of phytoplankton bloom
 168 degradation.
 169 After washing the meiofauna samples onto a 38 μm sieve, fauna were extracted with Ludox
 170 TM 40 (Somerfield and Warwick, 1996). The extraction was repeated three times. The
 171 extracts were sub-sampled following the method described by Somerfield and Warwick
 172 (1996). Sub-samples of 5 % of the extracted material (containing 85 – 448 nematodes,
 173 average = 226 nematodes) were evaporated slowly in anhydrous glycerol and mounted on
 174 slides for further analysis. The numbers of individuals identified should give reasonable, and
 175 comparable, estimates. Nematodes consistently comprised > 95 % of total meiofauna

abundance in all samples. Temporal dynamics of nematode communities in the Silver Pit were investigated by analysing changes in their taxonomic and trophic composition as well as changes in their size distribution.

2.3.1. Taxonomic analysis

All nematodes in each sub-sample were identified to genus or species level, using the taxonomic keys of Platt and Warwick (1983, 1988) and Warwick et al. (1998) and the online information system NeMys (Deprez, 2005; www.nemys.ugent.be). These are based on a phylogenetic classification scheme, where the structure of the taxonomic tree approximates the evolutionary relationships among organisms. Changes in phylogenetic diversity of species assemblages, however, are not explicitly linked to changes in functional diversity and so their ecological significance can be difficult to assess (Walker et al., 1999; Warwick and Clarke, 2001)

2.3.2. Trophic analysis

Despite the uniform organisation of free-living nematodes, a great diversity in mouth structure exists which reflects the ecological role of the species (Heip et al., 1985). Based on these morphological characteristics, Wieser (1953) devised a classification of feeding types including selective deposit feeders consuming bacteria and small-sized organic particles, non-selective deposit feeders also feeding on organic deposits but targeting larger-sized particles, epigrowth feeders scraping food off surfaces or feeding on diatoms and microalgae and predators feeding on nematodes and other small invertebrates. The disadvantage of this scheme is that it confines species to a single trophic role when many non-selective deposit feeders are known to be omnivores. For example, while the omnivores may feed on particles of a restricted size, their food can include bacteria, small flagellates and non-living aggregates. Moreover, many selective deposit and epigrowth feeders feed on diatoms but these are only numerous in shallow sediments and little is known about their diet in deeper sediments (Heip et al., 1985) and some genera classed as predators may scavenge dead

organic material as juveniles. Thus various authors have revised and modified Wieser's classification (Romeyn and Bouwman, 1983; Jensen, 1987; Moens and Vincx, 1997; Moens et al., 2004). However, in the absence of experimental evidence to describe the feeding habits of most species, Wieser's scheme remains the most feasible trophic classification of free-living marine nematodes from a variety of habitats.

2.3.3. Size-based analysis

Many biological and ecological processes are related to body size (Peters, 1983; Brown et al., 2004). In contrast to taxonomic and functional approaches, size-based analyses assume that all organisms of the same size, independent of their taxonomic or functional identity, will behave in approximately the same way in bioenergetic terms.

Individual body sizes were determined for nematodes in three of the four faunal replicates. One percent of all nematodes in each replicate (corresponding to a total of 2454 individuals), irrespective of species identity, were randomly selected for the calculation of individual body mass from allometric relationships linking body dimensions to mass. Nematode length (except long, filiform tails where present) and maximum width were measured using image analysis (Image Pro Plus version 4.5). The body mass of individuals was then calculated with the formulae of Andrassy (1956). Although more laborious, this volumetric, as opposed to a gravimetric, method was chosen since it is non-destructive and keeps measured specimens intact for taxonomic identification and the analysis of size distributions (Udalov et al., 2005). Nematode size distributions were described for each sample by assigning individuals to log₂ body-mass classes and calculating the proportion of all individuals in each class.

The similarities percentages procedure (SIMPER; Clarke, 1993) was used to identify species that consistently contributed to spatio-temporal differences between assemblages. These included the epigrowth feeder *Spilophorella paradoxa*, the non-selective deposit feeder *Sabatieria punctata* and the selective deposit feeder *Leptolaimus elegans* and individual body mass was therefore also estimated for these species separately. The percentages of organisms measured were 3 % of all *Sabatieria punctata* (corresponding to a total of 1479 individuals)

and 5 % of all *Spilophorella paradoxa* and *Leptolaimus elegans* (corresponding to a total of 372 and 1129 individuals respectively).

2.4. Coupling of sediment and water column chlorophyll *a*

An archive of surface chlorophyll *a* concentration images, derived from MODIS-Aqua satellite ocean colour data using a modified version of the case-II water algorithm 'OC5' (Gohin et al., 2002), was processed and supplied by Ifremer, France. This regional algorithm differs from generic, global satellite chlorophyll products in that additional sensor bands are used to correct for the presence of suspended sediments and the chromophoric (coloured) fraction of dissolved organic matter (cDOM). The surface chlorophyll maps cover the north-west Atlantic shelf at a resolution of 1 km² and were made available for the time period June 2002 to present by the EU/ESA project MarCoast (Marine and Coastal Environmental Information Services; www.gmes-marcoast.com). Maps in the archive were queried with the 'Zonal Statistics' function of ArcMap version 9.2 in order to extract discrete chlorophyll *a* values for the surface waters of the sampling stations A, B and C (Fig. 1). This approach provided a time-series of surface chlorophyll *a* estimates for each station. Valid, cloud-free estimates of chlorophyll *a* were obtained on approximately 23 % of all days during the time period of the study. The pelagic chlorophyll *a* values were then assembled, converted to monthly means and compared with chlorophyll *a* concentrations in the sediment.

2.5. Data processing

2.5.1. Univariate methods

Spatial and temporal patterns of environmental variables were investigated using Kruskal-Wallis one-way analysis by ranks where measures are transformed to ranks, ignoring group membership. The ranks were used to test the null-hypothesis (H_0) that there was no significant

shift in the centre of the groups between stations and over time. There was no statistically significant co-variation between the environmental measures included in the analyses.

Total nematode abundance, species diversity of the entire community and diversity within the four feeding types (H') were calculated from the species abundance data (Shannon and Weaver, 1949):

$$H' = - \sum_{i=1}^S p_i (\ln p_i)$$

where p_i is the number of individuals in the i^{th} species (n_i) divided by the total number of individuals (N).

The Index of Trophic Diversity (ITD), based on the proportion of each of four feeding types was calculated following Heip et al. (1985). ITD values range from 0.25 (highest trophic diversity with the four trophic groups accounting for 25 % each) to 1.0 (lowest trophic diversity when a single feeding type is present):

$$\text{ITD} = \sum \theta^2$$

where θ is the percentage contribution of each of four feeding types according to Wieser (1953).

Average individual body mass was calculated from the body dimensions across all species present in the total nematode community and for populations of *Spilophorella paradoxa*, *Sabatieria punctata* and *Leptolaimus elegans*.

Bartlett's and Cochran's tests were used to test for homogeneity of variance before spatio-temporal variations in all community attributes were analysed by two-way analysis of variance (ANOVA). Following the detection of significant 'station' and sampling 'month'

effects, the Tukey HSD multiple comparisons test was used in pair-wise comparisons of samples. When the interaction term station x month was significant, the two-way ANOVA was broken down to a one-way ANOVA.

All univariate statistical analyses were performed using the software package Statgraphics Plus version 3.3.

2.5.2. Multivariate methods

Non-metric multi-dimensional scaling (MDS) ordination using the Bray-Curtis similarity measure (Clarke et al., 2006) was applied to the relative abundance (i.e. proportion) of species, feeding types and geometric size classes (Clarke and Warwick, 2001). The statistical significance of spatial differences (averaged over the 12-month study period) and changes in distribution patterns over time (averaged over the three stations) were assessed by two-way crossed analysis of similarities (ANOSIM; Clarke, 1993), a non-parametric analogue of the univariate ANOVA.

BIO-ENV (Clarke and Ainsworth, 1993) and BVSTEP (Clarke and Warwick, 1998) were used to investigate relationships between environmental variables and the species, feeding type and size composition of nematode communities. Spearman rank correlations (R) between resemblance matrices derived from faunal data (based on Bray-Curtis similarity) and matrices derived from various sub-sets of environmental data (based on normalised Euclidean Distance) were used to identify suites of variables that best explained observed community patterns. The significance of the correlations was determined using a permutation procedure (Stoffels et al., 2005).

All multivariate statistical analyses were performed using Primer version 6.1.5 (Clarke and Gorley, 2006).

3. Results

3.1. Benthic habitat in the Silver Pit

314

315 All stations comprised soft bottom, muddy habitat. Stations were comparable with respect to
 316 water depth (A: 63 m, B: 64 m, C: 55 m) and sediment granulometry, although grain size at
 317 station C was slightly smaller than at station A and B (Table 1). The silt-clay content of the
 318 sediment did not differ significantly between stations and seasons, nor did the content of total
 319 organic carbon (Table 2). A temporal trend in chlorophyll *a* and phaeopigment concentrations
 320 in the sediment was apparent with a distinct peak in May 2005 and to a lesser extent in
 321 September 2005 (Tables 1 and 2, Fig. 2).

322

323 *3.2. Coupling of sediment and water chlorophyll a*

324

325 Pelagic chlorophyll *a* concentrations were typically higher to the north and to the south of the
 326 Silver Pit than over the central Silver Pit (see Fig. 1 for a monthly composite image of April
 327 2005). Time series analysis showed that the spring bloom started in the first week of April in
 328 both 2005 and 2006. Water chlorophyll *a* concentration was low in summer in both years,
 329 especially at station C (Fig. 3). It is likely that a deep chlorophyll maximum is present during
 330 this period of strongly stratified conditions (Olesen, 1995; Richardson et al., 2000), but this
 331 cannot be detected by the satellite. There was an autumn bloom in both years peaking during
 332 late September and early October. The spring bloom of 2006 was more pronounced than in
 333 2005.

334 The most productive station was station A with a mean water chlorophyll *a* concentration of
 335 $0.75 (\pm 0.60) \text{ mg m}^{-3}$ over the study period. Station B was intermediate with $0.64 (\pm 0.44) \text{ mg}$
 336 m^{-3} and station C was lowest with a mean value of $0.55 (\pm 0.37) \text{ mg m}^{-3}$. Hence there was a
 337 gradient in pelagic productivity through the Silver Pit from west (high) to east (low), with
 338 station A being closest to the areas of high primary production to the north and south of the
 339 Silver Pit (Fig. 1). Higher phytoplankton abundance at the surface was reflected in higher
 340 pigment concentrations in the upper layers of the sediment, particularly during the spring

bloom of both years (Fig. 3). However, the relationship between surface production and benthic deposition differed considerably between 2005 and 2006 (Fig. 3). It is unknown to what extent the settling particles are buried in the sediment at the place they arrive on the seafloor and how much is lost to other areas by advection near the sediment surface.

The temporal variability of pulses of primary production reaching the seafloor along the gradient of pelagic productivity allowed us to investigate the relationship between nematode taxonomic, trophic and size structure and changes in food quality and quantity.

3.3. Nematode species and feeding types

The analysis of spatio-temporal differences in community-related measures (Table 3) showed that density, species diversity and trophic diversity were higher at station A and B than at station C. Similar trends were recorded for species diversity within the four feeding types (data not shown). Spatial differences in univariate community attributes were generally more pronounced than temporal trends. With the exception of trophic diversity (ITD), the magnitude and direction of spatial differences were maintained throughout the 12-month study period. ITD did not change significantly over time at station A and B and decreased within the first 9 months at station C ($p = 0.059$). Communities at station C were most trophically diverse in February 2006.

Across all three stations, proportions of the epigrowth feeder *Spilophorella paradoxa* were consistently lower in winter (November 2005) than in spring (May 2005 and April 2006, Table 3, Fig. 4). The non-selective deposit feeder *Sabatieria punctata* dominated nematode communities at station B. Spatial distributions of *Leptolaimus elegans* were not consistent over time nor were temporal dynamics of this species consistent among stations. While relative abundance remained temporally stable at station B and C, values at station A were significantly increased in November 2005 following the autumn bloom (Table 3).

As shown in Table 4, the relative abundance of selective deposit feeders ($F = 78.56$, $p < 0.001$) and predators ($F = 15.43$, $p < 0.001$) were lowest at station C whereas communities at

station A were characterised by relatively low proportions of epigrowth feeders ($F = 26.75$, $p < 0.001$). Lowest proportions of non-selective deposit feeders were recorded at station B ($F = 6.12$, $p = 0.004$). Temporal patterns of nematode feeding types were inconsistent amongst stations and generally not statistically significant at $p > 0.05$.

There were significant spatial differences in distribution patterns of nematode species ($R = 0.583$, $p = 0.001$) and feeding guilds ($R = 0.492$, $p = 0.001$) and these generally dominated temporal dissimilarities (species assemblages: $R = 0.213$, $p = 0.001$, trophic groups $R = 0.065$, $p = 0.089$; Fig. 5).

Distribution patterns of nematode species were correlated with grain size, total organic carbon content in the sediment and sedimentary chlorophyll *a*:phaeopigment ratios ($R = 0.533$, $p = 0.010$). Spatial differences in species distribution patterns were most marked in May 2005 when station-specific differences in chlorophyll *a*:phaeopigment ratios were highest (Table 1). A combination of grain size and total organic carbon content in the sediment best explained the trophic structure of assemblages ($R = 0.424$, $p = 0.010$). Spatial differences in the trophic structure of nematode assemblages were most pronounced in April 2006 and coincided with low levels of total organic carbon in the sediment (Table 1).

3.4. Nematode body size

Spatially significant differences in individual body mass across the entire community included a higher average individual body mass at station C than at station A and B whereas the opposite trend was observed for populations of the epigrowth feeder *Spilophorella paradoxa* and the non-selective deposit feeder *Sabatieria punctata* (Table 5). Average individual body mass remained stable over the 12-month study period for the total community and for *Sabatieria punctata* and *Leptolaimus elegans* populations. Individual body mass of *Spilophorella paradoxa*, in contrast, peaked in September 2005 at all stations (Fig. 6).

In contrast to functional analyses, the investigation of the distribution of both absolute (not shown) and relative nematode abundance across body size classes revealed significant

temporal differences. Conversely, spatial differences were less pronounced when the total community was taken into account (Table 6). We observed a shift from larger to smaller individuals over the 12-month production cycle but this could not be linked to any of the measured environmental variables ($p > 0.05$).

At all three sampling stations, populations of the epigrowth feeder *Spilophorella paradoxa* were dominated by mature adults, as indicated by an abundance peak in size class -1 (Fig. 7). Significant spatial and temporal dissimilarities in the body size distribution (Table 6) were clearly related to differences in the total organic carbon content and the progression of chlorophyll *a* decomposition (i.e. chlorophyll *a*:phaeopigment ratio) in the sediment throughout the annual production cycle ($R = 0.368$, $p = 0.040$). There was a clear temporal succession in the size distribution of individuals in the population of *Spilophorella paradoxa* (Fig. 8) following deposition of the spring phytoplankton bloom in May 2005 and its decomposition over the following 11 months. Relatively high levels of both fresh (i.e. high chlorophyll *a*:phaeopigment ratios) and refractory (i.e. total organic carbon content) organic material in the sediment in May 2005 coincided with approximately equal proportions of juveniles and adults in the population, indicating continuous asynchronous breeding. Adults clearly dominated in July and September 2005 when sub-surface temperatures peaked and food availability was still high. The abundance of small individuals increased and that of larger individuals decreased over the winter months (November 2005 to February 2006) as both sub-surface temperatures and carbon resources in the sediment declined. The distribution of individuals among size classes was most homogeneous in April 2006, prior to the deposition of material from the next spring bloom.

4. Discussion

It has been estimated that sedimentation of organic matter accounts for approximately half of net primary production in stratified parts of UK shelf seas (Jago et al., 2007). Here we show from paired observations of pelagic and benthic pigment concentration that the degree of

transfer between surface and sediment is not constant, and can differ greatly between locations and years. The approximately two-fold higher concentration of phytoplankton in the water column over the Silver Pit during April 2006 did not result in enhanced benthic chlorophyll or carbon concentrations. This could be due to the combined effects of differences in (1) the rate of pelagic consumption (e.g. grazing), (2) local hydrodynamics, resulting in sedimentation of the phytoplankton at other locations and (3) the timing of benthic sample collection (i.e. time elapsed following the pelagic phytoplankton bloom). In the absence of light measurements and live observation of microalgal cells, *in situ* benthic primary production in the Silver Pit cannot be ruled out (McGee et al., 2008). However, investigating sediments from the Dogger Bank in the central North Sea, Reiss et al. (2007) showed that stations below 40 m water depth were characterised by a phytopigment composition originating from pelagic phytoplankton settled on the sea floor.

Nematode assemblages were studied at three stations in the Silver Pit area of the central North Sea during a 12-month production cycle to identify the relative importance of different environmental factors (including the quality and quantity of organic material reaching the seafloor) in controlling their taxonomic, trophic and biomass distribution patterns. Our data suggest that spatio-temporal differences in nematode species composition were partly a response to the sedimentation of pelagic production and appeared to reflect species-specific responses to its nutritional quality. These correlative relationships provide valuable pointers to the mechanisms potentially causing the observed variation.

4.1. Does a local increase in food promote consistent temporal changes in the taxonomic, trophic and size structure of nematode communities?

The taxonomic and trophic structure of nematode communities differed significantly among stations over relatively short distances, potentially masking some of the temporal dynamics. Nematode assemblages appeared to respond to spatial gradients of declining grain size and pelagic productivity in the Silver Pit from west to east as evidenced by decreasing density,

species diversity and trophic diversity. Taking the entire community into account, average individual body size increased with decreasing grain size and pelagic productivity. This result was counter to the analysis of Udalov et al. (2005) who investigated the relationship between mean nematode individual weight, sediment granulometry and food availability from 120 worldwide locations in the depth range 0 – 8260 m. In their analysis, nematode size increased with particle diameter and the concentration of sediment-bound chloroplastic pigments. However, the overall relationships masked more complex patterns within habitat types. Variability of generally more diverse, coarser sediments at shallower depths compared to fluid clays and oozes in the deep sea were mainly responsible for the changes in the nematode size structure from shelf to slope. Thus, the relationships between sediment granulometry and nematode size were strong for intertidal and shallow subtidal habitats but weaker for deeper subtidal areas. Equally, although Udalov et al. (2005) found evidence for decreases in the size of nematodes with decreasing food availability, the correlation with nematode size was significant for oligotrophic areas only and weakened when organic input was sufficient to cover the higher energy consumption rates of larger species (Warwick and Price, 1979). Low pelagic productivity and small grain size at the easternmost station in the Silver Pit resulted in high proportions of large-bodied non-selective deposit and epigrowth feeders in the nematode community whereas the density and diversity of their smaller-sized bacteria feeding congeners was reduced.

Although spatial differences in species composition dominated temporal dynamics, significant temporal changes were recorded over the 12-month production cycle. These were linked to changes in the quality and quantity of organic material reaching the seabed, as indicated by the chlorophyll *a*:phaeopigment ratio and the total organic carbon content. Davies and Payne (1984) provided evidence of a large settlement of diatoms immediately after the spring bloom in the northern North Sea, based on changes in the biochemical composition of material collected in moored sediment traps. The significant relationship between species distribution patterns and the state of phytoplankton remineralisation might thus point towards a species-specific response to the nutritional quality of sedimenting

organic material and the biochemical changes in the sediment associated with its decomposition (e.g. oxygenation, alterations in decomposer communities such as bacteria etc.).

There was no consistent temporal relationship between the trophic composition of nematode communities and spring chlorophyll *a* or carbon sedimentation, most likely a result of the trophic plasticity of most feeding types. Thus different species belonging to one trophic guild may be able to forage on different food sources and/or exploit their food at different rates (De Mesel et al., 2004; Postma-Blaauw et al., 2005). Investigating the uptake of a radio-labelled diatom by meiofauna species in a laboratory experiment, Ólafsson et al. (1999) found that nematodes with similar buccal cavities and of similar size showed surprisingly large differences in uptake, suggesting a strong selection both for and against the diatom among epigrowth and non-selective deposit feeders. Likewise, also in a laboratory study, Moens et al. (1999) demonstrated a highly species-specific response of bacteria feeding nematodes to different types, densities and age structures of bacterial strains.

4.2. Do selected nematode species, representing the main taxonomic and trophic categories available to the phylum, respond in a differential way to a pulsed food supply?

Phytoplankton sedimentation in stratified temperate marine waters is distinctly seasonal, but the majority of studies in subtidal environments found no unequivocal influence of sedimentation on the density and biomass of meiofaunal nematodes (Heip et al., 1985; Ólafsson et al., 1999). However, the apparent absence of a community response may be misleading. Phytoplankton during and after bloom conditions may settle quickly to the seafloor and these fresh cells may be largely intact and rich in nutrients (Graf, 1992; Richardson et al., 2000). Sedimented phytoplankton might be directly ingested or contribute to the microbial loop, ultimately increasing meiofaunal food supply. Fleege et al. (1989) hypothesised that this variation in phytoplankton settlement might affect meiofaunal survivorship and reproductive success. Our results suggest that populations of species,

509 assumed to feed on fresh organic material such as *Spilophorella paradoxa*, can exhibit
 510 temporal patterns in their abundance and size distribution. Given the seasonal cycle of detrital
 511 sedimentation, the balance of benthic food input and demand may shift seasonally, with
 512 spring surpluses followed by deficits later in the year. Across all three stations, relative
 513 abundance of *Spilophorella paradoxa* declined following the spring phytoplankton bloom in
 514 May 2005 and reached the lowest values the following November. This suggests that food
 515 deficits may be particularly acute for those species that feed at the sediment-water interface
 516 and depend on fresh inputs, although other unmeasured factors, such as competition for food,
 517 are also likely to influence population dynamics.

518 Temporal dynamics of *Spilophorella paradoxa* populations were clearly linked to the quality
 519 and quantity of organic material present in the sediment throughout an annual production
 520 cycle, confounded by annual fluctuations in sub-surface temperature. Temperature may
 521 trigger or terminate reproductive activities of nematodes and determine the development time
 522 of juveniles. Generally, higher ambient temperatures result in shorter development times;
 523 however, little is known of the development time of most species (Warwick and Price, 1979
 524 and references therein; Moens and Vincx, 2000a). High concentrations of fresh and refractory
 525 organic material in the sediment and rising sub-surface temperatures in the Silver Pit in May
 526 2005 resulted in approximately equal proportions of juveniles and adults in the population.
 527 Developmental time decreased as ambient temperatures increased, leading to adult-dominated
 528 populations in July and September 2005 when food availability was still sufficient to cover
 529 their metabolic demands. Declining food availability and temperatures slowed growth and,
 530 potentially, reproduction in subsequent months, leading to low numbers of individuals across
 531 the entire size spectrum in November 2005. The abundance peak of adults in summer 2005
 532 was followed by an increase of juveniles between February and April 2006 when
 533 temperatures in the sediment began to increase prior to the deposition of the spring
 534 phytoplankton bloom.

535 Many deeper dwelling, deposit-feeding nematodes, including populations of *Sabatieria*
 536 *punctata* and *Leptolaimus elegans* may not be able to use sedimenting phytoplankton as food

directly and they may rely on the microbial loop to stimulate bacterial production from sedimented material. Results from experimental studies, investigating the importance of freshly produced phytoplankton detritus in the diet of meiofauna are conflicting. Rudnick (1989) hypothesised that, in contrast to surface-dwelling nematodes, the food resources of sub-surface deposit feeders are more stable because they have access to and assimilate a portion of the buried pool of older organic matter. Consequently, one might not expect consistent temporal changes in their density and size distribution. Ólafsson et al. (1999) contradicted this hypothesis by revealing that sub-surface dwelling nematode species can assimilate as much phytoplankton carbon per unit biomass as surface dwellers. Recently, Franco et al. (2008) showed that *Sabatieria*-species can benefit from sedimenting phytoplankton by moving to the sediment surface. Patterns of food availability for deposit feeding organisms can be spatially and temporally complex and total organic carbon measurements might be inadequate to elucidate the mechanisms by which food abundance and variety regulate the organisation of deposit-feeding assemblages (Whitlatch, 1981). Whilst such measures provide information on the total standing stock of organic carbon, they generally overestimate its labile, readily utilisable fraction. At the same time, all living components (e.g. bacteria, microphytobenthos, protozoans) represent a high-quality food source but generally account for a minor fraction of the sediment organic carbon pools (Fabiano and Danovaro, 1994; Danovaro and Gambi, 2002).

5. Conclusions

Biomass-size distributions showed the most consistent temporal patterns in nematode assemblages. When applied to the whole community, the size-based approach has significant advantages over taxon-based and functional approaches through its simplicity, since it avoids distinguishing between taxonomic groups or defining distinct trophic guilds (Soetaert et al., 2002). Our results from the central North Sea show that the relationship between

phytoplankton sedimentation and meiofaunal response is complex and that community responses are weaker than those of some of the component species. Hence, size-based analyses provided the most insightful results when performed on selected species.

The differential responses within communities reflect different life histories and feeding modes and the changes in the resources available during the seasonal production cycle. The rate of response indicates that the life histories of meiofauna allow their populations to track the availability of suitable food resources. Since the feeding habits of most meiofauna species are not well known, the majority of previous studies on the structure and connectedness of aquatic food webs have ignored meiofaunal taxa or aggregated them into broad functional groups. Existing food web data are therefore strongly biased towards trophically higher organisms. Our findings suggest that future investigations of the relationship between size-structure of meiofauna populations (Vanaverbeke et al., 2004b) and potential food sources (Dauwe and Middelburg, 1998), factors regulating meiofauna fecundity (e.g. food, temperature; Moens and Vincx, 2000b) as well as novel information on species-specific feeding habits and preferences (Moens and Vincx, 1999) are crucial to the understanding of their role in food webs.

Acknowledgements

We thank the scientists, officers and crew of RV 'Cefas Endeavour' for practical assistance in the collection of samples. Claire Mason conducted the particle size analyses and Olga Andres carried out phytopigment analyses. We are grateful to Paul Whomersley and Tracy Maxwell for processing the meiofauna samples. The work was supported by the Department for Environment, Food and Rural Affairs (research projects MF10 and ME4118).

References

- 592 Andrassy, I., 1956. Die Rauminhalts- und Gewichtsbestimmung der Fadenwürmer
593 (Nematoden). *Acta Zoologica* 2, 1–14.
- 594 Boon, A.R., Duineveld, G.C.A., 1998. Chlorophyll *a* as a marker for bioturbation and carbon
595 flux in southern and central North Sea sediments. *Marine Ecology Progress Series* 162,
596 33–43.
- 597 Boon, A.R., Duineveld, G.C.A., Berghuis, E.M., van der Weele, J.A., 1998. Relationships
598 between benthic activity and the annual phytopigment cycle in near-bottom water and
599 sediments in the southern North Sea. *Estuarine, Coastal and Shelf Science* 46, 1–13.
- 600 Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M., West, G.B., 2004. Towards a
601 metabolic theory of ecology. *Ecology* 85, 1771–1789.
- 602 Chalcraft, D.R., Resetarits, W.J., 2003. Mapping functional similarity of predators on the
603 basis of trait similarities. *American Naturalist* 162, 390–402.
- 604 Chardy, P., Dauvin, J.-P., 1992. Carbon flows in a subtidal fine sand community from the
605 western English Channel: a simulation analysis. *Marine Ecology Progress Series* 81, 147–
606 161.
- 607 Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure.
608 *Australian Journal of Ecology* 18, 117–143.
- 609 Clarke, K.R., Ainsworth, M., 1993. A method of linking multivariate community structure to
610 environmental variables. *Marine Ecology Progress Series* 92, 205–219.
- 611 Clarke, K.R., Warwick, R.M., 1998. Quantifying structural redundancy in ecological
612 communities. *Oecologia* 113, 278–289.
- 613 Clarke, K.R., Warwick, R.M., 2001. Change in marine communities: an approach to statistical
614 analysis and interpretation. 2nd edition, Primer-E Ltd, Plymouth, UK, 172 pp.
- 615 Clarke, K.R., Gorley, R.N., 2006. PRIMER v6. User manual. Primer-E Ltd., Plymouth, UK,
616 190 pp.
- 617 Clarke, K.R., Somerfield, P.J., Chapman, M.G., 2006. On resemblance measures for
618 ecological studies, including taxonomic dissimilarities and zero-adjusted Bray-Curtis

- 619 coefficient for denuded assemblages. Journal of Experimental Marine Biology and
620 Ecology 330, 55–80.
- 621 Colebrook, J.M., Robinson, G.A., 1961. The seasonal cycle of the plankton in the North Sea
622 and the north-eastern Atlantic. Journal du Conseil Permanent International pour
623 l'Exploration de Mer 26, 156–165.
- 624 Danovaro, R., Gambi, C., 2002. Biodiversity and trophic structure of nematode assemblages
625 in seagrass systems: evidence for a coupling with changes in food availability. Marine
626 Biology 141, 667–677.
- 627 Danovaro, R., Scopa, M., Gambi, C., Fraschetti, S., 2007. Trophic importance of subtidal
628 metazoan meiofauna: evidence from *in situ* exclusion experiments on soft and rocky
629 substrates. Marine Biology 152, 339–350.
- 630 Dauwe, B., Middelburg, J.J., 1998. Amino acids and hexosamines as indicators of organic
631 matter degradation state in North Sea sediments. Limnology and Oceanography 43, 782–
632 789.
- 633 Davies, J.M., Payne, R., 1984. Supply of organic matter to the sediment in the northern North
634 Sea during a spring phytoplankton bloom. Marine Biology 78, 315–324.
- 635 De Mesel, I., Derycke, S., Moens, T., Van der Gucht, K., Vincx, M., Swings, J., 2004. Top-
636 down impact of bacterivorous nematodes on the bacterial community structure: a
637 microcosm study. Environmental Microbiology 6, 733–744.
- 638 Deprez, T., 2005. NeMys. World Wide Web electronic publication, www.nemys.ugent.be,
639 version (2/2008).
- 640 Dyer, K.R., 1986. Coastal and estuarine sediment dynamics. John Wiley & Sons, Chichester,
641 UK, 342 pp.
- 642 Fabiano, M., Danovaro, R., 1994. Composition of organic matter in sediments facing a river
643 estuary (Tyrrhenian Sea): relationships with bacteria and microphytobenthic biomass.
644 Hydrobiologia 277, 71–84.
- 645 Findlay, S.E.G., 1982. Effect of detrital nutritional quality on population dynamics of a
646 marine nematode (*Diplolaimella chitwoodi*). Marine Biology 68, 223–227.

- 647 Fleege, J.W., 1985. Meiofaunal densities and copepod species composition in a Louisiana,
648 USA, estuary. *Transactions of the American Microscopical Society* 104, 321–332.
- 649 Fleege, J.W., Shirley, T.C., Ziemann, D.A., 1989. Meiofaunal responses to sedimentation
650 from an Alaskan spring bloom. I. Major taxa. *Marine Ecology Progress Series* 57, 137–
651 145.
- 652 Franco, M.A., De Mesel, I., Demba Diallo M., Van der Gucht, K., Van Gansbeke, D., van
653 Rijswijk, P., Costa, M.J., Vincx, M., Vanaverbeke, J., 2007. Effect of phytoplankton
654 bloom deposition on benthic bacterial communities in two contrasting sediments in the
655 southern North Sea. *Aquatic Microbial Ecology* 48, 241–254.
- 656 Franco, M.A., Soetaert, K., Van Oevelen, D., Van Gansbeke, D., Costa, M.J., Vincx, M.,
657 Vanaverbeke, J., 2008. Density, vertical distribution and trophic responses of metazoan
658 meiobenthos to phytoplankton deposition in contrasting sediment types. *Marine Ecology*
659 *Progress Series* 358, 51–62.
- 660 Giere, O., 1993. *Meiobenthology - the microscopic fauna in aquatic sediments*. Springer
661 Verlag, Berlin, Germany, 328 pp.
- 662 Gohin, F., Druon, J.N., Lampert, L., 2002. A five channel chlorophyll concentration
663 algorithm applied to Sea WiFS data processed by SeaDAS in coastal waters. *International*
664 *Journal of Remote Sensing* 23, 1639–1661.
- 665 Graf, G., Bengtsson, J., Diesner, U., Schulz, R., Theede, H., 1982. Benthic response to
666 sedimentation of a spring phytoplankton bloom: process and budget. *Marine Biology* 67,
667 201–208.
- 668 Graf, G., 1992. Benthic-pelagic coupling: a benthic view. *Oceanography and Marine Biology:*
669 *an Annual Review* 30, 149–190.
- 670 Henrichs, S.M., Doyle, A.P., 1986. Decomposition of ^{14}C -labelled organic substances in
671 marine sediments. *Limnology and Oceanography* 31, 765–778.
- 672 Heip, C., Vincx, M., Vranken, G., 1985. The ecology of marine nematodes. *Oceanography*
673 *and Marine Biology: an Annual Review* 23, 399–489.

- 674 Jago, C.F., Kennaway, G.M., Novarino, G., Jones, S.E., 2007. Size and settling velocity of
675 suspended flocs during a *Phaeocystis* bloom in the tidally stirred Irish Sea, NW European
676 shelf. Marine Ecology Progress Series 345, 51–62.
- 677 Jensen, P., 1987. Feeding ecology of free-living aquatic nematodes. Marine Ecology Progress
678 Series 35, 187–196.
- 679 McGee, D., Laws, R.A., Cahoon, L. B., 2008. Live benthic diatoms from the upper
680 continental slope: extending the limits of marine primary production. Marine Ecology
681 Progress Series, 356, 103–112.
- 682 Mikola, J., 1998. Effects of microbivore species composition and basal resource enrichment
683 on trophic-level biomasses in an experimental microbial-based soil food web. Oecologia
684 117, 396–403.
- 685 Moens, T., Vincx, M., 1997. Observations on the feeding ecology of estuarine nematodes.
686 Journal of the Marine Biological Association of the United Kingdom 77, 211–227.
- 687 Moens, T., Vincx, M., 1999. Do meiofauna consume primary production? About many
688 questions and how to answer them, in: Dehairs, B.J.F., Goeyens, L. (Eds.), Integrated
689 Marine Systems Analysis, European Network for Integrated Marine Systems Analysis,
690 Minutes of the First Network Meeting, Brugge, Belgium, pp. 188–202.
- 691 Moens, T., Verbeek, L., de Maeyer, A., Swings, J., Vincx, M., 1999. Selective attraction of
692 marine bacterivorous nematodes to their bacterial food. Marine Ecology Progress Series
693 176, 165–178.
- 694 Moens, T., Vincx, M., 2000a. Temperature and salinity constraints on the life cycle of two
695 brackish-water nematode species. Journal of Experimental Marine Biology and Ecology,
696 243, 115–135.
- 697 Moens, T., Vincx, M., 2000b. Temperature, salinity and food thresholds in two brackishwater
698 bacterivorous nematode species: assessing niches from food absorption and respiration
699 experiments. Journal of Experimental Marine Biology and Ecology 243, 137–154.
- 700 Moens, T., Yeates, G.W., De Ley, P., 2004. Use of carbon and energy sources by nematodes.
701 Nematology Monographs and Perspectives 2, 529–545.

- Ólafsson, E., Elmgren, R., 1997. Seasonal dynamics of sublittoral meiobenthos in relation to phytoplankton sedimentation in the Baltic Sea. *Estuarine, Coastal and Shelf Science* 45, 149–164.
- Ólafsson, E., Modig, H., van de Bund, W.J., 1999. Species specific uptake of radio-labelled phytodetritus by benthic meiofauna from the Baltic Sea. *Marine Ecology Progress Series* 177, 63–72.
- Olesen, M., 1995. Comparison of the sedimentation of a diatom spring bloom and of a subsurface chlorophyll maximum. *Marine Biology* 121, 541–547.
- Peters, R.H., 1983. The ecological implications of body size. Cambridge University Press, Cambridge, UK, 329 pp.
- Platt, H.M., Warwick, R.M., 1983. Free-living marine nematodes. Part I. British enoploids. Cambridge University Press, Cambridge, UK, 307 pp.
- Platt, H.M., Warwick, R.M., 1988. Free-living marine nematodes. Part II. British chromadorids. Cambridge University Press, Cambridge, UK, 502 pp.
- Pope, J.G., Shepherd, J., Webb, J., 1994. Successful surf-riding on size spectra: the secret of survival in the sea. *Philosophical Transactions of the Royal Society* 343, 41–49.
- Postma-Blaauw, M.B., de Vries, F.T., de Goede, R.G.N., Bloem, J., Faber, J.H., Brussaard, L., 2005. Within-trophic group interactions of bacterivorous nematode species and their effects on the bacterial community and nitrogen mineralization. *Oecologia* 142, 428–439.
- Reiss, H., Wiekling, G., Kröncke, I., 2007. Microphytobenthos of the Dogger Bank: a comparison between shallow and deep areas using phytopigment composition of the sediment. *Marine Biology* 150, 1061–1071.
- Richardson, K., Visser, A.W., Pedersen, F.B., 2000. Subsurface phytoplankton blooms fuel pelagic production in the North Sea. *Journal of Plankton Research* 22, 1663–1671.
- Romeyn, K., Bouwman, L.A., 1983. Food selection and consumption by estuarine nematodes. *Hydrobiological Bulletin* 17, 103–109.
- Rudnick, D.T., 1989. The lags between the deposition and meiobenthic assimilation of phytodetritus. *Marine Ecology Progress Series* 50, 231–240.

- 730 Schmid-Araya, J.M., Hildrew, A.G., Robertson, A., Schmid, P.E., Winterbottom, J., 2002.
731 The importance of meiofauna in food webs: evidence from an acid stream. *Ecology* 83,
732 1271–1285.
- 733 Schratzberger, M., Dinmore, T.A., Jennings, S., 2002. Impacts of trawling on the diversity,
734 biomass and structure of meiofauna assemblages. *Marine Biology* 140, 83–93.
- 735 Schratzberger, M., Warr, K., Rogers, S.I., 2007. Functional diversity of nematode
736 communities in the southwestern North Sea. *Marine Environmental Research* 63, 368–
737 389.
- 738 Shannon, C.E., Weaver, W., 1949. The mathematical theory of communication. University of
739 Illinois Press, Urbana, USA, 144 pp.
- 740 Soetaert, K., Muthumbi, A., Heip, C., 2002. Size and shape of ocean margin nematodes:
741 morphological diversity and depth-related patterns. *Marine Ecology Progress Series* 242,
742 179–193.
- 743 Somerfield, P.J., Warwick, R.M., 1996. Meiofauna in marine pollution monitoring
744 programmes. A laboratory manual. Ministry of Agriculture, Fisheries and Food,
745 Directorate of Fisheries Research, Lowestoft, UK, 71 pp.
- 746 Stoffels, R.J., Clarke, K.R., Closs, G.P., 2005. Spatial scale and benthic community
747 organisation in the littoral zones of large oligotrophic lakes: potential for cross-scale
748 interactions. *Freshwater Biology* 50, 1131–1145.
- 749 Tett, P., 1987. Plankton, in: Baker, J.M., Wolff, W.M. (Eds.), *Biological surveys of estuaries*
750 *and coasts*. Cambridge University Press, Cambridge, UK, pp. 280–341.
- 751 Udalov, A.A., Azovsky, A.I., Mokievsky, V.O., 2005. Depth-related pattern in nematode size:
752 what does the depth itself really mean? *Progress in Oceanography* 67, 1–23.
- 753 Vanaverbeke, J., Steyaert, M., Soetaert, K., Rousseau, V., van Gansbeke, D., Parent, J.-Y.,
754 Vincx, M., 2004a. Changes in structural and functional diversity of nematode
755 communities during a spring phytoplankton bloom in the southern North Sea. *Journal of*
756 *Sea Research* 52, 281–292.

- 757 Vanaverbeke, J., Soetaert, K., Vincx, M., 2004b. Changes in morphometric characteristics of
758 nematode communities during a spring phytoplankton bloom deposition. *Marine Ecology*
759 *Progress Series* 273, 139–146.
- 760 Walker, B., Kinzig, A., Landridge, J., 1999. Plant attribute diversity, resilience, and
761 ecosystem function: the nature and significance of dominant and minor species.
762 *Ecosystems* 2, 95–113.
- 763 Warwick, R.M., Price, R., 1979. Ecological and metabolic studies on free-living nematodes
764 from an estuarine mud-flat. *Estuarine and Coastal Marine Science* 9, 257–271.
- 765 Warwick, R.M., Platt, H.M., Somerfield, P.J., 1998. Free-living marine nematodes. Part III.
766 British monohysterids. Field Studies Council, Shrewsbury, UK, 296 pp.
- 767 Warwick, R.M., Clarke, K.R., 2001. Practical measures of marine biodiversity based on
768 relatedness of species. *Oceanography and Marine Biology: an Annual Review* 39, 207–
769 231.
- 770 Whitlatch, R.B., 1981. Animal-sediment relationships in intertidal marine benthic habitats:
771 some determinants of deposit-feeding species diversity. *Journal of Experimental Marine*
772 *Biology and Ecology* 53, 31–45.
- 773 Wieser, W., 1953. Die Beziehung zwischen Mundhöhlengestalt, Ernährungsweise und
774 Vorkommen bei freilebenden marinen Nematoden. *Arkiv för Zoologi* 4, 439–484.

775 Table 1

776 Sediment properties (mean $\pm$ SE) at the three stations in the Silver Pit over a 12-month production cycle. TOC = total organic carbon, Chl *a* = chlorophyll *a*,

777 Phaeo = phaeopigment

Station	Date	Grain size [μm]	Silt-clay [%]	TOC [%]	Chl <i>a</i> [mg m^{-2}]	Phaeo [mg m^{-2}]	Chl <i>a</i> :Phaeo
A	21 – 26 May 2005	106	21	0.38	89 ± 16	111 ± 40	1.2 ± 0.5
	12 – 16 Jul 2005	98	24	0.49	21 ± 10	46 ± 2	0.5 ± 0.2
	29 Sep – 4 Oct 2005	99	26	0.56	31 ± 7	73 ± 3	0.4 ± 0.1
	5 – 10 Nov 2005	104	22	0.44	7 ± 1	22 ± 2	0.3 ± 0.1
	13 – 17 Feb 2006	86	28	0.51	21 ± 2	51 ± 8	0.5 ± 0.1
	26 Apr – 5 May 2006	84	29	0.20	16 ± 5	67 ± 13	0.2 ± 0.1
B	21 – 26 May 2005	99	21	0.31	26 ± 4	113 ± 10	0.2 ± 0.1
	12 – 16 Jul 2005	83	29	0.36	9 ± 4	25 ± 8	0.3 ± 0.1
	29 Sep – 4 Oct 2005	85	28	0.39	29 ± 4	67 ± 13	0.5 ± 0.1
	5 – 10 Nov 2005	90	26	0.36	8 ± 1	29 ± 2	0.3 ± 0.1
	13 – 17 Feb 2006	92	22	0.33	12 ± 1	46 ± 2	0.3 ± 0.0
	26 Apr – 5 May 2006	97	24	0.33	12 ± 3	53 ± 4	0.2 ± 0.0
C	21 – 26 May 2005	71	26	0.47	72 ± 38	54 ± 27	2.5 ± 1.6
	12 – 16 Jul 2005	78	25	0.34	11 ± 1	37 ± 6	0.3 ± 0.0
	29 Sep – 4 Oct 2005	69	28	0.34	18 ± 6	54 ± 4	0.3 ± 0.1
	5 – 10 Nov 2005	67	29	0.35	11 ± 1	25 ± 3	0.4 ± 0.0
	13 – 17 Feb 2006	73	24	0.44	6 ± 1	32 ± 2	0.2 ± 0.0
	26 Apr – 5 May 2006	74	25	0.33	8 ± 3	42 ± 20	0.2 ± 0.1

778

779 Table 2

780 Results from Kruskal-Wallis one-way analysis by ranks: Spatio-temporal differences in
 781 sediment properties

Source	Test statistic	<i>p</i> -value
Grain size		
Station	11.94	0.003
Month	1.09	0.950
Silt-clay		
Station	0.92	0.630
Month	4.13	0.530
Total organic carbon		
Station	3.82	0.148
Month	5.88	0.318
Chlorophyll <i>a</i>		
Station	3.43	0.180
Month	27.41	< 0.001
Phaeopigment		
Station	4.95	0.084
Month	25.80	< 0.001
Chlorophyll <i>a</i> :Phaeopigment		
Station	2.53	0.282
Month	8.72	0.121

782

Table 3
Results from two-way analysis of variance (ANOVA): Spatio-temporal differences in
nematode community-related measures and the relative abundance of discriminating species

Source	Sum of squares	Df	Mean square	F-Ratio	p-value
Total abundance (Log)					
Station	0.2681	2	0.1341	8.90	0.001
Month	0.1532	5	0.0306	2.03	0.088
Station x Month	0.1523	10	0.0152	1.01	0.446
Species diversity (H')					
Station	2.9267	2	1.4634	40.36	< 0.001
Month	0.1543	5	0.0309	0.85	0.520
Station x Month	0.4608	10	0.0461	0.05	0.270
Trophic diversity (ITD)					
Station	0.0627	2	0.0331	90.93	< 0.001
Month	0.0034	5	0.0008	2.28	0.059
Station x Month	0.0084	10	0.00084	2.42	0.018
<i>Spilophorella paradoxa</i> [%]					
Station	8.7173	2	4.3587	1.41	0.253
Month	47.5376	5	9.5075	3.08	0.016
Station x Month	38.3231	10	3.8323	1.24	0.288
<i>Sabatieria punctata</i> [%]					
Station	406.4300	2	203.2150	8.97	< 0.001
Month	121.2810	5	24.2562	1.07	0.387
Station x Month	256.3180	10	25.6318	1.13	0.357
<i>Leptolaimus elegans</i> [%]					
Station	452.2720	2	226.1360	33.76	< 0.001
Month	150.2450	5	30.0491	4.49	0.002
Station x Month	137.5660	10	13.7566	2.05	0.045

787 Table 4

788 Mean ($\pm$ SE) relative abundance of nematode feeding types (sensu Wieser, 1953) at the three
 789 stations in the Silver Pit over a 12-month production cycle

Station	Date	Selective deposit feeders [%]	Non- selective deposit feeders [%]	Epigrowth feeders [%]	Predators [%]
A	21 – 26 May 2005	21 $\pm$ 1	43 $\pm$ 1	29 $\pm$ 1	6 $\pm$ 0
	12 – 16 Jul 2005	18 $\pm$ 1	38 $\pm$ 2	37 $\pm$ 3	7 $\pm$ 1
	29 Sep – 4 Oct 2005	22 $\pm$ 1	42 $\pm$ 2	31 $\pm$ 0	5 $\pm$ 1
	5 – 10 Nov 2005	29 $\pm$ 2	37 $\pm$ 1	28 $\pm$ 4	7 $\pm$ 2
	13 – 17 Feb 2006	24 $\pm$ 1	42 $\pm$ 2	28 $\pm$ 2	6 $\pm$ 1
	26 Apr – 5 May 2006	20 $\pm$ 2	41 $\pm$ 2	32 $\pm$ 1	7 $\pm$ 1
B	21 – 26 May 2005	17 $\pm$ 2	35 $\pm$ 4	40 $\pm$ 5	8 $\pm$ 1
	12 – 16 Jul 2005	15 $\pm$ 2	35 $\pm$ 1	46 $\pm$ 2	5 $\pm$ 1
	29 Sep – 4 Oct 2005	18 $\pm$ 2	37 $\pm$ 6	42 $\pm$ 5	4 $\pm$ 0
	5 – 10 Nov 2005	15 $\pm$ 2	44 $\pm$ 1	36 $\pm$ 3	5 $\pm$ 1
	13 – 17 Feb 2006	17 $\pm$ 1	34 $\pm$ 3	44 $\pm$ 3	5 $\pm$ 1
	26 Apr – 5 May 2006	18 $\pm$ 1	33 $\pm$ 3	45 $\pm$ 4	4 $\pm$ 1
C	21 – 26 May 2005	8 $\pm$ 2	44 $\pm$ 4	45 $\pm$ 4	2 $\pm$ 1
	12 – 16 Jul 2005	10 $\pm$ 1	46 $\pm$ 2	42 $\pm$ 3	2 $\pm$ 0
	29 Sep – 4 Oct 2005	11 $\pm$ 1	43 $\pm$ 4	44 $\pm$ 4	2 $\pm$ 1
	5 – 10 Nov 2005	11 $\pm$ 0	39 $\pm$ 4	45 $\pm$ 4	5 $\pm$ 0
	13 – 17 Feb 2006	15 $\pm$ 3	40 $\pm$ 4	40 $\pm$ 2	5 $\pm$ 1
	26 Apr – 5 May 2006	11 $\pm$ 2	41 $\pm$ 5	42 $\pm$ 5	6 $\pm$ 1

790

791 Table 5

792 Results from two-way analysis of variance (ANOVA): Spatio-temporal differences in average
 793 individual body mass in the total nematode community and for populations of discriminating
 794 species

Source	Sum of squares	Df	Mean square	F-Ratio	p-value
Total community					
Station	0.280737	2	0.140369	3.45	0.043
Month	0.320326	5	0.0640652	1.57	0.193
Station x Month	0.490307	10	0.0490307	1.2	0.321
<i>Spilophorella paradoxa</i>					
Station	0.322737	2	0.161369	4.15	0.024
Month	0.481504	5	0.0963007	2.48	0.050
Station x Month	0.339041	10	0.0339041	0.87	0.566
<i>Sabatieria punctata</i>					
Station	0.271078	2	0.135539	4.23	0.022
Month	0.105911	5	0.0211822	0.66	0.656
Station x Month	0.316011	10	0.0316011	0.99	0.473
<i>Leptolaimus elegans</i>					
Station	0.00149259	2	0.000746296	0.41	0.667
Month	0.0101259	5	0.00202519	1.11	0.372
Station x Month	0.0280407	10	0.00280407	1.54	0.166

795

796 Table 6

797 Results from two-way crossed analysis of similarities (ANOSIM): Spatio-temporal
 798 differences in the size distribution of the total nematode community and populations of
 799 discriminating species

	Station		Month	
	Spearman R	<i>p</i> -value	Spearman R	<i>p</i> -value
Total community	0.111	0.074	0.123	0.032
<i>Spilophorella paradoxa</i>	0.159	0.009	0.282	0.001
<i>Sabatieria punctata</i>	0.126	0.060	0.033	0.312
<i>Leptolaimus elegans</i>	0.050	0.221	0.001	0.471

800

Figure captions

Fig. 1. Sampling stations in the Silver Pit (A, B and C), with background showing the mean monthly water chlorophyll *a* concentration for April 2005. The 25 m depth contour is shown as a solid white line, and the 50 m contour as a solid black line.

Fig. 2. Temporal changes in the mean ($\pm$ SE) sediment chlorophyll *a* and phaeopigment concentrations across the three stations in the Silver Pit.

Fig. 3. Comparison of mean pelagic and benthic chlorophyll *a* concentrations in each of the sampling months during 2005 and 2006 coded by sampling station. Shaded areas identify the spring bloom period showing the gradient in pelagic primary production throughout the Silver Pit from west (station A) to east (station C).

Fig. 4. Temporal changes in the mean ($\pm$ SE) relative abundance of *Spilophorella paradoxa* across the three stations in the Silver Pit.

Fig. 5. Non-metric multi-dimensional scaling (MDS) ordination of mean relative abundance of nematode species and feeding types (sensu Wieser 1953) at the three stations in the Silver Pit over a 12-month production cycle. Day of year in parentheses.

Fig. 6. Temporal changes in the mean ($\pm$ SE) individual body mass of *Spilophorella paradoxa* across the three stations in the Silver Pit.

Fig. 7. Spatial differences in the mean ($\pm$ SE) absolute abundance of *Spilophorella paradoxa* over a 12-month production cycle plotted against geometric size class.

828 Fig. 8. Temporal changes in the mean ($\pm$ SE) absolute abundance of *Spilophorella paradoxa*
829 in geometric size class -4 to class 0 across the three stations in the Silver Pit.

ACCEPTED MANUSCRIPT

Figure

Fig. 1

Fig. 3

Fig. 5

Fig. 7

Fig. 8