Accepted Manuscript

A weight of evidence approach for quality assessment of sediments impacted by an oil spill: The role of a set of biomarkers as a line of evidence

Carmen Morales-Caselles, Inmaculada Riba, Carmen Sarasquete, T. Ángel DelValls

PII: S0141-1136(08)00219-5
DOI: [10.1016/j.marenvres.2008.10.003](https://doi.org/10.1016/j.marenvres.2008.10.003)
Reference: MERE 3295

To appear in: *Marine Environmental Research*

Received Date: 3 April 2008
Revised Date: 13 October 2008
Accepted Date: 15 October 2008

Please cite this article as: Morales-Caselles, C., Riba, I., Sarasquete, C., DelValls, T., A weight of evidence approach for quality assessment of sediments impacted by an oil spill: The role of a set of biomarkers as a line of evidence, *Marine Environmental Research* (2008), doi: [10.1016/j.marenvres.2008.10.003](https://doi.org/10.1016/j.marenvres.2008.10.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**A WEIGHT OF EVIDENCE APPROACH FOR QUALITY ASSESSMENT OF
SEDIMENTS IMPACTED BY AN OIL SPILL: THE ROLE OF A SET OF
BIOMARKERS AS A LINE OF EVIDENCE**

Carmen Morales-Caselles^{1,2,*}, Inmaculada Riba^{1,2}, Carmen Sarasquete¹, T. Ángel
DelValls^{1,2}

¹ Unidad Asociada de Calidad Ambiental y Patología (CSIC & UCA) instituto de
Ciencias Marinas de Andalucía (ICMAN-CSIC), Avda. República Saharaui s/n, Puerto
Real 11510, Cádiz, Spain

² UNESCO UNITWIN/WiCop, Facultad de Ciencias del Mar y Ambientales,
Universidad de Cádiz, Polígono Río San Pedro s/n, Puerto Real 11510, Cádiz, Spain.

Corresponding author: Carmen Morales Caselles

carmen.morales@uca.es

Phone: +34 956016740

Fax: +34 956016040

Abstract

In an attempt to incorporate both line of evidence (LOE) and classical weight of evidence (WOE) approaches for the assessment of sediment quality, a set of biomarkers were analyzed in target tissues of two invertebrate species after 28 days of exposure to sediments impacted by oil (derived from the tanker *Prestige*, 2002). The integration of biomarkers with sediment contamination, acute toxicity and benthic alteration parameters provides an “early warning” tool which not only indicates the environmental quality of an area, but also constitutes an advisory tool for potential ecological risks. The selected biomarkers provide information about the first biological responses due to the presence of contaminants in the environment providing predictable reports about further effects to the ecosystem. The present study demonstrates that the use of a set of biomarkers as part of a WOE approach designed to assess contaminated sediments contributes added value to the classical LOE and allows characterisation of the environmental status of the studied area in a more precise and accurate way.

Keywords: PAHs, contamination, toxicity, sublethal, WOE, bioassay, clam, crab.

45

46 **1. INTRODUCTION**

47 Chemical analyses normally provide the main tool in sediment quality assessment, even
48 though chemical concentrations alone are inadequate for the prediction of biological
49 consequences. Biological effects can be established using laboratory tests that determine
50 toxic responses and field surveys of the communities living within impacted sediments,
51 i.e. observing pollution-induced degradation of the biota (Chapman et al., 1991).

52 Weight of evidence (WOE) investigations determine the possible ecological impacts of
53 chemicals or other stressors based on multiple lines of evidence (LOE) (Chapman,
54 2007). They have been widely used in recent years to assess sediment quality around the
55 world (Chapman, 2000; Borgmann et al., 2001), including different areas within the
56 Iberian Peninsula (DelValls and Chapman, 1998; Caeiro, 2004; Riba et al., 2004;
57 Martín-Díaz, 2004; Morales-Caselles, 2008 a).

58 The sinking of the tanker *Prestige* (2002) spilt about 63,000 tonnes of heavy fuel oil (a
59 mixture of saturated hydrocarbons, aromatic hydrocarbons, resins, and asphaltenes, with
60 most of the PAHs being of medium to high molecular weight) mainly affecting the
61 Galician Coast, NW Spain. Several investigations have focused on determining the
62 biological and environmental effects of this dramatic episode by following single lines
63 of evidence, such as chemical analyses (CSIC, 2003a; Franco et al., 2006; González et
64 al., 2006), toxicity (Mariño-Balsa et al., 2003; Martínez-Gómez et al., 2006; Marigómez
65 et al., 2006; Morales-Caselles et al., 2006) or benthic alteration (Junoy et al., 2005;
66 Serrano et al., 2006). Recently, authors presented a report (Morales-Caselles et al., 2008
67 a) where a classical WOE approach, the Sediment Quality Triad (SQT), based on three
68 lines of evidence (physicochemical characterization of the sediments, determination of

69 acute toxicity and benthic alteration) was carried out in the Galician Coast. This study
70 suggested a general recovery of the environmental health within the area. However,
71 there were signs (Morales-Caselles et al., 2008 a) that other sources of contaminants
72 apart from the *Prestige* oil spill could be producing some environmental stress. The aim
73 of the present study is to use a biomarker line of evidence (LOE) with the weight of
74 evidence (WOE) approach to further improve the sediment quality assessment of the
75 Galician Coast. Chapman and Hollert (2006) suggested that future applications of the
76 SQT should consider specific LOE in terms of risk assessment, ensuring that both
77 exposure and assessment of biological effects are adequately addressed, as both are of
78 causation and ecological relevance. Thus, the LOE selected in this study constitutes an
79 improvement of the classical WOE laboratory and field studies based on biomarker
80 determinations. Biomarkers can act as an important early warning system by indicating
81 whether environmental pollutants are present at sufficiently high concentrations to cause
82 an effect (Walker et al., 2006). In addition, biomarker responses are specific to certain
83 groups of contaminants, providing information related to possible causation (Galloway
84 et al. 2004). Chemicals such as the Polycyclic Aromatic Hydrocarbons (PAHs) have a
85 very short biological half-life in most species, yet nevertheless, may have long-term
86 effects (Walker et al., 2006). In this sense, some compounds may not produce acute
87 toxic effects, however sublethal effects are likely to be observed; carefully selected
88 biomarkers can highlight these sublethal effects, thus providing increased sensitivity
89 and robustness to WOE assessments. Thus, the aim of this research was to examine the
90 feasibility and viability of incorporating a biomarker LOE within the classical WOE
91 approach as applied to the assessment of oil contaminated sediments. This new
92 approach is demonstrated by examining the extent of biological impact and identity of

contaminants within sediments influenced by the accidental oil spill of the *Prestige* (2002) on the Galician coast.

2. METHODOLOGY

2.1 Approach

The study was performed in two areas of the Galician coast (NW Spain) affected by the *Prestige* oil spill in 2002 (Figure 1): Cies Island (A, B, C) within the Atlantic Island National Park (AINP) and the Bay of Corme-Laxe (D, E, F). Cies Island acted as a natural barrier protecting the rias from entry by the fuel. The Bay of Corme-Laxe is also considered a place with high ecological relevance, with a low anthropogenic and industrial influence (fishing and farming being the main economic activities).

The 4 LOE employed within the WOE approach were:

(a) Sediment contamination; physicochemical characterization of sediments by analyzing PAHs (acenaphthalene, acenaphthylene, anthracene, benzo(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, benzo(g,h,i) perylene, benzo(k)fluoranthene, chrysene, dibenzo(a,h) anthracene, fenanthrene, fluoranthene, fluorene, indene (1,2,3,cd)pyrene, naphthalene, and pyrene) using GC-MS with selected ion monitoring and trace metals (Zn, Pb, Cu, Ni and Hg) with anodic stripping voltammetry (Morales-Caselles et al., 2006);

(b) Acute toxicity and bioaccumulation; by performing sediment bioassays such as the commercial assay Microtox[®] (Morales-Caselles et al., 2007), the amphipod mortality test with *Corophium volutator* (Morales-Caselles et al., 2007), the polychaetes mortality assay (Casado-Martínez et al., in press) and bioaccumulation experiments with *Arenicola marina* (Morales-Caselles et al., 2008 b);

(c) 'In situ alteration'; benthic alteration was selected and determined by measuring parameters in situ based on taxonomic identification and community descriptive statistics (abundance-biomass analysis, species richness, diversity, dominance and proportions of the major taxonomic groups) (DeIvals and Chapman, 1998);

(d) Laboratory and field studies based on biomarkers; using two invertebrate species the crab *Carcinus maenas* and the clam *Ruditapes philippinarum*, and a set of biomarkers (Morales-Caselles et al., 2008 c): mixed function oxygenase activity, which is the first mode of detoxification in many organic pollutants, was measured using the adapted Ethoxyresorufin-O-deethylase (EROD) assay; the Ferric Reducing Ability of Plasma (FRAP) assay allows a measure of the antioxidant capacity; antioxidant Glutathione-S-transferase (GST) activity was determined by monitoring the rate of conjugation of glutathione (GSH) to 1-chloro-2,4-dinitrobenzene (CDNB) at 340 nm; oxidation of 1 mM NADPH by Glutathione Reductase activity (GR) in the presence of 10 mM oxidized glutathione was also monitored at 340 nm; phase II metabolizing enzyme Glutathione Peroxidase activity (GPX); all the biomarker responses were normalized with the total protein content (n=5).

2.2 Data integration

The data obtained from the different LOE were integrated through a multivariate analysis approach based on linking all the variables obtained (Riba et al., 2004) and a pie chart representation of comparisons between sites of multivariate factors (Riba et al., 2004; Morales-Caselles et al., 2008 a). The multivariate analysis was performed using principal component analysis (PCA) as the extraction procedure, which is a multivariate statistical technique to explore variable distributions (Riba et al., 2003). The original data set used in the analysis included the variables obtained from the 4-LOE and its objective was to derive a reduced number of new variables as linear combinations of the

original variables. This provides a description of correlations between the set of the original variables and their meaning in each study site with a minimum loss of information. For the representation of the pie charts the new factors obtained from the PCA were submitted to ANOVA and Tukey tests which identified significant differences in sensitivity among stations and controls for each factor (Morales-Caselles et al., 2008 a). Every study site has a pie chart divided into the obtained factors which use different colours depending on the level of significant differences in relation to the reference.

3. RESULTS AND DISCUSSION

Table 1 shows the summarized results of the different parameters analyzed. Sediment Hg and Pb content was significantly higher in all Corme-Laxe sites; The highest metal contents were observed at stations A and F. There was no observed general pattern in toxicity and benthic parameters between the sampling sites. Factor analysis revealed that the original variables could be grouped into three principle components (factors) that explained 79 % of the total variance (Table 2). The multivariate analysis is a tool that allows us to interpret large groups of different variables by grouping them using correlations; in addition it indicates the importance of each factor in every single study site.

3.1 Factor #1

Factor 1 explained 35.2 % of the variance and was influenced by the relationship between different variables related with chemicals, sublethal responses, bioaccumulation and benthic alteration. Sediment PAHs, Pb and Hg contents were positively correlated to the bioaccumulation of PAHs in *Arenicola marina* (under laboratory conditions), but negatively correlated to their mortality in the acute assays. A

165 set of antioxidant and detoxification biomarkers analyzed in crabs (activity of GPX,
166 GR, GST, EROD and FRAP) and the digestive gland of clams (GR and GST activity)
167 under laboratory conditions were also correlated with Factor #1. EROD and FRAP of
168 organisms exposed to sediments *in situ* were not significantly correlated with other
169 variables (Table 2). The aforementioned contaminants and the toxicity variables are
170 slightly connected to the benthic alteration explained by alteration of the specific
171 richness and an increase of the polychaete population, while a positive development of
172 crustaceans was detected. Thus, Factor #1 may be interpreted as representing
173 contamination by PAHs, Hg and Pb, which does not produce lethal effects; although
174 PAH bioaccumulation and sub-lethal responses in organisms (under laboratory and field
175 conditions) are associated with a slight alteration of the *in situ* benthic community.
176 Environmental alterations due to these contaminants have been reported by other
177 authors (DelValls et al., 1998). Factor 1 scores for stations E (1.0) and F (1.4) located in
178 Corme-Laxe (Figure 2) were rather high, suggesting an accumulation of contaminants
179 including fuel oil from the *Prestige* (Labandeira et al., 2001; Prego et al., 2006).
180 Although initial studies did not detect the presence of Pb and Hg in emulsified samples
181 of the *Prestige* fuel (with 54–59% water) (CSIC, 2003b), the presence of Pb was later
182 corroborated by other authors (Andrade et al., 2004). Thus, the contamination by PAHs,
183 Hg and Pb observed at Corme-Laxe appears to be caused by the remnants of the
184 *Prestige* (Morales-Caselles et al., 2008 b).
185 Although no lethal effects were observed, the bioaccumulation of PAHs by *A. marina*
186 demonstrates the high bioavailability of these compounds. The induction of different
187 biomarkers in the hepatopancreas of crabs and in the digestive glands of clams was also
188 correlated with the presence of these contaminants (PAHs, Hg and Pb), suggesting that
189 the deployed organisms suffered stress due to the presence of these substances in the

190 sediments (Morales-Caselles et al., 2008c). The correlation observed among the
191 biomarkers and the different variables defined by Factor #1 was stronger under
192 laboratory conditions, implying that field deployments result in less sensitivity (Astley
193 et al 1999). The flushing action of the open water environment may act to reduce the
194 impacts of sediment contaminants when compared with laboratory incubations.

195 Although, a good correlation was observed between the biomarkers induced in both
196 invertebrate species, the crab *Carcinus maenas* was more sensitive i.e., showed stronger
197 responses, than the clam *Ruditapes philippinarum* to this kind of pollution. On the
198 other hand some of the variables related to the benthic alteration present in Factor #1
199 corroborate the effects observed in the sub-lethal experiments.

200 **3.2 Factor #2**

201 The second factor, Factor #2 (24.5 % of the variance) connects the set of biomarkers
202 measured under field conditions (EROD and FRAP activity in crabs and clams, and
203 GPX and GST in clams), the mortality of *Arenicola* in the acute experiment, no toxicity
204 for amphipods, the alteration in the number of species and the decrease of the
205 Polychaete population. Positive and negative correlations for a few biomarkers were
206 also detected under laboratory conditions and no concordance with the amphipod
207 toxicity test was observed. The relationships between the biological responses identified
208 in Factor #2 are not correlated with any of the chemicals analyzed suggesting that a
209 contaminant or group of compounds that was not analysed, is the cause of the biomarker
210 responses. Taking into account that the acute toxicity observed in *Arenicola* and the rest
211 of the bioassays was relatively low (less than 30% mortality) (Morales-Caselles et al.,
212 2008 b), a source of contaminant not related to the sediment is the most likely cause of
213 these effects. Station A (0.1) located on Cies Island and site D (2.0) in Bay of Corme-
214 Laxe had the highest positive correlation with Factor #2 (Figure 2). The good

correlation observed between the biomarkers in both crabs and clams incubated under field conditions suggests that these locations suffer the stress of non-measured variables which, in the case of site D, could be related to the proximity of aquaculture infrastructures (mussel culture). Other authors (Otero et al., 2006), have described the negative impacts of these mussel aquaculture, including: the discharge of a large volume of bio-deposits containing high concentrations of nutrients; the release of drugs and pesticides into the environment; an increase in sedimentation and accumulation of organic matter and an increase in the concentration of nutrients in sediments and waters (mainly N and P). Negative effects on wild populations of animals have also been observed, ranging from genetic interaction and disease transmission, to changes in the composition of the structure of benthic fauna due to a change from oxic to anoxic conditions (Otero et al., 2006).

3.3 Factor #3

The third factor, Factor #3 explained 19.2 % of the variance and was correlated with sediments contaminated by the metals Zn, Cu and Ni, and PAHs; these contaminants were correlated with the antioxidant responses under laboratory assays with crabs and clams (GR) and the acute toxicity as determined by the amphipods assay, which was not significant (the toxicity detected was not enough to consider sediment samples as toxic according to this acute assay). Samples where the mortality rate of the amphipods was 20 % higher than the mortality recorded in the reference and showed significantly different ($*p < 0.05$) results compared to those obtained in the reference are considered as toxic (Morales-Caselles et al., 2008 b) The effects on the benthic community were manifested as an alteration in the diversity and percentage of molluscs. This suggests the presence of a stress related to the source of metals (Zn, Cu and Ni) and organic compounds (PAHs), which is different than the source explained by Factor #1. Sites A

(1.2) in the AINP and F (1.3) in the Bay of Corme-Laxe had positive Factor #3 values (Figure 2). Previous studies in Bay of Corme-Laxe detected severe contamination of the sediments by Cu (Prego et al., 2006), however a fuel oil origin was unlikely, with the major source of Cu being antifouling paints from the hulls of fishing vessels (Cobelo-García and Prego, 2003). Although contamination by Cu and Zn was observed in the uppermost sediment layers of the *Prestige* shipwreck area (Prego and Cobelo-García, 2004; Cobelo-García et al., 2005), this contamination is unlikely not be related to the shipwreck, because levels of Cu in the fuel oil carried by the *Prestige* were relatively low (3.39 mg kg^{-1}) and previous studies have shown that inputs from terrestrial sources are probably higher than from the spilled fuel oil (Carballeira et al., 1997; Villares et al., 2007). Some of the studied variables demonstrate the stress of these contaminants and the effects on the benthic community; however, the results mostly point to chronic contamination with low bioavailability, and potential, but unconfirmed biological risk. Thus, despite the high trace metal content of sediments from the Galician coast, their reactivity and bioavailability were very low (Otero et al., 2005), which maybe related to the high degree of pyritization found for some of the most toxic trace metals, favouring their release by oxidation of the sulphides that they form, thus making them bioavailable to benthic fauna.

3.4 Significant differences among stations

Station C, with an absence of PAH contamination and the lowest biological effects, was considered as the most suitable site to use as a reference station. Factor loadings obtained in the MAA were submitted to ANOVA and post-hoc examination using the Tukey test (Figure 3). Factor # 1 scores of sediments from Cies Island in the AINP were not significantly different, suggesting that the effects of the *Prestige* oil spill were no longer detectable

265 within the area. However, Factors # 2 and # 3 scores for station A were significantly
266 different ($p < 0.01$) from the reference station (C), suggesting the influence of non-
267 measured stressors from sources other than the *Prestige*. The factor #3 score of station
268 B was also significantly different ($p < 0.05$) from the reference station, suggesting the
269 presence of some contaminants considered a potential risk within the area, although in
270 general, the environmental quality of the sediments was relatively good. On the other
271 hand, Factor #1 scores of sediments from Corne-Laxe D, E and F were significantly
272 different ($p < 0.01$) from the reference station, suggesting the remaining contaminants
273 from the *Prestige* oil spill are still producing sub-lethal effects to the biota of the bay. In
274 addition, the significant difference ($p < 0.01$) in Factor #2 score observed at station D
275 suggests other unknown sources of contaminants responsible for biological stress in the
276 study site. In the case of station F a mixture of metals and PAHs from different sources
277 could be considered a potential risk in the area as shown by Factor #3 ($p < 0.01$).
278 Biomarkers are useful “early warning” indicators of the health status of animals from
279 impacted areas (Galloway et al., 2004; Montserrat et al., in press), including systems
280 impacted by oil (Anderson and Lee 2006), where complex mixtures of pollutants are
281 usually present. The use of a set of biomarkers, as part of a WOE approach designed to
282 assess contaminated sediments, contributed added value to the classical LOE concept
283 and allows for the characterization of the environmental status of the studied area in a
284 more precise and accurate way. In addition, the inclusion of chronic bioassays with two
285 invertebrate species, not only under laboratory conditions but also in field deployments,
286 elucidate different sources of contaminants apart from the sediments permitting a more
287 realistic approach to the original situation of the ecosystem, and the potential ecological
288 risks.

289 4. CONCLUSIONS

Multiple lines of evidence suggest that 4 years after the impact of the *Prestige* oil spill, the fuel is not producing acute toxic effects within the environment (Morales-Caselles et al., 2008 a); however, the use of a biomarker LOE within the WOE approach demonstrated that sub-lethal responses may occur in the area of Corme-Laxe, related to PAHs, Pb and Hg. No effects of oil spill were observed in the study sites located in AINP although contamination by metals, especially Zn, Cu and Ni was observed in some sites on Cíes Island. The inclusion of biomarkers in the SQT indicates that coastal anthropogenic influences (a mixture of pollutants that act as environmental stressors) are evident in both areas and should be considered a potential risk. In the case of Corme-Laxe, a possible impact related to the aquaculture of mussels was detected. Biomarkers demonstrated higher sensitivity when compared to acute toxicity approaches providing a more comprehensive and integrated weight-of-evidence approach (Morales-Caselles et al., 2008 a), that as demonstrated, can be easily incorporated as a fourth line of evidence into the classical SQT. Furthermore, this study confirms the suitability of biomarkers as a tool to assess the sources and potential ecological risk of metallic and petrogenic contaminants within sediments.

5. ACKNOWLEDGEMENTS

This work was supported by the projects VEM2003-20563/INTER, and CTM2005-07282-C03-01/TECNO financed by the Spanish Education and Science Ministry and by Centro Investigaciones Subacuáticas (CIS) funded by the Ministry of Environment. C Morales-Caselles thanks the Ministry of Education and Science for funding her research fellowship (FPU). The authors would like to thank the members of the CIS for their support and help in the chemical analysis and the benthic community information; special thanks are given to L Martín, N Fernández, A César, P Vidal and A Moreno.

314 **6. REFERENCES**

- 315 Anderson, J.W., Lee, R.F. 2006. Use of biomarkers in oil spill risk assessment in the
316 marine environment. *Human and Ecology Risk Assessment* 12, 1192-122
- 317 Andrade, M.L., Covelo, E.F., Vega, F.A., Marcet, P. 2004. Effect of the Prestige oil
318 spill on salt marsh soils on the coast of Galicia (Northwestern Spain). *Journal of*
319 *Environmental Quality* 33, 2103–2110
- 320 Astley, K.N., Meigh, H.C., Glegg, G.A., Braven, J., Depledge, M.H. 1999. Multi-variate
321 Analysis of Biomarker Responses in *Mytilus edulis* and *Carcinus maenas* from the Tees
322 Estuary (UK). *Marine Pollution Bulletin* 39, 145-154
- 323 Borgmann, U., Norwood, W.P., Reynoldson, T.B., Rosa, F. 2001. Identifying cause in
324 sediment assessments: bioavailability and the Sediment Quality Triad. *Can. Journal of*
325 *Fish Aquatic Science* 58, 950-969
- 326 Caeiro, S. 2004. Environmental data management in the Sado estuary: weight of
327 evidence to assess sediment quality. Ph.D. Thesis. University Nova of Lisbon. 383 pp
- 328 Carballeira, A., Carral, E., Puente, X.M., Villares, R. 1997. Estado de conservación de
329 la costa de Galicia. Nutrientes y metales pesados en sedimentos y organismos
330 intermareales. Universidad de Santiago de Compostela. 105 pp
- 331 Casado-Martínez, M.C., Forja, J.M., DelValls, T.A. Comparative toxicity assessment
332 using the amphipod *Corophium volutator* and the polychaete *Arenicola marina* for
333 dredged material management. *Environmental Toxicology* (in press)

- 334 Chapman, P.M., Power, E.A., Dexter, R.N., Andersen, H.B. 1991. Evaluation of effects
335 associated with an oil platform, using the sediment quality triad. *Environmental*
336 *Toxicology and Chemistry* 10, 407–424
- 337 Chapman, P.M. 2000. The Sediment Quality Triad: then, now and tomorrow.
338 *International Journal of Environmental Pollution* 13, 351-356
- 339 Chapman, P.M., Hollert, H. 2006. Should the sediment quality triad become a tetrad, a
340 pentad, or possibly even a hexad? *Journal of Soil and Sediment* 6, 4-8
- 341 Chapman, P.M. 2007. Determining when contamination is pollution – Weight of
342 evidence determinations for sediments and effluents. *Environment International* 33,
343 492-501
- 344 Cobelo-García, A., Labandeira, A., Prego, R.. 2005. Two opposite cases of metal
345 accumulation in ria sediments: Ferrol and Corme-Laxe (Galicia, NW Iberian Peninsula).
346 *Ciencias Marinas* 31, 653-659
- 347 Cobelo-García, A., Prego, R.. 2003. Land inputs, behaviour and contamination levels of
348 copper in a ria estuary (NW Spain). *Marine Environmental Research* 56, 403-422
- 349 Centro Superior Investigaciones Científicas (CSIC). 2003a. Heavy metal presence in the
350 shipwreck area of Prestige tanker and metal contents in the emulsified fuel (in
351 Spanish). Technical report no. 2, <http://csicprestige.iim.csic.es/>
- 352 Centro Superior Investigaciones Científicas (CSIC). 2003b. Caracterización del vertido
353 y evolución preliminar en el medio. CSIC Technical Report, vol. 2,
354 <http://csicprestige.iim.csic.es/>

- 355 DelValls, T.A., Chapman, P.M. 1998. Site-specific sediment quality values for the golf
356 of Cádiz (Spain) and San Francisco Bay (USA), using the sediment quality triad and
357 multivariate analysis. *Ciencias Marinas* 24, 313-336
- 358 DelValls, T.A., Conradi, M., García-Adiego, E., Forja, J.M., Gómez-Parra, A. 1998.
359 Analysis of macrobenthic community structure in relation to different environmental
360 sources of contamination in two littoral ecosystems from the Gulf of Cádiz (SW Spain).
361 *Hydrobiologia* 385, 59-70
- 362 Franco, M.A., Viñas, L., Soriano, J.A., de Armas, D., González, J.J., Beiras, R., Salas,
363 N., Bayona, J.M., Albaigés, J. 2006. Spatial distribution and ecotoxicity of petroleum
364 hydrocarbons in sediments from the Galicia continental shelf (NW Spain) after the
365 Prestige oil spill. *Marine Pollution Bulletin* 53, 260-271
- 366 Galloway, T.S., Brown, R.J., Browne, M.A., Dissanayake, A., Lowe, D., Jones, M.B.,
367 Depledge, M.H. 2004. A multibiomarker approach to environmental assessment.
368 *Environmental Science and Technology* 38, 1723-1731
- 369 González, J.J., Viñas, L., Franco, M.A., Fumiga, J., Soriano, J.A., Grueiro, Muniategui,
370 S., López-Mahía, P., Prada, D., Bayona, J.M., Alzaga, R., Albaigés, J. 2006. Spatial and
371 temporal distribution of dissolved/dispersed aromatic hydrocarbons in seawater in the
372 area affected by the Prestige oil spill. *Marine Pollution Bulletin* 53, 250-259
- 373 Junoy, J., Castellanos, C., Viéitez, J.M., de la Huz, M.R., Lastra, M. 2005. The
374 macroinfauna of the Galician sandy beaches (NW Spain) affected by the Prestige oil-
375 spill. *Marine Pollution Bulletin* 50, 526-536
- 376 Labandeira, A., Cobelo-García, A., Prego, R. 2001. La Ría de Corme y Lage: Revisión
377 de su conocimiento. *Monogr Quím Oceanogr* 1, 1-34

- 378 Marigómez, J., Soto, M., Cancio, I., Orbea, A., Garmendia, L., Cajaraville, M.P. 2006.
379 Cell and tissue biomarkers in mussel, and histopathology in hake and anchovy from Bay
380 of Biscay after the Prestige oil spill (Monitoring Campaign 2003). *Marine Pollution*
381 *Bulletin* 53, 287-304
- 382 Mariño-Balsa, J.C., Pérez, P., Estévez-Blanco, P., Saco-Álvarez, L., Fernández, E.,
383 Beiras, R. 2003. Assessment of the toxicity of sediment and seawater polluted by the
384 Prestige fuel spill using bioassays with clams (*Venerupis pollastra*, *Tappes decussates*
385 and *Venerupis rhomboideus*) and the microalga *Skeletonema costatum*. *Ciencias*
386 *Marinas* 29, 115–122
- 387 Martín-Díaz, L. 2004. Determinación de la calidad ambiental de sistemas litorales y de
388 estuario de la Península Ibérica utilizando ensayos de campo y laboratorio. Doctoral
389 thesis. University of Cádiz. 330 pp
- 390 Martín-Díaz, M.L., Blasco, J., Sales, D., DelValls, T.A. Biomarkers study for sediment
391 quality assessment in spanish ports using the crab *Carcinus maenas* and the clam
392 *Ruditapes philippinarum*. *Archives of Environmental Contamination and Toxicology*
393 (in press)
- 394 Martínez-Gómez, C., Campillo, J.A., Benedicto, J., Fernández, B., Valdés, J., García, I.,
395 Sánchez, F. 2006. Monitoring biomarkers in fish (*Lepidorhombus boscii* and
396 *Callionymus lyra*) from the northern Iberian shelf after the Prestige oil spill. *Marine*
397 *Pollution Bulletin* 53, 305-314
- 398 Montserrat, J.M., Martínez, P.E., Geracitano, L.A., Amado, L.L., Martinez Gaspar
399 Martins, C., Lopes Leaes Pinho, G., Soares Chaves, I., Ferreira-Cravo M, Ventura-

- 400 Lima, J., Bianchini, A. Pollution biomarkers in estuarine animals: Critical review and
401 new perspectives. *Comparative Biochemistry Physiology C* (in press)
- 402 Morales-Caselles, C., Jiménez-Tenorio, N., González de Canales M.L., Sarasquete, C.,
403 DelValls, T.A. 2006. Ecotoxicity of sediments contaminated by the oil spill associated
404 with the tanker “Prestige” using juveniles of the fish *Sparus aurata*. *Archives of*
405 *Environmental Contamination and Toxicology* 51, 652–660
- 406 Morales-Caselles, C., Kalman, J., Riba, I., DelValls, T.A. 2007. Comparing sediment
407 quality in spanish littoral areas affected by acute (Prestige, 2002) and chronic (Bay of
408 Algeciras) oil spills. *Environmental Pollution* 146, 233-240
- 409 Morales-Caselles, C., Riba, I., Sarasquete, C., DelValls, T.A. 2008a. Using a classical
410 weight-of-evidence approach for 4-years monitoring of the impact of an accidental oil
411 spill on sediment quality. *Environment International* 34, 514-523
- 412 Morales-Caselles, C., Kalman, J., Micaelo, A.C., Ferreira, A.M., Vale, C., Riba, I.
413 DelValls, T.A. 2008b. Sediment contamination, bioavailability and toxicity of sediments
414 affected by an acute oil spill. Four years after the sinking of the tanker Prestige (2002).
415 *Chemosphere* 71, 1207-1213
- 416 Morales-Caselles, C., Martín-Díaz, M.L., Riba, I., Sarasquete, C., DelValls, T.A. 2008c.
417 The role of biomarkers to assess oil-contaminated sediment quality using toxicity tests
418 with clams and crabs. *Environment Toxicology and Chemistry* 27, 1309-1316
- 419 Otero, X.L., Calvo de Anta, R.M., Macías, F. 2005. Trace elements in biodeposits and
420 sediments from mussel culture in the Ría de Arousa (Galicia, NW Spain).
421 *Environmental Pollution* 136, 119-134

- 422 Otero, X.L, Calvo de Anta, R.M., Macías, F. 2006. Sulphur partitioning in sediments
423 and biodeposits below mussel rafts in the Ría de Arousa (Galicia, NW Spain). *Marine*
424 *Environmental Research* 61, 305-325
- 425 Prego, R., Cobelo-García, A., Marmolejo-Rodríguez, J., Santos-Echeandía, J. 2006.
426 Trace elements in the Prestige fuel-oil spill: Levels and influence on Laxe Ria
427 sediments (NW Iberian Peninsula). *Ciencias Marinas* 32, 179-186
- 428 Prego, R., Cobelo-García, A. 2004 Cadmium, copper and lead contamination of the
429 seawater column on the Prestige shipwreck (NE Atlantic Ocean). *Analytica Chimica*
430 *Acta* 524, 23–26
- 431 Riba, I., Forja, J.M., Gómez-Parra, A., DelValls, T.A. 2004. Sediment quality in littoral
432 regions of the Gulf of Cádiz: a triad approach to address the influence of mining
433 activities. *Environmental Pollution* 132, 341-353
- 434 Riba, I., Zitko, V., Forja, J.M., DelValls, T.A. 2003. Deriving sediment quality
435 guidelines in the Guadalquivir estuary associated with the Aznalcóllar mining spill, a
436 comparison of different approaches. *Ciencias Marinas*, 29
- 437 Serrano, A., Sánchez, F., Preciado, I., Parra, S., Frutos, I. 2006. Spatial and temporal
438 changes in benthic communities of the Galician continental shelf after the Prestige oil
439 spill. *Marine Pollution Bulletin* 53, 315-331
- 440 Villares, R., Real, C., Fernández, J.A., Aboal, J., Carballeira, A. 2007. Use of an
441 environmental specimen bank for evaluating the impact of the Prestige oil spill on the
442 levels of trace elements in two species of *Fucus* on the coast of Galicia (NW Spain).
443 *Science of the Total Environment* 374, 379–387

- 444 Walker, C.H., Hopkin, S.P., Sibly, R.M., Peakall, D.B. 2006. Biomarkers. In: Principles
445 of Ecotoxicology. CRC Press Boca Raton, Florida. 315 pp

446 **FIGURE CAPTIONS**

447 Figure 1. Map of the coastal area of Galicia (NW Spain) showing the sampling sites in
448 the Atlantic Island National Park (A, B, C) and the area of Corme-Laxe (D, E and F).

449 Figure 2. Estimated factor scores for the three factors in each of the 6 cases. The factor
450 scores quantify the prevalence of each factor for every station and are used to establish
451 the definition of each factor.

452 Figure 3. Pie charts which represent the significant differences of the factors score in
453 every study site related to the reference site C (heavy shading: $p < 0.01$; light shading: p
454 < 0.05 ; no shading: no significant differences, $p > 0.05$).

455

456

1

2 Figure 2.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19 Figure 3.

20

21

22 **FIGURE CAPTIONS**

23 Figure 1. Map of the coastal area of Galicia (NW Spain) showing the sampling sites in
24 the Atlantic Island National Park (A, B, C) and the area of Corme-Laxe (D, E and F).

25 Figure 2. Estimated factor scores for the three factors in each of the 6 cases. The factor
26 scores quantify the prevalence of each factor for every station and are used to establish
27 the definition of each factor.

28 Figure 3. Pie charts which represent the significant differences in the factors' score in
29 every study site related to the reference site C (high density shading: $p < 0.01$; low
30 density shading: $p < 0.05$; no shading: no significant differences, $p > 0.05$).

31

32

33

34 Table 1. Summarized results and SD of chemical analysis (mg kg^{-1} for metals, $\mu\text{g kg}^{-1}$
35 for PAHs) the acute toxicity tests (*Corophium volutator* and *Arenicola marina*: %
36 mortality; Microtox: IC50; bioaccumulation of PAHs: $\mu\text{g kg}^{-1}$), biomarker responses
37 under field and laboratory conditions (glutathione peroxidase activity GPX: nmol min^{-1}
38 mg prot^{-1} , glutathione transferase GST activity $\text{nmol min}^{-1} \text{mg prot}^{-1}$, glutathione
39 reductase GR activity $\text{nmol min}^{-1} \text{mg prot}^{-1}$, ferric reducing ability of plasma FRAP
40 activity $\mu\text{M mg}^{-1} \text{min}^{-1}$ and EROD activity $\text{pmol mg}^{-1} \text{min}^{-1}$) and the alteration
41 parameters for sediments from the Atlantic Islands National Park (AINP) (A, B, C) and
42 Corme-Laxe (D, E, F). n.d.: not detected; n.a.: not available.

43

44

45

1

		A	B	C	D	E	F
Chemical analyses	Zn	377±36.06	91±1.34	164±11.31	25±1.41	19.9±1.34	271±16.26
	Pb	1.5±0.14	0.9±0.00	0.85±0.01	3.7±0.21	7.3±0.07	5.9±0.14
	Cu	5.2±0.35	1.4±0.14	1.4±0.21	0.7±0.21	0.43±0.05	4.2±0.21
	Ni	13.3±1.91	2.4±0.21	4.5±0.14	1.7±0.14	1.5±0.00	5.7±0.57
	Hg	0.7±0.21	0.8±0.07	0.6±0.07	2±0.14	2.1±0.28	3.4±0.28
	PAH	108±7.78	67±16.33	n.d.	38±2.47	52±0.71	323±13.44
Toxicity tests	Corophium	23±1.18	20±1.53	17±0.02	10±1.77	17±1.18	20±3.54
	Arenicola	28±1.96	28±1.97	22±1.96	39±0.79	17±0.02	17±0.24
	Microtox	5631±134	9422±301	1801±101	3977±214	21041±407	4398±282
	Bioaccumulation PAH	2927±1.48	2573±1.48	2666±1.24	2616±2.40	3911±0.71	3285±0.71
Biomarkers (laboratory)	GPX-crab-lab	11.6±0.59	9.7±0.55	8.2±1.72	19.3±0.00	19.5±2.88	15.9±1.18
	GPX-clam-lab	2.1±0.01	2.9±0.59	4.5±0.04	6.1±2.36	3.1±1.47	4.2±2.42
	GR-crab-lab	1.1±0.06	0.7±0.08	0.9±0.12	0.9±0.18	0.6±0.01	1.5±0.04
	GR-clam-lab	2.1±0.65	1.6±0.00	2.3±0.36	3.4±0.43	11.7±3.17	3.9±0.03
	GST-crab-lab	139.6±52.3	218.1±44.2	407.3±8.97	429.6±36.1	684.1±422	1070.9±198
	GST-clam-lab	1292.8±339	838.6±0.00	1624.5±209	1199.1±174	910.2±566	847.7±293
	EROD-crab-lab	0.05±0.00	0.06±0.00	0.06±0.00	0.04±0.00	0.10±0.00	0.09±0.00
	EROD-clam-lab	0.26±0.00	0.27±0.00	0.42±0.00	0.41±0.00	0.44±0.00	0.22±0.00
	FRAP-crab-lab	3.85±0.92	2.08±1.01	2.56±0.47	2.86±0.43	2.93±0.56	1.56±0.30
	FRAP-clam-lab	10.58±3.8	7.77±0.5	4.04±0.70	13.72±1.9	12.09±2.6	6.43±0.30
Biomarkers (field)	GPX-crab-field	17.8±0.07	23.1±2.12	15.9±4.10	41.4±2.83	193.1±0.1	125.7±11
	GPX-clam-field	10.5±0.49	3.6±0.14	4.0±0.21	25.5±0.35	3.2±0.49	7.0±2.40
	GR-crab-field	0.7±0.21	1.4±0.92	1.4±0.99	9.9±0.71	9.5±2.83	23.4±1.06
	GR-clam-field	2.9±0.00	1.3±0.00	3.8±0.00	9.7±0.85	14.7±0.07	8.1±1.77
	GST-crab-field	1097.8±571	1563.8±224	690.3±10.6	1489.0±219	7523.4±1129	6072.9±0.0
	GST-clam-field	2060.8±926	371.6±130	1198.7±45	3365.6±250	130.5±2.5	1557.9±192
	EROD-crab-field	0.13±0.00	2.96±0.00	0.02±0.00	8.51±0.00	0.39±0.00	0.48±0.00
	EROD-clam-field	0.15±0.00	0.07±0.00	0.09±0.00	0.63±0.00	0.07±0.00	0.13±0.00
	FRAP-crab-field	2.72±0.35	n.a.	n.a.	2.42±0.32	n.a.	n.a.
	FRAP-clam-field	10.39±1.4	3.09±0.53	2.63±0.91	23.58±2.5	2.01±0.65	6.58±1.01
Benthic alterations	Number species	28.5	33.9	42.4	28.6	32.1	48.2
	specific richness	5.1	5	4.3	3	3	2.9
	Diversity	15.3	28.4	39.1	30	40.1	15.4
	Dominance	0.50	0.10	0.06	0.15	0.19	0.20
	% Mollusca	15.3	28.4	39.1	30.0	40.1	15.4
	% Polychaete	20.0	21.5	21.7	20.0	22.2	23.1
	% Crustacea	37.0	41.0	39.1	50.0	51.4	61.5

2

3

4

Table 2. Sorted rotated factor loadings of 36 variables for the three principal factors resulting from the multivariate analysis of the single results obtained from the chemical analysis, the acute toxicity tests, the suite of biomarkers and the alteration parameters for the study of the sediments quality in the Galician Coast. Chemicals: loadings are related to the concentration of contaminants in sediments; Acute effects: loadings explain the toxicity detected by the acute assays and the bioaccumulation of PAHs in *Arenicola marina*. Sublethal effects: loadings are related to the induction of biomarkers. Benthic alteration: loadings are related to alteration of the biota (decrease of number of species, specific richness, diversity, diminution in the percentage of molluscs and crustacea and increase in the polychaete population. The group of variables selected for the interpretation represented a loading of 0.40 or higher for a good association between an original variable and a factor.

42

		Factor 1 35.2	Factor 2 24.5	Factor 3 19.2
Chemicals	Zn	—	—	0.90
	Pb	0.95	—	—
	Cu	—	—	0.93
	Ni	—	—	0.76
	Hg	0.95	—	—
	PAH	0.59	—	0.76
Acute effects	Corophium	—	-0.52	0.75
	Arenicola	-0.41	0.87	—
	Microtox	—	—	—
	Bioaccumulation	0.73	—	—
Sublethal effects (laboratory & field)	GPX-crab-lab	0.48	-0.77	—
	GPX-clam-lab	—	0.53	—
	GR-crab-lab	0.41	—	0.78
	GR-clam-lab	0.65	—	0.49
	GST-crab-lab	0.85	—	—
	GST-clam-lab	-0.53	—	—
	EROD-crab-lab	0.62	-0.78	—
	EROD-clam-lab	—	—	-0.89
	FRAP-crab-lab	-0.46	—	—
	FRAP-clam-lab	—	0.64	—
	GPX-crab-field	0.88	—	—
	GPX-clam-field	—	0.99	—
	GR-crab-field	0.91	—	—
	GR-clam-field	0.77	—	-0.46
	GST-crab-field	0.89	—	—
	GST-clam-field	—	0.75	—
	EROD-crab-field	—	0.88	—
	EROD-clam-field	—	0.98	—
	FRAP-crab-field	—	0.76	—
	FRAP-clam-field	—	0.99	—
Benthic alteration	Number of species	—	0.56	—
	specific richness	0.91	—	—
	Diversity	—	—	0.97
	% Mollusca	—	—	0.96
	% Polychaeta	0.62	-0.68	—
	% Crustacea	-0.96	—	—

43

44