



## The effect of 3-pentadecylphenol on DPPC bilayers ATR-IR and P NMR studies

Katarzyna Cieřlik-Boczula, Aleksander Koll

### ► To cite this version:

Katarzyna Cieřlik-Boczula, Aleksander Koll. The effect of 3-pentadecylphenol on DPPC bilayers ATR-IR and P NMR studies. Biophysical Chemistry, Elsevier, 2009, 140 (1-3), pp.51. 10.1016/j.bpc.2008.11.009 . hal-00505534

**HAL Id: hal-00505534**

**<https://hal.archives-ouvertes.fr/hal-00505534>**

Submitted on 24 Jul 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

The effect of 3-pentadecylphenol on DPPC bilayers ATR-IR and  $^{31}\text{P}$  NMR studies

Katarzyna Cieřlik-Boczula, Aleksander Koll

PII: S0301-4622(08)00253-6  
DOI: doi: [10.1016/j.bpc.2008.11.009](https://doi.org/10.1016/j.bpc.2008.11.009)  
Reference: BIOCHE 5191

To appear in: *Biophysical Chemistry*

Received date: 10 September 2008  
Revised date: 15 November 2008  
Accepted date: 17 November 2008



Please cite this article as: Katarzyna Cieřlik-Boczula, Aleksander Koll, The effect of 3-pentadecylphenol on DPPC bilayers ATR-IR and  $^{31}\text{P}$  NMR studies, *Biophysical Chemistry* (2008), doi: [10.1016/j.bpc.2008.11.009](https://doi.org/10.1016/j.bpc.2008.11.009)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Corresponding author: Katarzyna Cieřlik-Boczula

Address: Faculty of Chemistry, University of Wrocław, Joliot-Curie 14, 50-383 Wrocław,  
Poland

Tel.: +48 71 375 7249, Fax.: 071 328-23-48

e-mail address: kasia\_cieslik0@yahoo.com

# **The effect of 3-pentadecylphenol on DPPC bilayers**

## **ATR-IR and $^{31}\text{P}$ NMR studies**

Katarzyna Cieřlik-Boczula\*, Aleksander Koll

Faculty of Chemistry, University of Wrocław, Joliot-Curie 14, 50-383 Wrocław, Poland

**Keywords:** phenolic lipid; DPPC; liposome; ATR-IR;  $^{31}\text{P}$  NMR; phase transition

**Abstract**

The influence of 3-pentadecylphenol (PDP) on the structure and physicochemical properties of the lipid bilayers of DPPC liposomes was studied using ATR-IR and  $^{31}\text{P}$  NMR methods. On the basis of analysis of the bands assigned to the  $\text{CH}_2$  stretching,  $\text{CH}_2$  scissoring,  $\text{C}=\text{O}$  stretching, and  $\text{PO}_2^-$  stretching vibrations it was revealed that PDP influences both the hydrophobic and hydrophilic parts of the DPPC liposome bilayer. Analysis of the  $^{31}\text{P}$  NMR line-shape indicated a lamellar to non-lamellar phase transition in PDP-doped DPPC dispersions. It was shown that PDP/DPPC isotropic aggregates have similar  $\nu\text{C}=\text{O}$  and  $\nu_{\text{as}}\text{PO}_2^-$  band positions and lower *gauche* populations in the hydrophobic chain region compared with the DPPC bilayer in the liquid-crystal phase.

## Introduction

3-pentadecylphenol (PDP) belongs to the phenolic lipids group. It is a large group of compounds of natural origin [1-5] which can interact with proteins, DNA, and biomembranes and they are antibacterial, fungicidal, and cytotoxic agents [2,6,7]. Phenolic lipids compose of an aromatic headgroup related to phenol or dihydroxybenzene connected to a hydrophobic hydrocarbon chain. The PDP compound has a phenol ring with one pentadecyl (C15) chain, see Fig. 1 A. PDP compounds are derived after the hydrogenation of unsaturated fractions of alkylphenolic oil internationally called "cashew nut shell liquid" (CNSL) [8]. CNSL is derived from the shell of the cashew nut (*Anacardium occidentale L.*). The CNSL oil is composed of some amount of cardanol which mainly consists of PDP with the unsaturated C15 hydrocarbon chain and small amount of saturated one [8]. The low cost of obtaining PDP from CNSL (world-wide cashew nut production is nearly 500 000 tons per year) increases the interest in this compound by industry and science. The synthesis of hydrophobic Au<sub>core</sub>Ag<sub>shell</sub> nanoparticles in toluene proceeds by way of the interfacial reduction of silver ions by 3-pentadecylphenol followed by their deposition on hydrophobized Au nanoparticles. This leads to the formation of phase of stable and pure Au<sub>core</sub>Ag<sub>shell</sub> nanoparticles in toluene [9]. PDP molecules are used in the formation of oriented block copolymers with anisotropic proton conduction [10]. The lamellar nanostructure present in this system is formed by PDP compounds and is responsible for characteristic organization of copolymer structures in which proton conductivity is different in three macroscopic directions. 3-Pentadecylphenol itself has various biological properties. PDP can effectively function as an anti-obesity agent by inhibiting the activity of  $\alpha$ -glucosidase enzyme and thereby regulate animal body weight [11]. It also has quite promising antioxidant activity resulting from the presence of phenol OH group [12].

**Figure 1.**

Most of the biological functions of phenolic lipids are related to modification of the physical properties of biomembranes [2]. As an amphiphilic compound, PDP can easily incorporate into the lipid bilayer, change the physicochemical properties of biomembranes, and alter the activities of different membrane proteins. The character of interactions between lipids and PDP compounds on molecular level is necessary to understand the membrane-connected biological activities of PDP molecules.

Our previous ATR-IR studies showed the influence of PDP compounds on dipalmitoylphosphatidylcholine (DPPC) multilayers in dry or partially hydrated lipid films [13]. However, the level of hydration has a crucial influence on the type of organization of the lipid bilayers and on the interactions between the lipid and doped compound [14]. The studies of liposome suspensions presented in that paper allowed us to mimic the hydration condition of natural membranes.

In this paper we focus on characterizing the effect of PDP on dipalmitoylphosphatidylcholine (DPPC) bilayers using ATR-IR (Attenuated Total Reflection Infrared Spectroscopy) and  $^{31}\text{P}$  NMR methods. Phosphatidylcholines (PC) are the most prevalent phospholipids among those constituting the basic structure of eukaryotic cell membranes. DPPC lipid is one of the most numerous members of PC lipid group and is very often used as a model for mimic lipid-biomembrane properties. The structure of DPPC molecule is present in Fig. 1B. The ATR-IR method supplies information about the PDP-lipid interaction and characterize the influence of the new compound on the structure of the lipid bilayer under fully hydrated conditions.  $^{31}\text{P}$  NMR provides information regarding the macroscopic architecture of the membrane phospholipids. The  $^{31}\text{P}$  NMR spectral line-shape is diagnostic of bilayer, hexagonal  $\text{H}_{\text{II}}$ , or isotropic (micellar, cubic or rhombic) lipid

arrangements in the fully hydrated state [15,16]. The isotropic phase in the PDP/DPPC mixtures was observed. The presence of non-lamellar structures in the PDP-doped lipid bilayer can extend the biological functions of biomembrane; for example the fusion phenomena and many other related processes are facilitated by the formation of non-bilayer lipid aggregates.

## Materials and Methods

### *Materials*

All compounds were used of the highest available purity, without further purification. The lipid dipalmitoylphosphatidylcholine (DPPC, of purity >99.8%) and 3-pentadecylphenol (PDP, of purity >95%) were obtained from Sigma-Aldrich, Germany.

### *Preparation of liposomes*

The chloroform solution of DPPC and PDP was dried under a stream of nitrogen. Dry lipid films were hydrated by addition of 1 ml of 10 mM phosphate buffer, pH 7.2. The final concentration of DPPC was 10 mg/ml (for ATR-IR measurement) and 40 mg/ml (for  $^{31}\text{P}$  NMR measurement). The mixtures were kept at a temperature of 10°C above the respective gel-liquid crystalline phase-transition temperature and gently shaken until optical homogeneity was obtained. The liposomal suspensions were treated 10 times in the cooling/heating process (liposomes were incubated for 10 min. at 4°C, then heated to a temperature 10 C° higher than the  $T_m$  of the doped liposomes, and incubated at this temperature, also for 10 min.). Finally, the solution was extruded through a filter of 100nm pore size (LiposoFast with polycarbonat filter, Avestin, Canada).

### *ATR-IR (Attenuated Total Reflection Infrared) measurements*



The ATR-IR infrared spectra were recorded on a Nicolet Avatar 360 FT-IR spectrometer. The 256 scans were collected at a resolution of  $2\text{ cm}^{-1}$ . The liposome suspensions were prepared according to procedure described above and spread on one surface of ZnSe-ATR crystal (face angle:  $45^\circ$ , 6 reflections, Specac). The dry DPPC/PDP films were prepared by spreading  $200\text{ }\mu\text{l}$  of chloroform solution of DPPC/PDP mixture on one surface of ZnSe-ATR crystal and evaporating the solvent under a stream of nitrogen. The concentration of DPPC in chloroform solution was  $10\text{ mg/ml}$ . The spectra of dry film and liposome suspensions were recorded in a heating cycle from  $15^\circ\text{C}$  to  $80^\circ\text{C}$ . The sample temperature was equilibrated for 5 min. before acquisition of each spectrum. The data were analyzed by Grams software.

#### *$^{31}\text{P}$ NMR measurements*

The  $^{31}\text{P}$  NMR spectra were recorded on a Bruker 300 NMR Fourier Transform spectrometer operating at  $121.49\text{ MHz}$ . The PDP/DPPC suspensions were prepared according to the procedure described above in a  $\text{D}_2\text{O}$  solution and were put into 5mm thin-walled NMR tubes. The relaxation delay time was 3 s. The spectra were recorded over a large spectral width (36 kHz) using broad-band proton decoupling. The data were analyzed by Grams software.

## Results and discussions

### ATR-IR measurements

#### *The alkyl chain vibration region in dry DPPC/PDP film*

The ATR-IR studies of the PDP/DPPC dry films showed the interaction of the PDP compounds with the DPPC multilayer structure [13]. PDP influences both the hydrophobic and hydrophilic regions of DPPC layers. In dry DPPC film there is no phase transition in the investigated temperature range from 25 to 80°C [17]. However, the incorporation of PDP molecules into the DPPC layers caused the appearance of a phase transition under dry conditions [13]. Analysis of the positions of the stretching and scissoring CH<sub>2</sub> bands, commonly used vibration indicators of conformational and alkyl chain packing changes, clearly indicates the nature of the phase transition in the PDP/DPPC mixtures as a chain-melting phase transition. For an equimolar PDP/DPPC mixture at a temperature of around 48°C there is a characteristic shift of the stretching CH<sub>2</sub> bands to the higher frequency region associated with a sharp increase the amount of *gauche* conformers of the hydrocarbon chains in mixed DPPC layers (Fig. 2A). During this phase transition, conformational disordering is introduced and causes changes in lateral alkyl chain packing. The intensity, bandwidth, and position of the methylene deformation mode ( $\delta$ CH<sub>2</sub>) are sensitive to the type of lateral alkyl chain packing [18-21]. In a gel phase this band is relatively sharp and intensive, with the maximum at around 1468 cm<sup>-1</sup>, indicating hexagonal packing. During the melting-chain phase transition the increase in conformational disordering disturbs the hexagonal-like lipid chain packing, which is shown by an increase in bandwidth and decrease in intensity and position of the  $\delta$ CH<sub>2</sub> band. Figure 2B shows the shift of the maximum position of the  $\delta$ CH<sub>2</sub> band as a function of temperature in dry PDP/DPPC (50/50mol%) multilayer film as an example. The sharp shift of this band to the lower frequency region indicates the phase transition

accompanied by a marked increase in *gauche* conformers and a loss of hexagonal chain packing. The temperature of the phase transition in dry PDP-doped DPPC films, derived from analysis of the  $\nu_s\text{CH}_2$  and  $\delta\text{CH}_2$  bands, commonly used indicators of phase transition in lipid layers, was the same.

According to numerous previous studies on the influence of hydration on phase-transition temperature in DPPC bilayers, we did not observe any phase transition in the investigated temperature range in the dry DPPC multilayers (Fig. 2).

**Figure 2**

*The alkyl chain region in DPPC/PDP liposomes*

The ATR-IR method allowed us to study the liposomal suspensions. After subtraction of the phosphate buffer's spectrum from that of the liposomal suspension, we were able to analyze some of the most important vibrational bands. The temperature dependencies of the  $\text{CH}_2$  symmetric stretching frequency in the DPPC and PDP/DPPC (50/50mol%) liposomes are shown in Figure 3. The shift of the  $\nu_s\text{CH}_2$  bands to higher frequencies arises from the increased number of hydrocarbon *gauche* conformers in the investigated liposomes induced by the increase in temperature. The line of the  $\nu_s\text{CH}_2$  band positions as a function of temperature adopts for DPPC and PDP/DPPC liposomes a characteristic sigmoidal shape. The temperature of the main phase transition ( $T_m$ ) of DPPC liposomes determined by this method corresponds very well with the  $T_m$  derived by *many* other techniques and was 41.9°C. The presence of PDP molecules in the DPPC liposome wall caused an increase in  $T_m$  proportional to the increase in PDP concentration. For PDP/DPPC (50/50mol%) liposomes, the temperature of the observed phase transition rose to 50.9°C (Fig. 3). The width of the transition region in PDP-doped DPPC liposomes is sharp and comparable to that in DPPC

liposomes. The gel phase in both systems has a similar *trans/gauche* population, the positions of the CH<sub>2</sub> stretching bands for both the doped and undoped liposomes were approximately equal. Small differences occurred only at higher temperatures, where the PDP/DPPC mixtures seem to be more rigid (Fig. 3).

**Figure 3**

At the phase transition temperature of the DPPC and PDP/DPPC liposomes, a sharp decrease in intensity of the methylene CH<sub>2</sub> deformation band ( $\delta$ CH<sub>2</sub>) was observed, indicating drastic changes in lateral chain packing. In the PDP-doped liposomes, the loss of hexagonal chain packing, shown by the shift to a lower frequency region and decrease in intensity of the  $\delta$ CH<sub>2</sub> band, takes place in the observed phase transition.

#### *The interfacial region in dry DPPC/PDP film*

The interfacial region of DPPC assemblies is represented by ester group vibrations. The C=O stretching band is in the region between 1750 cm<sup>-1</sup> and 1700 cm<sup>-1</sup>. After spectral deconvolution or second-derivative calculations, two or more components of the  $\nu$ C=O mode appear. The splitting of the  $\nu$ C=O band is associated with the conformational nonequivalence of the C<sub>1</sub>-C<sub>2</sub> bonds of the *sn*-1 (*trans*) and *sn*-2 (*gauche*) chains in DPPC molecules [22-24]. The band at 1742 cm<sup>-1</sup> originates from the *sn*-1 C=O group, while the corresponding band for the *sn*-2 C=O group is found at 1725 cm<sup>-1</sup> in the spectrum of dry DPPC layers. These two conformations of ester carbonyl groups cause differences in the extent of hydration of the ester part. In general, in hydrated samples a broad band contour is found in a lower frequency region.

In the dry DPPC system the maximum of  $\nu\text{C=O}$  is centered around  $1736.7\text{ cm}^{-1}$  and is almost stable in the  $10$  to  $80^\circ\text{C}$  temperature range (Fig. 4A). The presence of PDP molecules shifts the maximum of the  $\nu\text{C=O}$  band to a higher frequency. In the PDP/DPPC (50/50mol%) dry mixture this band is around  $1739\text{ cm}^{-1}$  and shifts slightly to a higher frequency region with increasing temperature, as shown in Fig. 4A. The phase transition observed in the PDP-doped DPPC dry film (Fig. 2) is not connected with any distinct changes in the interaction and conformation state of  $\text{C=O}$  groups (Fig. 4A).

#### Figure 4

##### *The interfacial region in DPPC/PDP liposomes*

In buffer, the DPPC bilayer of liposomes has a fully hydrated state. Under this condition the increase in hydration of the DPPC bilayer takes place during the main phase transition and causes the characteristic shift of the  $\nu\text{C=O}$  band position to lower frequencies (Fig. 4B). The fluidization of the bilayer leads to more water molecules reaching the deeper part of the lipid membrane and hydration of the ester groups increases. Therefore, by monitoring the positions of the  $\nu\text{C=O}$  band as a function of temperature, it is possible to observe the main phase transition in DPPC liposome dispersion. The presence of PDP molecules in DPPC liposomes influences the ester group's vibrations. At a temperature lower than the phase transition the midpoint of the broad  $\text{C=O}$  stretching band is around  $1738.5\text{ cm}^{-1}$  and is higher than in the gel state of DPPC liposomes, where the  $\nu\text{C=O}$  band lies near  $1736\text{ cm}^{-1}$  (Fig. 4B). The phase transition observed in PDP/DPPC liposomes is connected with a drastic increase in the hydration of the ester groups of DPPC molecules. However, the jump in  $\nu\text{C=O}$  vibrations at around the temperature of the phase transition is much deeper in doped than in undoped DPPC liposomes, which suggests greater changes in the hydration of  $\text{C=O}$

moieties in the PDP/DPPC system (see Fig. 4B). In the higher temperature region, after the temperature of the phase transition, both the DPPC and PDP/DPPC systems have more similar  $\nu\text{C}=\text{O}$  band positions.

*The phosphate headgroup absorption region in dry DPPC/PDP film*

In the spectrum of DPPC, the phosphate moiety of the headgroup part gives rise to several characteristic vibrations in the infrared region. Asymmetric and symmetric stretching modes for the  $\text{PO}_2^-$  groups are found near  $1250\text{ cm}^{-1}$  and  $1085\text{ cm}^{-1}$ , respectively [25,26]. Weaker single-bond P-O stretching bands are in the  $900\text{-}800\text{ cm}^{-1}$  region [25]. The R-O-P-O-R' stretching vibration is a shoulder on the  $\nu_{\text{as}}\text{PO}_2^-$  band centered near  $1060\text{ cm}^{-1}$  [26]. The band most sensitive to the state of hydration of the phospholipid bilayer is the  $\nu_{\text{as}}\text{PO}_2^-$  band. In dry DPPC film it is centered at  $1260\text{ cm}^{-1}$ , and the formation of a hydrogen bond during hydration results in a band-shift towards lower frequencies. In dry PDP/DPPC film a strong interaction by the hydrogen bond between the phenolic OH group and the  $\text{PO}_2^-$  moiety was observed. In the dry, equimolar PDP/DPPC mixture the  $\nu_{\text{as}}\text{PO}_2^-$  band position was centered in a lower frequency region, at  $1244\text{ cm}^{-1}$  [13]. The scheme of interaction between PDP and DPPC molecules can be proposed on the basis of theoretical calculations [27]. The structure of the PDP-DPPC complex is presented in Figure 5. The formation of PDP-DPPC complex gave  $14.5\text{ kcal/mol}$  (PM3 method) gain of interaction energy resulting from both van der Waals forces and H-bond interactions [27]. Thus, in DPPC bilayer PDP molecules are located parallel between DPPC molecules. The phenol groups of PDP molecules locate between hydrophilic head group of DPPC molecules. The hydrocarbon chains of both molecules lays parallel to each other and form the hydrophobic region of lipid bilayer.

**Figure 5**

*The phosphate headgroup region in DPPC/PDP liposomes*

In fully hydrated DPPC liposomes, the  $\nu_{\text{as}}\text{PO}_2^-$  band reaches  $1220\text{ cm}^{-1}$ . The  $\nu_{\text{as}}\text{PO}_2^-$  frequencies are insensitive to the main phase transition in DPPC liposomes. The phase transition observed in PDP-doped DPPC liposomes is also insensitive to this band position and is centered around  $1220\text{ cm}^{-1}$ . In the presence of water molecules, PDP is still most probably able to form a hydrogen bond with the phosphate group, but this interaction has no influence on  $\nu_{\text{as}}\text{PO}_2^-$  band position. The DFT and semi-empirical calculations showed that in the presence of water molecule the H-bond between phenol and phosphate group can still exist [27].

 **$^{31}\text{P}$  NMR measurements**

$^{31}\text{P}$  NMR is a useful analytical technique for the study of the polymorphic phase behavior of hydrated phospholipids in excess water. Lipid phosphorus exhibits large chemical shift anisotropy.  $^{31}\text{P}$  NMR line-shape is sensitive to different types of motion of lipid molecules. Dynamics of phospholipid bilayer systems consist primarily of the rapid rotation of the lipid molecules about their long axes and they have a characteristic broad spectrum with a high-field peak and low-field shoulder [15,16] (Fig. 6A). In non-bilayer configurations, lateral diffusion takes place and in the micellar, cubic, or rhombic phase isotropic motion leads to a narrow, symmetric  $^{31}\text{P}$  NMR spectrum [15,16] (Fig. 6B).

**Figure 6**

$^{31}\text{P}$  NMR may be usefully applied to both model and biological membrane systems. In PDP-doped DPPC dispersions, a lamellar-nonlamellar phase transition appears. The  $^{31}\text{P}$  NMR

spectra obtained from aqueous ( $D_2O$ ) dispersions of the equimolar PDP/DPPC mixture at different temperatures are illustrated in Fig. 7A,B. At a temperature below the temperature of the phase transition, the PDP/DPPC (50/50mol%) dispersion adopts a lamellar structure. At 25°C the  $^{31}P$  NMR spectrum is very broad, with a low-field shoulder and high-field peak (Fig. 7A). This line-shape is characteristic of the lamellar (anisotropic) phase of phospholipids. At a higher temperature (60°C), above the temperature of the observed phase transition, this system adopts an isotropic phase. The  $^{31}P$  NMR spectrum exhibits a symmetric and narrow peak centered at 0 ppm (Fig. 7B). This isotropic signal indicates the presence of regions of high curvature in which lipid diffusion can result in the complete motional averaging of the chemical shift anisotropy. Thus the isotropic phase can be represented by micellar, cubic, or rhombic structures, but the  $^{31}P$  NMR technique does not allow us to determine which one the PDP/DPPC mixture adopts. The lamellar phase is present for DPPC liposomes both at 25 °C and 60°C, as was shown in Fig.7 C, D, respectively. The narrowing of the lamellar  $^{31}P$  NMR line-shape of DPPC liposomes present at higher temperature (60°C) derives from the increase of anisotropic motion accompanied by the rise of lipid bilayer fluidity after the main phase transition.

**Figure 7**

## Summary

As an amphiphilic compound, 3-pentadecylphenol (PDP) easily incorporates into the DPPC bilayer. The interactions of PDP with the DPPC bilayer occur through a combination of hydrogen-bonding and hydrophobic and conformational interactions. Characteristic of PDP is that it causes the appearance of a phase transition in dry DPPC film [13]. It was a chain-



melting phase transition accompanied by an increase in *gauche* conformers in the hydrocarbon tails and by the disorder of hexagonal chain packing. It is commonly known that the level of hydration determines the structure and physicochemical properties of a lipid bilayer. In order to mimic the physiological conditions of a biomembrane it is necessary to study liposomal suspensions. In PDP/DPPC liposome dispersions the phase transition was connected with changes in both the hydrophobic and hydrophilic regions of lipid aggregates. This process was accompanied by an increase in *gauche* conformers, a disturbance of hexagonal chain packing, and a rise in the hydration of the ester groups of the DPPC molecules. The sharp increase of  $\nu_{as}CH_2$  and decrease of  $\nu C=O$  and  $\delta CH_2$  frequency positions were observed around the phase transition temperature.

It is quite interesting that the PDP compounds did not form hydrogen bonds with the ester part of the dry DPPC bilayer. However, in dry PDP/DPPC (50/50mol%) film the  $\nu C=O$  band was shifted to a higher frequency region compared with that in pure DPPC layers, which suggests that changes occurred in the conformational state of these groups. The same higher-frequency shift of the  $\nu C=O$  band was observed in the fully hydrated state of the PDP/DPPC (50/50mol%) bilayer wall of liposomes at lower temperatures, before the phase transition temperature. The state of hydration of the lipid bilayer does not influence the type of interaction between the phenolic OH group and the ester part of DPPC.

Under dry lipid film conditions, PDP compounds form hydrogen bonds with the phosphate groups of DPPC lipids, which was shown by the lower-frequency shift of the  $\nu_{as}PO_2^-$  band [13] and by DFT and semi-empirical calculations [27]. In PDP/DPPC (50/50mol%) liposomes, where the PDP-concentration is very high, the  $\nu_{as}PO_2^-$  vibrations are centered around  $1220\text{ cm}^{-1}$ , which is characteristic of the fully hydrated state of phosphate groups. Most probably, the  $PO_2^-$  groups of DPPC molecules in the liposome suspension

interact by a hydrogen bond with both water and PDP molecules. Nevertheless, the presence of doped compounds does not influence  $\nu_{\text{as}}\text{PO}_2^-$  band position in a hydrated DPPC bilayer.

In DPPC liposomes at 41.9°C, the lamellar structures go from a gel phase to a liquid-crystal phase. In PDP-doped DPPC liposomes, a lamellar-nonlamellar phase transition takes place. The  $^{31}\text{P}$  NMR experiment showed the presence of an isotropic phase at a temperature above that of the phase transition of PDP/DPPC (50/50mol%) liposomes. The data at 60°C for PDP/DPPC (50/50 mol%) liposomes showed a sharp and pure isotropic peak centered at 0 ppm. Several lipid-assembly architectures will produce such a spectrum including micellar, cubic, and orthorhombic phases. Many long-hydrocarbon-chain phenolic lipids can induce in higher concentrations non-layered structures of the micellar or  $\text{H}_{\text{II}}$  type in lipid membranes [28]. This is responsible for the increase in permeability of a biomembrane in the presence of phenolic lipids and it changes the activities of membrane proteins [2]. PDP molecules create an isotropic phase represented by regions of high curvature. Such structures can be involved in membrane fusion phenomena (including related processes such as exo- and endocytosis) [2,15] and transbilayer transport processes (including lipid flip-flop and facilitation of transport). The advantage of PDP over other membrane fusion compounds is the low cost of obtaining it from the commonly accessible shell of the cashew nut. Incorporation of PDP into the lipid membrane of liposomes can facilitate the fusion of liposomes with cell membrane and enable the release of drug molecules from the interior of liposomes to target cells.

The phase transition of lamellar structures to isotropic ones in PDP/DPPC (50/50mol%) dispersion is accompanied by a larger shift of the  $\nu\text{C}=\text{O}$  band position than in DPPC liposomes. The isotropic phase is characterized by a lower *gauche* conformer population in the hydrocarbon chains, which makes the isotropic structure more rigid than the DPPC bilayer. Isotropic PDP/DPPC aggregates reveal a character of vibrations of the phosphate and carbonyl groups closely related to that of a DPPC bilayer, which suggests a

similar conformational and hydration state of the hydrophilic region in the pure and doped DPPC structures.

At lower temperatures, before the phase transition, a bilayer structure of PDP/DPPC mixtures was observed. The PDP/DPPC bilayer had a conformational state of the hydrophobic part similar to that of DPPC. The position of the CH<sub>2</sub> stretching vibrations was the same for both undoped DPPC and doped DPPC liposomes. More significant changes were observed in the hydrophilic part and they were connected with different conformational states of the carbonyl groups in the DPPC molecules.

## References

- [1] A. Kozubek and J.H.P. Tyman, Cereal grain resorcinolic lipids: mono and dienolic homologues are present in rye grains, *Chem. Phys. Lipids*. 78 (1995) 29-35.
- [2] A. Kozubek and J.H.P. Tyman, Resorcinolic lipids, the natural nonisoprenoid phenolic amphiphiles and their biological activity, *Chem. Rev.* 99 (19[21] R. G. Snyder, Vibrational spectra of crystalline n-paraffins : II. Intermolecular effects, *J. Mol. Spectrosc.*, 7 (1961) 116-144.
- [3] E. Wenckert, E. Loeser, S.N. Mahapatra, F. Schenker and E.M.J. Wilson, Wheat Bran Phenols, *J. Org. Chem.* 29 (1964) 435-439.
- [4] P.N. Reusch and H.L.J. Sadoff, 5-n-Alkylresorcinols from encysting *Azotobacter vinelandii*: isolation and characterization, *J. Bacteriol.* 139 (1979) 448-453.
- [5] C. Bandypadhyay, A. Gholap and V.J.R. Mamdapur, Characterization of alkenylresorcinol in mango (*Mangifera indica* L.) latex, *Agric. Food Chem.* 33 (1985) 377-379.
- [6] H. Itakowa, N. Totsuka, K. Nakahara, M. Maezuru, K. Takeya, M. Kondo, M. Inamatsu and H. Morita, A quantitative structure-activity relationship for antitumor activity of long-chain phenols from *Ginkgo biloba* L., *Chem. Pharm. Bull.* 37 (1989) 1619-1621.
- [7] A. Kozubek, M. Nietubyc and A.F. Sikorski, Modulation of the activities of membrane enzymes by cereal grain resorcinolic lipids, *Z. Naturforsch.* 47c (1992) 41-46.
- [8] I. Kubo, K. Komatsu and M. Ochi, Molluscicides from the cashew *Anacardium occidentale* and their large-scale isolation, *J. Agric. Food Chem.* 34 (1986) 970-973.
- [9] S.P. Chandran, J. Ghatak P.V. Satyam and M. Sastry, . Interfacial deposition of Ag on Au seeds leading to  $\text{Au}_{\text{core}}\text{Ag}_{\text{shell}}$  in organic media. *J. Colloid Interf. Scien.* 312 (2007) 498-505.
- [10] T. Kato, N. Mizoshita and K. Kishimoto, Functional liquid-Crystalline Assemblies: Self-Organized Soft Material, *Angew. Chem. Int. Ed.* 45 (2006) 38-68.
- [11] T. Nakatsu and Z. Chen, United States Patent No. 07/685,285, 15.05.1991

- [12] R. Amorati, G.F. Pedulli, L. Valgimigli, O.A. Attanasi, P. Filippone, Ch. Fiorucci and R. Saladino, Absolute rate constants for the reaction of peroxy radicals with cardanol derivatives, *J. Chem. Soc., Perkin Trans. 2* (2001) 2142-2146.
- [13] K. Ciesik, A. Koll and J. Grdadolnik, Structural characterization of a phenolic lipid and its derivative using vibrational spectroscopy, *Vib. Spectrosc.* 41 (2006) 14-20.
- [14] H.H. Mantsch and D. Chapman, *Infrared Spectroscopy of Biomolecules* (Wiley-Liss, New York, 1996).
- [15] P.R. Cullis and B. De Kruijff, Lipid Polymorphism and functional roles of lipids in biological membranes, *Biochim. Biophys. Acta* 559 (1979) 399-420.
- [16] P.R. Cullis and B. De Kruijff, The polymorphic phase behaviour of phosphatidylethanolamines of natural and synthetic origin. A  $^{31}\text{P}$  NMR study, *Biochim. Biophys. Acta* 513 (1978) 31-42.
- [17] M. Kodoma, M. Kuwabara and S. Seki, Successive phase-transition phenomena and phase diagram of the phosphatidylcholine-water system as revealed by differential scanning calorimetry 689 (1982) 567-570.
- [18] K. Cieřlik-Boczula, B. Czarnik-Matusewicz, M. Perevozkina, A. Filarowski, N. Boens, W. De Borggraeve and A. Koll, ATR-IR spectroscopic study of the structural changes in the hydrophobic region of ICPAN/DPPC bilayers, *J.Mol. Struct.* 878 (2008) 162-168.
- [19] D. G. Cameron, H. L. Casal, E. F. Gudgin and H. H. Mantsch, The gel phase of dipalmitoyl phosphatidylcholine. An infrared characterization of the acyl chain packing, *Biochim. Biophys. Acta* 596 (1980) 463-467.
- [20] D. G. Cameron, H. L. Casal and H. H. Mantsch, Characterization of the pretransition in 1,2-dipalmitoyl-sn-glycero-3-phosphocholine by Fourier transform infrared spectroscopy, *Biochem.* 19 (1980) 3665-3672.

- [21] R. G. Snyder, Vibrational spectra of crystalline n-paraffins : II. Intermolecular effects, J. Mol. Spectrosc. 7 (1961) 116-144.
- [22] S.F. Bush, H. Levin and I.W. Levin, Cholesterol-lipid interactions: An infrared and raman spectroscopic study of the carbonyl stretching mode region of 1,2-dipalmitoyl phosphatidylcholine bilayers, Chem. Phys. Lipids 27 (1980) 101-111.
- [23] I.W. Levin, E. Mushayakarara and R. Bittman, Vibrational assignment of the sn-1 and sn-2 chain carbonyl stretching modes of membrane phospholipids, J. Raman Spectrosc. 13 (1982) 231-234.
- [24] E. Mushayakarara and W.I. Levin, Determination of acyl chain conformation at the lipid interface region: Raman spectroscopic study of the carbonyl stretching mode region of dipalmitoyl phosphatidylcholine and structurally related molecules, J. Phys. Chem. 86 (1982) 2324-2327.
- [25] H.L. Casal and H.H. Mantsch, Polymorphic phase behaviour of phospholipid membranes studied by infrared spectroscopy, Biochim.Biophys. Acta 779 (1984) 381-401.
- [26] J.L.R. Arronda and F.M. Macarulla, Infrared spectroscopy of phosphatidylcholines in aqueous suspension a study of the phosphate group vibrations, Biochim.Biophys. Acta 794 (1984) 165-168.
- [27] K. Cieřlik-Boczula, A. Koll, Characterisation of the structure of PDP-DPPC bilayers by DFT and PM3 calculations, Polish J. Chem. 81 (2007) 1081-1093.
- [28] A. Kozubek, R.A. Demel, The effect of 5-(n-alk(en)yl) resorcinols from rye on membrane structure, Biochim. Biophys Acta 642 (1981) 242-251.

**Figure legends:**

**Figure 1.** Schematic representation of the structure of 3-pentadecylphenol (PDP) (left side) and dipalmitoylphosphatidylcholine (DPPC) molecule (right side). The PDP and DPPC are amphiphilic compounds with hydrophilic head-group and hydrophobic hydrocarbon-chain region.

**Figure 2.** (A) The maxima of the  $\nu_s\text{CH}_2$  band at different temperatures for the PDP/DPPC (50/50mol%) (■) and DPPC (○) dry multilayer film. (B) Frequency of methylene deformation modes ( $\delta\text{CH}_2$ ) of the aliphatic tails of PDP/DPPC (50/50mol%) (■) and DPPC (○) dry multilayer film as a function of temperature.

**Figure 3.** Wavenumber displacements of the bands due to the symmetric  $\text{CH}_2$  stretching vibration compared for DPPC liposomes (○) and PDP/DPPC (50/50mol%) (■) depending on temperature.

**Figure 4.** Temperature dependence of the position of the peak maximum of the  $\text{C}=\text{O}$  stretching band contour in the infrared spectra of dry DPPC (○) and PDP/DPPC (50/50 mol%) (■) film (A) and of buffer dispersion of DPPC (○) and PDP/DPPC (50/50mol%) (■) liposomes (B).

**Figure 5.** The model structure of dipalmitoylphosphatidylcholine (DPPC) and 3-pentadecylphenol (PDP) complex derived from PM3 calculation. There is H-bond between  $\text{P}=\text{O}$  and phenolic OH group with 2,734 Å of  $\text{O}(\text{H})\cdots\text{O}$  distance. C-blue, H-grey, O-red, P-yellow, N-dark blue [27].

**Figure 6.** Schematic representation of  $^{31}\text{P}$  NMR line-shape for different lipid aggregates.

**Figure 7.** Proton-decoupled  $^{31}\text{P}$  NMR spectra obtained from aqueous ( $\text{D}_2\text{O}$ ) dispersions of PDP/DPPC (50/50mol%) at a temperature before (25°C) (A) and after (60°C) (B) the phase transition and for DPPC liposomes at 25°C (C) and at 60°C (D).






Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7