Nicotinic actions on neuronal networks for cognition: General principles and long-term consequences

Rogier B. Poorthuis, Natalia A. Goriounova, Jonathan J. Couey, Huibert D.
Mansvelder

► To cite this version:

Rogier B. Poorthuis, Natalia A. Goriounova, Jonathan J. Couey, Huibert D. Mansvelder. Nicotinic actions on neuronal networks for cognition: General principles and long-term consequences. *Biochemical Pharmacology*, 2009, 78 (7), pp.668. 10.1016/j.bcp.2009.04.031 . hal-00509503

HAL Id: hal-00509503

<https://hal.science/hal-00509503>

Submitted on 13 Aug 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Nicotinic actions on neuronal networks for cognition:
General principles and long-term consequences

Authors: Rogier B. Poorthuis, Natalia A. Goriounova,
Jonathan J. Couey, Huibert D. Mansvelder

PII: S0006-2952(09)00338-4
DOI: doi:10.1016/j.bcp.2009.04.031
Reference: BCP 10169

To appear in: *BCP*

Received date: 1-4-2009
Accepted date: 27-4-2009

Please cite this article as: Poorthuis RB, Goriounova NA, Couey JJ, Mansvelder HD, Nicotinic actions on neuronal networks for cognition: General principles and long-term consequences, *Biochemical Pharmacology* (2008), doi:10.1016/j.bcp.2009.04.031

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Research Update for **Special Issue** ‘Nicotinic Receptor-based Therapeutics’

Nicotinic actions on neuronal networks for cognition:

General principles and long-term consequences

Rogier B. Poorthuis^{1,2}, Natalia A. Goriounova^{1,2}, Jonathan J. Couey^{1,2,3}, Huibert D.
Mansvelder¹

¹Department of Integrative Neurophysiology, CNCR, Neuroscience Campus, VU
University Amsterdam, The Netherlands.

²These authors contributed equally to this review article

³Present address: NTNU-Kavli Institute for Systems Neuroscience, Centre for the
Biology of Memory, Trondheim, Norway.

Running title: Nicotinic actions on neuronal networks for cognition

Corresponding Author:

Huibert D. Mansvelder

Department of Integrative Neurophysiology, CNCR,

Neuroscience Campus Amsterdam, VU University Amsterdam,

De Boelelaan 1085

1081 HV, Amsterdam,

The Netherlands.

e-mail: huibert.mansvelder@cncr.vu.nl

Number of words

Abstract: 198

Body text: 5409

Number of references: 134

Abstract

Nicotine enhances cognitive performance in humans and laboratory animals. The immediate positive actions of nicotine on learning, memory and attention are well-documented. Several brain areas involved in cognition, such as the prefrontal cortex, have been implicated. Besides acute effects on these brain areas and on brain function, a picture is emerging showing that long-term consequences of nicotine exposure during adolescence can be detrimental for cognitive performance. The majority of adult smokers started the habit during adolescence. Our knowledge on the types of nicotinic receptors in the brain areas that are candidates for mediating nicotine's effects is increasing. However, much less is known about the underlying cellular mechanisms. A series of recent studies have uncovered exciting features of the mechanisms by which nicotine alters prefrontal cortex neuronal activity, synaptic plasticity, gene expression and cognitive function, and how these changes may have a lasting effect on the developing brain. In this review, we discuss these exciting findings and identify several common principles by which nicotinic receptor activation modulates cortical circuits involved in cognition. Understanding how nicotine induces long-term changes in neuronal circuits and alters plasticity in the prefrontal cortex is essential to determining how these mechanisms interact to alter cognition.

Key words: Acetylcholine, Nicotine, Prefrontal cortex, Neuronal networks, Synaptic plasticity, Development

Nicotine and cognitive function

It has long been recognized that nicotine, the addictive substance in cigarettes, can have stimulating effects on brain function. The link to the psychoactive effects lies in the fact that nicotine stimulates nicotinic acetylcholine receptors (nAChRs) that are normally activated by the endogenous neurotransmitter acetylcholine (ACh) and interfere with cholinergic signalling. By boosting signal-to-noise ratio, the cholinergic system in the brain is important for a variety of cognitive functions, such as learning, memory and attention processes that involve many different brain regions [1]. Specifically in prefrontal cortex (PFC) cholinergic signals are involved in attention [2]. The entire cortex is innervated by projections from the basal forebrain that release ACh [1]. Nicotinic AChRs, with their highly dense and widespread distribution in neocortical and subcortical areas [3], are an important component of the ACh system. It comes therefore as no surprise that nicotine can affect cognitive processes both in humans and rodents [4-7].

The strong involvement of cholinergic signalling in cognitive functions was shown in studies where selective depletion of ACh in target areas or lesions of ACh projections was applied (reviewed in [2]). In tests that assess attention behaviour, which strongly relies on prefrontal cortex function, selective lesions in the basal forebrain cholinergic system as well as depletion of ACh directly in PFC result in a reduction in attention performance [8-10]. These data agree with studies showing a positive correlation between cortical acetylcholine release and attentional demand [11]. Attention deficits induced by lesions can be ameliorated by pharmacological agents that augment cholinergic signalling in the PFC [12]. In view of these findings, it is not surprising that activation of nAChRs can affect cognitive functions and, in particular, attentional performance. With a few exceptions, nAChRs agonists enhance cognitive

performance, while antagonists have the adverse effect [7]. For instance, nicotinic agonists improve working memory function [13] and overcome deficits induced by lesioning cholinergic innervation of the hippocampus [14, 15]. In contrast, nicotinic antagonists for different nicotinic subtypes applied to the hippocampus impair working memory function in the radial arm maze [16]. Some aspects of this enhancement appear to rely particularly on nicotinic receptor signalling in the prefrontal cortex [17]. Also in humans, nicotine was shown to enhance activity in the prefrontal cortex [18]. Finally, the enhancement of attention performance by nicotinic agonists was shown in a number of studies [7, 12, 17, 19-22].

Very little is known about the mechanisms underlying nicotine's power as cognitive enhancer. Even less is known about the long-term consequences of nicotine exposure for cognitive performance. Our understanding of how nicotinic compounds affect cortical circuits involved in cognition is far from complete, but in this review recently uncovered important new insights in these mechanisms will be highlighted.

Nicotinic receptors

Nicotinic AChRs belong to the cys-loop ligand-gated ion-channel family [23]. This group of pentameric transmembrane proteins form a water-filled pore upon binding of neurotransmitter after which charged ions can flow over the membrane. Twelve genes have been identified encoding neuronal nicotinic receptors [for review see 24]. Each gene encodes a subunit of the receptor that can be classified into α -type subunits and non- α -type subunits, based on the presence or absence, respectively, of a pair of cysteine amino acids [23-25, reviewed in 26, 27-29]. This cysteine pair is important for agonist binding and it has been thought therefore that α -subunits at least in part regulate agonist binding. In the central nervous system 9 α -subunits ($\alpha 2$ – $\alpha 10$)

1 encoded by CHRNA2–10) and 3 β -type subunits ($\beta 2$ – $\beta 4$; CHRNB2–4) are expressed.
 2 These subunits assemble in different stoichiometries to form the pentameric channel,
 3
 4 and the subunit composition of nAChRs varies depending on the brain region [for
 5
 6 review see 23, 24, 30, 31–34]. Nicotinic receptors can either assemble as homomeric
 7
 8 or heteromeric channels. Heteromeric channels are formed by a combination between
 9
 10 α and β subunits, whereas some α subunits can assemble into a homomeric channel.
 11
 12 When opened, the nicotinic receptor is a cation selective channel which permits flow
 13
 14 of Na^+ , K^+ and Ca^{2+} across the membrane. At normal resting membrane potential this
 15
 16 leads to a depolarizing current. The impact of nAChR activation on neuronal function
 17
 18 strongly depends on the subunit composition of the nAChRs. Each subunit
 19
 20 combination has its own activation and desensitization characteristics and has
 21
 22 different single channel conductance and agonist selectivity, potentially leading to
 23
 24 different kinetics of depolarizing currents in the target cell [15, 23].

25
 26
 27
 28
 29
 30
 31 The two most abundant nicotinic receptors in the brain are receptors that
 32
 33 contain $\alpha 4\beta 2$ or $\alpha 7$ subunits. $\alpha 4\beta 2^*$ receptors have a very high affinity for nicotine
 34
 35 and desensitize at low concentrations of nicotine, corresponding to blood
 36
 37 concentrations experienced by smokers [23, 35]. In contrast, the homomeric channels
 38
 39 containing $\alpha 7$ subunits have a low affinity for nicotine but do not desensitize at low
 40
 41 nicotinic concentrations [36, 37]. These phenomena will have significant impact on
 42
 43 how these receptors are activated in neuronal networks (see below). Another
 44
 45 distinctive feature of these receptors is their permeability for calcium. Amino acids
 46
 47 lining the pore of the protein largely determine the ion-selectivity of the channel. $\alpha 7^*$
 48
 49 nicotinic receptors are highly permeable for calcium compared to $\alpha 4\beta 2^*$ receptors
 50
 51 [38]. They serve, therefore, a distinguished role because this calcium influx can
 52
 53 influence cellular processes like neurotransmitter release and synaptic plasticity
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

directly. It has been suggested that $\alpha 7^*$ nicotinic receptors perform a complementary role to NMDA receptors. These channels are also calcium permeable, but are only opened at a more depolarized membrane potential [39].

Nicotinic AChRs in the PFC

There is ample evidence that nAChR activation in general affects attention performance [4, 7, 40, 41], but much less is known about the nAChR subtypes and brain areas involved. Several studies point to a specific role of cholinergic signalling in the medial prefrontal cortex and attention performance [9, 17]. However, only a limited number of studies have addressed the role of nAChR subtypes in the PFC and their role in attention behavior. Infusion of α -bungarotoxin, an $\alpha 7^*$ nicotinic receptor antagonist, into the prefrontal cortex impairs performance in a delayed response task, which requires effortful processing for response selection [42]. $\beta 2$ -containing nAChRs have also been implicated in mediating effects of nicotine on attention performance. Nicotine decreased response latency and reduced incorrect responses in the 5-choice serial reaction time task [43], a test that assesses sustained attention performance [44]. These effects of nicotine were completely antagonized by dihydro- β -erythroidine (DH β E), a specific blocker for $\beta 2$ -containing nicotinic receptors. In this study, methyllycaconitine (MLA), a somewhat selective blocker of $\alpha 7$ -containing nAChRs, did not alter the effects of nicotine [43]. However, genetic approaches assessing the role of nAChRs have shown that $\alpha 7^*$ receptors do have a role in attention. Knockout mice lacking the gene for the $\alpha 7$ nAChR subunits showed impaired task acquisition and a higher rate of omissions in the 5-choice serial reaction time task [45, 46]. Since the mice used for these studies lacked $\alpha 7^*$ nAChR expression throughout their brains, it is not known whether the impairment in

attention performance was attributable to the lack of $\alpha 7^*$ receptors specifically in the PFC, let alone whether the effects were attributable to a specific type of cells in the PFC neuronal circuits. If nicotinic compounds are to be designed as cognitive enhancers for therapeutic use, a detailed understanding of how nAChR activation affects PFC microcircuits will be indispensable.

Nicotine's modes of action

The large body of evidence demonstrating that nAChRs can affect cognitive processes offers an enticing chance to link protein function to complex behaviour [4, 5, 7, 47]. However, to understand the mechanisms involved at the level of neuronal networks, there are several bridges yet to be built. Typically, a cortical microcircuit consists of a set of excitatory and inhibitory neurons that are interconnected using highly dynamic connections. To understand how nAChR activation in the prefrontal cortex affects cognitive behaviour, an understanding is needed of how prefrontal cortical microcircuits generate output from the inputs they receive. One of the challenges is to distinguish the cell types and the connectivity patterns that are present in the prefrontal cortex circuitry. The next step is to understand how nicotine alters the functionality of these neuronal circuits. General principles on how nicotine affects microcircuits in the brain will depend on (i) which cell types in the circuits express nicotinic receptors and (ii) from which subunits these receptors are made of. The latter strongly determines activation and desensitization kinetics, agonist sensitivity and ion-specific channel conductance. (iii) The sub-cellular location of the receptor will determine what stage of information processing is affected, since nicotinic receptors can be found on dendritic, somatic, axonal and presynaptic compartments. Nicotinic AChRs expressed in axons or axon terminals can alter release of neurotransmitter at

specific sites, even independent of action potential depolarization [48-52]. This often results in an increased probability of release at these sites, changing how the information carried by these synapses enters and is processed by the cortex. Alternatively, nicotinic AChRs can alter whole neuron functions by changing resting membrane potentials. These nicotinic currents can drastically alter the availability of Na and K channels available for action potential generation (through inactivation), affect resting membrane potential (depolarization), and even potentially alter regional voltage signals via shunting [36, 53-55].

(iv) Given the fact that nAChR can be continuously activated by endogenous ACh, nAChR desensitization can affect ongoing neuronal activity just as nAChR activation [36, 37, 56, 57]. Therefore, the dynamics of endogenous cholinergic signalling will play an important role in the effects of exogenously administered nicotinic compounds. The dynamics of cholinergic signalling in the cortex during ongoing behaviour has long remained enigmatic. Due to low temporal resolution of classical techniques such as microdialysis, cholinergic modulation was considered to occur on a timescale of minutes [11, 58]. However, recently it was shown that cholinergic signalling involved in attention has a much faster temporal dynamics, suggesting that cholinergic signals convey more information than just arousing a network into an 'excited' state [2, 59]. How the interplay between nAChR activation and deactivation on a subsecond time scale by endogenous ACh and exogenously applied agonists will affect cortical neuronal network activity remains to be elucidated.

(v) Finally, it will be of significance whether nAChRs are activated by direct synaptic contact or via a slower process like volume transmission. Spill over and

diffusion of ACh will result in different activation and desensitization profiles compared to targeted fast synaptic release.

These factors will combine to alter neuronal network properties and will be central to understanding nicotine's effects on higher cognitive functions. Although we are only beginning to understand how nicotine is affecting neuronal circuits in the prefrontal cortex, several features have now been uncovered, some of which show similarities to cholinergic modulation of other cortical areas, emphasizing that common principles may exist guiding nicotinic modulation of cortical circuits.

Nicotinic modulation of thalamocortical communication

One of the first recognized functions for nAChRs in the central nervous system was its role in enhancing neurotransmitter release [48]. As first described in chicken medial habenula-interpeduncular synapses and later in the mossy fiber synapse in the rat hippocampus, nicotine augments synaptic release of glutamate via presynaptic receptors [48, 60]. The facilitating effect of nicotine was dependent on the extracellular calcium concentration, and nicotine application leads to a higher calcium signal in mossy fiber boutons. Depending on the subunit composition and precise location, nicotinic receptors can enhance presynaptic neurotransmitter release either through depolarization or direct calcium influx or both [61].

Nicotinic AChRs on axonal projections play a key role in regulating the transmission of thalamic information to the cortex [50, 52, 62-64]. Transdermal administration of nicotine to non-smokers does not affect cochlear activity but does affect the neural transmission of acoustic information [65]. A similar result was observed in the rat auditory cortex [66]. Critically, in this study antagonists of nAChRs reduced the evoked signal in the cortex, suggesting that endogenous ACh

acts through nAChRs to regulate thalamic transmission. The barrel cortex of the rat is perhaps the best studied model of thalamocortical transmission, and here too nicotinic agonists can alter cortical processing. Topical application of nicotinic agonist to the exposed cortex *in-vivo* increased the size of a whisker's functional representation in the cortex [67]. Earlier recordings in thalamocortical slices from the barrel cortex support this result by demonstrating that thalamic synapses, unlike intracortical synapses, are modulated by nAChRs [68]. Even in the visual cortex, nicotine can increase responsiveness to visual stimuli [18, 69].

Although the prefrontal cortex is thought to be a higher order processing area, it receives thalamic input from the dorsal medial nucleus of the thalamus. Thalamocortical glutamatergic transmission to the prefrontal cortex is augmented by the activation of nAChRs [52, 55, 63, 70, 71]. Autoradiographic labelling of nAChRs was reduced after lesions in the medial dorsal thalamus (MDT). This suggested that nAChRs are present on thalamocortical terminals and could potentially alter thalamic information processing in the PFC. When neurons in the MDT are stimulated *in vivo* action potentials are elicited in the prefrontal cortex. Infusing nicotine locally into the PFC enhanced the response elicited in the prefrontal cortex. Microdialysis experiments showed that nicotine induced glutamate release in the PFC which could be blocked by DH β E [63]. Lesioning the MDT strongly reduced the augmentation of glutamatergic inputs to layer V pyramidal neurons by nicotine [52]. This suggests that among the glutamatergic inputs received by layer V pyramidal neurons, nicotine selectively stimulates thalamic inputs (Figure 1). As with thalamocortical inputs to somatosensory cortex, the nAChRs responsible for the augmentation by nicotine were located away from the presynaptic terminal, most likely on the axons themselves [50, 52]. These nicotinic mechanisms differ from the mechanisms by which nicotine

1 increases excitatory transmission in the hippocampus and VTA, where activation of
2 $\alpha 7^*$ receptors leads to a direct stimulation of glutamate release [51, 60]. Support for
3
4 modulatory effects of presynaptic nAChRs activation in the PFC comes from a variety
5
6 of approaches including electrophysiological recordings and assay of release from
7
8 isolated nerve terminals [72-74]. A recent study testing the relative contribution of
9
10 $\beta 2^*$ nAChRs vs. $\alpha 7^*$ nAChRs on glutamatergic synaptosomes from PFC [72]
11
12 demonstrated that both $\alpha 7^*$ and non $\alpha 7^*$ nAChRs appear to be important although
13
14 each modulates excitatory amino acid (EAA) release via distinct mechanisms. Taken
15
16 together, these data suggest that nicotine selectively increases activity of inputs from
17
18 the thalamus to the cortex over other glutamatergic synapses.
19
20
21
22
23

24 Understanding how nAChRs can affect the function of cortical pyramidal
25
26 neurons is essential to understanding nicotine's effects on cognition. However, the
27
28 task is significantly more complicated. While so many aspects of pyramidal cell
29
30 function are well described, nAChRs are rarely found on pyramidal cells in the cortex
31
32 [55, 75]. A recent study has found nAChRs on layer VI pyramidal cells [71] (Figure
33
34 1). This specific population represents pyramidal cells projecting back to the
35
36 thalamus. Despite the fact that nAChRs are rarely found on pyramidal cells, nicotine
37
38 can still affect their function in many ways. In addition to glutamatergic inputs to
39
40 layer V pyramidal neurons, glutamatergic inputs to several types of layer V
41
42 interneurons were also excited by nicotine with a similar pharmacological profile
43
44 [55]. Although it was not shown in the study, it is tempting to speculate that these
45
46 nicotine-sensitive glutamatergic inputs to interneurons were of thalamic origin, but
47
48 this waits further testing. Nicotinic regulation of excitatory inputs to inhibitory
49
50 interneurons could serve to balance excitation and inhibition in the prefrontal cortex,
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

which is thought to be crucial for cortical functioning and information processing [76].

Cortical interneurons and nicotinic actions

Inhibitory neurons of the neocortex comprise a comparatively more diverse population of cells than excitatory cells. At least two types of interneurons are recognized to be morphologically and functionally distinct classes: fast spiking cells (FS) and low-threshold spiking cells (LTS) [76-81]. Fast spiking cells (FS) are physiologically equipped for high frequency firing, show little adaptation, and have been shown to synapse on or near the somata of their target cells [81-84]. As such, they occupy an ideal functional and morphological position to regulate the input window of pyramidal cells [85]. At least one study in the somatosensory cortex has demonstrated this functional position for FS cells [86]. Their functional role also appears to extend to regulating plasticity in this microcircuit [87]. This association with thalamic inputs has also been confirmed in the PFC [88]. While there is some disagreement as to whether this FS cells express nAChRs, this discrepancy appears to be species specific. Studies in rodents have failed to find nAChRs on FS cells in the cortex [55, 75] (Figure 1). In contrast, in at least one study FS cells in human cortex appear to express nAChRs [89]. FS cells appear to be important in regulating the precise timing of information coming into the cortex, and there is emerging consensus evidence that LTS interneurons play a role in shaping feedforward inhibition between excitatory cells [90, 91]. Like FS cells, LTS cells target specific dendritic subdomains of their target pyramids. In contrast to FS cells which are thought to regulate target cell activation and activity, LTS cells appear to regulate specific inputs to pyramidal cell apical dendrites, as well as mediating interlaminar feedforward inhibition in the

cortex. These interneurons express large nicotinic currents, and excitatory input to these cells is also enhanced by nicotine [55, 75, 92]. A third class of interneurons, identified based on their firing properties in response to depolarizing current steps, regular-spiking non-pyramidal neurons, were also excited by nicotine [55] (Figure 1).

Nicotinic AChR activation and synaptic plasticity

Synaptic plasticity is critically important for cognitive function, and in particular, synaptic plasticity in the PFC has been directly associated with attention and working memory [93]. The relative timing of action potentials in pre- and postsynaptic neurons has a profound impact on the induction of long-term potentiation or depression. When a presynaptic spike precedes a postsynaptic spike within a short time window of several tens of milliseconds, LTP is induced. The reverse order of spike-timing results in long term depression (LTD) [94, 95]. In mouse PFC, nicotine strongly affects this timing-dependent synaptic plasticity, which is called spike-timing-dependent plasticity (STDP). Stimulation of nicotinic AChRs in PFC modifies STDP induced by pairing stimulation of the excitatory inputs to PFC layer 5 pyramidal neurons with postsynaptic spikes elicited 5 ms after each synaptic response [55]. This coordinated stimulation induced robust LTP; however, when the same stimulus paradigm was applied in the presence of nicotine concentrations experienced by smokers, LTP was eliminated and a depression of the excitatory inputs to these cells was observed. Which nAChRs on what neurons are responsible for this effect? As discussed above, mouse PFC layer 5 pyramidal neurons do not express nicotinic receptors themselves. Rather, the nAChRs involved in the nicotinic modulation of LTP induction increase inhibitory GABAergic inputs to the pyramidal cells, as the nicotinic modulation of plasticity was abolished by inhibitors of GABA type A

(GABA_A) receptors. As described above, LTS and RSNP GABAergic interneurons found in the PFC layer 5 express nAChR subunits on their soma that activate these neurons when nicotine is present. FS interneurons are excited indirectly by nAChRs that increase glutamatergic excitation of those cells. Thus, nicotine exposure enhances inhibitory input to the layer V pyramidal neurons through both direct and indirect excitation of inhibitory GABA interneurons (Figure 1).

Studies in other cortical areas indicate that increases in postsynaptic calcium concentration are critical for the induction of synaptic plasticity [96-98]. Using two-photon imaging of intracellular calcium levels, it was found that action potentials that propagated from the soma into the dendrites of layer 5 pyramidal cells elicited increases in dendritic calcium concentration. Nicotine enhanced the GABA input to the same dendrites, resulting in less calcium entry, likely due to failure of action potential back-propagation from the soma. Thus, nicotine suppresses postsynaptic calcium changes, thereby altering the conditions necessary for synaptic potentiation. Burst-like stimulation of the pyramidal cell in the presence of nicotine could restore postsynaptic calcium to concentrations comparable to those seen in the absence of nicotine, as well as the STDP, indicating that strong postsynaptic stimulation could overcome the nicotinic modulation [55, 99].

The activation of distributed nAChRs provides the PFC neuronal network with a wide range of computational possibilities, but the functional consequences of this modulation are hard to predict from these data alone. Nicotine alters the rules for synaptic plasticity resulting from timed presynaptic and postsynaptic activity and increases LTP threshold by reducing dendritic calcium signals. As such, the function of the medial PFC network will most likely change in the presence of nicotine. Most likely, distal apical dendrite of layer 5 pyramidal neurons in superficial layers will be

more quantitatively affected by the nicotinic mechanisms we found to block STDP than the synapses that are located closer to the cell body. By reducing dendritic action potential propagation in apical dendrites, nicotine hampers communication between cell body and distal synapses in layer 5 pyramidal neurons. This potentially could strongly affect information processing in the neuronal network of the medial PFC as a whole, and will alter the output of the PFC. At the same time, increased activity in pyramidal neurons restores the conditions for STDP to occur. The presence of nicotine and increased threshold for STDP could reduce cognitive performance in healthy naive rodents [100]. Alternatively, since PFC neuronal activity could be increased during PFC-based cognitive behavior, nicotine may provide conditions under which signal-to-noise ratio in PFC information processing is enhanced, thereby improving cognitive performance [41, 100]. It is possible that enhancing signal-to-noise for phasic activity within the PFC, rather than simply increasing excitability, could be an effective mechanism for cognition-enhancing drugs.

Through the mechanisms discussed above, nicotinic AChR activation directly affects activity in cortical circuits involved in cognition. However, these same cellular and synaptic mechanisms also affect long-term synaptic plasticity, the effects of which outlast nAChR activation [51, 55, 101]. Thereby, nicotine may exert lasting effects on cognition. Thus far, a very limited number of studies have addressed this hypothesis, and mechanisms underlying long-term effects of nAChR activation have not been addressed. The majority of adult smokers started the habit during adolescence [102, 103], and evidence is accumulating that the adolescent brain is vulnerable for precipitating lasting changes upon nicotine exposure. In the following

sections, we will discuss the evidence for lasting effects of nAChR activation on cortical networks and cognitive function upon adolescent nicotine exposure.

Cortical development and nAChRs

Acetylcholine and nAChRs play critical roles in virtually all phases of brain maturation, during embryogenesis as well as postnatal development [reviewed in 104]. During postnatal development of sensory cortices there is a dramatic, transient increase in the expression of AChE [105]. Concurrently, nAChR γ 7 subunit gene expression also transiently increases in sensory cortices. Binding of [^{125} I] γ BgTx - to assess nAChR γ 7 levels - starts at birth in rat sensory cortex, peaks at postnatal day 10, and then declines to adult concentrations by postnatal day 20 [106]. Expression of γ 7 subunit mRNA follows a similar time course [107]. Moderate to high levels of messenger RNA are maintained into the first postnatal week, followed by a decline into adulthood [107]. The increase in cortical γ 7 mRNA precedes by the arrival of AChE-labeled thalamocortical afferents and preventing these afferents from reaching the cortex strongly reduces the γ 7 subunit mRNA and [^{125} I] γ BgTx binding in layers IV and VI [108], suggesting that the expression of γ 7 subunit-containing nAChRs is regulated by thalamic inputs. These γ 7 subunit-containing nAChRs could be located post-synaptically as well as pre-synaptically, on thalamic afferents. It has been hypothesized that pre-synaptic γ 7 nAChRs in primary auditory cortex are involved in the maturation of glutamate synapses by facilitating the conversion of 'silent synapses', containing only NMDA receptors into mature AMPA and NMDA receptor containing synapses [64, 109]. The authors suggest that through this mechanism of nAChR-induced maturation of glutamate synapses, the expression of γ 7 subunits in the auditory cortex could define a critical period of sensory cortex development in

which synaptic refinement of cortical circuitry and tuning to sensory inputs takes place.

In developing hippocampus, nAChRs containing $\gamma 7$ subunits can also activate 'silent' synapses that show a low probability of being active and turn them into high probability synapses [110]. Schaffer collateral to CA1 synapses that have a high probability of being active during development can be down-regulated by $\gamma 7$ and $\delta 2$ subunits-containing nAChRs [111]. In the rat PFC, there are strong developmental changes in nicotinic signalling of pyramidal neurons in layer VI. The nicotinic currents recorded from these neurons mediated by $\alpha 5$ -containing nAChRs peak at postnatal week 3 [71]. In human brain, the levels of nAChRs also show region-specific changes in development. Towards birth nicotine receptor binding is at its highest density, then nAChR levels start declining with the rate dependent on brain area: there is an apparent rapid decline in the hippocampus, whereas in cortical areas nicotine binding falls more gradually [112]. These findings suggest that nicotinic receptors play a role during development of excitatory glutamatergic connections and can contribute to shaping cortical neuronal circuitry.

In humans, as well as in rodents and other mammals, brain development is far from complete at birth, and many neuronal systems mature in response to the continuous interaction with the changing environment. This is especially true for the areas of the brain involved in higher cognitive functions, such as PFC, that shows dynamic changes in grey and white matter proceeding late into adolescence [113, 114]. This delayed maturation of PFC involves on the cellular level active rewiring of the intrinsic circuitry, particularly pyramidal –pyramidal connections within PFC and the local inhibitory circuitry [115, 116]. Also PFC depending cognitive performance such

as working memory and processing speed, voluntary response suppression, level off only by late adolescence [117-119].

Cholinergic signalling through nAChRs plays an important role in brain maturation. Similar to development of the PFC, the cholinergic system innervating it follows a more prolonged developmental time period [120]. When cholinergic innervation is disrupted during early postnatal development, delayed cortical neuronal development and permanent changes in cortical cytoarchitecture and cognitive behaviors are observed [121]. Thus, nAChRs appear to play multiple functional roles in brain development and their expression profiles coincide with important phases in cortical maturation. Since PFC maturation occurs later than other cortical areas and continues into late adolescence, the lasting effects of nicotine exposure on prefrontal cortex function may be especially strong during the adolescent period.

Long-term consequences of nicotine exposure during adolescence

An ever-growing amount of evidence shows that nicotine exposure during adolescence not only has direct effects on prefrontal cortical function but can also lead to adaptations in this brain area that last into adulthood. Adolescent smoking strongly correlates with cognitive and behavioural impairments during later life [122-124]. Functional MRI studies show that during working memory and attention tasks adolescent smokers have reduced PFC activation, less efficiency and altered functional coordination than in abstinent adolescents [125, 126]. Importantly, the history of smoking duration in years is correlated with the extent of diminished PFC activity, suggesting that nicotine exerts long-lasting effects on PFC function [126]. Though studies in humans reveal strong correlations between adolescent smoking and cognitive impairments during later life, genetic variability and diverse social

environment make it almost impossible to disentangle the causal relationships.

Animal models with a highly uniform genetic and environmental background between individuals offer the opportunity to directly address lasting prefrontal adaptations in response to nicotine exposure.

In rodents, nicotine exposure during adolescence induces stronger changes in gene expression in the PFC than during other periods of development and adulthood [127-129]. In PFC after chronic nicotine treatment, the maximal regulation of genes involved in vesicle release, signal transduction, cytoskeleton dynamics and transcription was observed at postnatal day 35, suggesting the role of nicotine in initiating long-term structural and functional adaptations in adolescent PFC [129]. The activity of specific early response genes (*arc*) was found to be elevated in adolescent PFC after acute nicotine exposure [127]. In addition, *c-fos* expression in the PFC in response to nicotine exposure is maximal during adolescence [102]. The expression of key molecules involved in plasticity is also altered in the PFC by adolescent nicotine exposure. Acute nicotine induces increases in the expression of the dendritically targeted dendrin mRNA in PFC of adolescent but not adult animals. Dendrin is an important component of cytoskeletal modifications at the synapse and therefore can lead to unique plasticity changes in the adolescent PFC [128]. Lasting synaptic adaptations involve activation of intracellular signalling pathway and such enzymes as extracellular regulated protein kinase (ERK) and cAMP response element binding protein (CREB). Specifically in the PFC, increases in phosphorylation of both these enzymes were found after repeated nicotine exposure [130]. Also changes in macromolecular constituents indicative of cell loss (reduced DNA) and altered cell size (protein/DNA ratio) can be seen in cortical regions of rodents after adolescent nicotine treatment [131].

Though these results describe direct changes after nicotine exposure, altered expression of genes involved in neuroplasticity can lead to structural changes in PFC neurons that last into adulthood. And indeed, repeated nicotine exposure also changes the structure of neurons in medial PFC: it increases both dendritic length and spine density [132]. Long-term changes have been observed in dendritic morphology of specific subpopulations of pyramidal neurons and these structural changes depended on the age of drug exposure [133]. Adolescent nicotine pretreatment produced an increase in basilar dendritic length in complex but not simple cells, while after adult nicotine exposure similar effect was seen in simple cells but not in complex [133].

Thus nicotine induces significant changes in gene expression and neuronal morphology in PFC specifically during the adolescent period. The key question now is of course, does adolescent nicotine exposure result in lasting altered cognitive function? Recently, this question was addressed by Counotte et al. [134]. Rats were trained in the 5 choice serial reaction time task [44] and were injected with either nicotine or saline for 10 days during adolescence (postnatal days 34 – 43) and attention performance was tested 5 weeks after the animals received the last injection with nicotine. Animals that received nicotine during adolescence showed a doubling in premature responses and a reduction in correct responses, suggesting increased impulsive behaviour and reduced attention performance [134]. Animals that received nicotine as adults did not show changes in impulsivity nor in attention performance [134]. What the mechanisms are underlying these lasting effects of nicotine on cognitive performance is at this point unknown.

Conclusions

Nicotine's effects on cognition imply a vital role for nAChRs in cortical function. While several studies point to nAChRs in the PFC as central to nicotine's effects on attention performance, general principles on nAChR regulation of thalamic inputs may apply to the neocortex. In general, the data thus far point to nicotinic actions on two different information streams: glutamatergic input from the thalamus to the neocortex is excited by $\beta 2$ subunit-containing nAChRs that are located on axons, but most likely not on the presynaptic terminals [52]. Within the neocortex, GABAergic transmission is enhanced by nicotine. Most of these nicotinic receptors are of the $\beta 2^*$ or $\alpha 7^*$ type and are located on the cell bodies of interneurons, or on the excitatory inputs to these interneurons [55]. Recently, important new features of nicotinic modulation of cortical circuits have been uncovered. Pyramidal neurons of layer VI contain $\alpha 5$ subunit-containing nAChRs that directly excite these neurons [71]. Despite these findings, large gaps remain in our understanding of nicotinic modulation of cortical circuits involved in cognition. For instance, little is known about the type of interneurons that are present in layer VI and how they are modulated by nicotine. PFC layers II and III are also still 'terra incognita' when it comes to nicotinic mechanisms. It will be exciting to learn whether in these layers common features can be identified that apply to other cortical areas.

From animal experiments, the long-term consequences of adolescent nicotine exposure for prefrontal function in adult life are emerging. Our understanding of the mechanisms is still fragmentary, but the field of research on the role of nicotinic signalling in neuronal network development and cognition function is opening up. Depending on the developmental stage, the magnitude of effects induced by nicotine and the specific targets affected by nicotine vary. In adolescence, when brain development is still ongoing, the PFC appears to be a vulnerable target for lasting

1 nicotinic actions. In rodents, even a short exposure to nicotine during this period can
2 induce a cascade of intracellular signalling, gene expression profiles and structural
3 changes that last into adulthood and may induce permanent deficiencies in attention
4 and cognitive control. The challenge that lies ahead is to uncover the exact
5 mechanisms underlying these long-term consequences of nicotine-induced cognitive
6 impairments.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

We thank Marjolijn Mertz and Dr. Lorna Role for stimulating discussions. H.D.M. was supported by grants from The Netherlands Council for Scientific Research (NWO 917.76.360, 912.06.148).

References

- [1] Everitt BJ, Robbins TW. Central cholinergic systems and cognition. *Annu Rev Psychol* 1997;48:649-84.
- [2] Parikh V, Sarter M. Cholinergic mediation of attention: contributions of phasic and tonic increases in prefrontal cholinergic activity. *Ann N Y Acad Sci* 2008;1129:225-35.
- [3] Perry EK, Court JA, Johnson M, Piggott MA, Perry RH. Autoradiographic distribution of [3H]nicotine binding in human cortex: relative abundance in subicular complex. *J Chem Neuroanat* 1992;5:399-405.
- [4] Mansvelder HD, van Aerde KI, Couey JJ, Brussaard AB. Nicotinic modulation of neuronal networks: from receptors to cognition. *Psychopharmacology* 2006;1-14.
- [5] Picciotto MR. Nicotine as a modulator of behavior: beyond the inverted U. *Trends in Pharmacological Sciences* 2003;24:493-9.
- [6] Paterson D, Nordberg A. Neuronal nicotinic receptors in the human brain. *Prog Neurobiol* 2000;61:75-111.
- [7] Levin ED, McClernon FJ, Rezvani AH. Nicotinic effects on cognitive function: behavioral characterization, pharmacological specification, and anatomic localization. *Psychopharmacology (Berl)* 2006;184:523-39.
- [8] Muir JL, Everitt BJ, Robbins TW. AMPA-induced excitotoxic lesions of the basal forebrain: a significant role for the cortical cholinergic system in attentional function. *J Neurosci* 1994;14:2313-26.
- [9] McGaughy J, Dalley JW, Morrison CH, Everitt BJ, Robbins TW. Selective behavioral and neurochemical effects of cholinergic lesions produced by intrabasis infusions of 192 IgG-saporin on attentional performance in a five-choice serial reaction time task. *J Neurosci* 2002;22:1905-13.
- [10] Dalley JW, Theobald DE, Bouger P, Chudasama Y, Cardinal RN, Robbins TW. Cortical cholinergic function and deficits in visual attentional performance in rats following 192 IgG-saporin-induced lesions of the medial prefrontal cortex. *Cereb Cortex* 2004;14:922-32.
- [11] Himmelheber AM, Sarter M, Bruno JP. The effects of manipulations of attentional demand on cortical acetylcholine release. *Brain Res Cogn Brain Res* 2001;12:353-70.
- [12] Muir JL, Everitt BJ, Robbins TW. Reversal of visual attentional dysfunction following lesions of the cholinergic basal forebrain by physostigmine and nicotine but not by the 5-HT₃ receptor antagonist, ondansetron. *Psychopharmacology (Berl)* 1995;118:82-92.
- [13] Levin ED, Kaplan S, Boardman A. Acute nicotine interactions with nicotinic and muscarinic antagonists: working and reference memory effects in the 16-arm radial maze. *Behavioural pharmacology* 1997;8:236-42.
- [14] Yu ZY, Wang W, Fritschy JM, Witte OW, Redecker C. Changes in neocortical and hippocampal GABA_A receptor subunit distribution during brain maturation and aging. *Brain Res* 2006;1099:73-81.
- [15] McGehee DS, Role LW. Physiological diversity of nicotinic acetylcholine receptors expressed by vertebrate neurons. *AnnuRevPhysiol* 1995;57:521-46.
- [16] Felix R, Levin ED. Nicotinic antagonist administration into the ventral hippocampus and spatial working memory in rats. *Neuroscience* 1997;81:1009-17.

- [17] Hahn B, Shoaib M, Stolerman IP. Involvement of the prefrontal cortex but not the dorsal hippocampus in the attention-enhancing effects of nicotine in rats. *Psychopharmacology* 2003;168:271-9.
- [18] Lawrence NS, Ross TJ, Stein EA. Cognitive mechanisms of nicotine on visual attention. *Neuron* 2002;36:539-48.
- [19] Corbetta M, Shulman GL. Control of goal-directed and stimulus-driven attention in the brain. *Nature reviews* 2002;3:201-15.
- [20] Muir JL, Everitt BJ, Robbins TW. The cerebral cortex of the rat and visual attentional function: dissociable effects of mediofrontal, cingulate, anterior dorsolateral, and parietal cortex lesions on a five-choice serial reaction time task. *Cereb Cortex* 1996;6:470-81.
- [21] Stolerman IP, Mirza NR, Shoaib M. Nicotine psychopharmacology: addiction, cognition and neuroadaptation. *Med Res Rev* 1995;15:47-72.
- [22] Rezvani AH, Bushnell PJ, Levin ED. Effects of nicotine and mecamylamine on choice accuracy in an operant visual signal detection task in female rats. *Psychopharmacology* 2002;164:369-75.
- [23] Millar NS, Gotti C. Diversity of vertebrate nicotinic acetylcholine receptors. *Neuropharmacology* 2009;56:237-46.
- [24] Le Novère N, Corringer PJ, Changeux JP. The diversity of subunit composition in nAChRs: Evolutionary origins, physiologic and pharmacologic consequences. *Journal Of Neurobiology* 2002;53:447-56.
- [25] Chavez-Noriega LE, Crona JH, Washburn MS, Urrutia A, Elliott KJ, Johnson EC. Pharmacological Characterization of Recombinant Human Neuronal Nicotinic Acetylcholine Receptors $\alpha 2\beta 2$, $\alpha 2\beta 4$, $\alpha 3\beta 2$, $\alpha 3\beta 4$, $\alpha 4\beta 2$, $\alpha 4\beta 4$ and $\alpha 7$ Expressed in *Xenopus* Oocytes. *J Pharmacol Exp Ther* 1997;280:346-56.
- [26] Changeux JP, Edelstein SJ. Allosteric mechanisms in normal and pathological nicotinic acetylcholine receptors. *Current Opinion In Neurobiology* 2001;11:369-77.
- [27] Hogg RC, Raggenbass M, Bertrand D. Nicotinic acetylcholine receptors: from structure to brain function. *Reviews Of Physiology, Biochemistry And Pharmacology*, Vol 147 2003, 2003. p. 1-46.
- [28] Gotti C, Clementi F. Neuronal nicotinic receptors: from structure to pathology. *Progress in Neurobiology* 2004;74:363-96.
- [29] Gotti C, Zoli M, Clementi F. Brain nicotinic acetylcholine receptors: native subtypes and their relevance. *Trends in Pharmacological Sciences* 2006;27:482-91.
- [30] Grady SR, Murphy KL, Cao J, Marks MJ, McIntosh JM, Collins AC. Characterization of nicotinic agonist-induced [^{3} H] dopamine release from synaptosomes prepared from four mouse brain regions. *Journal Of Pharmacology And Experimental Therapeutics* 2002;301:651-60.
- [31] McGehee DS. Nicotinic receptors and hippocampal synaptic plasticity...it's all in the timing. *Trends in Neurosciences* 2002;25:171.
- [32] Wonnacott S, Sidhpura N, Balfour DJK. Nicotine: from molecular mechanisms to behaviour. *Current Opinion in Pharmacology* 2005;5:53.
- [33] Mineur YS, Picciotto MR. Genetics of nicotinic acetylcholine receptors: Relevance to nicotine addiction. *Biochemical Pharmacology* 2008;75:323-33.
- [34] Alkondon M, Albuquerque EX. The nicotinic acetylcholine receptor subtypes and their function in the hippocampus and cerebral cortex. *Prog Brain Res* 2004;109-20.

- [35] Abumaria N, Rygula R, Havemann-Reinecke U, Ruther E, Bodemer W, Roos C, et al. Identification of genes regulated by chronic social stress in the rat dorsal raphe nucleus. *Cell Mol Neurobiol* 2006;26:145-62.
- [36] Mansvelder HD, Keath JR, McGehee DS. Synaptic mechanisms underlie nicotine-induced excitability of brain reward areas. *Neuron* 2002;33:905-19.
- [37] Wooltorton JR, Pidoplichko VI, Broide RS, Dani JA. Differential desensitization and distribution of nicotinic acetylcholine receptor subtypes in midbrain dopamine areas. *J Neurosci* 2003;23:3176-85.
- [38] Fucile S. Ca²⁺ permeability of nicotinic acetylcholine receptors. *Cell Calcium* 2004;35:1-8.
- [39] Dingledine R, Borges K, Bowie D, Traynelis SF. The glutamate receptor ion channels. *Pharmacol Rev* 1999;51:7-61.
- [40] Mirza NR, Stolerman IP. The role of nicotinic and muscarinic acetylcholine receptors in attention. *Psychopharmacology* 2000;148:243-50.
- [41] Mirza NR, Stolerman IP. Nicotine enhances sustained attention in the rat under specific task conditions. *Psychopharmacology (Berl)* 1998;138:266-74.
- [42] Granon S, Poucet B, Thinus-Blanc C, Changeux JP, Vidal C. Nicotinic and muscarinic receptors in the rat prefrontal cortex: differential roles in working memory, response selection and effortful processing. *Psychopharmacology* 1995;119:139-44.
- [43] Blondel A, Sanger DJ, Moser PC. Characterisation of the effects of nicotine in the five-choice serial reaction time task in rats: antagonist studies. *Psychopharmacology* 2000;149:293-305.
- [44] Robbins TW. The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology* 2002;163:362-80.
- [45] Young JW, Finlayson K, Spratt C, Marston HM, Crawford N, Kelly JS, et al. Nicotine improves sustained attention in mice: evidence for involvement of the alpha7 nicotinic acetylcholine receptor. *Neuropsychopharmacology* 2004;29:891-900.
- [46] Hoyle E, Genn RF, Fernandes C, Stolerman IP. Impaired performance of alpha7 nicotinic receptor knockout mice in the five-choice serial reaction time task. *Psychopharmacology* 2006;189:211-23.
- [47] Maskos U. Emerging concepts: novel integration of in vivo approaches to localize the function of nicotinic receptors. *Journal of Neurochemistry* 2007;100:596-602.
- [48] McGehee DS, Heath MJ, Gelber S, Devay P, Role LW. Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science* 1995;269:1692-6.
- [49] Lena C, Changeux JP, Mulle C. Evidence for "preterminal" nicotinic receptors on GABAergic axons in the rat interpeduncular nucleus. *J Neurosci* 1993;13:2680-8.
- [50] Kawai H, Lazar R, Metherate R. Nicotinic control of axon excitability regulates thalamocortical transmission. *Nat Neurosci* 2007;10:1168-75.
- [51] Mansvelder HD, McGehee DS. Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 2000;27:349-57.
- [52] Lambe EK, Picciotto MR, Aghajanian GK. Nicotine Induces Glutamate Release from Thalamocortical Terminals in Prefrontal Cortex. *Neuropsychopharmacology* 2003;28:216-25.

- [53] Pidoplichko VI, DeBiasi M, Williams JT, Dani JA. Nicotine activates and desensitizes midbrain dopamine neurons. *Nature* 1997;390:401-4.
- [54] Dani JA, De Biasi M. Cellular mechanisms of nicotine addiction. *PharmacolBiochemBehav* 2001;70:439-46.
- [55] Couey JJ, Meredith RM, Spijker S, Poorthuis R, Smit AB, Brussaard AB, et al. Distributed Network Actions by Nicotine Increase the Threshold for Spike-Timing-Dependent Plasticity in Prefrontal Cortex. *Neuron* 2007;54(1):73-87.
- [56] Mansvelder HD, McGehee DS. Cellular and synaptic mechanisms of nicotine addiction. *J Neurobiol* 2002;53:606-17.
- [57] Picciotto MR, Addy NA, Mineur YS, Brunzell DH. It is not "either/or": Activation and desensitization of nicotinic acetylcholine receptors both contribute to behaviors related to nicotine addiction and mood. *Prog Neurobiol* 2008;84:329-42.
- [58] Arnold HM, Burk JA, Hodgson EM, Sarter M, Bruno JP. Differential cortical acetylcholine release in rats performing a sustained attention task versus behavioral control tasks that do not explicitly tax attention. *Neuroscience* 2002;114:451-60.
- [59] Parikh V, Kozak R, Martinez V, Sarter M. Prefrontal acetylcholine release controls cue detection on multiple timescales. *Neuron* 2007;56:141-54.
- [60] Gray R, Rajan AS, Radcliffe KA, Yakehiro M, Dani JA. Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature* 1996;383:713-6.
- [61] Dajas-Bailador F, Wonnacott S. Nicotinic acetylcholine receptors and the regulation of neuronal signalling. *Trends in pharmacological sciences* 2004;25:317-24.
- [62] Clarke PB. Nicotinic modulation of thalamocortical neurotransmission. *Prog Brain Res* 2003;145:253-60.
- [63] Gioanni Y, Rougeot C, Clarke PB, Lepouse C, Thierry AM, Vidal C. Nicotinic receptors in the rat prefrontal cortex: increase in glutamate release and facilitation of mediodorsal thalamo-cortical transmission. *EurJNeurosci* 1999;11:18-30.
- [64] Metherate R. Nicotinic acetylcholine receptors in sensory cortex. *Learn Mem* 2004;11:50-9.
- [65] Harkrider AW, Champlin CA, McFadden D. Acute effect of nicotine on non-smokers: I. OAEs and ABRs. *Hearing Research* 2001;160:73-88.
- [66] Liang K, Poytress BS, Weinberger NM, Metherate R. Nicotinic modulation of tone-evoked responses in auditory cortex reflects the strength of prior auditory learning. *Neurobiology of Learning and Memory* 2008;90:138-46.
- [67] Penschuck S, Chen-Bee CH, Prakash N, Frostig RD. In vivo modulation of a cortical functional sensory representation shortly after topical cholinergic agent application. *J Comp Neurol* 2002;452:38-50.
- [68] Gil Z, Connors BW, Amitai Y. Differential Regulation of Neocortical Synapses by Neuromodulators and Activity. *Neuron* 1997;19:679-86.
- [69] Disney AA, Aoki C, Hawken MJ. Gain Modulation by Nicotine in Macaque V1. *Neuron* 2007;56:701-13.
- [70] Vidal C, Changeux JP. Nicotinic and muscarinic modulations of excitatory synaptic transmission in the rat prefrontal cortex in vitro. *Neuroscience* 1993;56:23-32.

- [71] Kassam SM, Herman PM, Goodfellow NM, Alves NC, Lambe EK. Developmental Excitation of Corticothalamic Neurons by Nicotinic Acetylcholine Receptors. *J Neurosci* 2008;28:8756-64.
- [72] Dickinson JA, Kew JNC, Wonnacott S. Presynaptic $\alpha 7$ and $\beta 2$ -containing nicotinic acetylcholine receptors modulate excitatory amino acid release from rat prefrontal cortex nerve terminals via distinct cellular mechanisms. *Mol Pharmacol* 2008;mol.108.046623.
- [73] Lubin M, Erisir A, Aoki C. Ultrastructural Immunolocalization of the $\alpha 7$ nAChR Subunit in Guinea Pig Medial Prefrontal Cortex. *Ann NY Acad Sci* 1999;868:628-32.
- [74] Wang BW, Liao WN, Chang CT, Wang SJ. Facilitation of glutamate release by nicotine involves the activation of a Ca^{2+} /calmodulin signaling pathway in rat prefrontal cortex nerve terminals. *Synapse* 2006;59:491-501.
- [75] Gullledge AT, Park SB, Kawaguchi Y, Stuart GJ. Heterogeneity of phasic cholinergic signalling in neocortical neurons. *J Neurophysiol* 2007.
- [76] Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C. Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* 2004;5:793-807.
- [77] Gupta A, Wang Y, Markram H. Organizing Principles for a Diversity of GABAergic Interneurons and Synapses in the Neocortex. *Science* 2000;287:273-8.
- [78] Kawaguchi Y, Kubota Y. Physiological and morphological identification of somatostatin- or vasoactive intestinal polypeptide-containing cells among GABAergic cell subtypes in rat frontal cortex. *J Neurosci* 1996;16:2701-15.
- [79] Wang Y, Gupta A, Toledo-Rodriguez M, Wu CZ, Markram H. Anatomical, Physiological, Molecular and Circuit Properties of Nest Basket Cells in the Developing Somatosensory Cortex. *Cereb Cortex* 2002;12:395-410.
- [80] Kawaguchi Y. Groupings of nonpyramidal and pyramidal cells with specific physiological and morphological characteristics in rat frontal cortex. *J Neurophysiol* 1993;69:416-31.
- [81] Kawaguchi Y, Kondo S. Parvalbumin, somatostatin and cholecystokinin as chemical markers for specific GABAergic interneuron types in the rat frontal cortex. *Journal of Neurocytology* 2002;31:277-87.
- [82] Gonzalez-Burgos G, Krimer LS, Povysheva NV, Barrionuevo G, Lewis DA. Functional Properties of Fast Spiking Interneurons and Their Synaptic Connections With Pyramidal Cells in Primate Dorsolateral Prefrontal Cortex. *J Neurophysiol* 2005;93:942-53.
- [83] Kawaguchi Y, Kubota Y. Neurochemical features and synaptic connections of large physiologically-identified GABAergic cells in the rat frontal cortex. *Neuroscience* 1998;85:677-701.
- [84] Angulo MC, Staiger JF, Rossier J, Audinat E. Distinct Local Circuits Between Neocortical Pyramidal Cells and Fast-Spiking Interneurons in Young Adult Rats. *J Neurophysiol* 2003;89:943-53.
- [85] Klyachko VA, Stevens CF. Excitatory and Feed-Forward Inhibitory Hippocampal Synapses Work Synergistically as an Adaptive Filter of Natural Spike Trains. *PLoS Biol* 2006;4:e207.
- [86] Sun QQ, Huguenard JR, Prince DA. Barrel cortex microcircuits: thalamocortical feedforward inhibition in spiny stellate cells is mediated by a small number of fast-spiking interneurons. *J Neurosci* 2006;26:1219-30.

- [87] Bacci A, Huguenard JR. Enhancement of spike-timing precision by autaptic transmission in neocortical inhibitory interneurons. *Neuron* 2006;49:119-30.
- [88] Rotaru DC, Barrionuevo G, Sesack SR. Mediodorsal thalamic afferents to layer III of the rat prefrontal cortex: Synaptic relationships to subclasses of interneurons. *The Journal of Comparative Neurology* 2005;490:220-38.
- [89] Krenz I, Kalkan D, Wevers A, de Vos RA, Steur EN, Lindstrom J, et al. Parvalbumin-containing interneurons of the human cerebral cortex express nicotinic acetylcholine receptor proteins. *J Chem Neuroanat* 2001;21:239-46.
- [90] Kapfer C, Glickfeld LL, Atallah BV, Scanziani M. Supralinear increase of recurrent inhibition during sparse activity in the somatosensory cortex. *Nat Neurosci* 2007;10:743-53.
- [91] Silberberg G, Markram H. Disynaptic inhibition between neocortical pyramidal cells mediated by Martinotti cells. *Neuron* 2007;53:735-46.
- [92] Xiang Z, Huguenard JR, Prince DA. Cholinergic Switching Within Neocortical Inhibitory Networks. *Science* 1998;281:985-8.
- [93] Laroche S, Davis S, Jay TM. Plasticity at hippocampal to prefrontal cortex synapses: dual roles in working memory and consolidation. *Hippocampus* 2000;10:438-46.
- [94] Markram H, Lubke J, Frotscher M, Sakmann B. Regulation of synaptic efficacy by coincidence of postsynaptic APs and EPSPs. *Science* 1997;275:213-5.
- [95] Bi GQ, Poo MM. Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. *J Neurosci* 1998;18:10464-72.
- [96] Koester HJ, Sakmann B. Calcium dynamics in single spines during coincident pre- and postsynaptic activity depend on relative timing of back-propagating action potentials and subthreshold excitatory postsynaptic potentials. *Proc Natl Acad Sci U S A* 1998;95:9596-601.
- [97] Magee JC, Johnston D. A synaptically controlled, associative signal for Hebbian plasticity in hippocampal neurons. *Science* 1997;275:209-13.
- [98] Sjöström PJ, Nelson SB. Spike timing, calcium signals and synaptic plasticity. *Current Opinion in Neurobiology* 2002;12:305-14.
- [99] McGehee DS. Nicotine and Synaptic Plasticity in Prefrontal Cortex. *Sci STKE* 2007;2007:pe44-.
- [100] Day M, Pan JB, Buckley MJ, Cronin E, Hollingsworth PR, Hirst WD, et al. Differential effects of ciproxifan and nicotine on impulsivity and attention measures in the 5-choice serial reaction time test. *Biochemical Pharmacology* 2006;In Press, Corrected Proof.
- [101] Ji D, Lape R, Dani JA. Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron* 2001;31:131-41.
- [102] Leslie FM, Loughlin SE, Wang R, Perez L, Lotfipour S, Belluzzia JD. Adolescent development of forebrain stimulant responsiveness: insights from animal studies. *Ann N Y Acad Sci* 2004;1021:148-59.
- [103] Breslau N, Peterson EL. Smoking cessation in young adults: age at initiation of cigarette smoking and other suspected influences. *Am J Public Health* 1996;86:214-20.
- [104] Mansvelder HD, Role LW. Neuronal receptors for nicotine: functional diversity and developmental changes. In: Miller M, editor. *Brain Development Normal processes and the effects of alcohol and nicotine*. Oxford: Oxford University Press, 2006. p. 341-62.

- [105] Robertson RT, Mostamand F, Kageyama GH, Gallardo KA, Yu J. Primary auditory cortex in the rat: transient expression of acetylcholinesterase activity in developing geniculocortical projections. *Brain Res Dev Brain Res* 1991;58:81-95.
- [106] Fuchs JL. [125I]alpha-bungarotoxin binding marks primary sensory area developing rat neocortex. *Brain Res* 1989;501:223-34.
- [107] Broide RS, O'Connor LT, Smith MA, Smith JA, Leslie FM. Developmental expression of alpha 7 neuronal nicotinic receptor messenger RNA in rat sensory cortex and thalamus. *Neuroscience* 1995;67:83-94.
- [108] Broide RS, Robertson RT, Leslie FM. Regulation of alpha7 nicotinic acetylcholine receptors in the developing rat somatosensory cortex by thalamocortical afferents. *J Neurosci* 1996;16:2956-71.
- [109] Metherate R, Hsieh CY. Regulation of glutamate synapses by nicotinic acetylcholine receptors in auditory cortex. *Neurobiol Learn Mem* 2003;80:285-90.
- [110] Maggi L, Le Magueresse C, Changeux JP, Cherubini E. Nicotine activates immature "silent" connections in the developing hippocampus. *Proc Natl Acad Sci U S A* 2003;100:2059-64.
- [111] Maggi L, Sola E, Minneci F, Le Magueresse C, Changeux JP, Cherubini E. Persistent decrease in synaptic efficacy induced by nicotine at Schaffer collateral-CA1 synapses in the immature rat hippocampus. *J Physiol* 2004;559:863-74.
- [112] Court JA, Martin-Ruiz C, Graham A, Perry E. Nicotinic receptors in human brain: topography and pathology. *J Chem Neuroanat* 2000;20:281-98.
- [113] Gogtay N, Giedd JN, Lusk L, Hayashi KM, Greenstein D, Vaituzis AC, et al. Dynamic mapping of human cortical development during childhood through early adulthood. *Proc Natl Acad Sci U S A* 2004;101:8174-9.
- [114] Casey BJ, Tottenham N, Liston C, Durston S. Imaging the developing brain: what have we learned about cognitive development? *Trends Cogn Sci* 2005;9:104-10.
- [115] Cruz DA, Eggan SM, Lewis DA. Postnatal development of pre- and postsynaptic GABA markers at chandelier cell connections with pyramidal neurons in monkey prefrontal cortex. *J Comp Neurol* 2003;465:385-400.
- [116] Woo TU, Pucak ML, Kye CH, Matus CV, Lewis DA. Peripubertal refinement of the intrinsic and associational circuitry in monkey prefrontal cortex. *Neuroscience* 1997;80:1149-58.
- [117] Luna B, Garver KE, Urban TA, Lazar NA, Sweeney JA. Maturation of cognitive processes from late childhood to adulthood. *Child Dev* 2004;75:1357-72.
- [118] Luciana M, Nelson CA. The functional emergence of prefrontally-guided working memory systems in four- to eight-year-old children. *Neuropsychologia* 1998;36:273-93.
- [119] Lewis DA. Development of the prefrontal cortex during adolescence: insights into vulnerable neural circuits in schizophrenia. *Neuropsychopharmacology* 1997;16:385-98.
- [120] Mechawar N, Descarries L. The cholinergic innervation develops early and rapidly in the rat cerebral cortex: a quantitative immunocytochemical study. *Neuroscience* 2001;108:555-67.

- [121] Hohmann CF, Berger-Sweeney J. Cholinergic regulation of cortical development and plasticity. New twists to an old story. *Perspect Dev Neurobiol* 1998;5:401-25.
- [122] Wiers RW, Bartholow BD, van den Wildenberg E, Thush C, Engels RC, Sher KJ, et al. Automatic and controlled processes and the development of addictive behaviors in adolescents: a review and a model. *Pharmacol Biochem Behav* 2007;86:263-83.
- [123] Ernst M, Pine DS, Hardin M. Triadic model of the neurobiology of motivated behavior in adolescence. *Psychol Med* 2006;36:299-312.
- [124] Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *Am J Psychiatry* 2003;160:1041-52.
- [125] Jacobsen LK, Mencl WE, Constable RT, Westerveld M, Pugh KR. Impact of smoking abstinence on working memory neurocircuitry in adolescent daily tobacco smokers. *Psychopharmacology (Berl)* 2007;193:557-66.
- [126] Musso F, Bettermann F, Vucurevic G, Stoeter P, Konrad A, Winterer G. Smoking impacts on prefrontal attentional network function in young adult brains. *Psychopharmacology (Berl)* 2007;191:159-69.
- [127] Schochet TL, Kelley AE, Landry CF. Differential expression of arc mRNA and other plasticity-related genes induced by nicotine in adolescent rat forebrain. *Neuroscience* 2005;135:285-97.
- [128] Schochet TL, Bremer QZ, Brownfield MS, Kelley AE, Landry CF. The dendritically targeted protein Dendrin is induced by acute nicotine in cortical regions of adolescent rat brain. *Eur J Neurosci* 2008;28:1967-79.
- [129] Poleskaya OO, Fryxell KJ, Merchant AD, Locklear LL, Ker KF, McDonald CG, et al. Nicotine causes age-dependent changes in gene expression in the adolescent female rat brain. *Neurotoxicol Teratol* 2007;29:126-40.
- [130] Brunzell DH, Russell DS, Picciotto MR. In vivo nicotine treatment regulates mesocorticolimbic CREB and ERK signaling in C57Bl/6J mice. *J Neurochem* 2003;84:1431-41.
- [131] Trauth JA, Seidler FJ, Slotkin TA. An animal model of adolescent nicotine exposure: effects on gene expression and macromolecular constituents in rat brain regions. *Brain Res* 2000;867:29-39.
- [132] Brown RW, Kolb B. Nicotine sensitization increases dendritic length and spine density in the nucleus accumbens and cingulate cortex. *Brain Res* 2001;899:94-100.
- [133] Bergstrom HC, McDonald CG, French HT, Smith RF. Continuous nicotine administration produces selective, age-dependent structural alteration of pyramidal neurons from prelimbic cortex. *Synapse* 2008;62:31-9.
- [134] Counotte DS, Spijker S, Van de Burgwal LH, Hogenboom F, Schoffelman AN, De Vries TJ, et al. Long-lasting cognitive deficits resulting from adolescent nicotine exposure in rats. *Neuropsychopharmacology* 2009;34:299-306.

Figure Legend

Figure 1.

Schematic model of the prefrontal cortex microcircuit and the identified nAChR subunits found to be expressed by the different cell types. The assignment of nAChR subunits to cell types and projections is mainly based on references [52, 55, 71], but are in line with findings in other cortical areas [64]. P: pyramidal neuron; FS: Fast Spiking interneuron; LTS: Low threshold spiking interneuron; RSNP: Regular spiking non-pyramidal interneuron.