Synthesis of partially-bridged phosphonate and thiophosphonate resorcinarenes

Barbara Cantadori, Paolo Betti, Francesca Boccini, Chiara Massera, Enrico Dalcanale

► To cite this version:

Barbara Cantadori, Paolo Betti, Francesca Boccini, Chiara Massera, Enrico Dalcanale. Synthesis of partially-bridged phosphonate and thiophosphonate resorcinarenes. *Supramolecular Chemistry*, 2008, 20 (01-02), pp.29-34. 10.1080/10610270701798803 . hal-00513526

HAL Id: hal-00513526

<https://hal.science/hal-00513526>

Submitted on 1 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis of partially-bridged phosphonate and thiophosphonate resorcinarenes

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2007-0098.R1
Manuscript Type:	Special Issue Paper
Date Submitted by the Author:	21-Aug-2007
Complete List of Authors:	Cantadori, Barbara; University of Parma, Organic and Industrial Chemistry Betti, Paolo; University of Parma, Organic and Industrial Chemistry Bocchini, Francesca; University of Parma, Organic and Industrial Chemistry Massera, Chiara; University of Parma, General, Inorganic, Analytical, and Physical Chemistry Dalcanale, Enrico; University of Parma, Organic and Industrial Chemistry
Keywords:	Thiophosphonated resorcinarenes, phosphonated resorcinarenes, Partially-bridged cavitands
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
figure 1.cdx scheme 1.cdx scheme 2.cdx

Synthesis of partially-bridged phosphonate and thiophosphonate resorcinarenes

BARBARA CANTADORI^a, PAOLO BETTI^a, FRANCESCA BOCCINI^a, CHIARA MASSERA^b, and ENRICO DALCANALE^{*a}

^a Dipartimento di Chimica Organica ed Industriale, Università di Parma and INSTM

UdR Parma, viale G. Usberti 17/A, 43100 Parma, Italy

^b Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica e Chimica Fisica,

Università di Parma, viale G. Usberti 17/A, 43100 Parma, Italy

* Corresponding author. E-mail: enrico.dalcanale@unipr.it

The synthesis of partially-bridged thiophosphonate and phosphonate resorcinarenes is reported in this paper. Resorcinarenes functionalized at the upper rim with 2 or 3 P=X (X=S,O) bridges oriented towards the molecular cavity are obtained from the treatment of the corresponding [iiii] tetrasubstituted cavitand with **1,2-Dihydroxybenzene** and base in DMF. This synthetic route selectively promoted the formation of one of the two disubstituted resorcinarenes, the AC-isomer. The crystal structure of the AC-dithiophosphonated resorcinarene **4** has been determined.

Keywords: Thiophosphonated resorcinarenes; phosphonated resorcinarenes; Partially-bridged cavitands

Dedicated to Prof. D. N. Reinhoudt on occasion of his 65th birthday.

INTRODUCTION

Cavitands are particularly attractive as host molecules because of the wide choice of suitable bridging groups. In fact, the appropriate structural preorganization of the host in combination with the appropriate set of complementary binding sites leads to the formation of highly stable complexes with neutral and charged guests [1]. The most common bridging groups employed for cavitands are alkylenedioxy [2] and derivatives [3], dialkylsilicon [4,5], heterophenylene [6,7], and phosphoryl [8,9]. In particular, phosphoryl derivatives are able to form strong complexes with cationic guests, such as metal ions [10,11], alkali metal, and ammonium cations [12] and are efficient hydrogen bond acceptors in the complexation of alcohols [13–15]. Because of their high binding capabilities, P=O groups have been included in preorganized structures to increase the stability of the complexes thus formed. Dutasta and co-workers reported the synthesis and the study of the binding properties of tetrabridged phosphonated cavitands having all inward configuration of the P=O groups [16]. To ensure strong binding, the P=O groups must be oriented toward the cavity (indicated as [iiii]).

Partially-bridged thiophosphonate and phosphonate resorcinarenes (Figure 1) are attractive molecules as intermediates in the preparation of **preorganized** and multidentate cavitand ligands.

[Insert figure 1]

In particular, we are interested in AC-disubstituted resorcinarenes **4** and **7** for their potential use as ligand precursors in the self-assembly of coordination cages capable of molecular recognition. A synthetic study of these partially-bridged building blocks is presented herein as well as the X-ray crystallographic investigation of the AC-dithiophosphonated resorcinarene **4**.

RESULTS AND DISCUSSION

Synthesis of resorcinarenes bearing three and two P=X (X=S,O) bridges at the upper rim

The starting cavitand **1** (Scheme 1) was prepared in good yields according to the protocols reported in the literature [10] starting from the ethyl-footed resorcinarene. Briefly, 4.05 equivalents of dichlorophenylphosphine were added to the resorcinarene solution under basic conditions to give a tetraphosphonito cavitand. The intermediate was treated with sulphur to obtain the desired tetraphosphonate cavitand **1**. This synthetic procedure **allows** the formation of the [iiii] stereoisomer with the four P=S oriented towards the molecular cavity as major product. In order to synthesise partially-

bridged thiophosphonate resorcinarenes, we performed the same protocol **reducing the amount of** bridging agent. When 2.8 equivalents of dichlorophenylphosphine were used, cavitand **1** was obtained as the major product in low yield (14%), while the tri-bridged compound **3** was also formed (6%) and only traces of the AC- and AB- di-bridged resorcinarenes were found (**4** and **5**). Other attempts to obtain one of the partially bridged resorcinarene in higher yields by varying the stoichiometric amount of phosphine derivative were unsuccessful. Therefore, this approach proved to be inadequate for the preparation of suitable amounts of partially bridged resorcinarenes.

A novel approach for the selective excision of one and two quinoxaline units from tetraquinoxaline cavitand has been developed by Gutierrez-Tunstad and co-workers [17]. After the treatment of tetraquinoxaline cavitand with **1,2-Dihydroxybenzene** under basic conditions they reported a overall yield of ca 56% for triquinoxaline resorcinarene and ca 48% for AC-diquinoxaline resorcinarene. The AB-diquinoxaline isomer was obtained in a maximum yield of 11%. Following this approach, we investigated the behaviour of [iiii] tetrathiophosphonate cavitand in the presence of **1,2-Dihydroxybenzene** under basic conditions.

In order to prepare the [iii] trisubstituted resorcinarene **3**, the [iiii] tetrathiophosphonate cavitand **1** was reacted with 1 equivalent of **1,2-Dihydroxybenzene** in the presence of K_2CO_3 as base (Scheme 1).

[Insert Scheme 1]

Under these conditions, we obtained [iii] trithiophosphonated resorcinarene **3** as major product in a yield of 52%. When 2 equivalents of **1,2-Dihydroxybenzene** were added to

the reaction, a mixture of the [iii] trithiophosphonated resorcinarene **3** (21%), of the [ii] AC-dithiophosphonated isomer **4** (28%) and only traces of the [ii] AB-dithiophosphonated isomer **5** were isolated. Higher yields of the desired product **4** were not achieved when the amounts of **1,2-Dihydroxybenzene** was increased. Moreover, during the reaction, the stereochemistry of the residual P=S groups did not change. Therefore, this synthetic route allows a good control of the products distribution between the di-bridged resorcinarenes without altering the P=S stereochemistry.

The stability of phosphonate derivatives was also investigated under basic conditions in the presence of **1,2-Dihydroxybenzene**. Tetraphosphonate cavitand **2** is readily available following Dutasta's synthetic procedure [16]. In analogy to [iiii] tetrathio phosphonate cavitand **1**, the treatment of compound **2** with **1,2-Dihydroxybenzene** and K₂CO₃ in DMF led to the abstraction of a variable number of P=O bridges depending on the number of equivalents of nucleophile used (Scheme 2).

[Insert scheme 2]

When 1 equivalent of **1,2-Dihydroxybenzene** was added to a mixture of cavitand **2** and K₂CO₃ in DMF the [iii] triphosphonate resorcinarene **6** was obtained in good yields (65%). Under these conditions, a small amount of the [ii] AC-diphosphonate isomer **7** was isolated, but neither the corresponding [ii] AB isomer nor the starting compound were observed. Similarly, when 2 equivalents of **1,2-Dihydroxybenzene** were used we obtained the [ii] AC-diphosphonate isomer **7** as the major product (62% yields) without the presence of the [ii] AB-diphosphonate isomer **8**.

In comparison to [iiii] tetrathiophosphonate cavitand **1**, the selective excision of one or two phosphonate groups from [iiii] tetraphosphonate cavitand **2** is more sensitive to the amount of nucleophile used. In particular, the reaction of 2 equivalents of **1,2-Dihydroxybenzene** with [iiii] tetrathiophosphonate cavitand **1** leads to the formation of similar amount of [ii] and [iii] resorcinarenes, whereas that with [iiii] tetraphosphonate cavitand **2** produces exclusively the AC isomer. Also in this case, the nucleophilic attack of **1,2-Dihydroxybenzene** to the P=O bridges does not change the stereochemistry of the remaining bridges. This last aspect is crucial to retain the recognition proprieties in the partially bridged resorcinarenes bearing P=O groups.

X-ray crystallographic investigation

The stereochemistry of the [ii] AC-disubstituted isomer has been confirmed also by the X-ray crystal structure of resorcinarene **4** (Figures 2 and 3) [18]. Crystals of **4** were obtained by slow evaporation of a solution of **4** in acetone. Compound **4** crystallizes in the space group P21/n as two **independent** resorcinarenes A and B. The structure of each resorcinarene shows indeed the presence of two P=S groups pointing inside the cavity and of four C₂H₅ chains at the lower rim. The cavity is stabilized by a network of intramolecular O-H...O hydrogen bonds among the OH groups on the phenyl rings at the upper rim, with the O...O distances spanning from 2.691(3) to 2.784(3) Å and the O-H...O angles going from 167.2(2)° to 175.2(2)°. Each resorcinarene crystallizes with two acetone molecules, one dispersed in the lattice and one with a CH₃ group inside the cavity, giving rise to two weak CH...π interactions (resorcinarene A: C...centroid distances of 3.061(2) and 3.294(3) Å, with C-H...centroid angles of 147.6(4)° and

147.6(3)° respectively; resorcinarene B: distances of 3.001(2) and 3.105(3) Å, with angles of 157.5(2)° and 143.1(4)° respectively, Figure 2).

[Insert figure 2]

In the lattice the resorcinarenes A and B face each other to give a supramolecular dimer stabilized by a network of hydrogen bonds involving the acetone molecules inside the cavity and the OH groups at the upper rim [O3D...O3S 2.675(3) Å, O3D-H3D...O3S 168.8(2)°; O3E...O1S 2.695(3) Å, O3E-H3E...O1S 166.1(2)°] (Figure 3).

[Insert figure 3]

CONCLUSIONS

An effective protocol for the preparation of partially-bridged thiophosphonate and phosphonate resorcinarenes is reported. Different amounts of **1,2-Dihydroxybenzene** were used to perform the abstraction of one or two P=X (X=S, O) bridges from the corresponding tetrabridged cavitands in good yields. This synthetic route allowed a good control of the product distribution producing mainly the [ii] AC-dibridged isomer and the [iii] trisubstituted resorcinarene. Moreover, during the excision of the P=X groups, the configuration of the remaining P=X (X=S, O) bridges on the molecules is retained. These compounds are useful precursors for the preparation of differently functionalized cavitands having specific molecular recognition properties [19].

EXPERIMENTAL

Materials and methods

All commercial reagents were ACS grade. All solvents were dried over 3 Å and 4 Å molecular sieves. ^1H NMR spectra were recorded on a **Bruker** Avance 300 MHz spectrometer at 300 K and all chemical shifts were reported in parts per million (ppm) in relation to the proton resonances resulting from incomplete deuteration of the NMR solvents. ^{31}P NMR spectra were recorded on a **Bruker** 162 MHz spectrometer and all chemical shifts were reported in ppm relative to external 85% H_3PO_4 at 0.00 ppm. Electrospray ionisation mass spectrometry (ESI-MS) experiments were performed on a Waters ZMD spectrometer equipped with an electrospray interface. Column chromatography was performed using silica gel 60 (Merck 70-230 mesh) as stationary phase. Cavitands **1** [10] and **2** [16] were prepared according to the protocols reported in literature.

Synthesis of the [iii] trithiophosphonated resorcinarene **3** and the [ii] AC-dithiophosphonated resorcinarene **4**

A mixture of cavitand **1** (0.500 g, 0.43 mmol), K_2CO_3 (0.599 g, 4.34 mmol), and freshly **recrystallized 1,2-Dihydroxybenzene** (0.095 g, 0.87 mmol) in 10 mL of dry DMF was stirred in a sealed tube at 80 °C for 5 h. The reaction mixture was then cooled and poured into ice-cold brine. The solid formed was then filtered, washed with water, and dried. The crude product was purified by column chromatography (SiO_2 ; CH_2Cl_2 /ethyl

acetate = 95/5) to give **3** in a 21% yield (0.095 g, 0.09 mmol) followed by **4** in a 28% yield (0.108 g, 0.12 mmol).

3

¹H NMR (CDCl₃, 300 MHz): δ 8.17 (m, 6H, P-ArH), 7.55 (m, 9H, P-ArH), 7.32 (s, 2H, ArH_{down}), 7.20 (s, 2H, ArH_{down}), 6.68 (d, 2H, ArH_{up}, ⁴J_{HP} = 2.2 Hz), 6.49 (s, 2H, ArH_{up}), 4.60 (m, 3H, ArCH), 4.29 (t, 1H, ArCH, J = 7.9 Hz), 2.40-2.23 (m, 8H, ArCH-CH₂), 1.09-0.91 (m, 12H, CH₂CH₃); ESI-MS: m/z 1037 [M+Na]⁺; methanol.

4

¹H NMR (Acetone-d₆, 300 MHz): δ 8.22 (dd, 4H, P-ArH, ³J_{PH} = 14.8 Hz, J_{HH} = 6.9 Hz), 7.78 (s, 4H, ArH_{down}), 7.75 (m, 2H, P-ArH), 7.67 (m, 4H, P-ArH), 6.44 (s, 4H, ArH_{up}, ⁴J_{HP} = 2.2 Hz), 4.63 (t, 2H, ArCH, J = 8.1 Hz), 4.34 (t, 2H, ArCH, J = 8.0 Hz), 2.51 (m, 4H, ArCH-CH₂), 2.40 (m, 4H, ArCH-CH₂), 1.05 (t, 6H, CH₂CH₃, J = 7.2 Hz), 0.92 (t, 6H, CH₂CH₃, J = 7.2 Hz); ³¹P NMR (CDCl₃, 162 MHz): δ 81.2 (s, 2P); ESI-MS: m/z 877 [M+H]⁺, 899 [M+Na]⁺; acetone.

Crystal structure determination of compound 4

The molecular structure of compound C₄₈H₄₆O₈S₂P₂·2C₃H₆O **4** was determined by single-crystal X-ray diffraction methods. Crystallographic and experimental details are summarized in Table 1.

[Insert table 1]

Intensity data were collected using a Mo K α radiation on a Bruker AXS Smart 1000 single-crystal diffractometer equipped with a CCD area detector at 293(2) K. The structure was solved by direct methods with the SIR97 program [20] and refined on F_o^2 by full-matrix least-squares procedures with the SHELXL-97 program [21]. The data reduction for **4** was performed using the SAINT [22] and SADABS [23] programs. All the non-hydrogen atoms were refined with anisotropic atomic displacements. The hydrogen atoms were included in the refinement at idealized geometries (C–H 0.95 Å) and refined “riding” on the corresponding parent atoms. The weighting scheme used in the last cycle of refinement was $w = 1 / [\sigma^2 F_o^2 + (0.0695P)^2 + 4.56219P]$. Geometric calculations and molecular graphics were performed with the PARST97 program [24] and the DS ViewerPro 5.0 package. Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publications no. CCDC-655076. These data can be obtained free of charge at www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; Fax: +44-1223/336-033; E-mail: deposit@ccdc.cam.ac.uk].

Synthesis of the [iii] triphosphonated resorcinarene **6**

66 mg (0.6 mmol) of freshly **recrystallized 1,2-Dihydroxybenzene** were dissolved in ca. 50 mL of DMF and 828 mg of K₂CO₃ (ca. 10 eq) were added. After the addition of 650 mg (0.6 mmol) of cavitand **2** the reaction mixture was heated to 80 °C and stirred for 5 hours. The reaction mixture was then cooled to room temperature and DMF was

completely evaporated in *vacuum*. The solid was dissolved in CH₂Cl₂ and three extractions with acidic water were performed. The final product was purified by silica gel chromatography with CH₂Cl₂–EtOH as eluent (90/10 v/v). The yield of the reaction was 65 % (375 mg).

¹H NMR (CDCl₃, 300 MHz): δ 9.56 (bs, 2H, Ar-OH), 8.05 (m, 6H, P-ArH), 7.59 (m, 9H, P-ArH), 7.36 (s, 2H, ArH_{down}), 7.17 (s, 2H, ArH_{down}), 6.93 (s, 2H, ArH_{up}), 6.80 (s, 2H, ArH_{up}), 4.60 (m, 3H, ArCH), 4.37 (t, 1H, ArCH, J= 7.9 Hz), 2.38 (m, 6H, ArCH-CH₂), 2.26 (m, 2H, ArCH-CH₂), 1.08 (t, 6H, CH₂CH₃, J=7.2 Hz), 1.00 (t, 3H, CH₂CH₃, J=7.2 Hz), 0.94 (t, 3H, CH₂CH₃, J=7.2 Hz); ESI-MS: m/z 989 [M+Na]⁺; methanol

Synthesis of the [ii] AC-diphosphonated resorcinarene **7**

K₂CO₃ (0.511 g, 3.69 mmol) and freshly **recrystallized 1,2-Dihydroxybenzene** (0.081 g, 0.74 mmol) were added to a solution of cavitand **2** (0.402 g, 0.37 mmol) in 12 mL of dry DMF. The reaction mixture was stirred at 80° C for 5 h, cooled at room temperature and then poured into ice-cold brine. The solid formed was filtered, washed with water, and dried. The desired product **7** (183 mg) was collected by filtration in a 62% yield after the treatment of the crude with diethyl ether.

¹H NMR (CDCl₃, 300 MHz): δ 7.94 (m, 4H, P-ArH), 7.60 (m, 2H, P-ArH), 7.48 (m, 4H, P-ArH), 7.01 (s, 4H, ArH_{down}), 6.37 (s, 4H, ArH_{up}), 4.40 (t, 2H, ArCH, J= 7.0 Hz), 4.28 (t, 2H, ArCH, J= 7.8 Hz), 2.26 (m, 4H, ArCH-CH₂), 2.17(m, 4H, ArCH-CH₂),

0.96 (t, 6H, CH₂CH₃, J= 6.9 Hz), 0.87 (t, 6H, CH₂CH₃, J= 6.9 Hz); ESI-MS: m/z 868 [M+Na]⁺, 884 [M+K]⁺; methanol.

Acknowledgements

This work was supported by MUR through PRIN 2005. The instrumentation facilities at the Centro Interfacoltà Misure G. Casnati of the University of Parma were used.

REFERENCES

- [1] Cram, D. J. *Angew. Chem. Int. Ed. Engl.* **1986**, *98*, 1041.
- [2] Cram, D. J.; Karbach, S.; Kim, H.-E.; Knobler, C. B.; Maverich, E. F.; Ericson, J. L.; Helgeson, R. C. *J. Am. Chem. Soc.* **1988**, *110*, 2229.
- [3] Xi, H.; Gibb, C. L. D.; Stevens, E. D.; Gibb, B. C. *Chem. Commun.* **1998**, 1743.
- [4] Cram, D. J.; Stewart, K. D.; Goldberg, I.; Trueblood, K. N. *J. Am. Chem. Soc.* **1985**, *107*, 2574.
- [5] Tucker, J. A.; Knobler, C. B.; Trueblood, K. N.; Cram, D. J. *J. Am. Chem. Soc.* **1989**, *111*, 3688.
- [6] Moran, J. R.; Ericson, J. L.; Dalcanale, E.; Bryant, J. A.; Knobler, C. B.; Cram, D. J. *J. Am. Chem. Soc.* **1991**, *113*, 5707.
- [7] Cram, D. J.; Choi, H. J.; Bryant, J. A.; Knobler, C. B. *J. Am. Chem. Soc.* **1992**, *114*, 7748.
- [8] Lippmann, T.; Wilde, H.; Dalcanale, E.; Mavilla, L.; Mann, G.; Heyer, U.; Spera, S. *J. Org. Chem.* **1995**, *60*, 235.
- [9] Dalcanale, E.; Jacopozzi, P.; Ugozzoli, F.; Mann, G. *Supramol. Chem.* **1998**, *9*, 305.

- [10] Bibal, B.; Tinant, B.; Declercq, J.-P.; Dutasta, J.-P. *Chem. Commun.* **2002**, 432.
- [11] Bibal, B.; Declercq, J.-P.; Dutasta, J.-P.; Tinant, B.; Valade, A.-G. *Tetrahedron* **2003**, *59*, 5849.
- [12] Dutasta, J.-P. *Top. Curr. Chem.*, **2004**, *232*, 55.
- [13] Ventola, E.; Vainiotalo, P.; Suman, M.; Dalcanale E. *J. Am. Soc. Mass Spectrom.* **2006**, *17*, 213.
- [14] Pinalli, R.; Suman, M.; Dalcanale, E. *Eur. J. Org. Chem.*, **2004**, *3*, 451.
- [15] Suman, M.; Freddi, M.; Massera, C.; Ugozzoli, F.; Dalcanale E. *J. Am. Chem. Soc.* **2003**, *125*, 12068.
- [16] Delangle, P.; Mulatier, J.-C.; Tinant, B.; Declercq, J.-P.; Dutasta, J.-P. *Eur. J. Org. Chem.* **2001**, *16*, 3695.
- [17] Castro, P.; Zhao, G.; Masangkay, G. A.; Hernandez, C.; Gutierrez-Tunstad, L. M. *Organic Letters* **2004**, *6*, 333.
- [18] The crystal structure of a relevant AC-diphosphonater resorcinarene with undecyl feet has been reported by Dutasta's group: B. Dubessy, PhD thesis **2006**, École Normale Supérieure de Lyon.
- [19] Pirondini, L.; Dalcanale E. *Chem. Soc. Rev.* **2007**, *36*, 695.
- [20] Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, R. *J. Appl. Crystallogr.* **1999**, *32*, 115.
- [21] Sheldrick, G. M.; *SHELXL97. Program for Crystal Structure Refinement*, University of Göttingen: Göttingen, Germany, 1997.
- [22] *SAINT, Software Users Guide*, 6.0; Bruker Analytical X-ray Systems: 1999.

[23] Sheldrick, G. M.; *SADABS Area-Detector Absorption Correction*, 2.03; University of Göttingen: Göttingen, Germany, 1999.

[24] Nardelli, M. *J. Appl. Crystallogr.* **1996**, 29, 296.

Captions to Figures, Schemes and Table

FIGURE 1 Partially-bridged thiophosphonate and phosphonate resorcinarenes: tri- and di-substituted molecules.

SCHEME 1 Preparation of the [iii] tri- and the [ii] AC-dithiophosphorylated resorcinarenes. (a) **1,2-Dihydroxybenzene**, K_2CO_3 , dry DMF, 80° C, 5 h.

SCHEME 2 Abstraction of one or two phosphoryl bridges from the tetraphosphonate cavitand 2. (a) 1 eq. **1,2-Dihydroxybenzene**, K_2CO_3 , dry DMF, 80° C, 5 h. (b) 2 eq. **1,2-Dihydroxybenzene**, K_2CO_3 , dry DMF, 80° C, 5 h.

FIGURE 2 Molecular structure of resorcinarene A showing the intramolecular hydrogen bonds as yellow dotted lines (above) and the CH- π interactions as green dotted lines (below; only the hydrogens involved in the interactions are shown). Color code: carbon, grey; hydrogen, white; oxygen, red; phosphorus, orange; sulphur, yellow; centroid, green.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE 3 The dimer formed by resorcinarene A and B through a network of hydrogen bonds involving the acetone guests. Color code: carbon, grey; hydrogen, white; oxygen, red; phosphorus, orange; sulphur, yellow.

TABLE 1 Crystallographic data and refinement details for resorcinarene 4

For Peer Review Only

FIGURE 2 Molecular structure of resorcinarene A showing the intramolecular hydrogen bonds as yellow dotted lines (above) and the CH-n interactions as green dotted lines (below; only the hydrogens involved in the interactions are shown). Color code: carbon, grey; hydrogen, white; oxygen, red; phosphorus, orange; sulphur, yellow; centroid, green.
100x124mm (150 x 150 DPI)

FIGURE 3 The dimer formed by resorcinarene A and B through a network of hydrogen bonds involving the acetone guests. Color code: carbon, grey; hydrogen, white; oxygen, red; phosphorus, orange; sulphur, yellow.
107x91mm (150 x 150 DPI)

Resorcinarene 4	
Formula	C ₅₄ H ₅₈ O ₁₀ P ₂ S ₂
Formula weight	993.06
Crystal system	Monoclinic
Space group	<i>P21/n</i>
<i>a</i> /Å	11.611(1)
<i>b</i> /Å	50.654(5)
<i>c</i> /Å	17.557(2)
β /°	97.519(1)
<i>V</i> /Å ³	10237(2)
<i>Z</i>	8
<i>D</i> /g cm ⁻³	1.289
<i>F</i> (000)	4192
μ /mm ⁻¹	0.224
$\theta_{\min, \max}$ /°	2.83, 26.84
Reflections collected	83251
Independent reflections	19682 (<i>R</i> _{int} = 0.0389)
Obs. refl. [<i>F</i> _o >4σ(<i>F</i> _o)]	14100
Data / restr. / param	19682 / 0 / 1255
R indices [<i>F</i> _o >4σ(<i>F</i> _o)] ^a	<i>R</i> 1 = 0.0558, <i>wR</i> 2 = 0.1372
R indices (all data)	<i>R</i> 1 = 0.1848, <i>wR</i> 2 = 0.1481
$\Delta\rho_{\min, \max}$ /e Å ⁻³	-0.311, 0.232
<i>S</i> ^b	1.025

^a*R*₁ = $\sum ||F_o| - |F_c|| / \sum |F_o|$, *wR*₂ = $[\sum w(F_o^2 - F_c^2)^2 / \sum wF_o^4]^{1/2}$. ^bGoodness-of-fit *S* = $[\sum w(F_o^2 - F_c^2)^2 / (n - p)]^{1/2}$, where *n* is the number of reflections and *p* the number of parameters.

TABLE 1 Crystallographic data and refinement details for resorcinarene **4**