Monopyrrolotetrathiafulvalenium Dication and its Complexation with 1,5 Dinaphtho[38]crown-10

Jan O. Jeppesen

► To cite this version:

Jan O. Jeppesen. Monopyrrolotetrathiafulvalenium Dication and its Complexation with 1,5 Dinaphtho[38]crown-10. *Supramolecular Chemistry*, 2009, 21 (01-02), pp.157-163. 10.1080/10610270802549725 . hal-00513550

HAL Id: hal-00513550

<https://hal.science/hal-00513550>

Submitted on 1 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Monopyrrolotetrathiafulvalenium Dication and its Complexation with 1,5 Dinaphtho[38]crown-10

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2008-0120.R1
Manuscript Type:	Special Issue Paper
Date Submitted by the Author:	10-Oct-2008
Complete List of Authors:	Jeppesen, Jan; Univeristy of Southern Denmark, Chemistry
Keywords:	Crown ether, Host-Guest chemistry, Oxidation, Supramolecular chemistry, Tetrathiafulvalenes
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p> <p>Scheme 1.cdx Scheme 2.cdx Figure 1.cdx Figure 2.cdx Figure 3.cdx Figure 4.cdx Figure 5.cdx Graphical Abstract.cdx</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Submission to the ISMSC 2008 Special Issue of Supramolecular Chemistry

Monopyrrolotetrathiafulvalenium Dication and its Complexation with 1,5-Dinaphtho[38]crown-10

BO M. PETERSEN, MONICA JØRGENSEN, PAUL C. STEIN and
JAN O. JEPPESEN*

Department of Physics and Chemistry, University of Southern Denmark, Campusvej 55, DK-5230, Odense M, Denmark

* Corresponding author. E-mail: joj@ifk.sdu.dk

Graphical Abstract

The synthesis and characterisation of a monopyrrolotetrathiafulvalene (MPTTF) dication 1^{2+} , isolated both as its perchlorate (ClO_4^-) and hexachloroantimonate (SbCl_6^-) salt, is described. The complexation between the π -electron acceptor 1^{2+} and the π -electron donating macrocyclic polyether 1,5-dinaphtho[38]crown-10 (1/5DN38C10) has been investigated using ^1H NMR spectroscopy and a binding constant (K_a) of $3.7 \pm 0.3 \times 10^2 \text{ M}^{-1}$ was obtained for the 1^{2+} ⋯1/5DN38C10 complex (as its perchlorate salt) in acetonitrile at 298 K. The results reveal that the strength of the binding between 1^{2+} and 1/5DN38C10 is one order of magnitude smaller than that for the binding between the parent tetrathiafulvalene (TTF) dication (*i.e.*, TTF^{2+}) and 1/5DN38C10.

Keywords: Crown ether; Host-Guest chemistry; Oxidation; Supramolecular chemistry; Tetrathiafulvalenes

INTRODUCTION

The redox-active tetrathiafulvalene (TTF) unit and its derivatives have been intensively studied in the past three decades [1], because they can exist in three stable forms, specifically $\text{TTF}(0)$, TTF^+ , and TTF^{2+} and for this reason TTF derivatives have found widespread use in materials chemistry [2]. In recent years, however, the growing utility of TTF derivatives as building blocks in macrocyclic and supramolecular chemistry has revealed that the TTF unit is useful beyond the field of materials chemistry and the unique π -electron donor properties of TTF and its derivatives have been used to prepare a number of intriguing molecular and supramolecular systems, including chemical sensors [3], charge separating ligands [4],

porphyrins [5], artificial muscles [6], molecular shuttles [7], chemical springs [8], and synthetic switches [9]. Previous studies [10] have shown that TTF and its derivatives are excellent recognition sites for the π -electron accepting tetracationic cyclophane [11], cyclobis(paraquat-*p*-phenylene) (CBPQT⁴⁺). Although a considerable range of catenanes [12,13] and rotaxanes [7,13,14], incorporating a TTF unit in conjunction with CBPQT⁴⁺, have been reported, the use of the TTF dication (*i.e.*, TTF²⁺) as an π -electron acceptor in supramolecular chemistry has only been rarely described in the literature. In fact, to the best of our knowledge, only one example has been reported in the literature [15], namely the use of the parent TTF dication as a guest for the π -electron donating macrocyclic polyether 1,5-dinaphtho[38]crown-10 (1/5DN38C10). The binding constant (K_a) for the TTF²⁺⊂1/5DN38C10 complex was determined to be $4.1 \times 10^3 \text{ M}^{-1}$ in acetonitrile at 298 K, indicating a rather strong association between 1/5DN38C10 and the TTF dication. Therefore, we decided to elaborate on this fundamental finding. In this paper, we describe the synthesis and characterisation of a monopyrrolotetrathiafulvalene (MPTTF) dication isolated both as its perchlorate (ClO₄⁻) and hexachloroantimonate (SbCl₆⁻) salt followed by our complexation studies between the MPTTF dication and 1/5DN38C10.

RESULTS AND DISCUSSIONS

The MPTTF derivative investigated in this work, namely 5-methyl-2-[4,5-bis(pentylthio)-1,3-dithiole-2-ylidene]-1,3-dithiolo[4,5-*c*]pyrrole (**1**) was prepared according to [a slightly modified \(cf. Experimental\)](#) literature procedure [16]. The corresponding monopyrrolotetrathiafulvalenium dication **1**²⁺ was subsequently synthesized as outlined in Scheme 1. Addition of excess of iron(III)perchlorate (Fe(ClO₄)₃) dissolved in 1,2-dimethoxyethane to a solution of **1** in 1,2-dimethoxyethane produced a blue precipitate, which was filtered affording **1**•(ClO₄)₂ as an analytically dark blue powder in 75% yield. The

Deleted: the

1
2 bis(hexachloroantimonate) salt of $\mathbf{1}^{2+}$ (i.e., $\mathbf{1} \cdot (\text{SbCl}_6)_2$) was synthesized by addition of
3
4 2.3 equivalents of the chemical oxidant tris(4-bromophenyl)ammoniumyl
5
6 hexachloroantimonate to a solution of $\mathbf{1}$ in dichloromethane providing $\mathbf{1} \cdot (\text{SbCl}_6)_2$ as a dark
7
8 solid in 97% yield after precipitation using petroleum ether and filtration.
9

10 The absorption spectra (Fig. 1) of $\mathbf{1} \cdot (\text{ClO}_4)_2$ and $\mathbf{1} \cdot (\text{SbCl}_6)_2$ recorded in acetonitrile at room
11
12 temperature both shows a strong absorption band centred around 630 nm, which is
13
14 characteristic of structures containing the doubly-oxidised MPTTF $^{2+}$ unit [9c]. A comparison
15
16 (Fig. 2) of the ^1H NMR spectra of the neutral $\mathbf{1}$ and the dicationic $\mathbf{1} \cdot (\text{SbCl}_6)_2$ revealed
17
18 significant changes in the chemical shift of the signals associated with the protons located
19
20 close to the MPTTF unit, by virtue of the deshielding arising from the two sets of cationic
21
22 6π -electron aromatic rings. Consequently, the signals for the NMe, SCH_2 , and pyrrolo protons
23
24 are shifted downfield with 0.55, 0.67, and 1.60 ppm, respectively, whereas the remaining
25
26 protons on the pentyl chains only shows small changes in their chemical shift values upon
27
28 oxidation of the MPTTF unit to MPTTF $^{2+}$.
29

30 Formation of the MPTTF dication $\mathbf{1}^{2+}$ (as is its perchlorate salt) was also followed *in situ*
31
32 using absorption spectroscopy in the region from 350 to 1000 nm. The titrational addition of
33
34 1.0 equivalent of iron(III)perchlorate to the yellow MPTTF compound $\mathbf{1}$ in acetonitrile at
35
36 room temperature leads to the formation of the MPTTF radical cation $\mathbf{1}^{+\cdot}$ as evidenced by the
37
38 production of an intense green-coloured solution and the appearance (Fig. 3A) of two
39
40 absorption bands centred on 432 and 761 nm, respectively, in the absorption spectra. The
41
42 titration of the green-coloured acetonitrile solution containing the MPTTF radical cation $\mathbf{1}^{+\cdot}$
43
44 with an additional 1.0 equivalent of iron(III)perchlorate produce an intense blue-coloured
45
46 solution and results in the disappearance (Fig. 3B) of the 432 and 761 nm band, respectively,
47
48 and the appearance of a more narrow absorption band at 637 nm consistent with the formation
49
50 of the MPTTF dication $\mathbf{1}^{2+}$ (*vide supra*). Consequently, the doubly-oxidised MPTTF unit $\mathbf{1}^{2+}$
51
52
53
54
55
56
57
58
59
60

can be readily distinguished from the neutral MPTTF unit **1** (pale yellow) and the mono-oxidised MPTTF unit **1**²⁺ (green) by its intense blue colour.

The π -electron donating macrocyclic polyether 1/5DN38C10 was envisioned as a suitable host for the MPTTF dication **1**²⁺. By employing ¹H NMR spectroscopy **1**²⁺ was shown to form a complex with 1/5DN38C10. A comparison of the ¹H NMR spectra (CD₃CN, 298 K) of the non-complexed **1**²⁺ and **1**²⁺ in the presence of 1/5DN38C10 reveals significant chemical shift differences for the resonances associated with the aromatic *H*-2, *H*-3, and *H*-4 protons (Scheme 2) located on the 1/5DN38C10 macrocycle (upfield shifts, $\Delta\delta = -0.07$ to -0.23 ppm) indicating that **1**²⁺ is complexed by 1/5DN38C10. A continuous variation ¹H NMR experiment was carried out to determine the stoichiometry of the binding between **1**²⁺ and 1/5DN38C10. The Job plot obtained (Fig. 4) exhibit a maximum at a mole ratio of approximately 0.5, thus indicating that the 1/5DN38C10 receptor forms a 1:1 complex with **1**²⁺. To determine the binding constant (K_a) for the 1:1 complex between **1**²⁺ and the 1/5DN38C10 receptor ¹H NMR titration experiments (Fig. 5) were carried out in CD₃CN at 298 K. By following the changes (Fig. 5) in the chemical shift for the resonances associated with the three non-equivalent aromatic protons (*i.e.*, *H*-2, *H*-3, and *H*-4) in the 1/5DN38C10 macrocycle, upon addition of an increasing amount of **1**²⁺ to the receptor 1/5DN38C10, the binding constant for this 1:1 interaction could be determined using nonlinear curve fitting [17]; this yielded an average K_a value of $3.7 \pm 0.3 \times 10^2 \text{ M}^{-1}$ for the interaction between the **1**²⁺ dication and 1/5DN38C10. A comparison of the K_a values for the **1**²⁺⊂1/5DN38C10 and TTF²⁺⊂1/5DN38C10 ($K_a = 4.1 \times 10^3 \text{ M}^{-1}$, *vide supra*) complexes reveal that the strength of the binding between **1**²⁺ and 1/5DN38C10 is one order of magnitude smaller than that for the binding between TTF²⁺ and 1/5DN38C10. This observation can most likely be accounted for by the fact that the MPTTF unit is more steric hindered as compared to the TTF unit which lead to the formation of a less stable complex in the former case.

CONCLUSIONS

In conclusion, we have shown that it is possible to oxidise the MPTTF compound **1** to its corresponding dication 1^{2+} using either iron(III)perchlorate or tris(4-bromophenyl)ammoniumyl hexachloroantimonate. The resulting salts $1 \cdot (\text{ClO}_4)_2$ and $1 \cdot (\text{SbCl}_6)_2$ were isolated and characterised. It was shown that 1^{2+} forms a 1:1 complex with the π -electron donating macrocyclic polyether 1/5DN38C10 and a K_a value of $3.7 \pm 0.3 \times 10^2 \text{ M}^{-1}$ was obtained for the $1^{2+} \subset 1/5\text{DN38C10}$ complex (as its perchlorate salt) in acetonitrile at 298 K. These findings are undoubtedly not unimportant when it comes to future design and construction of supramolecular systems based on monopyrrolotetrathiafulvalenium dications.

EXPERIMENTAL

Chemicals and solvents were purchased from Aldrich and were used as received, except 2-(4,5-bis(pentylthio)-1,3-dithiole-2-ylidene)-1,3-dithiolo[4,5-*c*]pyrrole [18] and 1,5-dinaphtho[38]crown-10 (1/5DN38C10) [19] which were synthesised according to literature procedures. All reactions were carried out under an anhydrous nitrogen atmosphere. Thin layer chromatography (TLC) was carried out using aluminium sheets pre-coated with silica gel 60F (Merck 5554). The plates were inspected under UV light and, if required, developed in I_2 vapour. Column chromatography was carried out using silica gel 60F (Merck 9385, 0.040–0.063 mm). Melting points were determined on a Büchi melting point apparatus and are uncorrected. ^1H and ^{13}C NMR spectra were recorded at 298 K on a Varian Gemini-300BB (300 MHz) or on Varian Unity (500 MHz) instrument using tetramethylsilane (TMS) or the residual solvent as the internal standard. All chemical shifts are quoted on a δ scale in ppm, and all coupling constants (J) are expressed in Hertz (Hz). Samples were prepared using CDCl_3 or CD_3CN purchased from Cambridge Isotope Labs. Absorption spectra were recorded

Deleted: (1)

Deleted: 16

Formatted: Superscript

on a Shimadzu 1601 spectrophotometer, while elemental analyses were performed by the Atlantic Microlab, Inc., Atlanta, Georgia.

5-Methyl-2-[4,5-bis(pentylthio)-1,3-dithiole-2-ylidene]-1,3-dithiolo[4,5-c]pyrrole (1)

2-(4,5-Bis(pentylthio)-1,3-dithiole-2-ylidene)-1,3-dithiolo[4,5-c]pyrrole (0.84 g, 1.88 mmol) was dissolved in anhydrous DMF (15 mL) and degassed (N₂, 15 min). NaH (60% suspension in mineral oil, 0.75 g, 18.8 mmol) was added in one portion, resulting in an immediate color change from yellow to brown. The reaction mixture was stirred for 12 min at room temperature before iodomethane (2.6 mL, 41.8 mmol) was added. Stirring was continued for another 1 h before the reaction mixture was poured into brine (150 mL) and extracted with CH₂Cl₂ (2 × 100 mL). The combined organic phases were carefully washed with brine (2 × 100 mL), dried (MgSO₄), and the solvent removed in vacuo. The resulting red-brown oil was purified by column chromatography (300 mL SiO₂, Ø = 5 cm, CH₂Cl₂/cyclohexane 1:1 v/v). The yellow band was collected and concentrated to give the title compound **1** as a yellow oil which solidified upon standing. Yield: 0.89 g (quant.); ¹H NMR (300 MHz, CDCl₃): δ = 0.90 (t, *J* = 7.2 Hz, 6H, CH₂CH₃), 1.23–1.46 (m, 8H, CH₂CH₂CH₃), 1.63 (pentet, *J* = 7.2 Hz, 4H, SCH₂CH₂CH₂), 2.81 (t, *J* = 7.2 Hz, 4H, SCH₂CH₂), 3.62 (s, 3H, NCH₃), 6.40 ppm (s, 2H, pyrrole-*H*); ¹³C NMR (75 MHz, CDCl₃): δ = 14.1 (CH₂CH₃), 23.3 (CH₂CH₂CH₃), 29.6 (CH₂CH₂CH₃), 30.8 (SCH₂CH₂), 36.4 (SCH₂CH₂), 37.3 (NCH₃), 110.7 (fulvalene C=C), 113.6 (Pyrrole-α), 119.1 (Pyrrole-β), 120.1 (fulvalene C=C), 127.6 ppm (dithiole C=C).

Formatted: Font: Italic

5-Methyl-2-(4,5-bis(pentylthio)-1,3-dithiole-2-ylidene)-1,3-dithiolo[4,5-c]pyrrolonium bis(perchlorate) (1•(ClO₄)₂)

CAUTION! *The dry product is explosive. It will explode when heated to approximately 160 °C. Explosion can also occur when scraping the compound off a glass surface with a metal spatula.*

A solution of $\text{Fe}(\text{ClO}_4)_3 \cdot x\text{H}_2\text{O}$ (1.92 g) in anhydrous 1,2-dimethoxyethane (20 mL) was added dropwise to a solution of 5-methyl-2-(4,5-bis(pentylthio)-1,3-dithiole-2-ylidene)-1,3-dithiolo[4,5-*c*]pyrrole (**1**) (0.097 g, 210 μmol) in anhydrous 1,2-dimethoxyethane (3 mL) resulting in a colour change from yellow to green followed by a colour change from green to blue and the formation of a dark blue precipitate. The precipitate was collected by filtration and washed with a small amount of anhydrous 1,2-dimethoxyethane to give a blue solid, which was dissolved in a minimum of anhydrous MeCN. Evaporation of the solvent *in vacuo* gave the title compound **1**•(ClO_4)₂ as a dark blue powder. Yield: 0.104 g (75%); UV-Vis (MeCN, 293 K): λ_{max} (ϵ) = 404 nm ($7.2 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$), 633 nm ($1.4 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$); elemental analysis calcd (%) for $\text{C}_{19}\text{H}_{27}\text{Cl}_2\text{NO}_8\text{S}_6$: C 34.54, H 4.12, N 2.12, S 29.12; found: C 34.26, H 4.15, N 2.29, S 28.90.

5-Methyl-2-(4,5-bis(pentylthio)-1,3-dithiole-2-ylidene)-1,3-dithiolo[4,5-*c*]pyrrolium

bis(hexachloroantimonate) (**1**•(SbCl_6)₂)

5-Methyl-2-(4,5-bis(pentylthio)-1,3-dithiole-2-ylidene)-1,3-dithiolo[4,5-*c*]pyrrole (**1**) (0.075 g, 162 μmol) was dissolved in CH_2Cl_2 (50 mL) and the resulting yellow solution was degassed (N_2) by placing the solution in an ultrasound bath for 15 min. Tris(4-bromophenyl)ammoniumyl hexachloroantimonate (0.301 g, 369 μmol) was added in one portion causing an immediate colour change from yellow to dark green. The mixture was stirred for 5 min under N_2 before being poured into petroleum ether (500 mL). Stirring was continued for another 30 min before the dark precipitate was collected by filtration and dried *in vacuo* providing the title compound **1**•(SbCl_6)₂ as a dark solid. Yield: 0.178 g (97%); ^1H NMR (500 MHz, CD_3CN): δ = 0.94 (t, J = 7.2 Hz, 6H, CH_2CH_3), 1.40 (sextet, J = 7.2 Hz, 4H, CH_2CH_3), 1.47–1.53 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.89 (pentet, J = 7.2 Hz, 4H, SCH_2CH_2), 3.48 (t, J = 7.2 Hz, 4H, SCH_2CH_2), 4.17 (s, 3H, NCH_3), 8.00 ppm (s, 2H, pyrrole-*H*); UV-Vis (MeCN, 293 K): λ_{max} (ϵ) = 270 nm ($2.8 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$), 633 nm ($1.2 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$).

Acknowledgements

We gratefully acknowledge financial support provided by the Danish Natural Science Research Council through the SONS Programme of the European Science Foundation, which is also funded by the European Commission, Sixth Framework Programme and the Strategic Research Council (Project #2117-05-0115) in Denmark.

References

- [1] (a) Segura, J. L.; Martín, N. *Angew. Chem. Int. Ed.* **2001**, *40*, 1372; (b) Schukat, G.; Fanghänel, E. *Sulfur Rep.* **2003**, *24*, 1; (c) Becher, J.; Jeppesen, J. O.; Nielsen, K. *Synthetic Met.* **2003**, *133–134*, 309; (d) Otsubo, T.; Takimiya, K. *Bull. Chem. Soc. Jpn.* **2004**, *77*, 43; (e) Jeppesen, J. O.; Nielsen, M. B.; Becher, J. *Chem. Rev.* **2004**, *104*, 5115; (f) Gorgues, A.; Hudhomme, P.; Sallé, M. *Chem. Rev.* **2004**, *104*, 5151; (g) Frère, P.; Skabara, P. J. *Chem. Soc. Rev.* **2005**, *34*, 69.
- [2] Organic Superconductors including Fullerenes (Eds.: J. M. Williams, J. R. Ferraro, R. J. Thorn, K. D. Carlson, U. Geise, H. H. Wang, A. M. Kini, M.-H. Wangbo), Prentice-Hall, Engelwood Cliffs, 1992; (b) Bryce, M. R. *J. Mater. Chem.* **2000**, *10*, 589; (c) Nielsen, M. B.; Lomholt, C.; Becher, J. *Chem. Soc. Rev.* **2000**, *29*, 153.
- [3] (a) Heuzé, K.; Mézière, C.; Fourmigué, M.; Batail, P.; Coulon, C.; Canadell, E.; Auban-Senzier, P.; Jérôme, D. *Chem. Mater.* **2000**, *12*, 1898; (b) Trippe, G.; Levillain, E.; Derf, F. L.; Gorgues, A.; Sallé, M.; Jeppesen, J. O.; Nielsen, K.; Becher, J. *Org. Lett.* **2002**, *4*, 2461; (c) Nielsen, K. A.; Jeppesen, J. O.; Levillain, E.; Becher, J. *Angew. Chem. Int. Ed.* **2003**, *42*, 187; (d) Lyskawa, J.; Derf, F. L.; Levillain, E.; Mazari, M.; Salle, M.; Dubois, L.; Viel, P.; Bureau, C.; Palacin, S. *J. Am. Chem. Soc.* **2004**, *126*, 12194; (e) Trippe, G.; Derf, F. L.; Lyskawa, J.; Mazari, M.; Roncali, J.; Gorgues, A.; Levillain, E.; Salle, M. *Chem. Eur. J.* **2004**, *10*, 6497; (f) Nielsen, K. A.; Cho, W.-S.; Jeppesen, J. O.; Lynch, V.

Deleted: -1409

Deleted: -190

Deleted: t

Deleted: -315

Deleted: -58

Deleted: -5132

Deleted: -5184

Deleted: -98

Deleted: , pp. 1–210

Deleted: -598

Deleted: -164

Deleted: -1904

Deleted: -2464

Deleted: ,

Deleted: -191

Deleted: -12195

Deleted: -6509

- M.; Becher, J.; Sessler, J. L. *J. Am. Chem. Soc.* **2004**, *126*, 16296; (g) Zhao, B.-T.; Deleted: -16297
- Blesa, M.-J.; Mercier, N.; Derf, F. L.; Salle, M. *New J. Chem.* **2005**, *29*, 1164; (h) Lu, Deleted: -1167
- H.; Xu, W.; Zhang, D.; Chen, C.; Zhu, D. *Org. Lett.* **2005**, *7*, 4629; (i) Lu, H.; Xu, W.; Deleted: -4632
- Zhang, D.; Zhu, D. *Chem. Commun.* **2005**, 4777; (j) Nielsen, K. A.; Cho, W.-S.; Sarova, Deleted: -4779
- G. H.; Petersen, B. M.; Bond, A. D.; Becher, J.; Jensen, F.; Guldi, D. M.; Sessler, J. L.;
- Jeppesen, J. O. *Angew. Chem. Int. Ed.* **2006**, *45*, 6848; (k) Nielsen, K. A.; Cho, W.-S.; Deleted: -6853
- Lyskawa, J.; Levillain, E.; Lynch, V. M.; Sessler, J. L.; Jeppesen, J. O. *J. Am. Chem.*
- Soc.* **2006**, *128*, 2444; (l) Nielsen, K. A.; Sarova, G. H.; Martín-Gomis, L.; Fernández- Deleted: -2451
- Lázaro, F.; Stein, P. C.; Sanguinet, L.; Levillain, E.; Sessler, J. L.; Guldi, D. M.; Sastre-
- Santos, Á.; Jeppesen, J. O. *J. Am. Chem. Soc.* **2008**, *130*, 460; (m) Nielsen, K. A.; Deleted: -462
- Martín-Gomis, L.; Sarova, G. H.; Sanguinet, L.; Gross, D. E.; Fernández-Lázaro, F.;
- Stein, P. C.; Levillain, E.; Sessler, J. L.; Guldi, D. M.; Sastre-Santos, Á.; Jeppesen, J. O.
- Tetrahedron.* **2008**, *64*, 8449; Deleted: -8463
- [4] Martín, N.; Sánchez, L.; Guldi, D. M. *Chem. Commun.* **2000**, 113; Deleted: -114
- [5] (a) Becher, J.; Brimert, T.; Jeppesen, J. O.; Pedersen, J. Z.; Zubarev, R.; Bjørnholm, T.;
- Reitzel, N.; Jensen, T. R.; Kjaer, K.; Levillain, E. *Angew. Chem. Int. Ed.* **2001**, *40*,
- 2497; (b) Sadaïke, S.-i.; Takimiya, K.; Aso Y.; Otsubo, T. *Tetrahedron Lett.* **2003**, *44*,
- 161; (c) Li, H.; Jeppesen, J. O.; Levillain, E.; Becher, J. *Chem. Commun.* **2003**, 846; (d) Deleted: -2500
- Valentin, M. Di.; Bisol, A.; Agostini, G.; Liddell, P. A.; Kodis, G.; Moore, A. L.; Deleted: -165
- Moore, T. A.; Gust, D.; Carbonera, D. *J. Phys. Chem. B* **2005**, *109* 14401; (e) Xiao, X.; Deleted: -847
- Xu, W.; Zhang, D.; Xu, H.; Lu, H.; Zhu, D. *J. Mater. Chem.* **2005**, *15*, 2557; (f) Saha,
- S.; Flood, A. H.; Stoddart, J. F.; Impellizzeri, S.; Silvi, S.; Venturi, M.; Credi, A. *J. Am.*
- Chem. Soc.* **2007**, *129*, 12159; (n) Nielsen, K. A.; Levillain, E.; Lynch, V. M.; Sessler, J. Deleted: -12171
- L.; Jeppesen, J. O. *Chem. Eur. J.* **2008**, in press. Formatted: German (Germany)
- Formatted: German (Germany)
- Formatted: German (Germany)
- Formatted: German (Germany)
- Formatted: German (Germany)

- [6] Liu, Y.; Flood, A. H.; Bonvallet, P. A.; Vignon, S. A.; Northrop, B. H.; Tseng, H.-R.; Jeppesen, J. O.; Huang, T. J.; Brough, B.; Baller, M.; Magonov, S.; Solares, S. D.; Goddard, W. A.; Ho, C.-M.; Stoddart, J. F. *J. Am. Chem. Soc.* **2005**, *127*, 9745.

Deleted: -9759

- [7] (a) Pease, A. R.; Jeppesen, J. O.; Stoddart, J. F.; Luo, Y.; Collier, C. P.; Heath, J. R. *Acc. Chem. Res.* **2001**, *34*, 433; (b) Jeppesen, J. O.; Becher, J.; Stoddart, J. F. *Org. Lett.*

Deleted: -444

2002, *4*, 557; (c) Luo, Y.; Collier, C. P.; Jeppesen, J. O.; Nielsen, K. A.; Delonno, E.;

Deleted: -560

Ho, G.; Perkins, J.; Tseng, H.-R.; Yamamoto, T.; Stoddart, J. F.; Heath, J. R.

ChemPhysChem, **2002**, *3*, 519; (d) Yamamoto, T.; Tseng, H.-R.; Stoddart, J. F.;

Deleted: -525

Balzani, V.; Credi, A.; Marchioni, F.; Venturi, M. *Collect. Czech. Chem. Commun.*

2003, *68*, 1488; (e) Jeppesen, J. O.; Nygaard, S.; Vignon, S. A.; Stoddart, J. F. *Eur. J.*

Deleted: -1514

Org. Chem. **2005**, 196; (f) Nygaard, S.; Leung, K. C.-F.; Aprahamian, I.; Ikeda, T.;

Deleted: -220

Saha, S.; Laursen, B. W.; Kim, S.-Y.; Hansen, S. W.; Stein, P. C.; Flood, A. H.;

Stoddart, J. F.; Jeppesen, J. O. *J. Am. Chem. Soc.* **2007**, *129*, 960.

Deleted: -970

- [8] Nygaard, S.; Liu, Y.; Stein, P. C.; Flood, A. H.; Jeppesen, J. O. *Adv. Funct. Mat.* **2007**,

17, 751.

Deleted: -762

- [9] (a) Farren, C.; Christensen, C. A.; FitzGerald, S.; Bryce, M. R.; Beeby, A. *J. Org.*

Chem. **2002**, *67*, 9130; (b) Zhang, G.; Zhang, D.; Guo, X.; Zhu, D. *Org. Lett.* **2004**, *6*,

Deleted: -9139

1209; (c) Nygaard, S.; Laursen, B. W.; Flood, A. H.; Hansen, C. N.; Jeppesen, J. O.;

Deleted: -1212

Stoddart, J. F. *Chem. Commun.* **2006**, *2*, 144; (d) Choi, J. W.; Flood, A. H.; Steuerman,

Deleted: -146

D. W.; Nygaard, S.; Braunschweig, A. B.; Moonen, N. N. P.; Laursen, B. W.; Luo, Y.;

DeIonno, E.; Peters, A. J.; Jeppesen, J. O.; Xu, K.; Stoddart, J. F.; Heath, J. R. *Chem.*

Eur. J. **2006**, *12*, 261.

Deleted: -279

- [10] (a) Anelli, P.-L.; Asakawa, M.; Ashton, P. R.; Bissell, R. A.; Clavier, G.; Górski, R.; Kaifer, A. E.; Langford, S. J.; Mattersteig, G.; Menzer, S.; Philp, D.; Slawin, A. M. Z.;

- Spencer, N.; Stoddart, J. F.; Tolley, M. S.; Williams, D. J. *Chem. Eur. J.* **1997**, *3*, 1113; Deleted: -1135
- (b) Devonport, W.; Blower, M. A.; Bryce, M. R.; Goldenberg, L. M. *J. Org. Chem.* **1997**, *62*, 885; (c) Bryce, M. R.; Cooke, G.; Duclairoir, F. M. A.; Rotello, V. M. *Tetrahedron Lett.* **2001**, *42*, 1143; Deleted: -887
- (d) Nielsen M. B.; Jeppesen, J. O.; Lau, J.; Lomholt, C.; Damgaard, D.; Jacobsen, J. P.; Becher, J.; Stoddart, J. F. *J. Org. Chem.* **2001**, *66*, 3559; Deleted: -1145
- (e) Jeppesen, J. O.; Vignon, S. A.; Stoddart, J. F. *Chem. Eur. J.* **2003**, *9*, 4611; Deleted: -3563
- (f) Nygaard, S.; Hansen, C. N.; Jeppesen, J. O. *J. Org. Chem.* **2007**, *72*, 1617; Deleted: -4625
- [11] (a) Anelli, P.-L.; Ashton, P. R.; Ballardini, R.; Balzani, V.; Delgado, M.; Gandolfi, M. T.; Goodnow, T. T.; Kaifer, A. E.; Philp, D.; Pietraszkiewicz, M.; Prodi, L.; Reddington, M. V.; Slawin, A. M. Z.; Spencer, N.; Stoddart, J. F.; Vicent, C.; Williams, D. J. *J. Am. Chem. Soc.* **1992**, *114*, 193; Deleted: -1626
- (b) Asakawa, M.; Dehaen, W.; L'abbé, G.; Menzer, S.; Nouwen, J.; Raymo, F. M.; Stoddart, J. F.; Williams, D. J. *J. Org. Chem.* **1996**, *61*, 9591; Deleted: -218
- [12] (a) Asakawa, M.; Ashton, P. R.; Balzani, V.; Credi, A.; Hamers, G.; Mattersteig, G.; Montalti, N.; Shipway, A. N.; Spencer, N.; Stoddart, J. F.; Tolley, M. S.; Venturi, M.; White, A. J. P.; Williams, D. J. *Angew. Chem. Int. Ed.* **1998**, *37*, 333; Deleted: -9595
- (b) Balzani, V.; Credi, A.; Mattersteig, G.; Matthews, O. A.; Raymo, F. M.; Stoddart, J. F.; Venturi, M.; White, A. J. P.; Williams, D. J. *J. Org. Chem.* **2000**, *65*, 1924; Deleted: -337
- (c) Collier, C. P.; Mattersteig, G.; Wong, E. W.; Lou, Y.; Beverly, K.; Sampaio, J.; Raymo, F. M.; Stoddart, J. F.; Heath, J. R. *Science* **2000**, *289*, 1172; Deleted: -1936
- (d) Diehl, M. R.; Steuerman, D. W.; Tseng, H.-R.; Vignon, S. A.; Star, A.; Celestre, P. C.; Stoddart, J. F.; Heath, J. R. *ChemPhysChem* **2003**, *4*, 1335; Deleted: -1175
- (e) Kim, Y.-H.; Jang, S. S.; Jang, Y. H.; Goddard III, W. A. *Phys. Lett. Rev.* **2005**, *94*, 156801-1; Deleted: -1339
- (f) Nygaard, S.; Hansen, S. W.; Huffman, J. C.; Jensen, F.; Flood, A. H.; Jeppesen, J. O. *J. Am. Chem. Soc.* **2007**, *129*, 7354; Deleted: -156801-4
- Deleted: -7363

- [13] (a) Flood, A. H.; Peters, A. J.; Vignon, S. A.; Steuerman, D. W.; Tseng, H.-R.; Kang, S.; Heath, J. R.; Stoddart, J. F. *Chem. Eur. J.* **2004**, *10*, 6558; (b) Flood, A. H.; Stoddart, J. F.; Steuerman, D. W.; Heath, J. R. *Science* **2004**, *306*, 2055; (c) Mendes, P. M.; Flood, A. H.; Stoddart, J. F. *Appl. Phys. A* **2005**, *80*, 1197.

Deleted: -6564

Deleted: -2056

Deleted: -1209

- [14] (a) Jeppesen, J. O.; Perkins, J.; Becher, J.; Stoddart, J. F. *Org. Lett.* **2000**, *2*, 3547; (b) Jeppesen, J. O.; Perkins, J.; Becher, J.; Stoddart, J. F. *Angew. Chem. Int. Ed.* **2001**, *40*, 1216; (c) Jeppesen, J. O.; Nielsen, K. A.; Perkins, J.; Vignon, S. A.; Di Fabio, A.; Ballardini, R.; Gandolfi, M. T.; Venturi, M.; Balzani, V.; Becher, J.; Stoddart, J. F. *Chem. Eur. J.* **2003**, *9*, 2982; (d) Tseng, H.-R.; Vignon, S. A.; Celestre, P. C.; Perkins, J.; Jeppesen, J. O.; Di Fabio, A.; Ballardini, R.; Gandolfi, M. T.; Venturi, M.; Balzani, V.; Stoddart, J. F. *Chem. Eur. J.* **2004**, *10*, 155; (e) Kang, S.; Vignon, S. A.; Tseng, H.-R.; Stoddart, J. F. *Chem. Eur. J.* **2004**, *10*, 2555; (f) Lee, I. C.; Frank, C. W.; Yamamoto, T.; Tseng, H.-R.; Flood, A. H.; Stoddart, J. F.; Jeppesen, J. O. *Langmuir* **2004**, *20*, 5809; (g) Laursen, B. W.; Nygaard, S.; Jeppesen, J. O.; Stoddart, J. F. *Org. Lett.* **2004**, *6*, 4167; (h) Flood, A. H.; Laursen, B. W.; Nygaard, S.; Jeppesen, J. O.; Stoddart, J. F. *Org. Lett.* **2006**, *8*, 2205; (i) Nygaard, S.; Laursen, B. W.; Hansen, T. S.; Bond, A. D.; Flood, A. H.; Jeppesen, J. O. *Angew. Chem. Int. Ed.* **2007**, *46*, 6093.

Deleted: -3550

Deleted: ,

Deleted: -1221

Deleted: -3007

Deleted: -172

Deleted: -2564

Deleted: -5828

Deleted: -4170

Deleted: -2208

Deleted: -6097

- [15] Ashton, P. R.; Balzani, V.; Becher, J.; Credi, A.; Fyfe, M. C. T.; Mattersteig, G.; Menzer, S.; Nielsen, M. B.; Raymo, F. M.; Stoddart, J. F.; Venturi, M.; Williams, D. J. *J. Am. Chem. Soc.* **1999**, *121*, 3951.

Deleted: -3957

Deleted: -1260

- [16] Gopee, H.; Nielsen, K. A.; Jeppesen, J. O. *Synthesis* **2005**, 1251.

- [17] Binding Constants: The Measurement of Molecular Complex Stability. Connors, A. K. Wiley: New York, 1987.

- [18] (a) Jeppesen, J. O.; Takimiya, K.; Jensen, F.; Becher, J. *Org. Lett.* **1999**, *1*, 1291.;
(b) Jeppesen, J. O.; Takimiya, K.; Jensen, F.; Brimert, T.; Nielsen, K.; Thorup, N.;
Becher, J. *J. Org. Chem.* **2000**, *65*, 5794.

Deleted: -620

- [19] Hamilton, D. G.; Davies, J. E.; Prodi, L.; Sanders, J. H. M. *Chem. Eur. J.* **1998**, *4*, 608.

For Peer Review Only

Captions to SCHEMES and FIGURES

SCHEME 1 Synthesis of $\mathbf{1} \cdot (\text{ClO}_4)_2$ and $\mathbf{1} \cdot (\text{SbCl}_6)_2$.

SCHEME 2 Complexation of $\mathbf{1}^{2+}$ by 1/5DN38C10.

FIGURE 1 Absorption spectra recorded in MeCN at 293 K of A) $\mathbf{1} \cdot (\text{ClO}_4)_2$, B) $\mathbf{1} \cdot (\text{SbCl}_6)_2$, and C) the oxidant tris(4-bromophenyl)ammoniumyl $\cdot\text{SbCl}_6$.

FIGURE 2 A) ^1H NMR spectrum (300 MHz, 298 K) of the neutral MPTTF $\mathbf{1}$ recorded in CDCl_3 and B) ^1H NMR spectrum (500 MHz, 298 K) of the MPTTF dication $\mathbf{1}^{2+}$ (as its SbCl_6^- salt) recorded in CD_3CN .

FIGURE 3 Absorption spectra of MPTTF $\mathbf{1}$ recorded in MeCN at 293 K upon addition of increasing amounts of iron(III)perchlorate. A) 0–1 equivalent. B) A) 1–2 equivalent.

FIGURE 4 Job plot (CD_3CN , 298 K) corresponding to the interaction between the receptor 1/5DN38C10 and the substrate $\mathbf{1}^{2+}$ (as its hexachloroantimonate salt); these results are consistent with a 1:1 receptor-to-substrate binding. The *H*-4 proton in the naphthalene unit was used as the probe as this proton experience the largest changes in the chemical shift values upon mixing 1/5DN38C10 and $\mathbf{1}^{2+}$.

FIGURE 5 ^1H NMR (CD_3CN , 298 K) titration curves of the perturbation in the chemical shift of the three non-equivalent aromatic protons in the 1/5DN38C10 receptor upon addition of increasing amounts of $\mathbf{1}^{2+}$ (as its perchlorate salt). A) *H*-2, B) *H*-3, and C) *H*-4. The data were analyzed using standard curve fitting methods (*i.e.*, Connors 1:1 model) [17]; this gave the values shown in the insets from which an average K_a value of $3.7 \pm 0.3 \times 10^2 \text{ M}^{-1}$ for the interaction between the $\mathbf{1}^{2+}$ dication and 1/5DN38C10 was obtained.