Brillouin scattering study of the fast dynamics of m-toluidine

Lucia Comez, Monica Pietrella, Daniele Fioretto, Giulio Monaco, Filippo Scarponi, Roberto Verbeni, Luciano Palmieri

► To cite this version:

Lucia Comez, Monica Pietrella, Daniele Fioretto, Giulio Monaco, Filippo Scarponi, et al.. Brillouin scattering study of the fast dynamics of m-toluidine. Philosophical Magazine, 2007, 87 (3-5), pp.651-656. 10.1080/14786430601021686 . hal-00514355

HAL Id: hal-00514355

<https://hal.science/hal-00514355>

Submitted on 2 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Brillouin scattering study of the fast dynamics of m-toluidine

Journal:	<i>Philosophical Magazine & Philosophical Magazine Letters</i>
Manuscript ID:	TPHM-06-May-0161.R1
Journal Selection:	Philosophical Magazine
Date Submitted by the Author:	31-Jul-2006
Complete List of Authors:	Comez, Lucia; Università di Perugia, Dipartimento di Fisica; INFM CRS-SOFT, Università di Roma "La Sapienza" Pietrella, Monica; Università di Perugia, Fisica Fioretto, Daniele; Università di Perugia, Fisica; INFM CRS-SOFT, Università di Roma "La Sapienza" Monaco, Giulio; European Synchrotron Radiation Facility Scarponi, Filippo; Università di Perugia, Fisica Verbeni, Roberto; European Synchrotron Radiation Facility Palmieri, Luciano; Università di Perugia, Fisica
Keywords:	glass transition, phonons, supercooled liquids, X-ray spectroscopy
Keywords (user supplied):

Brillouin scattering study of the fast dynamics of m-toluidine

L. Comez^{1,2}, M. Pietrella¹, D. Fioretto^{1,2}, G. Monaco³, F. Scarponi¹, R. Verbeni³, and L. Palmieri¹

¹ *Dipartimento di Fisica, Università di Perugia, Via Pascoli, I-06123 Perugia, Italy.*

² *INFN CRS-SOFT, c/o Università di Roma "La Sapienza", P. A. Moro 2, I-00185 Roma, Italy.*

³ *European Synchrotron Radiation Facility B.P. 220 F-38043 Grenoble Cedex, France.*

Abstract

Brillouin light and x-ray scattering have been used to investigate the dynamic structure factor of supercooled and glassy m-toluidine in the temperature range between 34 and 303 K. In addition to the ubiquitous α relaxation, evidence is given of a fast relaxation process active in the ns-ps time region. Information on the activation energy of the fast process is obtained from the attenuation of longitudinal acoustic modes.

**corresponding author:*

Email: comez@fisica.unipg.it -Tel-Fax: +39 075 5852767

1. Introduction

The fast dynamics of liquids and glassy systems is dominated by different molecular motions. The most studied is the cooperative motion, commonly referred to as α -process, whose characteristic time progressively increases when decreasing the temperature, eventually diverging when approaching the liquid to glass transition. The mode coupling theory explains the nature of the α -relaxation process at the early stage of supercooling of a liquid as originating from nonlinear coupling between density correlators [1].

In addition to this relaxation process, one or more secondary relaxations are typically revealed in real liquids, which remain active also in the glassy state, whose nature is still not completely understood.

M-toluidine is a fragile glass former characterized by a spatial organization of the molecules induced by hydrogen bonds. A recent inelastic x-ray scattering (IXS) investigation of the dynamic structure factor of m-toluidine has given evidence for a cusp-like behaviour of the nonergodicity factor and of a critical nonergodicity parameter which oscillates in phase with the static structure factor, showing that local order at molecular level can coexist with the signatures of glass transition predicted by the idealized version of the mode-coupling theory [2].

M-toluidine has already been the subject of experimental investigations concerning its acoustic properties and dynamical processes by means of ultrasonic measurements [3-5] and Brillouin light scattering (BLS) experiments [6]. In this paper we contribute to this topic reporting new BLS measurements taken on m-toluidine, which extend the study of the dynamics of density fluctuations to the glassy phase ($T_g=187$ K). From the frequency position and linewidth of Brillouin lines, related to the velocity and the attenuation of longitudinal acoustic modes propagating in the system, and from a comparison with previous IXS data, we give evidence of a fast relaxation process active also below the glass transition temperature, we estimate its activation energy and discuss its nature.

2. Experimental details

M-toluidine ($\text{CH}_3\text{-C}_6\text{H}_4\text{-NH}_2$), purchased at 99 % from Merck, has been distilled in order to obtain a high-purity sample and prevent crystallization. The sample was transferred into a cylindrical pyrex cell, 1 mm inner diameter, which was sealed in dry nitrogen environment. A Cryomech ST405 cryostat was used to regulate temperature. A Coherent-Innova 300 model Ar^+

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

laser was operated with a typical power of 300 mW on a single mode of the 514.5 nm line. The light scattered from the sample in the backscattering geometry was analysed by means of a Sandercock-type (3+3)-pass tandem Fabry-Perot interferometer characterized by a finesse of about 100 and a contrast ratio higher than 10^9 . Spectra were collected in the temperature range between 34 and 303 K. IXS measurements were performed at the very high energy resolution IXS beamline ID16 [7] at the European Synchrotron Radiation Facility from 54 to 293 K. At each temperature, five different Q-values are typically detected, namely $Q=1, 2, 4, 7$, and 10 nm^{-1} , as described in Ref. [2]. Both experiments have been performed on cooling.

3. Results and discussion

Figure 1 shows typical BLS and IXS spectra of m-toluidine taken at values of the exchanged wavevector $Q = 0.038$ and 2 nm^{-1} , respectively, at ambient temperature. Both spectra show inelastic features, the so called Brillouin peaks, at about 10 and 600 GHz, respectively. The presence of relaxation processes coupled with density fluctuations can be revealed by the temperature and Q behaviours of velocity and attenuation of acoustic modes, proportional to the frequency position Ω and linewidth Γ of Brillouin lines. In all measured BLS spectra the linewidth of Brillouin peaks is larger than the instrumental function and a non-negligible quasielastic (Mountain) contribution is observed in the central part of the spectra, indicating the presence of relaxation processes. Among these, the structural relaxation is the most important for temperatures higher than T_g . For temperatures close and below the glass transition, however, the characteristic time of the structural relaxation becomes so long that it cannot be responsible for the hypersonic absorption. This result gives a direct evidence of the presence, in the ten GHz frequency region, of some secondary relaxation process which remains activated even in the glassy phase. This seems to be a peculiarity of fragile systems, as recently reported in Ref. [8].

[Insert Fig. 1 about here]

To be more quantitative, the values of Ω and Γ have been obtained by fitting the BLS spectra in a narrow frequency region around the Brillouin peak by means of the damped harmonic oscillator (DHO) model function

$$I(q, \omega) = \frac{I_0}{\pi} \frac{\Gamma \Omega^2}{(\omega^2 - \Omega^2)^2 + (\Gamma \omega)^2} \quad [1]$$

convoluted to the instrumental function. In the case of IXS spectra, a Lorentzian function centred at zero frequency was added to account for the central line and the whole spectrum was convoluted by the instrumental function after correction for the detailed balance [2]. The results of the fit are reported in Fig. 2 in the form of longitudinal sound velocity (ω/Q) (see bottom panel of Fig. 2) and apparent kinematic viscosity (Γ/Q^2) (see top panel of Fig. 2). Here, the exchanged momentum Q has been determined through the relationship $Q=4\pi n/\lambda_i$, where n is the refractive index of the medium and λ_i is the incident wavelength of the laser source. For the determination of the refractive index as a function of temperature, we have proceeded as follows: for the liquid phase ($T>T_g$) the refractive index was taken from the literature [9]; for temperatures $T<T_g$, it has been estimated using the Clausius–Mossotti relation, where the thermal variation of the density has been deduced from the thermal expansion coefficient determined from $\alpha V= 5.3 \times 10^{-4} \text{ cm}^3 \text{ K}^{-1}$ - an expression valid in the crystal and in proximity of the glass transition [10] - and from the density value $\rho(T_g)= 1.0745 \text{ gr cm}^{-3}$.

In figure 2, together with our results we also report previous BLS data [6], obtained in a narrow temperature range, showing a satisfactory agreement between two series of data. The unrelaxed limit of Γ/Q^2 , providing the longitudinal kinematic viscosity η_∞/ρ , can be reached either for a given temperature at very high frequency or for a given frequency at very low temperature. This condition is met by IXS spectra in the whole reported T range and by BLS only in the low temperature limit, where the values of Γ/Q^2 obtained by the two techniques tend to coincide. When increasing the temperature, the BLS values of Γ/Q^2 increase and become a factor 3 higher than the unrelaxed limit even for $T<T_g$. This behaviour is common in other glass-formers, studied by some of us (see, for instance, Refs. [11-15]), where this excess of acoustic attenuation has been proved to be related to the interplay of acoustic modes with internal degrees of freedom. On these bases also in case of m-toluidine, we believe that the fast process that emerges from this analysis - a relaxation of density fluctuations with a characteristic relaxation time in the ns-ps timescale - could be associated to a coupling among the translational and one or more (still unidentified) intramolecular degrees of freedom, possibly associated to the presence of the NH_2 and CH_3 groups.

[Insert Fig. 2 about here]

In the hypothesis that this relaxation process is thermally activated, i.e. its characteristic time follows the Arrhenius law $\tau(T) = \tau_0 \exp(E_a/k_B T)$, the value of the activation energy E_a can be estimated from the temperature dependence of Γ . In fact it can be shown [13] that, if the strength of the fast relaxation process is negligibly temperature dependent, the major contribution to the width of Brillouin lines comes from the temperature dependence of the relaxation time through the relationship:

$$\lim_{\omega\tau \gg 1} \left(\frac{\Gamma}{Q^2} \right) = \frac{\eta_\infty}{\rho} + \frac{\Delta}{\rho} \sin\left(\frac{\beta\pi}{2}\right) \Omega^{-(\beta+1)} \tau_0^{-\beta} \exp\left(-\frac{\beta E_a}{k_B T}\right) \quad [2]$$

where β is the stretching parameter whose value is 1 for single exponential relaxation and $0 < \beta < 1$ for stretched functions, such as Cole-Davidson or Cole-Cole functions. By a fitting procedure of Eq. (2) to the data of Fig. 2, in the 34-134 K temperature range, the value $\beta E_a = (611 \pm 80)$ K is obtained, corresponding to the solid line in the figure.

As order of magnitude, the quantity βE_a , evaluated for m-toluidine, well compares with the values calculated in other fragile glass formers of different molecular weight: orthoterphenyl $[(140 \pm 20)$ K], polybutadiene $[(260 \pm 20)$ K] and salol $[(400 \pm 40)$ K], where the existence of an intramolecular contribution in the spectrum of density fluctuations has been verified. [8,13,14,16]. Concerning this topic, a viscoelastic analysis of BLS spectra is currently in progress in order to gain further information on the temperature behaviour of the shape and of the characteristic time of this fast relaxation.

Finally, some additional comment would have to be reported with respect to the figure 2. Magnifying the image of the kinematic viscosity (Γ/Q^2) as a function of temperature, the appearance of two features can be identified, one at T_g where the slope drastically changes, and a second one at about 135 K. At the same time, observing the velocity (ω/Q) curves of fig. 2, this kink-like anomaly is present again in the BLS data. This feature in the acoustic properties at about 135 K could be related to an effect observed in the thermodynamic properties of m-toluidine. It is well known that the heat capacity undergoes a large drop in vicinity of the vitrification at the usual temperature of glass transition, T_g , named $T_{g\alpha}$. On further cooling, some systems exhibit another drop at a characteristic temperature, called $T_{g\beta}$. An exhausting explanation of this phenomenon still does not exist. A possible interpretation refers to fast local rearrangements that freeze out at a much lower temperature with respect to $T_{g\alpha}$ where long range diffusion ceases. First observed in

isopropylbenzene [17], this “secondary glass transition” has also been characterized for several molecular liquids by Oguni and co-workers [18], and experimentally observed in m-toluidine by Simionesco et al. [19]. In particular, $T_{g\beta}$ in m-toluidine turns out about to 130 K, a temperature surprisingly similar to the one where the observed change can be recognised in our BLS data. This evidence hints at a correspondence between dynamics and thermodynamics anomalies, which deserves more efforts of understanding. Also, new data in systems which present similar features would be precious.

4. Conclusions

In conclusion, a fast relaxation process has been reported to affect the dynamics of density fluctuations of m-toluidine in the ns-ps range explored by Brillouin light scattering. The presence of a secondary relaxation, beside the structural relaxation process, has been identified by examining the temperature behaviour of longitudinal kinematic viscosity in proximity of the glass transition, which shows an excess with respect to the unrelaxed value estimated through inelastic x-ray scattering. This behaviour is strongly resembling to that of other fragile glass-formers where the presence of an intramolecular contribution in the spectrum of density fluctuations has been recognized. Further, the activation energy for the fast relaxation process evaluated here well compares with those of the above mentioned glass-forming systems. This indication supports the idea that the coupling with internal degrees of freedom as a reasonable origin for the fast relaxation process revealed in m-toluidine in the BLS frequency region.

References

- [1] W. Götze and L. Sjögren, Rep. Prog. Phys. **55** 241 (1992).
- [2] L. Comez, S. Corezzi, G. Monaco, R. Verbeni, D. Fioretto, Phys. Rev. Lett. **94** 155702 (2005).
- [3] M. Cutroni and A. Mandanici, J. Chem. Phys. **114** 7124 (2001).
- [4] M. Cutroni and A. Mandanici, J. Chem. Phys. **116** 7326 (2002).
- [5] C. Dreyfus, R. Gupta, B. Bonello, C. Bousquet, A. Taschin, M. Ricci, G. Pratesi, J. Chem. Phys. **116** 7323 (2002).

- [6] A. Aouadi, C. Dreyfus, M. Massot, R. M. Pick, T. Berger, W. Steffen, A. Patkowski, C. Alba-Simionesco, *J. Chem. Phys.* **112** 9860 (2000).
- [7] ESRF, <http://www.esrf.fr/UsersAndScience/Experiments/>.
- [8] L. Comez, D. Fioretto, G. Monaco, G. Ruocco, *J. Non-Cryst. Solids* **307-310** 148-153 (2002).
- [9] J. Timmermans, in *Physico-Chemical Constants of Pure Organic Compounds*, (Elsevier, New York 1950).
- [10] C. Alba-Simionesco, D. Morineau, B. Frick, N. Higonenc, H. Fujimori, *J. Non-Cryst. Solids*, **235-237** 367 (1998).
- [11] G. Monaco, D. Fioretto, C. Masciovecchio, G. Ruocco, *Phys. Rev. Lett.* **82** 1776 (1999).
- [12] S. Mossa, G., Monaco, G. Ruocco, *Europhys Lett.* **60** 92 (2002).
- [13] D. Fioretto, C. Masciovecchio, M. Mattarelli, G. Monaco, L. Palmieri, G. Ruocco, F. Sette, *Phil. Mag. B* **82** 273 (2002);
- [14] D. Fioretto, M. Mattarelli, C. Masciovecchio, G. Monaco, G. Ruocco, *Phys. Rev. B* **65** 224205 (2002).
- [15] R. H. Gee and R. H. Boyd, *J. Chem. Phys.* **101** 8028 (1994).
- [16] G. Monaco, D. Fioretto, L. Comez, G. Ruocco, *Phys. Rev. E* **63** 061502 (2001).
- [17] K. Kishimoto, S. Suga, S. Seki, *Bull. Chem. Soc. Jpn.* **46** 3020 (1973).
- [18] H. Fujimori and M. Oguni, *Solid State Comm.* **94** 157 (1995).
- [19] C. Alba-Simionesco, J. Fan, C. A. Angell, 1999, *J. Chem. Phys.* **110** 5262 (1999).

Figure captions

Figure.1

BLS (top panel) and IXS (bottom panel) spectra of m-toluidine taken at ambient temperature and $Q=0.038 \text{ nm}^{-1}$ and $Q=2 \text{ nm}^{-1}$, respectively.

Figure.2

Apparent longitudinal kinematic viscosity (top panel) and longitudinal sound velocity of m-toluidine as obtained from X-ray scattering at $Q=2 \text{ nm}^{-1}$ (open squares) and from light scattering (full circles). The solid line in the top panel represents the result of the fitting procedure of Eq. (2)

to the experimental data. Open circles are from Ref. [6]. Inset: Enlarged view of the region where the fit has been performed to better visualize data and fitting curve.

For Peer Review Only

Figure 1
279x215mm (150 x 150 DPI)

Figure 2
279x215mm (150 x 150 DPI)