Herpesviruses In Respiratory Liquids Of Horses: Putative Implication In Airway Inflammation And Association With Cytological Features

Guillaume Fortier, Emmanuelle van Erck, Christine Fortier, Eric Richard,
Didier Pottier, Stéphane Pronost, Fabien Mischack, Etienne Thiry, Pierre
Lekeux

► To cite this version:

Guillaume Fortier, Emmanuelle van Erck, Christine Fortier, Eric Richard, Didier Pottier, et al..
Herpesviruses In Respiratory Liquids Of Horses: Putative Implication In Airway Inflammation
And Association With Cytological Features. *Veterinary Microbiology*, 2009, 139 (1-2), pp.34.
10.1016/j.vetmic.2009.04.021 . hal-00520653

HAL Id: hal-00520653

<https://hal.science/hal-00520653>

Submitted on 24 Sep 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Herpesviruses In Respiratory Liquids Of Horses:
Putative Implication In Airway Inflammation And Association
With Cytological Features

Authors: Guillaume Fortier, Emmanuelle van Erck, Christine
Fortier, Eric Richard, Didier Pottier, Stéphane Pronost, Fabien
Misczack, Etienne Thiry, Pierre Lekeux

PII: S0378-1135(09)00213-2
DOI: doi:10.1016/j.vetmic.2009.04.021
Reference: VETMIC 4425

To appear in:

Received date: 18-2-2009
Revised date: 31-3-2009
Accepted date: 14-4-2009

Please cite this article as: Fortier, G., van Erck, E., Fortier, C., Richard, E.,
Pottier, D., Pronost, S., Misczack, F., Thiry, E., Lekeux, P., Herpesviruses In
Respiratory Liquids Of Horses: Putative Implication In Airway Inflammation
And Association With Cytological Features, *Veterinary Microbiology* (2008),
doi:10.1016/j.vetmic.2009.04.021

This is a PDF file of an unedited manuscript that has been accepted for publication.
As a service to our customers we are providing this early version of the manuscript.
The manuscript will undergo copyediting, typesetting, and review of the resulting proof
before it is published in its final form. Please note that during the production process
errors may be discovered which could affect the content, and all legal disclaimers that
apply to the journal pertain.

HERPESVIRUSES IN RESPIRATORY LIQUIDS OF HORSES: PUTATIVE IMPLICATION IN AIRWAY INFLAMMATION AND ASSOCIATION WITH CYTOLOGICAL FEATURES

Guillaume Fortier ^{a-b,*}, Emmanuelle van Erck ^b, Christine Fortier ^a, Eric Richard ^{b-c}, Didier Pottier ^d, Stéphane Pronost ^a, Fabien Mischack ^a, Etienne Thiry ^e, Pierre Lekeux ^b.

^a Frank Duncombe Laboratory, 14053 CAEN cedex 4, France.

^b Department of Functional Sciences, Physiology, ^c Department of Infectious and Parasitic Diseases, Virology and viral Diseases, Faculty of Veterinary Medicine, University of Liège, B-4000 LIEGE, Belgium.

^e CIRALE, 14430- Goustranville, France

^d GRECAN, UFR des Sciences Pharmaceutiques, Université de Caen Basse-Normandie, 14000 Caen, France.

* Corresponding author: Guillaume Fortier,

tel.: +33 231 47 19 19; fax: +33 231 47 19 49.

email address: g.fortier@cgl4.fr

Abstract

The objectives of this study were to estimate the prevalence and the potential role of equine herpesviruses (EHVs) detection in both bronchoalveolar lavage (BAL) and tracheal wash (TW). The population included a control group (CTL; 37 TW and 25 BAL) and a pathological group (PAT; 259 TW and 387 BAL), including horses either suffering from respiratory diseases including syndrome of tracheal inflammation, inflammatory airway disease, recurrent airway obstruction, or submitted to respiratory investigation because of exercise intolerance or poor performance. Each respiratory liquid was submitted to a standardised cytological analysis, mentioning the morphological abnormalities of exfoliated epithelial cells (ECAb) and ciliocytophthoria (CCPh) as markers of potential viral infection, as well as PCR assays including a consensus PCR and virus-specific PCR for both equine alphaherpesviruses (EHV-

1; EHV-4) and gammaherpesviruses (EHV-2; EHV-5). The EHV infections were more prevalent in the TW of PAT group ($P = 0.004$), with the highest prevalence being for EHV-2 ($P = 0.006$). The EHV detection in BALs was not significantly different between groups. The EHV detection in TW was correlated to the polymorphonuclear neutrophil (PMN) counts in the respiratory liquid but not with CCPh or ECAb. CCPh or ECAb were associated with both consensus PCR and EHV-2 and EHV-5 virus-type PCR in the BAL only. The significant detection of EHV in the TW of PAT group in association with the PMN increased counts could lead to further investigations about their putative role in equine syndrome of tracheal inflammation.

Keywords: ciliocytophthoria, respiratory disease, horse, herpesvirus, PCR, gammaherpesvirus

1. Introduction

Respiratory diseases are a consensually recognised cause of wastage in the horse industry. Among respiratory diseases, airway inflammation is the most prevalent clinical feature and is largely encountered in the field. Recently, inflammatory airway disease (IAD) was redefined as a clinical syndrome causing a significant elevation of polymorphonuclear neutrophil (PMN) count in the bronchoalveolar lavage fluid (Couetil et al., 2007). Less is however known about the syndrome of tracheal inflammation (STI) even if viruses, bacteria or fungi are suspected as potential actors of this local neutrophilia (Couetil et al., 2007). Modifications in the cytological profiles of BAL or TW have sometimes been evoked in case of viral diseases but without clear demonstration (Whitwell and Greet, 1984; Viel, 2002). Transient neutrophilia in respiratory liquids has been also described after viral exposure but cannot be interpreted as a specific profile (McGorum and Dixon, 1994; Sutton et al., 1998). On the other hand, ciliocytophthoria is a criteria used for the differential diagnosis of viral disease in human pathology, especially for respiratory or genital cytology (Papanicolaou et al., 1961). Whitwell and Greet (1984) also described this anomaly in cases of viral diseases and more particularly in those caused by herpesvirus in horses with tracheal inflammation and mucosal erosion. Pointing out the putative role of viruses in inflammatory airway diseases, Viel (2002) emphasized the need of incorporation of exfoliated epithelial cells (EEC) in the BAL cytological analysis.

Due to their high prevalence and their capability of latency in various sites, viruses and particularly herpesviruses are frequently incriminated in equine poor performance syndromes (Wood et al., 2005). However, alphaherpesviruses such as equid herpesvirus 1 (EHV-1) or equid herpesvirus 4 (EHV-4) do not seem to be clearly associated with such syndromes (Newton et al., 2003; Wood et al., 2005). The alphaherpesviruses are recognised as causative agents of various epidemic diseases in the horse population (i.e; abortion,

myeloencephalopathy) and as well as of upper respiratory tract infections, but equine gammaherpesviruses have been less studied and have been characterized (Agius and Studdert, 1994) and reclassified (Davison et al., 2009) only recently. Their real pathogenic significance remains unclear because of their frequent isolation in healthy horse populations and their high seroprevalence worldwide (Bell et al., 2006). In mice, equid herpesvirus 2 (EHV-2) DNA can be retrieved from the lungs 30 days after experimental exposure and infection (Borchers et al., 2002). In foals or young race horses, this virus has been shown to induce conjunctivitis and mild respiratory diseases, (Bell et al., 2006). In addition, equid herpesvirus 5 (EHV-5) has been recently associated with equine pulmonary multinodular syndrome (Williams et al., 2007).

Performing both consensus PCR and virus-specific PCR assays, the purposes of this study were (1) to precise the DNA prevalence of EHV-1, 2, 4 and 5 in respiratory liquids of horses suffering from various respiratory diseases and compare it to a control group, and (2) to determine if particular cytological profiles of the sampled liquids could be associated with herpesviruses detection and to investigate within the different groups the correlation between EHV's detection and neutrophilia or epithelial cells abnormalities in both the TW and the BAL.

2. Materials and methods

2.1. Horse population and respiratory liquids

The study included 708 samples (647 horses aged 18 months - 13 years; Table 1) of various breeds and/or disciplines (Standardbred, Thoroughbred, Endurance and French saddle horses) harvested between October 2006 and February 2008. Horses from the control group (CTL) were referred to a sports medicine facility (CIRALE, Goustranville, France) for routine clinical and fitness follow-up using standardised protocols. Thorough clinical and

complementary exams were systematically performed, including haematological and biochemical evaluation at rest and post-exercise, Doppler echocardiography, standardised exercise tests on both treadmill and racetrack, locomotor evaluation, high-speed treadmill video-endoscopy of the upper respiratory tract, and collection of respiratory liquids including TW and BAL. Each CTL horse was free of any abnormality for all these clinical investigations, and particularly a scoring mucus ≤ 2 (Gerber et al., 2004), no signs of infection/inflammation at the haemato-biochemical evaluation (Couetil et al., 2007) and both TW and BAL cytological values within proposed references (Robinson, 2003). The pathological group (PAT) was composed of horses with signs of various respiratory disorders or “poor performance syndrome” and originated from 28 veterinary practices located close to training yards or race tracks in the Normandy or Ile de France regions. To be included, at least one of the following criteria had to be detected by the clinician: coughing, scoring mucus > 2 , TW or BAL neutrophilia, or poor performance suspected by the practitioner to be due at least partly to respiratory dysfunctions. Any horse exhibiting dyspnoea at the moment of sampling or clinical signs of either acute viral (i.e. serous nasal discharge, hyperthermia, ...) or bacterial infection (i.e. purulent nasal discharge, leukocytosis, ...), were excluded from the study.

2.2. Samples processing and cytological analysis

Each practice or referring veterinarian was expected to follow the main recommendations regarding sampling and shipment of TW or BAL (Viel, 2002; Pickles et al., 2002; Malikides et al., 2007). Samples of each group were excluded if a set of quality criteria was not met (i.e. presence of squamous pharyngeal cells, cell degradation, contaminating bacterial flora or lack of historical information). Aliquots of respiratory liquids were placed in EDTA- and serum tubes and respectively submitted for cytology and virology. May Grunwald Giemsa stain was applied to slides previously prepared by smooth cytocentrifugation (75 x g, 10 min) using a

Cytospin (Shandon-Thermoscientific, Waltham, MA, USA). Differential cell counts were performed by counting an average of 300 cells and the results of each cell type were expressed as a percentage of the total nucleated cells. Presence of exfoliated epithelial cells (EEC) were noted and defined as either normal or presenting morphological abnormalities (ECAb) when free cilia, ciliated plates or detached ciliated tufts were found in the liquid (Sutton et al., 1997; Viel, 2002) or ciliocytophthoria (CCPh) when the previous criteria was accompanied by cytoplasmic or nuclear modifications (Whitwell and Greet, 1984; Freeman et al., 1985). The EEC were initially included in the sample cell counts to determine if there were any significant differences in proportion between the CTL and the PAT group. Thereafter, EEC were excluded from the differential cell counts but ECAb and CCPh were taken into account in the statistical analysis as “present” or “absent”. Hematological and biochemical tests were performed with an Advia 120 (Bayer) and a Rx Imola (Randox,)

2.3 Nucleic acids extraction and PCR assays

Respiratory liquids were centrifuged at 2000 x g, 4 °C, during 15 min, and pellets were homogenised with 500 µl of supernatant. DNA was extracted from 140 µl of each sample, using the QIAamp DNA Mini Kit (Qiagen, France) according to the manufacturer’s instructions, and eluted in a 50 µl final volume. Concentrated DNA was stored at -80 °C until used. All extraction included a negative control where respiratory liquid was replaced with distilled water and a positive control.

The five PCR assays systematically carried out on each sample for this study are summarized in table 2. A consensus herpesvirus PCR assay was carried out as previously described by Leon et al. (2008). EHV-2 was detected by amplification of a sequence encoding a product homologous to human interleukin-10 (Leon, personal communication 2008; adapted from Borchers and Frolich, 1997) whereas EHV-5 was detected by a virus-specific EHV-5 PCR assay described by Dynon et al. (2001). Classical PCRs (consensus PCR and virus-specific

EHV-2 and EHV-5 PCR) were performed with pure Taq Ready to Go PCR beads (Amersham Biosciences, Europe GmbH, France); the final concentrations of each primer were 1 μ M consensus PCR, 1 μ M EHV-2 and 800 nM EHV-5. Classical PCRs were performed on GeneAmp PCR System 9700 (Applied Biosystems). Virus-specific real-time PCRs were performed as described by Diallo et al. (2007). Each test was used as a simplex assay in a total volume of 25 μ l containing 2x Taqman Universal PCR Master Mix (Applied Biosystems, France), 1 μ M of each primer and the optimum concentration of probe (between 100 and 400 nM as determined after titration). Real time amplifications were performed on Step One Real-Time PCR system (Applied Biosystems). Each different PCR test included a negative control where DNA was replaced with distilled water and a specific positive control where DNA from each virus was amplified. DNA from American Type Culture Collection reference strains for EHV-1 (Kentucky reference strain, KyA), EHV-2 (LK strain) and EHV-4 strain 405/76 and DNA from our laboratory for EHV-5 (Fortier et al. 2008, *in press*). Positive samples by consensual PCR which were negative with each virus-specific PCR assays (EHV-1, 2, 4 and 5), were sequenced after extraction of the PCR product from agarose using the QIAquick gel extraction kit (Qiagen, France). Both sense and anti-sense strands of the PCR products were sequenced (Biofidal, France). The nucleotide sequences were assembled and manually edited (CodonCode Aligner), and consensus PCR sequences were analysed for similarities on the National Center for Biotechnology Information (NCBI) website using databases and the Basic Local Alignment Search Tool program (BLASTn).

2.4 Statistical analysis

Differences in the results between the groups were compared using a Fisher's exact test. A Cochran Armitage trend test was used to evaluate the trend between PMN cell counts and herpesviruses DNA detection in the samples. According to the percentage of PMNs in the

liquids, PMN results were conventionally distributed into 6 different classes, from 0 to 20%, from 21 to 40%, from 41 to 60%, 61 to 80% and > 80%. Odds ratio (OR) were performed to evaluate the risk of being in the PAT group when herpesvirus DNA was detected. All tests were performed on SAS System (Cary, NC, USA). *P* values < 0.05 were considered significant.

3. Results

3.1. PCR assays in the respiratory fluids

In the TW of the CTL group, EHV-2 and EHV-4 were detected by virus-specific PCR whereas EHV-1 and EHV-5 were not. In this group, two samples were positive using the consensus PCR but were not confirmed by the specific tests and no clear identification was achieved using the sequencing. In the PAT group, all the targeted viruses were detected. Using PCR, 114 and 112 PAT TW were positive by virus-specific and consensus PCRs respectively (Table 3); 2 samples were detected positive for both EHV-2 and EHV-5. The prevalence of herpesviruses in the TW was significantly higher in the PAT group ($P=0.004$). Detection of EHV-2 was also higher in the PAT group compared to the CTL group ($P=0.006$). The highest level of detection for both groups was found for EHV-2, with 11% and 33% for the CTL group and PAT group respectively. In the BAL, only EHV-2 and EHV-4 were detected in the CTL group (Table 3), whereas in the PAT group all viruses were detected, as in the TW. The prevalence of these viruses in BAL was not significantly different between groups. The highest level of detection for both groups was found for EHV-2, with 36% and 21% for the CTL group and PAT group respectively. Whatever the group and sample, EHV-2 was the most frequently detected strain. A double detection (EHV-2 and EHV-4) was found in two BAL samples of the PAT group. Among the 708 samples, Asinine herpesvirus 5 (AHV-5) was identified in 24 PAT liquids (16 in the BAL and 8 in the TW of

by sequencing the amplified fragment of the consensus PCR (Table 3 and 4). Among them, a mild positive result was obtained in 8 samples at the second run using the virus-specific EHV-5 PCR. Only consensus PCR positive results were taken into account for the statistical analysis but not virus type specific results. For the two real-time PCR (EHV-1, EHV-4) the cycle threshold (C_t) was mentioned for each analysis. In the TW of PAT group, the C_t range was 22 - 42 for EHV-4 and 36.5 - 38.4 for EHV-1, whereas the C_t was 38 for the only EHV-4 positive sample in the CTL group.

In the BAL of the PAT group, the C_t range was 34 - 44 for EHV-4 and 35 - 40.5 for EHV-1, whereas the C_t was 39 for the only EHV-4 positive sample in the CTL group.

Inclusion in the PAT group was positively and significantly associated with the detection of herpesviruses by both consensus PCR (odds ratio= 3.26, 95% C.I = 1.38-7.04) and EHV-2 specific PCR assay (odds ratio= 4.03, 95% C.I = 1.37-11.74). No significant association was observed for any other herpesvirus neither in the TW nor in the BAL.

3.2. Association between groups, viral detection and cytology of the liquids

The proportion of EEC in the BAL (0.7 % vs 2.7 %) and TW (15.6 % vs 12.8 %) were not significantly different between the CTL group and the PAT group and ranged with normal values (Derksen et al., 1989). Percentages of CCPh or ECAb were not significantly associated with groups (Table 4). No morphological abnormality was detected among the BAL epithelial cells (ECAb and CCPh) in the CTL group. The proportion of CCPh was increased in the PAT group, but not significantly ($P = 0.32$).

As described in Table 5, the presence of one of the two positive criteria for epithelial cell alterations (ECAb, CCPh) was significantly associated with herpesvirus DNA detection for EHV-2 and EHV-5, but in the BAL only. No other significant association was found for EHV-4 in either liquid neither for any other herpesvirus in the TW. No significant association

were found between herpesviruses detection (in TW or BAL) and either lymphocytes, giant cells, mast cells, eosinophils, macrophages, or haemosiderophages. In the TW only, a significant association between PMN counts level and DNA detection with the consensus PCR or the EHV-2 virus-type specific PCR was observed (Table 6), whereas a tendency, at the limit of significance, was found for EHV-4 detection ($P = 0.05$). In the TW, the highest percentage of EHV-2 detection was in the sub-group of 40-60% of PMN and in the sub-group 80-100% of PMN for EHV-4 with the consensus PCR. The mean level of PMN was higher when herpesvirus DNA was detected with consensus PCR ($P = 0.003$) or EHV-2 specific PCR ($P = 0.04$). The results were not significant using EHV-5, EHV-1 and EHV-4. The trend test was not significant for any herpesvirus in the BAL.

No significant trend was found between ciliocytophthoria and the PMN cell counts in either BAL or TW, meanwhile the highest percentage of CCPh was found in the 40-60 % PMN class for TW.

Discussion

The precise role of equine herpesviruses in subclinical respiratory diseases has been documented mainly through the results of serologic assays or naso-pharyngeal sampling studies in young racing horses (Wood et al., 2005).

Previous studies evaluating viral causes of respiratory diseases in young horses in training mainly focused on alphaherpesviruses, which remain the most worldwide studied herpesviruses in the equine population (Newton et al., 2003; Wood et al., 2005). Gammaherpesviruses were seldom included and information lacked regarding their implication in larger populations of horses suffering from various respiratory disorders. Our study aimed at investigating this missing information by undertaking a systematic DNA detection of herpesviruses in TW and BAL respiratory liquids of horses suspected of

respiratory affections. Serologic tools were not implemented for EHV detection as they may not be a reliable method for investigating viral reactivation because of cross-reaction, poor humoral response and lack of seroconversion due to repeated exposures and vaccination. One of the limits in the PCR strategy used in the present study is that it could not allow discrimination between latent, non replicating and lytic states of the detected herpesviruses (Pusterla et al., 2008).

Among the alphaherpesviruses detected in the present study, EHV-4 was much more prevalent than EHV-1, and this can be related to its particular role in clinical respiratory disorders in the field. Even if the C_t values obtained for EHV-1 and EHV-4 were in previously described ranges (Diallo et al., 2007), taking into account the artificial dilution due to the sampling method for respiratory liquids, the detection of latent viruses could not be avoided. EHV-4 were sometimes retrieved from liquids with low C_t (high viral load) when a history of infectious disease of suspected viral origin was mentioned by the practitioner.

Equine herpesvirus DNA detection was clearly different in the two groups of horses, especially in the TW.

The high prevalence of EHV-2 and EHV-5 DNA shown in our study's pathological group is in accordance with previous observations, indicating their frequent detection in nasal swabs, yet from horses without clear clinical signs (Bell et al., 2006). Only few reports exist on EHV-2 and EHV-5 and airway diseases of horses (Allen and Murray, 2004) and most reports concern alphaherpesviruses and relate experimental viral infections with these strains in a lytic phase (Sutton et al., 1998).

EHV-2 is known to be latent in lymphocytes (Drummer et al., 1996) or macrophages (Borchers et al., 2002), however no correlation was found between DNA detection and lymphocyte or macrophage counts in both TW and BAL cytological analysis (data not shown). Detection of EHV-2 in the BAL of the CTL group, without EEC abnormalities

(ECAb or CCPh) could be due to latency states of the virus in the lymphocytes or macrophages of these liquids. Identification of AHV-5 by sequencing the PCR product of the consensus PCR, with only a low percentage of confirmation by virus-specific PCR tests (EHV-5) was also surprising. No history of concomitant grazing or neighbouring between horses and donkeys could be proven retrospectively. We hypothesised that the difference in detection between the consensus PCR and virus-type specific EHV-5 PCR could have been a consequence of the two different PCR methods, a nested PCR assay versus a standard PCR assay for EHV-5. The AHV-5 matching assessed with Genbank could be due to a lack of reported sequences for the EHV-5 virus. The EHV-5 DNA detection was therefore probably underestimated in the present work, mainly in the BAL samples.

Herpesvirus DNA detection was then considered in the light of other potential criteria of viral infection such as clinical suspicion or expression (group), epithelial cell morphological criteria or cytological neutrophilic profiles in the respiratory samples. The interpretation of high exfoliated epithelial cell counts or abnormal morphology in TW or BAL analysis is a recurrent preoccupation of clinicians and searchers and is often considered as a result of mechanical trauma during sampling procedures, leading to a wide range of published epithelial cell percentages between studies (Derksen et al., 1989). More interesting is the analysis of cell morphology during cytological examination, which takes into account the presence and integrity of the *ciliae*, nuclear damage (Whitwell and Greet, 1984; Viel, 2002), or the better defined ciliocytophthoria (Papanicolaou et al., 1961; Freeman et al., 1985).

The disparity between results of the TW and BAL in investigations of respiratory disorders and perhaps in virus DNA detection could be attributed to the intrinsic differences of these two liquids. The *trachea* is a “crossroad” connecting the upper and lower respiratory tracts and the TW is therefore recognised as potentially contaminated and of variable cytological quality compared to BAL, even in a healthy population. For this reason, IAD or RAO are

exclusively based on a BAL cytological diagnosis. This was recently confirmed by Couetil et al.(2007) who suggested that TW should be studied more in depth in terms of the mechanisms inducing tracheal inflammation (i.e; viruses, bacteria, fungi...). In the present study, ciliocytophthoria (CCPh) or epithelial cell abnormalities (ECAb) did not seem to be sensitive nonspecific “herpesvirus markers” in TW, whereas in the BAL theses epithelial alterations were significantly linked to the presence of both EHV-2 and EHV-5 whatever the group considered.

Detection of other viruses such as rhinovirus or influenza virus was not achieved in this study. However these viruses are known to potentially induce mild to severe airway inflammation in horses even a long time after an initial exposure (Willoughby et al., 1992; Sutton et al., 1997). Epithelial cell erosion in the *trachea* (CCPh or ECAb) may also be induced by these viruses explaining the lack of specificity the chosen criteria for positives samples with herpesviruses specifically. The tracheobronchial lymphoid associated tissues represent major latency sites (Allen and Murray, 2004), almost exclusive for herpesviruses and less exposed to other viruses. Epithelial cell damage is therefore more likely to be associated to herpesviruses pathogenicity. During an EHV-2 experimental infection, Borchers et al. (1998) retrieved infectious viruses in nasal or ocular swabs, and viral DNA in the lung and lymphoid tissues after an induced reactivation with dexamethasone (30 days post infection). One could also hypothesize that reactivation can take place after various stressful conditions (intense exercise, competition, transport,...) or corticotherapy and lead to viral shedding in the respiratory tract.

Conversely, in the TW, herpesvirus DNA detection was significantly associated with increased neutrophil counts and group. The causative effect of herpesviruses on inflammation after viral exposure remains to be elucidated, even being experimentally showed with EHV-1 (Smith et al., 2000). One of the EHV-2 G protein coupled receptor (GPCR) is a receptor for

interleukine 8 (IL-8) and the EHV-2 open reading frame E7 codes for IL-10, which is an immunosuppressive cytokine. The chemo-attractant effect of IL-8 for PMNs has previously been described in several clinical or experimental situations (Franchini et al., 2000). EHV-2 can be retrieved in the lung several weeks post infection in mice (Borchers et al., 2002). As EHV-2 replication occurs following the delivery of virus or virus genome by a cell-to-cell contact mechanism from lymphocytes to fully permissive epithelial cells, the detection of viral DNA could be associated with immunomodulation of the host pathway, epithelial cells modifications and neutrophilic recruitment a long time after natural exposure (Drummer et al., 1996). Schlocker et al. (1995) suggested early on that EHV-2 could play a role in equine chronic pulmonary disease. The significant association between CCPh or ECAb and DNA detection could also be explained by such a particular viral physiopathology.

Genomic variations (Browning and Studdert, 1987; Sharp et al., 2007) and concomitant infections with several field strains of EHV-2 have been demonstrated (Bell et al., 2006) and may determine a graduation in the clinical expression of the disease.

Conclusions

The purposes of this study were (1) to precise the DNA prevalence of four EHV-1s in respiratory liquids of horses suffering from various respiratory diseases in comparison with a control group, (2) to determine if particular cytological profiles of the liquids were associated with EHV-1s detection, by investigating, in the different groups, the correlation between EHV-1s detection and neutrophilia or epithelial cells abnormalities and/or ciliocytophthoria. Herpesviruses DNA was more frequently detected in the TW of the PAT group and associated with liquid neutrophilia. Among these viruses, EHV-2 was the most prevalent. No specific epithelial cells profile could be found in association with viral DNA detection in the TW but ciliocytophthoria or epithelial cells abnormalities were interesting features of the BAL significantly associated with the detection of EHV-2 and EHV-5. As respiratory tract

inflammation has already been demonstrated during experimental infection with EHV-1 or influenza virus in the horse, equine herpesviruses and particularly gammaherpesviruses may be implicated in the syndrome of tracheal inflammation.

Even if the precise status of the viral infection (lytic, non replicating, latent) could not be elucidated through the PCR assays strategy used in this study, an experimental infection with EHV-2 in “naïve” horses or induced reactivation should give more information about the potential role of this gammaherpesvirus in the horse. Other tools, such as interferon gamma detection or cytokine profile of the liquids could be useful to better understand or clarify the association with virus DNA detection and cytology of the liquids, particularly the TW.

Acknowledgements

The authors gratefully acknowledge the practitioners who participated to this study. The project received financial support from the Conseil general du Calvados and the French national Studs (Haras Nationaux).

References

- Agius, C.T., Studdert, M.J., 1994. Equine herpesviruses 2 and 5: comparisons with other members of the subfamily gammaherpesvirinae. *Adv.Virus Res.* 44, 357-379.
- Allen, G.P. and Murray, J.T., 2004. Equid herpesvirus 2 (EHV-2) and Equid herpesvirus (EHV-5). In: Coetzer,J.; Thomson,G.; Tustin,R. (Eds.), *Infectious disease of livestock*. Oxford Press, Capetown, UK, pp. 860-868.
- Bell, S.A., Balasuriya, U.B., Gardner, I.A., Barry, P.A., Wilson, W.D., Ferraro, G.L., MacLachlan, N.J., 2006. Temporal detection of equine herpesvirus infections of a cohort of mares and their foals. *Vet.Microbiol.* 116, 249-257.

- 330 Borchers, K., Brackmann, J., Kershaw, O., 2002. The mouse is not permissive for equine
331 herpesvirus 2 (EHV-2), however viral DNA persisted in lung and spleen depending on the
332 inoculation route. *Arch.Virol.* 147, 1437-1444.
- 333 Borchers, K., Frolich, K., 1997. Antibodies against equine herpesviruses in free-ranging
334 mountain zebras from Namibia. *J.Wildl.Dis.* 33, 812-817.
- 335 Borchers, K., Wolfinger, U., Ludwig, H., Thein, P., Baxi, S., Field, H.J., Slater, J.D., 1998.
336 Virological and molecular biological investigations into equine herpes virus type 2 (EHV-2)
337 experimental infections. *Virus Res.* 55, 101-106.
- 338 Browning, G.F., Studdert, M.J., 1987. Genomic heterogeneity of equine betaherpesviruses.
339 *J.Gen.Virol.* 68, 1441-1447.
- 340 Couetil, L.L., Hoffman, A.M., Hodgson, J., Buechner-Maxwell, V., Viel, L., Wood, J.L.,
341 Lavoie, J.P., 2007. Inflammatory airway disease of horses. *J.Vet.Intern.Med.* 21, 356-361.
- 342 Davison, A.J., Eberle, R., Ehlers, B., Hayward, G.S., McGeoch, D.J., Minson, A.C., Pellett,
343 P.E., Roizman, B., Studdert, M.J., Thiry, E., 2009. The order Herpesvirales. *Arch.Virol.* 154,
344 171-177.
- 345 Derksen, F.J., Brown, C.M., Sonea, I., Darien, B.J., Robinson, N.E., 1989. Comparison of
346 transtracheal aspirate and bronchoalveolar lavage cytology in 50 horses with chronic lung
347 disease. *Equine Vet.J.* 21, 23-26.
- 348 Diallo, I.S., Hewitson, G., Wright, L., Rodwell, B.J., Corney, B.G., 2006. Detection of equine
349 herpesvirus type 1 using a real-time polymerase chain reaction. *J.Virol.Methods* 131, 92-98.
- 350 Diallo, I.S., Hewitson, G., Wright, L.L., Kelly, M.A., Rodwell, B.J., Corney, B.G., 2007.
351 Multiplex real-time PCR for the detection and differentiation of equid herpesvirus 1 (EHV-1)
352 and equid herpesvirus 4 (EHV-4). *Vet.Microbiol.* 123, 93-103.
- 353 Drummer, H.E., Reubel, G.H., Studdert, M.J., 1996. Equine gammaherpesvirus 2 (EHV2) is
354 latent in B lymphocytes. *Arch.Virol.* 141, 495-504.
- 355 Dynon, K., Varrasso, A., Ficorilli, N., Holloway, S., Reubel, G., Li, F., Hartley, C., Studdert,
356 M., Drummer, H., 2001. Identification of equine herpesvirus 3 (equine coital exanthema
357 virus), equine gammaherpesviruses 2 and 5, equine adenoviruses 1 and 2, equine arteritis
358 virus and equine rhinitis A virus by polymerase chain reaction. *Aust.Vet.J.* 79, 695-702.
- 359 Fortier, G., Pronost, S., Mischczak, F., Fortier, C., Leon, A., Richard, E., Van Erck, E., Thiry,
360 E., Lekeux, P., 2008. Identification of equid herpesvirus-5 in respiratory liquids: A
361 retrospective study of 785 samples taken in 2006-2007. *Vet J.* *in press*
- 362 Franchini, M., Gill, U., von, F.R., Bracher, V.D., 2000. Interleukin-8 concentration and
363 neutrophil chemotactic activity in bronchoalveolar lavage fluid of horses with chronic
364 obstructive pulmonary disease following exposure to hay. *Am. J. Vet. Res.* 61, 1369-1374.
- 365 Freeman, K.P., Roszel, J.F., Slusher, S.H., 1985. Inclusions in equine cytologic specimens.
366 *J.Am.Vet.Med.Assoc.* 186, 359-364.

- 367 Gerber, V., Straub, R., Marti, E., Hauptman, J., Herholz, C., King, M., Imhof, A., Tahon, L.,
368 Robinson, N.E., 2004. Endoscopic scoring of mucus quantity and quality: observer and horse
369 variance and relationship to inflammation, mucus viscoelasticity and volume. *Equine Vet. J.*
370 36, 576-582.
- 371 Leon, A., Fortier, G., Fortier, C., Freymuth, F., Tapprest, J., Leclercq, R., Pronost, S., 2008.
372 Detection of equine herpesviruses in aborted foetuses by consensus PCR. *Vet.Microbiol.* 126,
373 20-29.
- 374 Malikides, N., Hughes, K.J., Hodgson, J.L., 2007. Comparison of tracheal aspirates before
375 and after high-speed treadmill exercise in racehorses. *Aust.Vet.J.* 85, 414-419.
- 376 McGorum, B.C., Dixon, P.M., 1994. The analysis and interpretation of equine
377 bronchoalveolar lavage fluid (BALF) cytology. *Equine vet.Educ.* 6, 203-209.
- 378 Newton, J.R., Wood, J.L., Chanter, N., 2003. A case control study of factors and infections
379 associated with clinically apparent respiratory disease in UK Thoroughbred racehorses.
380 *Prev.Vet.Med.* 60, 107-132.
- 381 Papanicolaou, G., Bridges, E., Railey, C., 1961. Degeneration of the ciliated cells of the
382 bronchial epithelium (ciliocytophthoria) in its relation to pulmonary disease. *Am*
383 *Rev.Respir.Dis.* 83, 641-659.
- 384 Pickles, K., Pirie, R.S., Rhind, S., Dixon, P.M., McGorum, B.C., 2002. Cytological analysis
385 of equine bronchoalveolar lavage fluid. Part 3: The effect of time, temperature and fixatives.
386 *Equine Vet.J.* 34, 297-301.
- 387 Pusterla, N., Mapes, S., Wilson, W.D., 2008. Use of viral loads in blood and nasopharyngeal
388 secretions for the diagnosis of EHV-1 infection in field cases. *Vet.Rec.* 162, 728-729.
- 389 Robinson, N.E., 2003. Inflammatory airway disease: defining the syndrome. Conclusions of
390 the Havemeyer workshop. *Equine Vet. Educ.* 15, 61-63.
- 391 Schlocker, N., Gerber-Bretscher, R., von Fellenberg, R., 1995. Equine herpesvirus 2 in
392 pulmonary macrophages of horses. *Am.J.Vet.Res.* 56, 749-754.
- 393 Sharp, E.L., Farrell, H.E., Borchers, K., Holmes, E.C., Davis-Poynter, N.J., 2007. Sequence
394 analysis of the equid herpesvirus 2 chemokine receptor homologues E1, ORF74 and E6
395 demonstrates high sequence divergence between field isolates. *J.Gen.Virol.* 88, 2450-2462.
- 396 Smith, K.C., Whitwell, K.E., Mumford, J.A., Hannant, D., Blunden, A.S., Tearle, J.P., 2000.
397 Virulence of the V592 isolate of equid herpesvirus-1 in ponies. *J.Comp Pathol.* 122, 288-297.
- 398 Sutton, G.A., Viel, L., Carman, P.S., Boag, B.L., 1997. Study of the duration and distribution
399 of equine influenza virus subtype 2 (H3N8) antigens in experimentally infected ponies in
400 vivo. *Can.J.Vet.Res.* 61, 113-120.
- 401 Sutton, G.A., Viel, L., Carman, P.S., Boag, B.L., 1998. Pathogenesis and clinical signs of
402 equine herpesvirus-1 in experimentally infected ponies in vivo. *Can.J.Vet.Res.* 62, 49-55.
- 403 VanDevanter, D.R., Warrenner, P., Bennett, L., Schultz, E.R., Coulter, S., Garber, R.L., Rose,
404 T.M., 1996. Detection and analysis of diverse herpesviral species by consensus primer PCR.
405 *J.Clin.Microbiol.* 34, 1666-1671.

- Viel, L., 2002. Significance of bronchoalveolar cytology in inflammation airway disease of horse. In: Hoffmann A., Robinson N. E. Wade J. F. Eds. (Eds), Inflammation airway disease. Havemeyer Foundation Monograph series No 9, Boston, USA, 52-55.
- Whitwell, K.E., Greet, T.R., 1984. Collection and evaluation of tracheobronchial washes in the horse. *Equine Vet.J.* 16, 499-508.
- Williams, K.J., Maes, R., del, P.F., Lim, A., Wise, A., Bolin, D.C., Caswell, J., Jackson, C., Robinson, N.E., Derksen, F., Scott, M.A., Uhal, B.D., Li, X., Youssef, S.A., Bolin, S.R., 2007. Equine multinodular pulmonary fibrosis: a newly recognized herpesvirus-associated fibrotic lung disease. *Vet.Pathol.* 44, 849-862.
- Willoughby, R., Ecker, G., McKee, S., Riddolls, L., Vernailen, C., Dubovi, E., Lein, D., Mahony, J.B., Chernesky, M., Nagy, E., 1992. The effects of equine rhinovirus, influenza virus and herpesvirus infection on tracheal clearance rate in horses. *Can.J.Vet.Res.* 56, 115-121.
- Wood, J.L., Newton, J.R., Chanter, N., Mumford, J.A., 2005. Association between respiratory disease and bacterial and viral infections in British racehorses. *J.Clin.Microbiol.* 43, 120-126.

Table 1

Description of the studied population

	Control group (CTL)		Pathological group (PAT)	
	Tracheal washes (TW)	Bronchoalveolar lavages (BAL)	Tracheal washes (TW)	Bronchoalveolar lavages (BAL)
Number of samples	37	25	259	387
Age (years) (mean $\pm$ s.d.)	5.6 $\pm$ 2.9	6.3 $\pm$ 3.05	5.8 $\pm$ 2.2	5.4 $\pm$ 3.4
PMNs count (%) (mean $\pm$ s.d.)	10.4 $\pm$ 6.9	5.7 $\pm$ 2.9	57.1 $\pm$ 34.7*	32.6 $\pm$ 29.1*

PMN: Polymorphonuclear neutrophils

* Statistically different from the corresponding CTL sample ($P < 0.0001$)

Table 2

Characteristics of the PCR assays used for the detection of equine herpesviruses.

PCR assay	Target gene	Primer and probe sequences (5'-3')	amplification program			Product size (bp)	PCR type	References
Consensual herpes	DNA pol	DFA: + GAYTTYGCNAGYYTNTAYCC	94°C	5min	x45	215-315	Classical Nested	Van Devanter et al. (1996)
		ILK: + TCCTGGACAAGCAGCARNYSGCNMTNAA	94°C	30sec				
		KG1: - GTCTTGCTCACCAGNTCNACNCCYTT	46°C	1min				
		TGV: + TGTAACTCGGTGTAYGGNTTYACNGGNGT*	72°C	1min				
		IYG: - CACAGAGTCCGTRTCNCRTADAT*	72°C	10min				
EHV1	gB	EHV1F1: + CATGTCAACGCACTCCCA	95°C	10min	x50	63	Real-time	Diallo et al.(2006)
		EHV1R1: - GGGTCGGGCGTTTCTGT	95°C	15sec				
		EHV1MGBPROBE: 6 FAM-CCCTACGCTGCTCC-MGB-NFQ	64°C	1min				
EHV2	« human interleukin 10 homologous sequence »	EHV2pS: + AACTCCCATCCCCAGAAGAC EHV2pAS: - TGTTTCCTCGCTAACCAACC	95°C	15min	x45	245	Classical One step	Leon (personal communication 2008)
			94°C	30sec				
			60°C	1min30				
			72°C	1min30				
			72°C	10min				
EHV4	gB	EHV4F1: + GGGCTATTGGATTACAGCGAGAT EHV4R1: - TAGAATCGGAGGGCGTGAAG EHV4MGBPROBE: VIC-CAGCGCCGTAACCAG-MGB-NFQ	95°C	10min	x50	59	Real-time	Diallo et al. (2007)
			95°C	15sec				
			64°C	1min				
EHV5	gB	EHV5-gB-S: + ATGAACCTGACAGATGTGCC EHV5-gB-AS: - CACGTTCACTATCACGTCGC	94°C	5min	x50	293	Classical One step	Dynon et al.(2001)
			94°C	30sec				
			60°C	45sec				
			72°C	1min30				
			72°C	8min				

+: forward primer, - : reverse primer

* second round primers of nested PCR

Table 3

PCR detections of herpesviruses in the tracheal washes (TW) and bronchoalveolar lavages (BAL) of the control (CTL) and pathological (PAT) groups.

	TW			BAL		
	CTL (n=37)	PAT (n=259)	<i>P</i> value ^a	CTL (n=25)	PAT (n=387)	<i>P</i> value ^a
PCR target	+	+		+	+	
EHV-1	0 [0%]	4 [2%] ^b	> 0.05	0 [0%]	4 [1%] ^c	>0.05
EHV-2	4 [11%]	85 [33%]	0.006 *	9 [36%]	82 [21%]	> 0.05
EHV-4	1 [3%]	17 [7%]	> 0.05	1 [4%]	16 [4%]	> 0.05
EHV-5	0 [0%]	8 [3%]	> 0.05	0 [0%]	16 [4%]	>0.05
EHV consensus PCR	7 [19%]	112 ^d [43%]	0.004 *	10 [40%]	133 ^e [34%]	>0.05

^a Fisher's exact test

^b 11 tests were not performed on this run for EHV-1 due to insufficient quantity of biological material.

^c 2 tests were not performed on this run for EHV-1 due to insufficient quantity of biological material.

* Statistically significant result

^d 8 samples were sequenced as Asinine herpesvirus 5 in the PAT group.

^e 16 samples were sequenced as Asinine herpesvirus 5 in the PAT group.

Table 4

Association between ciliocytophthoria or epithelial cells abnormalities with the status of horses and sample type (TW or BAL)

		CTL		PAT		<i>P</i> value ^a
		+	-	+	-	
BAL	ECAb	0/25 [0%]	25/25 [100%]	15/387 [4%]	372/387 [96%]	0.6
	CCPh	0/25 [0%]	25/25 [100%]	31/387 [8%]	356/387 [92%]	0.2
TW	ECAb	9/37 [24%]	28/37 [76%]	61/259 [23%]	198/259 [76%]	1
	CCPh	7/37 [19%]	30/37 [81%]	72/259 [27%]	187/259 [73%]	0.3

CTL = control group; PAT = pathological group; CCPh = ciliocytophthoria; ECAb = exfoliated epithelial cells abnormalities

^a Fisher's exact test

Note: a combination of the two criteria (CCPh and/or ECAb) is not significant in TW or BAL (0.24 and 0.70 respectively)

Table 5

Herpesvirus DNA detection by ciliocytophthoria or epithelial cells abnormalities in the BAL and TW

<i>P</i> values ^a	EHVs		EHV-2		EHV-5		EHV-4	
	TW	BAL	TW	BAL	TW	BAL	TW	BAL
Ciliocytophthoria	0.28	0.04*	0.39	0.17	0.31	0.02*	0.41	1
Epithelial cells abnormalities	0.2	0.01*	0.5	0.007*	0.16	0.45	1	0.47
CCPh or ECAb	0.30	0.06	0.33	0.13	0.52	0.034*	0.59	1

* Significant results ($P < 0.05$)^a Fisher's exact test

Table 6

Association between herpesvirus DNA detection and polymorphonuclear neutrophils cells (PMN) percentage in the TW for the control group (CTL) and the pathological group (PAT).

Herpesvirus		EHVs		EHV-2		EHV-4	
	PMN (%)	+	-	+	-	+	-
CTL	0-20	18.92	81.08	10.81	89.19	2.86	97.14
PAT	0-20	37.74	62.26	30.19	69.81	3.77	96.23
PAT	21-40	40.91	59.09	25.00	75.00	4.55	95.45
PAT	41-60	45.45	54.55	40.91	59.09	0.00	100
PAT	61-80	40.54	59.46	35.14	64.86	8.11	91.89
PAT	81-100	47.57	52.43	34.95	65.05	9.71	90.29
Cochran –Armitage trend test / two-sided $P > Z $		0.009		0.0165		0.0542	

Results for EHV-1 and EHV-5 were not significant