17β -Estradiol enhances α integrin subunit gene expression through ER α –Sp1 interaction and reduces cell motility and invasion of ER α -positive breast cancer cells

Diego Sisci, Emilia Middea, Catia Morelli, Marilena Lanzino, Saveria Aquila, Pietro Rizza, Stefania Catalano, Ivan Casaburi, Marcello Maggiolini, Sebastiano Andò

► To cite this version:

Diego Sisci, Emilia Middea, Catia Morelli, Marilena Lanzino, Saveria Aquila, et al.. 17β -Estradiol enhances α integrin subunit gene expression through ER α –Sp1 interaction and reduces cell motility and invasion of ER α -positive breast cancer cells. Breast Cancer Research and Treatment, 2010, 124 (1), pp.63-77. 10.1007/s10549-009-0713-6 . hal-00524848

HAL Id: hal-00524848

<https://hal.science/hal-00524848>

Submitted on 9 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

17 β -estradiol enhances α_5 integrin subunit gene expression through ER α -Sp1 interaction and reduces cell motility and invasion of ER α positive breast cancer cells.

Diego Sisci, Emilia Middea, Catia Morelli, Marilena Lanzino, Saveria Aquila, Pietro Rizza¹,

Stefania Catalano, Ivan Casaburi, Marcello Maggiolini, and Sebastiano Andò^{1*}

Department of Pharmaco-Biology - University of Calabria, Arcavacata di Rende, Italy

¹Department of Cell Biology - University of Calabria, Arcavacata di Rende, Italy

Running head: Estradiol enhances $\alpha_5\beta_1$ integrin expression.

***Corresponding author Mailing address:** Dipartimento di Biologia Cellulare, Università della Calabria, Arcavacata di Rende (Cosenza) 87030 ITALY. Phone: +39 0984 496201; FAX: +39 0984 496203; E-mail: sebastiano.ando@unical.it

Keywords: Cell adhesion, $\alpha_5\beta_1$ integrin, ER α , and Sp1.

Words count: Introduction, Results and Discussion, 2948

Materials and Methods, 2530

1 **Abstract**

2 In breast tumors the expression of estrogen receptor alpha ($ER\alpha$) is known to be associated with
3 a more favorable prognosis. $ER\alpha$ expression has been reported to reduce the metastatic potential of
4 breast cancer cells. Recently, we have observed that extracellular matrix proteins activate $ER\alpha$ and that
5 both liganded and unliganded receptor modulates cell invasiveness acting at nuclear level. To explain
6 the mechanisms by which $ER\alpha$ regulates cell adhesion, we have evaluated the expression of $\alpha_5\beta_1$
7 integrin, prevalently expressed in stationary cells, in response to 17β -estradiol (E2). Here we show that
8 E2/ $ER\alpha$ increases the expression of integrin $\alpha_5\beta_1$ through Sp1 mediated binding to a GC rich region
9 located upstream of an ERE half-site in the 5' flanking region of the α_5 gene forming a ternary $ER\alpha$ -
10 Sp1-DNA complex. Estrogen responsiveness of the α_5 gene promoter, as observed in HeLa cells,
11 underlies a general mechanism of regulation which is not strictly linked to the cell type.

12 Our data reveal novel insight into the molecular mechanisms sustaining the reduced
13 invasiveness of $ER\alpha$ expressing cells demonstrating that $\alpha_5\beta_1$ integrin expression is related to the
14 maintenance of the stationary status of the cells, counteracting E2/ $ER\alpha$ capability to enhance breast
15 cancer cell migration and invasion.

16 .

17

18

19

20

21

22

23

1 **Introduction**

2 In the mammary gland estrogen receptor alpha ($ER\alpha$) functions as a hormone-dependent transcription
3 factor regulating the expression of proto-oncogenes which control cell growth and differentiation [1-5].
4 In breast cancer, however, the stimulatory effect of 17β -estradiol (E_2)/ $ER\alpha$ signaling on tumor cell
5 growth strongly correlates with a lower invasiveness and reduced metastatic potential [6-8]. For
6 instance, previous studies and our recent findings demonstrate how the exposure of breast cancer cells
7 to extracellular matrix (ECM) proteins increases cell proliferation [9], enhances functional
8 transactivation of $ER\alpha$ and reduces cell invasiveness in $ER\alpha$ positive breast cancer cells [10].

9 On the other hand, it is well documented how E_2 / $ER\alpha$ up-regulates genes which reduce cell
10 invasiveness like α_1 antichymotrypsin [11] and down-regulates genes involved in the enzymatic
11 remodeling and degradation of ECM proteins like metalloproteinase-9 [12] and Collagenase type IV
12 [13].

13 Cell adhesion to ECM is mainly mediated by integrins which mostly control the physical strength of
14 cell substratum attachment [14]. Several studies have demonstrated that variations in cell surface
15 expression levels of integrin [15], or in adsorbed concentration of its substratum, or its ligand [16], or
16 in the integrin-ligand affinity [17], could affect the adhesion strength and the speed of cell movement
17 [14].

18 A role for E_2 / $ER\alpha$ in integrin expression derives from physiological circumstances consistent with
19 evidence indicating that $\alpha_5\beta_1$ levels double in breast tissues between puberty and sexual maturity and
20 drop dramatically after ovariectomy [9]. So, E_2 seems to exert a specific modulatory role on integrin
21 expression which mostly regulates the adhesion of tumor cells to substrates.

22 Fibronectin (Fn) mediated attachment and signaling in cells occurs in response to Fn binding to cell
23 surface integrins. Fn has been demonstrated to bind to different Fn specific integrins including
24 $\alpha_4\beta_1$, $\alpha_5\beta_1$, $\alpha_v\beta_1$, $\alpha_v\beta_3$, and $\alpha_v\beta_6$ [18]. However previous studies have shown that the α_4 subunit is not

present in mammary cells [19]. Furthermore immunohistochemical analyses revealed little or no α_v protein in mammary epithelia or in myoepithelial cells [20] at any stage of development examined. Other authors have reported that the α_5 is the key receptor subunit for Fn and that α_5 knockout mice most closely resemble Fn knockout mice [18, 21].

On the basis of these findings we reasonably focused our attention on the effects of E2 on the α_5 integrin subunit being this subunit fully representative of the $\alpha_5\beta_1$ integrin expression.

In the present study we elucidated the molecular mechanism by which E2/ER α signaling up-regulates the expression of the α_5 integrin subunit either in MCF-7 or HeLa cells ectopically expressing ER α . This event appears to occur through the formation of a transcriptionally-active ER α –Sp1 complex that binds to the GC-rich (Sp1) (-420 to -370) region located up-stream of ATG and very likely indicates a general modulatory mechanism induced by E2 which is not tightly linked to the cell type investigated.

Materials and Methods

Cell lines and cell culture conditions

Two ER α positive (ER+) human breast cancer epithelial cell lines, MCF-7 and T47D and an ER α negative (ER–) *cervicae* carcinoma cell line HeLa were used. MCF-7 and T47D cells were maintained in monolayer culture in Dulbecco's modified Eagle's/Ham's F-12 medium (1:1) (DMEM/F12; Gibco, USA) supplemented with 5 % calf serum (CS; Eurobio, F), 100 UI/ml penicillin (Eurobio, F), 100 ug/ml streptomycin (Eurobio, F) and 0.2 mM L-glutamine (Eurobio, F). HeLa cells were maintained in monolayer cultures in Minimal Essential Medium with Earle's salts (MEM; Gibco, USA) supplemented with 10% fetal bovine serum (Eurobio, F), 100 UI/ml penicillin, 100 ug/ml streptomycin and 0.2 mM L-glutamine. Cells were passaged weekly using trypsin-EDTA (Eurobio, F) and media were changed every two days. Both cell lines were cultured at 37°C in 5% CO₂ in a humidified atmosphere.

1 In all experiments, steroids and growth factors were deprived from cells growing them in
2 serum- and phenol red-free DMEM/F12 (SFM; Gibco, USA) containing 0.5 % BSA and 2 mM L-
3 glutamine for 24 hrs.

4 ***Fluorescence-Activated Cell Sorting.***

5 Serum starved MCF-7 cells were treated with 10 nM E2 for 4, 12, and 24 h or for 24h with
6 increasing concentration of E2 (1, 10, 100 nM). Cells were detached with EDTA solution (Versene),
7 washed and incubated with anti-integrin α_5 antibody (H-104, Santa Cruz Biotechnology, USA) or anti-
8 integrin α_4 antibody (7.2, Santa Cruz Biotechnology, USA) for 1 h at 22°C. Cells were then washed
9 twice in PBS and incubated with a fluorescein isothiocyanate (FITC)-conjugated secondary antibody
10 (Santa Cruz Biotechnology, USA) for 1 h at 37°C. After two washes in PBS, cells were fixed with 70%
11 ethanol and subjected to FACS analysis. At least 1×10^4 cells were measured in a FACS flow cytometer
12 (Becton Dickinson, USA). MCF-10 cells were used as control.

13 ***Western blotting***

14 To evaluate the influence of E2 on α_5 protein expression, MCF-7 and T47D cells, serum-
15 starved for 24 h, were incubated with 10 nM E2 (Sigma Aldrich, USA) for 16, 24, and 48 h. In some
16 experiments the cells were treated with 1 μ M of the E2 antagonist ICI 182,780 (ICI; Zeneca, UK). At
17 the end of each incubation time, cells were washed with ice-cold PBS and lysed for 1 min at 4 °C in
18 Triton lysis buffer (50 mM Hepes pH 7.5, 150 mM NaCl, 1.5 mM $MgCl_2$, 1mM $CaCl_2$, 1 % Triton X-
19 100, 0.2 mM Na_3VO_4 , 1 % PMSF, 1 % Aprotinin). Cell lysates were cleared by centrifugation (14000
20 rpm for 10 min at 4 °C) and the protein content was determined using Bradford Dye Reagent (Bio Rad,
21 USA) according to manufacturer's instructions. Cellular lysates (10 to 30 μ g of protein/lane) were
22 resolved by SDS-PAGE, transferred to nitrocellulose membranes and probed with either anti- α_5
23 integrin polyclonal antibody (H-104, Santa Cruz Biotechnology, USA) or anti-GAPDH polyclonal
24 antibody (FL-335, Santa Cruz Biotechnology, USA). The antigen-antibody complexes were detected

by incubation of the membranes with peroxidase-coupled anti-rabbit IgG and developed using the ECL Plus Western Blotting detection system (Amersham Pharmacia Biotech, UK).

RT-PCR

MCF-7 and T47D cells were incubated for 24 h in SFM and then treated with E2 10 nM for 24 h and/or 1 μ M ICI 182,780 (ICI; Zeneca, UK). At the end of the treatment total cellular RNA was extracted using RNeasy (Qiagen, USA) and reverse transcribed using Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase (Promega, USA). Briefly, reverse transcription (RT) was performed using 1 μ g of total RNA in a final volume of 10 μ l by incubation at 37 °C for 30 min with 200 U of M-MLV reverse transcriptase, 0.4 μ g oligo dT, 0.5 μ M deoxy-nucleotidetriphosphate (dNTP) and 24U RNAsin (Promega, USA), followed by heat denaturation for 5 min at 95 °C. Subsequent PCR analysis was performed on 1 μ l of the RT product in a final volume of 50 μ l. The following pairs of primers were used to amplify the 188 bp of α_5 integrin: 5'-CCCCGCTCGTGCCGCTGCTGTT-3' (α_5 forward) and 5'-CCCCCACCACC-ATCCCGTCTCC-3' (α_5 reverse). The PCR amplification was performed using 1.25U GoTaq DNA polymerase (Promega, usa), 1X PCR buffer (10mM Tris-HCl, 50mM KCl), 2.5mM MgCl₂, 0.2mM of each dNTP, and 1.5 pM of primers for α_5 or 36B4. PCR conditions were 30 seconds (sec) at 94 °C, 40 sec at 59 °C, and 60 sec at 72 °C for 30 cycles. PCR products (10 μ l) were separated on a 1.0 % agarose gel. Amplification of the 408 bp of ribosomal RNA 36B4 was performed as control using the following primers: 5'-CTCAACATCTCCCCCTTCTC-3' (36B4 forward) and 5'-CAAATCCCA-TATCCTCGTCC-3' (36B4 reverse).

Plasmid constructs

To create the Luciferase expression vector pGL3-p α_5 encoding the 5'-flanking region of the α_5 gene [22], genomic DNA was extracted from MCF-7 by TRIzol reagent following the manufacturer's instructions and amplified by PCR using the following primer pairs 5'-ACTCCTGGGTT-

1 TTCAATACTG-3' (forward) and 5'-CCCGCTCTTCCCTGTCC-3' (reverse). PCR amplification was
2 performed using the same reagents reported above. PCR conditions were 5 min at 95 °C followed by 1
3 min at 94 °C, 1 min at 58 °C, and 1 min at 72 °C for 30 cycles. The obtained fragment was inserted in
4 the pCR 2.1 plasmid using the TA cloning kit (Invitrogen, USA), sequenced, and cut with Hind III and
5 Xho I. The insert was then cloned into the pGL3 basic vector (Promega, USA). The deletion mutant in
6 positions -424 to -370 of the α_5 gene 5'-flanking sequence corresponding to a Sp1/ERE half-site motif
7 (Figure 4A) was created using PCR (pGL3-p α_5 - Δ Sp1/ERE). The oligonucleotides used as PCR
8 primers were as follows: 5'-ACTCCTGGGTTTCAATACTG-3' (forward), 5'-CCTTAAGAG-
9 GAAGCCTGTATATGTGTG-3' (reverse) and 5'-CGAATTCGATTTCCTTCCTCATTAGGA-3'
10 (forward), 5'-CCCGCTCTTCCC-TGTCC-3' (reverse). The PCR products were digested with EcoRI
11 (underlined in primers), ligated and inserted in the pCR 2.1 plasmid using the TA cloning kit. The
12 obtained plasmid was digested with Hind III and Xho I. The insert was then cloned into the pGL3 basic
13 vector. The pGL3-Sp1/ERE plasmid was obtained using two complementary oligonucleotides: 5'-
14 ATATGAGCTCCATATACAGGCTTCCTCCGCCACCAGAGGTGATTTCCTTCCTCATTAGG-
15 AAGCTTTATAT-3' and 5'-ATATAAAGCTTCCTAATGAGGAAAGGAATCACCTCTGGTGGGC-
16 GGAGGAAGCCTGTATATGGAGCT-CATAT-3'. Self annealing was obtained by boiling and
17 cooling down 10 μ g of each oligonucleotide for 5 min in water. The insert was digested with Sac I and
18 Hind III (underlined in primers) and successively cloned in pGL3 basic vector.

19 Site mutation of the Sp1 site (-404 to -399) up-stream to the half-ERE binding motif was generated
20 using PCR. The oligonucleotides used as PCR primers were as follows: 5'-AAGGTA-
21 CCACTCCTGGGTTT TCAATACTG-3' (forward), 5'-CTCTGGTGGATGGAGGAAGCCTGTAT-3'
22 (reverse) and 5'-ATACAGGCTTCCTCCatCCACCAGAG-3' (forward), 5'-AACTCGAGCCCG-
23 CTCTTCCCTGTCC-3' (reverse). PCR products were inserted into the pCR 2.1 plasmid using the TA

cloning kit. The obtained plasmid was digested with Kpn I and Xho I (underlined in primers). The insert was then cloned into the pGL3 basic vector pGL3-p α 5 (-Sp1).

Plasmids, transfections and ERE-luciferase assay

In addition to the plasmids described above, the following plasmids were used: the SV40 promoter-based pSG5 vector encoding ER α (wild-type) pSG5-HeG0 (HeG0, Tora et al, 1989); pCMV-hE241G, an ER α construct deleted in the nuclear localization signal region (NLS) (aa. 250-303) and inserted in the pCMV expression plasmid (E241G; a gift from Song R) and the *Renilla reniformis* luciferase expression vector pRL-Tk (Promega, USA).

To monitor the transcriptional activation of α ₅ by E₂, MCF-7 and HeLa cells (5x10⁴ density) were plated on a 24-well plates, grown in DMEM/F12 to an approximate confluence of 70-80 % and then co-transfected with either pGL3-p α 5 (encoding for luciferase under the control of the human α ₅ promoter) and pRL-Tk (MCF-7 and HeLa) or with pGL3-p α 5, HeG0, and pRL-Tk, or pGL3-p α 5, E241G, and pRL-Tk (HeLa). Specific deletion and mutations of the pGL3-p α 5 plasmid were generated as described above. The effects of each site mutation/deletion were examined in transfection assays in MCF-7 cells. Transfections were carried out using Fugene 6 (Roche, USA). Cells were transfected in growing medium and 6 h after transfection were washed twice with SFM and shifted to SFM for 24 h and then treated for an additional 24 h with 10 nM E₂ and/or 1 μ M ICI 182,780 (ICI; Zeneca, UK) and/or 100nM mithramycin A (Sigma, USA). Where indicated, cells were treated with mithramycin A for 1 h before stimulation with E₂.

Firefly luciferase and *Renilla reniformis* luciferase activities were determined using the Dual-Luciferase Reporter Assay System (Promega, USA) according to the manufacturer's instructions. Firefly luciferase activity was normalized to *Renilla reniformis* luciferase activity and expressed as relative luciferase units.

To evaluate the effects of ER α activity on α_5 integrin expression HeLa cells were transfected, as above with increasing concentrations of pSG5-HeG0 (1 and 2 μ g), as control HeLa cells were transfected with an empty pSG5 vector. Following transfection, cells were serum starved for 24 h and then treated for 24 h with 10 nM E2.

Knock-down of $\alpha_5\beta_1$ integrin gene expression

siRNA was used for effective depletion of $\alpha_5\beta_1$ integrin gene (Santa Cruz Biotech, USA). A scrambled siRNA (Invitrogen, USA) that lacks identity with known gene targets was used as control for non sequence-specific effects. MCF-7 and T47D cells were trypsinized, transfected in suspension with Lipofectamine 2000 (Invitrogen, USA), as suggested by the manufacturer, and then plated in six-well plates (3.5×10^5 cells per well). Briefly, cells were transfected with 100 pmol siRNAs in 3 ml total growing medium. siRNAs were diluted in 500 μ l of SFM without antibiotic; after 5 min lipofectamine 2000 was added to the mix and incubated at room temperature for 20 min. siRNA-Lipofectamine 2000 complex was then added to the cells and incubated at 37°C for 6 h, switched to fresh growing medium and further incubated for additional 18 h. Then cells were switched to SFM for other 24 h and treated or not with 10 nM E2 for 24 h before motility analysis. Forty eight hours of incubation with the siRNAs was chosen based on a time course performed to evaluate the time needed to obtain a knock down of at least 60% of α_5 integrin content.

Chromatin immunoprecipitation

MCF-7 cells were grown in 10 cm plates. Confluent cultures (90%) were shifted to SFM for 24 h and then treated with 10 nM E2 or left untreated in PRF-CT for 2 h. Following treatment, ChIP methodology was performed as described previously (Morelli et al., 2004). The precleared chromatin was precipitated with anti-ER α mAb (Santa Cruz, USA) for ER α , anti-Sp1 pAb (Upstate, USA) for Sp1 and anti-Polymerase II pAb (Santa Cruz, USA) for Pol II. A 4 μ l volume of each sample was used as template for PCR with specific primers. The following pairs of primers were used to amplify 312 bp

of the $\alpha 5$ promoter region containing the Sp1/half-ERE 5'-ACCCCAAAGTCTCCTCCCTCCCG-3' (forward); 5'-CCCGCGTCCCCCACCCCTAAG-3' (reverse). The PCR conditions were: 1 min at 94 °C, 45 sec at 59 °C, and 45 sec at 72 °C. The amplification products obtained in 28 cycles were analyzed in a 2 % agarose gel and visualized by ethidium bromide staining.

Gel Mobility Shift Assay

Nuclear protein extracts were prepared and collected as reported in DAPA. The probes were generated by annealing single-stranded oligonucleotides (reported below) labeled with [γ - 32 P]ATP and T4 polynucleotide kinase and then purified using Sephadex G50 spin columns. The oligonucleotides used as probes and as cold competitors (synthesized by Sigma Genosys, UK) are listed below. The protein binding reactions were carried out in 20 μ l of buffer (20 mM HEPES, pH 8, 1 mM EDTA, 50 mM KCl, 10 mM dithiothreitol, 10 % glycerol, 1 mg/ml BSA, 50 μ g/ml poly (dI-dC) with 50000 cpm of labeled probe, 20 μ g of MCF-7 nuclear protein or 15 ng of Sp1 and 0.18 pmol of purified ER α recombinant proteins (both purchased from Invitrogen, USA), and 5 μ g of poly (dI-dC)). The above-mentioned mixture was incubated at room temperature for 20 min in the presence or absence of unlabeled competitor oligonucleotide or specific anti-ER α (anti ER α F-10) or Sp1 (anti Sp1 1C6) antibodies, both purchased from Santa Cruz Biotechnology, USA. The entire reaction mixture was electrophoresed through a 6 % polyacrylamide gel for 3 h at 150 V. Gel was dried and subjected to autoradiography at -70 °C.

Oligonucleotides used (nucleotide motifs of interest are underlined): Probe - 5'-TTCCTCCGCCC-ACCAGAGGTGATTCC-3'; Mutated probe - 5'-TTCCTCCGCCCACCAGAaaTtATTCC-3'.

DNA affinity precipitation assay

DNA affinity precipitation assay was performed as described by Suzuki T. et al [23] with minor modifications. Nuclear extracts were prepared from MCF-7 cells. Sub-confluent cultures (70%) were shifted to SFM for 24 h and then treated with 10 nM E2 for 24 h, or left untreated in SFM. Following

treatment, cells were scraped in cold PBS, and resuspended in 400 μ l of cold buffer A (10 mM HEPES-KOH, pH 7.9, at 4 °C, 1.5 mM $MgCl_2$, 10 mM KCl, 0.5 mM dithiothreitol, 0.2 mM PMSF, 1 mM leupeptin). The cells were allowed to swell on ice for 10 min, centrifuged, and the pellet was resuspended in 50 μ l of cold Buffer B (20 mM HEPES-KOH, pH 7.9, 25% glycerol, 1.5 mM $MgCl_2$, 420 mM NaCl, 0.2 mM EDTA, 0.5 mM dithiothreitol, 0.2 mM PMSF, 1 mM leupeptin) and incubated on ice for 20 min for high salt extraction. Cellular debris was removed by centrifugation for 2 min at 4 °C, and the supernatant fraction (containing DNA-binding proteins) was stored at -70 °C. The yield was determined by Bradford method.

The 26 bp probe was prepared by annealing the biotin-treated sense oligonucleotide (5'-TTCTCCGCCACCAGAGGTGATTCC-3') and non-biotinated anti-sense oligonucleotide. A 3 pmol sample of the probe was added to 100 μ g of the protein nuclear lysates and placed on ice for 40 min. Then 20 μ l of streptavidin-coated beads (Promega) were added and the sample was incubated for a further 3 h at 4 °C with rotation. After separation of the beads from the supernatant by centrifugation, the beads were washed five times with binding buffer, suspended with 40 μ l of sample buffer, boiled and used for Western blotting as a DNA-bound fraction. The supernatant was mixed with an equal volume of 2 $\times$ sample buffer, boiled and used for Western blotting as an unbound fraction. For competition, a 10-fold excess of unlabelled probe was added to the nuclear lysate and incubated on ice for 30 min before addition of the labeled probe.

Motility assay

To evaluate the role of $\alpha 5 \beta 1$ integrin on cell motility we performed a knock down of $\alpha 5 \beta 1$ integrin, as reported in *plasmids and transfections* section, in both MCF-7 and T47D cell lines, before motility assays. After transfection, cells were maintained in SFM for 24 h and then treated with E2 for additional 24h. At the end, cells were dispersed with versene (Eurobio, F), washed twice, re-suspended in SFM, counted using a haemocytometer and plated on 24-well modified Boyden chambers containing

1 porous (8 μ m) polycarbonate membranes (Costar, USA). For cell invasion experiments the chambers
2 were coated on the internal surface, with 30 μ g/ml Fibronectin (Fn) by incubation at room temperature.
3 The lower chambers were loaded with 500 μ l of either SFM for migration assays or SFM containing
4 5% CS for invasion assays, while synchronized cells (1×10^5) suspended in 200 μ l of SFM with or
5 without 10 nM E2 were plated into the upper chambers. After 8 hrs of incubation in 5% CO₂ at 37° C,
6 the cells in the upper chamber were removed by a cotton swab, so that only cells that had migrated
7 through the membrane remained. The membranes were then fixed and stained in Coomassie blue
8 solution (0.25g Coomassie blue, 45ml water, 45ml methanol, 10ml glacial acetic acid) for 5 min, then
9 each well was rinsed three times with distilled water. The migrated cells were counted using an
10 inverted microscope.

11 *Statistical analysis*

12 All data were expressed as the mean $\pm$ SD (standard deviation) of at least three independent experiments.
13 Statistical significances were tested using Student's t test.

14

15 **RESULTS**

16 *α_5 integrin subunit expression in MCF-7 cells upon E2 exposure.*

17 The expression of the α_5 integrin subunit in response to E2 stimulation was evaluated in MCF-7
18 cells. We focused our attention on the α_5 integrin subunit, which is strongly indicative of integrin $\alpha_5\beta_1$
19 expression, since only the β_1 subunit protein has been demonstrated to form a heterodimer with the α_5
20 subunit, so the detection of the α_5 subunit has been reported to mean that the $\alpha_5\beta_1$ integrin was also
21 detected herein.

22 Serum-starved MCF-7 cells were treated with 10 nM E2 for 4, 12 and 24 hours (h) before
23 immunodetection of the α_5 integrin subunit by FACS analysis (see *Materials and Methods*). As shown

1 in figure 1A, the exposure to E2 induced an increase in α_5 integrin subunit expression that became
2 significant starting from 12 h of treatment. The significant up-regulation of α_5 integrin subunit levels
3 was also observed in serum starved MCF-7 cells treated for 24 h with increasing concentrations of E2
4 (1, 10, 100 nM) (Figure 1B). The increment of α_5 integrin subunit expression in response to E2 was
5 also observed in T47D cells (data not shown).

6 To further assess the specificity of E2-dependent α_5 integrin subunit expression, we evaluated
7 whether E2 was also able to regulate α_4 integrin subunit levels, since its expression has been reported
8 to be associated with a reduction of breast cancer cell invasiveness [24]. E2 administration induced a
9 dose-dependent increase of α_5 expression, but did not affect α_4 integrin subunit levels (Figure 1C,i).

10 ***E2 regulates the expression of α_5 integrin subunit***

11 To further investigate the mechanistic basis for the estrogen-dependent regulation of α_5 integrin
12 subunit expression, we carried out a time course analysis. MCF-7 cells were serum starved for 24 h and
13 then treated with 10 nM E2 for 16, 24 and 48 h. The α_5 integrin subunit expression was detected on
14 protein lysates. As shown in Figure 2A a marked decrease of α_5 integrin subunit was observed
15 following a prolonged starvation (48 h), while E2 stimulation induced a significant increase of its
16 expression. In addition, E2 induced a dose-dependent increase of α_5 integrin subunit expression (Figure
17 2C). The up-regulation observed was mediated by ER activation since the addition of ICI 182,780
18 abrogated it (Figure 2C).

19 The observed up-regulation was directly correlated to a significant increase in α_5 integrin
20 mRNA transcription (Figure 2B and D).

21 To demonstrate that the increased expression of the α_5 integrin subunit was specifically mediated by
22 ER α , we transfected HeLa cells with increased amounts of HeG0, treated with E2 (10 nM) for 24 h,
23 and protein lysates were used to detect the α_5 integrin subunit. E2 stimulation increased α_5 integrin

expression in ER α content related manner (Figure 2C), supporting the conclusion that E2 regulates the expression of $\alpha_5\beta_1$ integrin in ER α -positive cells. The increased expression of α_5 integrin subunit lead us to evaluate the role of E2 activated ER α on the regulation of α_5 integrin subunit expression.

The 5'-flanking region of α_5 gene is regulated by ER α

To investigate the mechanisms by which E2 regulates α_5 gene expression, we prepared a pGL3-p α_5 plasmid containing a construct bearing the 5'-flanking region of the α_5 gene [22] obtained by PCR, as described in *Materials and Methods*, conjugated with a luciferase reporter gene. This region has been previously shown to be active as a promoter in the control of α_5 integrin gene expression [22].

MCF-7 cells were transfected with the pGL3-p α_5 reporter plasmid and exposed to 10nM E2 and/or 1 μ M ICI for 24h before luciferase assay. Figure 3A shows a significant increase in luciferase expression induced by E2 that is abolished by ICI. The same results were obtained in HeLa cells transfected with HeG0, pGL3-p α_5 , and pRL-Tk (Figure 3C), while E2 failed to induce luciferase activity in HeLa cells transfected with the reporter plasmids alone (Figure 3B) and in the same cells transfected with pCMV-hE241G, a construct bearing ER α deleted in the nuclear localization signal (NLS), pGL3-p α_5 , and pRL-Tk (Figure 3D). All together, these data indicate the α_5 integrin subunit as an E2/ER α -regulated gene.

ER α and Sp1 bind to the 5'-flanking region of α_5 integrin subunit

The 5'-flanking region of the α_5 integrin subunit contains several elements located between positions -178 and -27 that are required for efficient basic transcription of the gene, such as AP1 sites, [22] and Sp1 and nuclear factor 1 (NF-1) sites which respond to cell adhesion on Fn [25]. The structural analysis of the α_5 promoter revealed the presence of an half ERE motif adjacent to a SP1 site which may potentially mediate the response to E2 treatment [26]. Since ER α can activate target promoters either in a classical way, by directly binding to specific ERE sequences, or indirectly,

1 through a functional interaction with other nuclear proteins such as Sp1 and subsequent binding to Sp1
2 responsive sequences [27-32], we assessed the occupancy of the 5'-flanking region of the α_5 integrin
3 subunit by ER α and Sp1 in MCF-7 cells using ChIP assay. As illustrated in Figure 4, unliganded ER α
4 is associated with the 5'-flanking region of the α_5 integrin gene containing the Sp1/half-ERE binding
5 sites and its binding is strongly enhanced by the addition of E2. The binding of ER α to this promoter
6 region is concomitant with the recruitment of Sp1. A small amount of Pol II was found to bind the
7 promoter in untreated cells, reflecting a basal regulation of the α_5 integrin gene expression (Figure 4).
8 E2 addition increased the occupancy of the 5'-flanking region of α_5 integrin gene by Pol II.

9 ***ER α regulates α_5 gene expression by binding to a Sp1/ERE half-site motif.***

10 On the basis of ChIP results, we hypothesized that both the half ERE motif and the SP1 site
11 may be involved in E2 induced α_5 integrin subunit expression. To confirm our hypothesis two reporter
12 plasmids were constructed as described in *Materials and Methods*, the pGL3-p α_5 - Δ Sp1/ERE, which
13 contains a specific deletion from positions -420 to -370 of the 5'-flanking sequence of α_5 gene
14 corresponding to a Sp1/half ERE motifs (Figure 5A), and a pGL3-Sp1/ERE which contains the putative
15 sequence responsible of E2 responsiveness (Figure 5A).

16 As reported in Figure 5B, the deletion of the Sp1/ERE half-site motif totally abolished the E2
17 responsiveness of the promoter. As expected, an increase in luciferase expression was observed in cells
18 transfected with the pGL3-Sp1/ERE construct in response to E2 that was abrogated by ICI (Figure 5C),
19 showing that the SP1/half-ERE element is the target for the regulation of the α_5 integrin subunit
20 expression in response to E2.

21 ***ER α enhances Sp1 binding to α_5 promoter.***

22 The results of our functional assays, which provide evidence for the involvement of the
23 Sp1/half-ERE binding site in mediating the effects of E2 on α_5 integrin expression, fit well with our gel

mobility shift assay. Nuclear extracts of MCF-7 cells were incubated in the presence of a ^{32}P -labeled oligonucleotide reproducing the Sp1/half-ERE motif (Figure 6A, lane 6). The protein-DNA complexes formed revealed four slower migrating bands in the untreated nuclear extracts (Figure 6A, lane 1) which are depleted in the presence of an excess of unlabeled probe (Figure 6A, lane 2) and markedly increased in nuclear extracts obtained from E2 treated cells (Figure 6A, lane 3). The specificity of the binding was demonstrated by immunodepletion induced by either anti-ER α (Figure 6A, lane 4) or anti-Sp1 (Figure 6A, lane 5) antibodies. Addressing the binding of the bipartite ER-Sp1 complex to the selected Sp1/half-ERE responsive region.

To determine whether ER α “per se” is able to bind to its own reported half-site and to cooperate with Sp1 binding, gel mobility shift assays were performed using both ER α and Sp1 purified recombinant proteins with the same ^{32}P -labeled oligonucleotide in a cell free system (Figure 6B). A single gel-shifted band was visible using Sp1 purified protein (Figure 6B, lane 2), the specificity of this band was demonstrated using an excess of unlabeled probe that caused a dramatic reduction of the band (Figure 6B, lane 3), while the addition of a cold oligonucleotide mutated in the ERE half site slightly reduced the formation of the protein/DNA complex (Figure 6B, lanes 3 and 4, respectively). Unexpectedly, no band was observed incubating the labeled oligonucleotide with ER α purified recombinant protein (Figure 6B, lanes 5, 6, and 7). It is worth to note that the gel-shifted band observed incubating the probe with Sp1 native protein became newly detectable in the presence of increasing amounts of ER α (Figure 6B, lanes 8 to 10) that resulted drastically reduced using an excess of unlabeled probe (Figure 6B, lanes 11 to 13).

Sp1 is required for E2 regulated expression of the α_5 integrin subunit.

To explore whether Sp1 contributes to the regulation of α_5 gene expression mediated by ER α , we inhibited the SP1 binding to DNA using mithramycin A, a GC-specific DNA binding drug. As

1 shown in figure 7A, mithramycin A treatment caused a significant decrease in luciferase expression of
2 the α_5 promoter construct in response to E2 stimulation in MCF-7 cells.

3 The requirement of Sp1 recruitment in the regulation of α_5 gene transcription in response to E2 was
4 also confirmed by DAPA (Figure 7B). A mutation in the Sp1 site caused a dramatic reduction of ER α
5 binding to a biotinated oligonucleotide bearing the Sp1 consensus sequence and the half-ERE site,
6 while Sp1 binding was completely abrogated (Figure 7B).

7 We confirmed these data using luciferase assays mutating the Sp1 site located at -404 to -399 of the 5'-
8 flanking region of α_5 gene (Figure 7C). All together our data indicate that the half-ERE motif is not
9 the target of the E2-dependent ER/Sp1 complex, suggesting that the regulation of α_5 gene transcription
10 and expression occurs through a functional interaction of the complex with the Sp1 responsive
11 sequences.

12 *$\alpha_5\beta_1$ integrin counteracts E2-induced cell motility in ER-positive breast cancer cells.*

13 Since we have previously demonstrated that ER α expression leads to a lower invasiveness and
14 to a reduced metastatic potential of breast cancer cells through a genomic action [10], we investigated
15 if $\alpha_5\beta_1$ integrin expression was involved in ER α mediated reduction of breast cancer cells migration
16 and invasion by knocking down $\alpha_5\beta_1$ integrin (Figure 8). A knock down of about 60% of $\alpha_5\beta_1$ integrin
17 expression was obtained after 48 h (Figure 8A). In agreement with previous studies [8, 10], E2
18 treatment produced a marked and significant reduction of both cell migration (Figure 8B) and invasion
19 (Figure 8C) in MCF-7 cells as well as in T47D cells (data not shown). Interestingly, a dramatic
20 increase of both cell migration (Figure 8B) and invasion (Figure 8C) was observed only in $\alpha_5\beta_1$
21 integrin silenced cells treated with E2, indicating that $\alpha_5\beta_1$ integrin counteract E2-induced cell motility
22 in ER-positive breast cancer cells.

1 **Discussion**

2 Previous findings have demonstrated that the concentration of the ECM protein Fn and $\alpha_5\beta_1$
3 integrin are developmentally and hormonally regulated by steroids in normal mouse mammary glands
4 [9]. The expression of the Fn-specific integrin receptor $\alpha_5\beta_1$, was found to increase between puberty
5 and sexual maturity and decrease in late pregnancy and lactation, indicating that Fn signaling may be
6 required for mammary gland proliferation in response to ovarian steroids [9]. Breast tumours, which
7 are regulated by ovarian steroids, are usually accompanied by striking changes in ECM components
8 and expression of integrins. While the epithelial basal membrane is intact in normal mammary tissues,
9 it is widely damaged in invasive carcinomas [33-34]. Particularly, in breast cancers exhibiting major
10 aggressiveness and invasiveness, ER α is no longer expressed. For instance “*in vitro*” invasiveness and
11 motility of ER α negative breast and ovarian cancer cells appear to be dramatically higher than ER α
12 positive cells [7-8, 35]. On the contrary, E2 has been found to enhance breast cancer cell migration and
13 invasion through a rapid and dynamic cytoskeleton rearrangement [36-37] and a reversible
14 epithelial/mesenchymal-like transition of ER-positive cells [38]. So, the role of E2 and ER α in breast
15 cancer cell motility and invasion remains still controversial.

16 We report here for the first time that E2/ER α modulates the expression of $\alpha_5\beta_1$ integrin in breast cancer
17 cells, through the direct activation of the regulatory region of α_5 gene.

18 Previously we demonstrated how cell adhesion on Fn induces ER α translocation from the
19 cytoplasm into the nucleus increasing the expression of estrogen responsive genes involved in the
20 down-regulation of cell motility and invasion [10]. The increased expression of α_5 integrin in response
21 to E2 stimulation reinforces the evidence that ER α may induce the expression of genes which enhance
22 cell adhesion and reduce cell migration. Some target genes have been described previously, like

1 fibulin-1, a Fn-binding ECM protein [35] that is up-regulated by ER α , metalloproteinase 9 [12] and
2 type IV collagenase [13] whose expression is negatively regulated by ER α .

3 On the basis of previous finding demonstrating that motile cells express high levels of integrin
4 $\alpha_3\beta_1$ and $\alpha_v\beta_1$ fibronectin receptor [39], while stationary cells prevalently express integrin $\alpha_5\beta_1$ [40],
5 the expression of the latter integrin seems to be a key indicator of the ability of cells to migrate. Indeed,
6 overexpression of $\alpha_5\beta_1$ on the surface of Chinese hamster ovary cells suppress their transformed
7 phenotype, causing the loss of their ability to grow in soft agar and to form tumors in nude mice [41].
8 On the contrary, $\alpha_5\beta_1$ integrin has been shown to contribute to the progression of breast cancer. For
9 instance, ligand binding to $\alpha_5\beta_1$ integrin, in the absence of $\alpha_4\beta_1$, activates invasion-stimulatory signals
10 in breast cancer cells [24, 42].

11 Virtually little is known about the molecular mechanisms controlling α_5 expression. We
12 demonstrated that the 5'-flanking region of the α_5 gene, previously reported to act as a promoter [22],
13 is a useful tool in the study of the molecular events controlling α_5 gene expression. The 5'-flanking
14 region lies within a CpG island, lacks TATA and CCAAT boxes, and contains consensus binding sites
15 for nuclear proteins such as the *ets* family of proto-oncogenes, Sp1, AP-2, and AP-1 [22]. We have
16 identified a half-ERE site (-393 to -388) located down-stream and in close proximity of a Sp1 binding
17 site (-404 to -399) in the 5'-flanking region of the α_5 gene. These adjacent binding sites are crucial
18 effectors of E2 action inducing α_5 gene expression as it emerges from our reporter assay studies. The
19 transcriptional regulation of the α_5 gene was further investigated in HeLa cells ectopically expressing a
20 wild type ER α and a mutant of the receptor containing a deletion in the nuclear localization signal. The
21 results obtained confirmed estrogen responsiveness of the α_5 gene promoter in HeLa cells expressing
22 ER α but not the deleted mutant, demonstrating how the activation of the α_5 promoter by E2/ER α
23 underlines a general mechanism of regulation not tightly linked to the cell type.

1 Theoretically, E2 regulation of the α_5 gene may be mediated by ER α binding to the 5'-
2 regulatory half-ERE motif and/or ER α may interact with Sp1 to transactivate the α_5 gene by binding to
3 the Sp1/ half-ERE binding site. Indeed, E2 has been found to activate the transcription of several ER α -
4 regulated genes in a no classical way either through the binding of ER α and the recruitment of Sp1 to
5 their own responsive element separated by 10-23 bases (Sp1(N)₁₀₋₂₃ ERE half-site) such as c-myc [43],
6 cathepsin D [44], retinoic acid receptor α_1 [45], progesteron receptor A [26] and IRS-1 [46] or through
7 its recruitment on E2-responsive GC-rich promoter motifs by interacting with Sp1 [27-32], as reported
8 for epidermal growth factor receptor [47], progesteron receptor B [48], *cad* [32], and cyclin D1 [49].

9 EMSA studies, using oligonucleotides reproducing the Sp1/half-ERE site contained in the 5'-
10 flanking region of the α_5 gene, revealed that α_5 gene transcription induced by E2-activated ER α is
11 dependent on GC-rich SP1 binding sites at -404 to -399 in the 5'-flanking region of the α_5 gene.

12 Overall, our results suggest that the responsiveness of the α_5 integrin subunit to E2 is mediated
13 by the enhanced binding of the Sp1/ER α complex to the GC-rich Sp1 region and not to the half ERE
14 site.

15 The control of α_5 expression by E2 may play an important role in regulating the different
16 biological processes that are dependent on $\alpha_5\beta_1$ integrin-Fn interactions in ER α positive cells, such as
17 the mechanisms that cells use to detach from and migrate out of the Fn-rich stroma in which they
18 mature.

19 Generally, $\alpha_5\beta_1$ integrin expression is also related to the maintenance of the differentiated and
20 stationary status of the cells, so we may argue how the expression of α_5 integrin subunit could
21 counteract E2/ER α capability to enhance “*per se*” breast cancer cell migration stimulating the
22 reversible epithelial/mesenchymal-like transition [38]. Indeed the loss of $\alpha_5\beta_1$ integrin expression is

1 concomitant with a dramatic increase of both breast cancer cell motility and invasion upon E2
2 exposure.

3 Our data are apparently in contrast to recent data demonstrating how $\alpha_4\beta_1$ integrin antagonizes
4 $\alpha_5\beta_1$ integrin mediated cell invasion in both normal and cancerous mammary cells [24, 42]. The
5 authors reported that activation of $\alpha_5\beta_1$ integrin signals increases cell invasion on type I collagen
6 matrix by up-regulating interstitial collagenase matrix metalloproteinase-1 expression. Since the
7 substratum used in our cell invasion experiments is Fn, it is not surprising that the receptors involved in
8 mediating cell invasion are distinct, and that the increased $\alpha_5\beta_1$ integrin expression enhances cell
9 adhesion on Fn coated membrane.

10 Here we demonstrate that in MCF-7 cells, not expressing $\alpha_4\beta_1$ integrin, E2 exposure induces a
11 strong increase of α_5 integrin expression concomitantly with a reduced cell migration as well as cell
12 invasion on Fn. The increased expression of $\alpha_5\beta_1$ integrin in response to E2 treatment appears to be
13 consistent with previously reported observations demonstrating that an increased expression of $\alpha_5\beta_1$
14 integrin, such as in ERB-B2 positive breast cancer cells, is responsible of the long term survival of
15 dormant micro-metastasis in bone marrow [50-51].

16 On the basis of all these observations, we may conclude that the up-regulatory effects induced
17 by E2 on $\alpha_5\beta_1$ integrin expression, may be crucially involved in the regulation of ER α -positive breast
18 cancer cell motility and invasiveness.

References

1. Altucci L, Addeo R, Cicatiello L, Dauvois S, Parker MG, Truss M, Beato M, Sica V, Bresciani F, Weisz A (1996) 17beta-Estradiol induces cyclin D1 gene transcription, p36D1-p34cdk4 complex activation and p105Rb phosphorylation during mitogenic stimulation of G(1)-arrested human breast cancer cells. *Oncogene* 12(11):2315-2324.
2. Duan R, Xie W, Burghardt RC, Safe S (2001) Estrogen receptor-mediated activation of the serum response element in MCF-7 cells through MAPK-dependent phosphorylation of Elk-1. *J Biol Chem* 276(15):11590-11598.
3. Planas-Silva MD, Shang Y, Donaher JL, Brown M, Weinberg RA (2001) AIB1 enhances estrogen-dependent induction of cyclin D1 expression. *Cancer Res* 61(10):3858-3862.
4. Sabbah M, Courilleau D, Mester J, Redeuilh G (1999) Estrogen induction of the cyclin D1 promoter: involvement of a cAMP response-like element. *Proc Natl Acad Sci U S A* 96(20):11217-11222.
5. Weisz A, Bresciani F (1993) Estrogen regulation of proto-oncogenes coding for nuclear proteins. *Crit Rev Oncog* 4(4):361-388.
6. Liotta LA, Stetler-Stevenson WG (1991) Tumor invasion and metastasis: an imbalance of positive and negative regulation. *Cancer Res* 51(18 Suppl):5054s-5059s.
7. Thompson EW, Paik S, Brunner N, Sommers CL, Zugmaier G, Clarke R, Shima TB, Torri J, Donahue S, Lippman ME *et al* (1992) Association of increased basement membrane invasiveness with absence of estrogen receptor and expression of vimentin in human breast cancer cell lines. *J Cell Physiol* 150(3):534-544.
8. Rochefort H, Platet N, Hayashido Y, Derocq D, Lucas A, Cunat S, Garcia M (1998) Estrogen receptor mediated inhibition of cancer cell invasion and motility: an overview. *J Steroid Biochem Mol Biol* 65(1-6):163-168.
9. Woodward TL, Mienaltowski AS, Modi RR, Bennett JM, Haslam SZ (2001) Fibronectin and the alpha(5)beta(1) integrin are under developmental and ovarian steroid regulation in the normal mouse mammary gland. *Endocrinology* 142(7):3214-3222.
10. Sisci D, Aquila S, Middea E, Gentile M, Maggiolini M, Mastroianni F, Montanaro D, Ando S (2004) Fibronectin and type IV collagen activate ERalpha AF-1 by c-Src pathway: effect on breast cancer cell motility. *Oncogene* 23(55):8920-8930.
11. Confort C, Rochefort H, Vignon F (1995) Insulin-like growth factors (IGFs) stimulate the release of alpha 1-antichymotrypsin and soluble IGF-II/mannose 6-phosphate receptor from MCF7 breast cancer cells. *Endocrinology* 136(9):3759-3766.
12. Crowe DL, Brown TN (1999) Transcriptional inhibition of matrix metalloproteinase 9 (MMP-9) activity by a c-fos/estrogen receptor fusion protein is mediated by the proximal AP-1 site of the MMP-9 promoter and correlates with reduced tumor cell invasion. *Neoplasia* 1(4):368-372.
13. Abbas Abidi SM, Howard EW, Dmytryk JJ, Pento JT (1997) Differential influence of antiestrogens on the in vitro release of gelatinases (type IV collagenases) by invasive and non-invasive breast cancer cells. *Clin Exp Metastasis* 15(4):432-439.
14. Lynch L, Vodyanik PI, Boettiger D, Guvakova MA (2005) Insulin-like growth factor I controls adhesion strength mediated by alpha5beta1 integrins in motile carcinoma cells. *Mol Biol Cell* 16(1):51-63.
15. Keely PJ, Fong AM, Zutter MM, Santoro SA (1995) Alteration of collagen-dependent adhesion, motility, and morphogenesis by the expression of antisense alpha 2 integrin mRNA in mammary cells. *J Cell Sci* 108 (Pt 2):595-607.

16. DiMilla PA, Stone JA, Quinn JA, Albelda SM, Lauffenburger DA (1993) Maximal migration of human smooth muscle cells on fibronectin and type IV collagen occurs at an intermediate attachment strength. *J Cell Biol* 122(3):729-737.
17. Huttenlocher A, Ginsberg MH, Horwitz AF (1996) Modulation of cell migration by integrin-mediated cytoskeletal linkages and ligand-binding affinity. *J Cell Biol* 134(6):1551-1562.
18. Yang JT, Bader BL, Kreidberg JA, Ullman-Cullere M, Trevithick JE, Hynes RO (1999) Overlapping and independent functions of fibronectin receptor integrins in early mesodermal development. *Dev Biol* 215(2):264-277.
19. Delcommenne M, Streuli CH (1995) Control of integrin expression by extracellular matrix. *J Biol Chem* 270(45):26794-26801.
20. Woodward TL, Xie J, Fendrick JL, Haslam SZ (2000) Proliferation of mouse mammary epithelial cells in vitro: interactions among epidermal growth factor, insulin-like growth factor I, ovarian hormones, and extracellular matrix proteins. *Endocrinology* 141(10):3578-3586.
21. Watt FM, Hlodivala KJ (1994) Cell adhesion. Fibronectin and integrin knockouts come unstuck. *Curr Biol* 4(3):270-272.
22. Birkenmeier TM, McQuillan JJ, Boedeker ED, Argraves WS, Ruoslahti E, Dean DC (1991) The alpha 5 beta 1 fibronectin receptor. Characterization of the alpha 5 gene promoter. *J Biol Chem* 266(30):20544-20549.
23. Suzuki T, Fujisawa JI, Toita M, Yoshida M (1993) The trans-activator tax of human T-cell leukemia virus type 1 (HTLV-1) interacts with cAMP-responsive element (CRE) binding and CRE modulator proteins that bind to the 21-base-pair enhancer of HTLV-1. *Proc Natl Acad Sci U S A* 90(2):610-614.
24. Jia Y, Zeng ZZ, Markwart SM, Rockwood KF, Ignatoski KM, Ethier SP, Livant DL (2004) Integrin fibronectin receptors in matrix metalloproteinase-1-dependent invasion by breast cancer and mammary epithelial cells. *Cancer Res* 64(23):8674-8681.
25. Larouche K, Leclerc S, Salesse C, Guerin SL (2000) Expression of the alpha 5 integrin subunit gene promoter is positively regulated by the extracellular matrix component fibronectin through the transcription factor Sp1 in corneal epithelial cells in vitro. *J Biol Chem* 275(50):39182-39192.
26. Petz LN, Nardulli AM (2000) Sp1 binding sites and an estrogen response element half-site are involved in regulation of the human progesterone receptor A promoter. *Mol Endocrinol* 14(7):972-985.
27. Safe S (2001) Transcriptional activation of genes by 17 beta-estradiol through estrogen receptor-Sp1 interactions. *Vitam Horm* 62:231-252.
28. Safe S, Kim K (2004) Nuclear receptor-mediated transactivation through interaction with Sp proteins. *Prog Nucleic Acid Res Mol Biol* 77:1-36.
29. Anderson I, Gorski J (2000) Estrogen receptor alpha interaction with estrogen response element half-sites from the rat prolactin gene. *Biochemistry* 39(13):3842-3847.
30. Wang F, Hoivik D, Pollenz R, Safe S (1998) Functional and physical interactions between the estrogen receptor Sp1 and nuclear aryl hydrocarbon receptor complexes. *Nucleic Acids Res* 26(12):3044-3052.
31. Porter W, Wang F, Wang W, Duan R, Safe S (1996) Role of estrogen receptor/Sp1 complexes in estrogen-induced heat shock protein 27 gene expression. *Mol Endocrinol* 10(11):1371-1378.
32. Khan S, Abdelrahim M, Samudio I, Safe S (2003) Estrogen receptor/Sp1 complexes are required for induction of cad gene expression by 17beta-estradiol in breast cancer cells. *Endocrinology* 144(6):2325-2335.

33. Nerlich AG, Wiest I, Wagner E, Sauer U, Schleicher ED (1997) Gene expression and protein deposition of major basement membrane components and TGF-beta 1 in human breast cancer. *Anticancer Res* 17(6D):4443-4449.
34. Ronnov-Jessen L, Petersen OW, Bissell MJ (1996) Cellular changes involved in conversion of normal to malignant breast: importance of the stromal reaction. *Physiol Rev* 76(1):69-125.
35. Hayashido Y, Lucas A, Rougeot C, Godyna S, Argraves WS, Rochefort H (1998) Estradiol and fibulin-1 inhibit motility of human ovarian- and breast-cancer cells induced by fibronectin. *Int J Cancer* 75(4):654-658.
36. Giretti MS, Fu XD, De Rosa G, Sarotto I, Baldacci C, Garibaldi S, Mannella P, Biglia N, Sismondi P, Genazzani AR *et al* (2008) Extra-nuclear signalling of estrogen receptor to breast cancer cytoskeletal remodelling, migration and invasion. *PLoS One* 3(5):e2238.
37. Saji S, Kawakami M, Hayashi S, Yoshida N, Hirose M, Horiguchi S, Itoh A, Funata N, Schreiber SL, Yoshida M *et al* (2005) Significance of HDAC6 regulation via estrogen signaling for cell motility and prognosis in estrogen receptor-positive breast cancer. *Oncogene* 24(28):4531-4539.
38. Planas-Silva MD, Waltz PK (2007) Estrogen promotes reversible epithelial-to-mesenchymal-like transition and collective motility in MCF-7 breast cancer cells. *J Steroid Biochem Mol Biol* 104(1-2):11-21.
39. Gehlsen KR, Dickerson K, Argraves WS, Engvall E, Ruoslahti E (1989) Subunit structure of a laminin-binding integrin and localization of its binding site on laminin. *J Biol Chem* 264(32):19034-19038.
40. Bronner-Fraser M (1985) Alterations in neural crest migration by a monoclonal antibody that affects cell adhesion. *J Cell Biol* 101(2):610-617.
41. Giancotti FG, Ruoslahti E (1990) Elevated levels of the alpha 5 beta 1 fibronectin receptor suppress the transformed phenotype of Chinese hamster ovary cells. *Cell* 60(5):849-859.
42. Woods Ignatoski KM, Grewal NK, Markwart S, Livant DL, Ethier SP (2003) p38MAPK induces cell surface alpha4 integrin downregulation to facilitate erbB-2-mediated invasion. *Neoplasia* 5(2):128-134.
43. Dubik D, Shiu RP (1992) Mechanism of estrogen activation of c-myc oncogene expression. *Oncogene* 7(8):1587-1594.
44. Krishnan V, Wang X, Safe S (1994) Estrogen receptor-Sp1 complexes mediate estrogen-induced cathepsin D gene expression in MCF-7 human breast cancer cells. *J Biol Chem* 269(22):15912-15917.
45. Rishi AK, Shao ZM, Baumann RG, Li XS, Sheikh MS, Kimura S, Bashirelahi N, Fontana JA (1995) Estradiol regulation of the human retinoic acid receptor alpha gene in human breast carcinoma cells is mediated via an imperfect half-palindromic estrogen response element and Sp1 motifs. *Cancer Res* 55(21):4999-5006.
46. Panno ML, Mauro L, Marsico S, Bellizzi D, Rizza P, Morelli C, Salerno M, Giordano F, Ando S (2006) Evidence that the mouse insulin receptor substrate-1 belongs to the gene family on which the promoter is activated by estrogen receptor alpha through its interaction with Sp1. *J Mol Endocrinol* 36(1):91-105.
47. Salvatori L, Ravenna L, Felli MP, Cardillo MR, Russo MA, Frati L, Gulino A, Petrangeli E (2000) Identification of an estrogen-mediated deoxyribonucleic acid-binding independent transactivation pathway on the epidermal growth factor receptor gene promoter. *Endocrinology* 141(6):2266-2274.
48. Schultz JR, Petz LN, Nardulli AM (2003) Estrogen receptor alpha and Sp1 regulate progesterone receptor gene expression. *Mol Cell Endocrinol* 201(1-2):165-175.

- 1 49. Castro-Rivera E, Samudio I, Safe S (2001) Estrogen regulation of cyclin D1 gene expression in
2 ZR-75 breast cancer cells involves multiple enhancer elements. J Biol Chem 276(33):30853-
3 30861.
- 4 50. Korah R, Boots M, Wieder R (2004) Integrin alpha5beta1 promotes survival of growth-arrested
5 breast cancer cells: an in vitro paradigm for breast cancer dormancy in bone marrow. Cancer
6 Res 64(13):4514-4522.
- 7 51. Spangenberg C, Lausch EU, Trost TM, Prawitt D, May A, Keppler R, Fees SA, Reutzel D, Bell
8 C, Schmitt S *et al* (2006) ERBB2-mediated transcriptional up-regulation of the alpha5beta1
9 integrin fibronectin receptor promotes tumor cell survival under adverse conditions. Cancer Res
10 66(7):3715-3725.

11
12
13 **Footnotes**

14 **Diego Sisci** and **Emilia Middea** contributed equally to this work.

15 **Financial support:** AIRC 2007, MIUR Ex 60%, and Regione Calabria

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

1 **Legends of figures**

2 **Figure 1 – Cell surface expression of α_5 integrin subunit in response to E2 treatment in MCF-7**

3 **cells.** α_5 integrin subunit expression was determined by Fluorescence activated cell sorting analysis
4 and western blotting. A) MCF-7 cells were serum starved for 24 h, treated with 10 nM E2 for 4, 12, and
5 24 h, and detached with EDTA solution. Cells were re-suspended in PBS and incubated with anti-
6 α_5 integrin antibody followed by fluorescein-conjugated anti-rabbit IgG. Cells were analyzed in a
7 FACS flow cytometer. The results indicate the percentage of positive cells compared to cells labeled
8 with an isotype matched control antibody; the results are representative of three independent
9 experiments and reported as mean $\pm$ SD. Statistical analysis was performed by Student's t-test (*p<0.05
10 *vs.* untreated).

11 B) MCF-7 cells serum-starved for 24 h were treated or left untreated with E2 (0, 1, 10, and 100 nM) for
12 24 h. Cells lysates were subjected to western blotting with an anti- α_5 integrin. GAPDH expression was
13 evaluated as loading control.

14 The graph represents the mean $\pm$ SD of the optical density relative increase of α_5 integrin subunit levels
15 with respect to untreated samples that were arbitrarily set at 1. The results are representative of three
16 independent experiments. Statistical analysis was performed by Student's t-test (*p<0.05 *vs.* untreated).

17 C) Serum-starved MCF-7 cells (i) were incubated or not with E2 (0, 10, and 100 nM) for 24 h, and
18 detached with EDTA solution. Cells were re-suspended in PBS and incubated with either an anti- α_5
19 (white bar) or an anti- α_4 (grey bar) integrin antibody followed by fluorescein-conjugated anti-rabbit or
20 anti-mouse IgG respectively. Cells were analyzed in a FACS flow cytometer. MCF-10 cells (ii) were
21 used as control. The results indicate the percentage of positive cells compared to cells labeled with an
22 isotype matched control antibody; the results are representative of three independent experiments and
23 reported as mean $\pm$ SD. Statistical analysis was performed by Student's t-test (*p<0.05 *vs.* untreated).

1 **Figure 2 – 17 β -estradiol increases the expression of the α_5 integrin subunit in MCF-7 cells.** MCF-
2 7 cells serum-starved for 24 h were treated or left untreated with 10 nM E2 for 16, 24, and 48 h (A) and
3 with 1, 10 and 100 nM of E2 and/or 1 μ M ICI 182,780 (C). Cells were lysed after incubation, and
4 proteins were subjected to western blotting with an anti-integrin α_5 antibody or an anti-GAPDH as
5 loading control. The results are representative of three independent experiments. The graph represents
6 the mean $\pm$ SD of the optical density of α_5 integrin subunit normalized versus GAPDH and represented
7 as relative increase with respect to untreated samples that were arbitrarily set at 1. Data were
8 statistically analyzed by Student's t-test (* p<0.05 vs untreated).

9 MCF-7 cells were treated for 24 and 48 h with 10 nM E2 (B) or with 1, 10 and 100 nM E2 and/or 1
10 mM ICI 182,780 (D) and total RNA was extracted. The abundance of α_5 integrin subunit mRNA was
11 detected by RT-PCR, as described in *Materials and Methods*. 36B4 mRNA levels were determined as
12 control. The results are representative of three independent experiments. The graph represents the
13 mean $\pm$ SD of the optical density of α_5 integrin subunit normalized versus 36B4 and reported as relative
14 increase with respect to untreated samples that were arbitrarily set at 1. Data were statistically analyzed
15 by Student's t-test (* p<0.05 vs untreated).

16 E) HeLa cells were transfected with increasing amount of pSG5-HeG0 (ER α : 1 μ g and 2 μ g) or pSG5
17 empty vector (Vector). Transfected cells were serum starved for 24 h and consecutively treated with E2
18 (10 nM). α_5 integrin and ER α expression was detected by WB. The graph represents the mean $\pm$ SD of
19 the optical density relative increase of α_5 integrin subunit levels with respect to untreated samples that
20 were arbitrarily set at 1. The results are representative of three independent experiments. Data were
21 statistically analyzed by Student's t-test (* p<0.05 vs untreated).

22 **Figure 3 – The 5'-flanking region of α_5 gene acts as promoter and is activated by E2/ER α .** The
23 transcriptional activity of ER α on the α_5 integrin subunit gene was evaluated using luciferase reporter

system, as described in *Materials and Methods*. MCF-7 cells (A), and HeLa cells (B) were transfected with a DNA mixtures containing 0,5 μ g pGL3-p α 5 reporter plasmid (detailed in *Materials and Methods*) and 50ng of pRL-Tk. Transfected cells were serum starved for 24 h and consecutively treated with E2 (10 nM) and/or ICI (1 μ M) for 24 h. C) HeLa cells transfected with 0,2 μ g pSG5-HeG0, 0,3 μ g pGL3-p α 5, and 50ng of pRL-Tk. D) HeLa cells were transfected with 0,2 μ g pCMV-hE241G, 0,3 μ g pGL3-p α 5, and 50ng of pRL-Tk. Firefly luciferase activity was internally normalized to Renilla luciferase activity and expressed as fold of increase with respect to untreated samples. Data were statistically analyzed by Student's t-test (* p<0.05 vs control). The results are expressed as mean $\pm$ SD from four independent experiments

Figure 4 – ER α is recruited to the 5'-flanking region of α 5 gene. Chromatin immunoprecipitations were carried out on serum-starved MCF-7 cells as described in *Materials and Methods*. Cells were treated or left untreated with E2 (10 nM) for 45 minutes, lysed and DNA-associated proteins were precipitated using either anti-ER α Ab, or anti-Sp1 Ab, or anti-Pol II Ab. The 5'-flanking sequence of α 5 gene containing both Sp1 and half-ERE sites was detected by PCR with specific primers listed in *Materials and Methods*. Inputs DNA were amplified as loading controls.

Figure 5 – The Sp1/ERE half-sites contained in the 5'-flanking region of the α 5 gene is responsible of E2/ER α activation. A specific deletion (-420/-370) in the 5'-flanking region of the α 5 gene was obtained as described in *Materials and Methods* (A). MCF-7 cells were transfected with a DNA mixtures containing 0.5 μ g pGL3-p Δ α 5 reporter plasmid (B), 0,5 μ g pGL3-Sp1/ERE (C), and 50ng of pRL-Tk. Transfected cells were serum starved for 24 h and then treated with E2 (10 nM) and/or ICI (1 μ M) for 24 h. Firefly luciferase activity was internally normalized to Renilla luciferase activity and expressed as fold of increase with respect to untreated samples. Results represent the

mean $\pm$ SD from several experiments. Data were statistically analyzed by Student's t-test (* $p < 0.05$ vs control).

Figure 6 - E2 enhances the binding of Sp1 to the 5'-flanking region of the α_5 gene containing the Sp1/half-ERE binding sites.

A) Nuclear extracts from MCF-7 cells treated for 24 h with E2 10 nM (lanes 3 to 5) or untreated (lane 1 and 2) were incubated with [γ - 32]P-labeled oligonucleotides containing the Sp1 and the half-ERE binding sites (reported in *Materials and Methods*) and subjected to electrophoresis in a 6% polyacrylamide gel. The specificity of the binding was tested by adding to the reaction mixture either anti-ER α (lane 4) or anti-Sp1 antibodies (lane 5). Competition experiments were performed by adding a 100-fold molar excess of unlabeled probe (lane 2). Lane 6 contains the probe.

B), The [γ - 32]P-labeled oligonucleotide (lane 1), described above, was incubated with either purified Sp1 protein (15 ng) (lane 2 to 4) or purified ER α protein (0,18 pmol) (lane 5 to 7). The presence of shifted complexes was also evaluated by incubating Sp1 (10 ng) with different amounts of the recombinant ER α (0; 0,18; 0,36 and 0,54 pmol). Competition experiments were performed by adding a 100-fold molar excess of either unlabeled probe (lane 3, 6, and 11 to 13) or the probe mutated in the half-ERE site (reported in *Materials and Methods*) (lane 4, 7).

Figure 7 - Sp1 is required for the expression of α_5 gene in response to E2. MCF-7 cells (A) were transfected with DNA mixtures containing 0,5 μ g pGL3-p α_5 reporter plasmid and 50 ng of pRL-Tk. Transfected cells were serum starved for 24 h and consecutively treated with E2 (10 nM) for 24 h and/or pre-treated with Mithramycin A (100 μ M). Firefly luciferase activity was internally normalized to Renilla luciferase activity and expressed as fold of increase with respect to untreated samples. Data were statistically analyzed by Student's t-test (* $p < 0.05$). The results are expressed as mean $\pm$ SD. from several independent experiments.

1 B) Nuclear extracts from E2-treated and untreated MCF-7 cells for 24 h were incubated with either a
2 biotinated oligonucleotide containing the Sp1 and the half-ERE binding sites or a biotinated
3 oligonucleotide mutated in the Sp1 consensus site and subjected to DNA affinity precipitation assay as
4 described in Materials and Methods. The specificity of the binding was tested by loading the unbound
5 fraction (C-). E2 treated MCF-7 cells lysate was used as positive control (C+). The results are
6 representative of three independent experiments.

7 C) MCF-7 cells were transfected with DNA mixtures containing 0,5 μ g pGL3-p α 5 reporter plasmid, or
8 the same plasmid mutated in the Sp1 site located at -404/-399 (detailed in *Materials and Methods*) and
9 50ng of pRL-Tk. Transfected cells were serum starved for 24 h and consecutively treated with E2 (10
10 nM) for 24 h. Firefly luciferase activity was internally normalized to Renilla luciferase activity and
11 expressed as fold of increase with respect to untreated samples. Data were statistically analyzed by
12 Student's t-test (* $p < 0.05$ vs control). The results are expressed as mean $\pm$ SD. from several independent
13 experiments.

14 **Figure 8 – $\alpha_5\beta_1$ integrin expression inhibits E2-mediated cell migration and invasion.** MCF-7 cells
15 were transfected in suspension with 100 pmol siRNAs/well (si $\alpha_5\beta_1$ integrin or a scrambled siRNA for
16 control samples), plated, starved for 24 h and treated with E2 (10 nM) for additional 24 h. Cytoplasmic
17 protein extracts (A) (25 μ g/lane) were analyzed by WB to evaluate $\alpha_5\beta_1$ integrin content. GAPDH was
18 evaluated as loading control. The graph represents the mean $\pm$ SD of the optical density of α_5 integrin
19 subunit normalized versus GAPDH. MCF-7 cells, transfected and treated as before, were detached,
20 plated and allowed to migrate for 16 hrs on porous membranes coated (C) or not (B) with 30 μ g/ml
21 Fibronectin (Fn) to evaluate the effects of Estradiol (10 nM) exposure on cell motility (B) and invasion
22 (C). Cells were then fixed, stained and the cells migrated to the lower surfaces of the membranes were
23 counted. The results represent the mean $\pm$ S.D. of three independent experiments. Data were statistically
24 analyzed by Student's t test (* $p < 0.05$ vs untreated; ■ $p < 0.05$ vs the corresponding treatment in

1 scrambled).

2

3

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8