

Higher-dimensional normalisation strategies for acyclicity

Yves Guiraud, Philippe Malbos

► **To cite this version:**

Yves Guiraud, Philippe Malbos. Higher-dimensional normalisation strategies for acyclicity. *Advances in Mathematics*, Elsevier, 2012, 231 (3-4), pp.2294-2351. <10.1016/j.aim.2012.05.010>. <hal-00531242v3>

HAL Id: hal-00531242

<https://hal.archives-ouvertes.fr/hal-00531242v3>

Submitted on 8 Aug 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HIGHER-DIMENSIONAL NORMALISATION STRATEGIES FOR ACYCLICITY

YVES GUIRAUD – PHILIPPE MALBOS

Abstract – We introduce *acyclic polygraphs*, a notion of complete categorical cellular model for (small) categories, containing generators, relations and higher-dimensional globular syzygies. We give a rewriting method to construct explicit acyclic polygraphs from *convergent* presentations. For that, we introduce *higher-dimensional normalisation strategies*, defined as homotopically coherent ways to relate each cell of a polygraph to its normal form, then we prove that acyclicity is equivalent to the existence of a normalisation strategy. Using acyclic polygraphs, we define a higher-dimensional homotopical finiteness condition for higher categories which extends Squier’s finite derivation type for monoids. We relate this homotopical property to a new homological finiteness condition that we introduce here.

Keywords – rewriting; polygraphic resolution; homology of small categories; identities among relations.

M.S.C. 2000 – 18C10; 18D05; 18G10; 18G20; 68Q42.

Affiliation – INRIA and Université de Lyon, Institut Camille Jordan, CNRS UMR 5208, Université Claude Bernard Lyon 1, 43 boulevard du 11 novembre 1918, 69622 Villeurbanne cedex, France.

CONTENTS

1	Introduction	2
2	Polygraphic resolutions	8
3	Normalisation strategies for polygraphs	15
4	Polygraphic resolutions from convergent presentations	25
5	Abelianisation of polygraphic resolutions	38
6	Examples	53

1. INTRODUCTION

1.1. An overview of Squier's theory

In the eighties, Squier has established a link between some computational, homological and homotopical properties of monoids, [41, 42]. This allowed him to answer an open question: does a finitely generated monoid with a decidable word problem always admit a finite convergent presentation?

1.1.1. The word problem and rewriting theory. Given a monoid \mathbf{M} , a generating set Σ_1 for \mathbf{M} provides a way to represent the elements of \mathbf{M} in the free monoid Σ_1^* , *i.e.*, as finite words written with the elements of Σ_1 . But, if \mathbf{M} is not free, there is no reason for an element of \mathbf{M} to have a single representative in Σ_1^* . The *word problem* for \mathbf{M} consists in finding a generating set Σ_1 and an algorithm that can determine whether or not any two elements of Σ_1^* represent the same element of \mathbf{M} .

One way to solve the word problem is to exhibit a finite presentation (Σ_1, Σ_2) of \mathbf{M} with a good computational property, called *convergence* in rewriting theory. There, one studies presentations where the relations in Σ_2 are not seen as equalities between the words in Σ_1^* , such as $u = v$, but, instead, as rewriting rules that can only be applied in one direction, like $u \Rightarrow v$, simulating a non-reversible computational process. Convergence is defined as the conjunction of the following two conditions:

- *termination, i.e.*, all the computations end eventually,
- *confluence, i.e.*, different computations on the same input lead to the same result.

A finite convergent presentation (Σ_1, Σ_2) of \mathbf{M} gives a solution to the word problem: the *normal form algorithm*. Given an element u in Σ_1^* , convergence ensures that all the applications of directed relations to u , in any possible manner, will eventually produce a unique result: an element \hat{u} of Σ_1^* where no directed relation applies anymore, called the *normal form* of u . And, by construction, two elements u and v of Σ_1^* represent the same element of \mathbf{M} if, and only if, their normal forms are equal in Σ_1^* . Finally, finiteness ensures that one can determine if an element of Σ_1^* is a normal form, by examining all the relations. (As far as rewriting is concerned, this article is self-contained, but this wider mathematical field is covered in more details by Book and Otto, [8], Baader and Nipkow, [3], and the group Terese, [45].)

Thus, if a monoid admits a finite convergent presentation, it has a decidable word problem. In the middle of the eighties, it was still unknown if the converse implication held. In [20], Kapur and Narendran had exhibited a monoid that admits a finite generating set for which the word problem was solvable, but that do not admit a finite convergent presentation with the same generators. However, this did not answer the original question, the generating set having been fixed.

1.1.2. From computational to homological properties. At that time, Squier linked the existence of a finite convergent presentation to a homological invariant of the monoid, the homological type left-FP₃, that is independent of the choice of a presentation of \mathbf{M} and, in particular, of a generating set. A monoid \mathbf{M} has homological type left-FP₃ when there exists an exact sequence

$$P_3 \longrightarrow P_2 \longrightarrow P_1 \longrightarrow P_0 \longrightarrow \mathbb{Z} \longrightarrow 0$$


of (left) modules over \mathbf{M} , where \mathbb{Z} denotes the trivial \mathbf{M} -module and each P_i is projective and finitely generated.

From a presentation (Σ_1, Σ_2) of \mathbf{M} , one can build an exact sequence of free \mathbf{M} -modules


$$\mathbb{Z}\mathbf{M}[\Sigma_2] \xrightarrow{J} \mathbb{Z}\mathbf{M}[\Sigma_1] \longrightarrow \mathbb{Z}\mathbf{M} \longrightarrow \mathbb{Z} \longrightarrow 0, \quad (1)$$

where $\mathbb{Z}\mathbf{M}[\Sigma_1]$ and $\mathbb{Z}\mathbf{M}[\Sigma_2]$ are the free \mathbf{M} -modules over Σ_1 and Σ_2 , respectively. The differential J , called the *Fox Jacobian* after [15], is defined on a directed relation $\alpha : u \Rightarrow v$ by $J(\alpha) = [u] - [v]$, where $[\cdot]$ is the unique derivation of Σ_1^* with values in $\mathbb{Z}\mathbf{M}[\Sigma_1]$ that extends the canonical inclusion of Σ_1 into $\mathbb{Z}\mathbf{M}[\Sigma_1]$.

In [41], Squier proved that, when (Σ_1, Σ_2) is a convergent presentation, its *critical branchings* form a generating set of the kernel of the Fox Jacobian. A critical branching of (Σ_1, Σ_2) is an overlapping application of two different directed relations on the same word u of Σ_1^* , where u has minimal size. For example, the relations $\alpha : xy \Rightarrow v$ and $\beta : yz \Rightarrow w$ generate a critical branching $(\alpha z, x\beta)$ on $u = xyz$:



Convergence of the presentation (Σ_1, Σ_2) ensures that any critical branching (f, g) can be completed as in the following diagram:



The boundary of such a branching is defined as the element $J'(f, g) = [f] - [g] + [f'] - [g']$, where $[\cdot]$ extends the canonical inclusion of Σ_2 into $\mathbb{Z}\mathbf{M}[\Sigma_2]$.

Squier proves that the set Σ_3 of critical branchings of (Σ_1, Σ_2) and the boundary J' extend the exact sequence (1) by one step:

$$\mathbb{Z}\mathbf{M}[\Sigma_3] \xrightarrow{J'} \mathbb{Z}\mathbf{M}[\Sigma_2] \xrightarrow{J} \mathbb{Z}\mathbf{M}[\Sigma_1] \longrightarrow \mathbb{Z}\mathbf{M} \longrightarrow \mathbb{Z} \longrightarrow 0. \quad (2)$$

Moreover, when (Σ_1, Σ_2) is finite, then Σ_3 is finite, proving that, if a monoid has a finite convergent presentation, then it is of homological type left-FP₃.

Finally, Squier exhibited a finitely generated monoid, with a decidable word problem, but that is not of homological type left-FP₃. This gave a negative answer to the aforementioned open question: there exist finitely generated monoids with a decidable word problem that do not admit a finite convergent presentation (for any possible finite set of generators).

1.1.3. From computational to homotopical properties. In [42], Squier links the existence of a finite convergent presentation to a homotopical invariant of monoids, called *finite derivation type* (FDT₃) and that is a natural extension of the properties of being finitely generated (FDT₁) and finitely presented (FDT₂).

1. Introduction

To define this invariant, for a monoid \mathbf{M} with a presentation (Σ_1, Σ_2) , Squier constructs a cellular complex $S(\Sigma_1, \Sigma_2)$ with one 0-cell, whose 1-cells are the elements of the free monoid Σ_1^* and whose 2-cells are generated by the relations of Σ_2 . More precisely, there is a 2-cell between every pair of words with shape wu_1w' and wv_1w' such that $u_1 = v_1$ is a relation in Σ_2 . Then, to get $S(\Sigma_1, \Sigma_2)$, one fills with 3-cells all the squares formed by independent applications of relations, such as the following one, where (u_1, v_1) and (u_2, v_2) are relations in Σ_2 :

$$\begin{array}{ccc}
 & u_1 = v_1 & wv_1w'u_2w'' & u_2 = v_2 \\
 & \parallel & & \parallel \\
 wu_1w'u_2w'' & & & wv_1w'v_2w'' \\
 & u_2 = v_2 & wu_1w'v_2w'' & u_1 = v_1
 \end{array}$$

If Σ_3 is a set of 3-cells over $S(\Sigma_1, \Sigma_2)$, then the set $\Sigma_1^*\Sigma_3\Sigma_1^*$ is the set of 3-cells $u\beta v$, with β in Σ_3 and u and v in Σ_1^* , and whose boundary is the one of β multiplied by u on the left and v on the right. A *homotopy basis* of (Σ_1, Σ_2) is a set Σ_3 of 3-cells such that $\Sigma_1^*\Sigma_3\Sigma_1^*$ makes the complex $S(\Sigma_1, \Sigma_2)$ contractible. A monoid is of *finite derivation type* (FDT₃) if it admits a finite presentation whose associated complex admits a finite homotopy basis or, in other words, whose “relations among the relations” are finitely generated.

Squier proves that, given a convergent presentation (Σ_1, Σ_2) , it is sufficient to attach one 3-cell to each 3-dimensional sphere corresponding to a critical branching to get a homotopy basis of (Σ_1, Σ_2) . Moreover, if Σ_2 is finite, the presentation (Σ_1, Σ_2) has finitely many critical branchings proving that, if a monoid admits a finite convergent presentation, then it is FDT₃. Squier used this result to give another proof that there exist finitely generated monoids with a decidable word problem that do not admit a finite convergent presentation.

1.1.4. Refinements of Squier’s conditions. By his results, Squier has opened two different directions, one homological and one homotopical, to explore in the quest for a complete characterisation of the existence of finite convergent presentations in the case of monoids. The corresponding invariants are related: FDT₃ implies left-FP₃, as proved by several authors, [13, 39, 24]. The converse implication is false in general, as already noted by Squier in [42], yet it is true in the special case of groups, [14], the latter result being based on the Brown-Hueschmann isomorphism between homotopical and homological syzygies, [10].

However, the invariants left-FP₃ and FDT₃ are not complete characterisations of the property to admit a finite convergent presentation: they are necessary, but not sufficient conditions, as already proved by Squier in [42]. Following this observation, various refinements of both invariants have been explored.

In the homological direction, thanks to the notion of Abelian resolution, one defines the more restrictive conditions left-FP_n, for every natural number $n > 3$, and left-FP_∞: a monoid \mathbf{M} has homological type left-FP_∞ when there exists a resolution of the trivial \mathbf{M} -module by finitely generated and projective \mathbf{M} -modules. In [21], a notion of n -fold critical branching is used to complete the exact sequence (2) into a resolution, obtaining the following implication: if a monoid admits a finite convergent presentation, then it is of homological type left-FP_∞, the converse implication still being false in general. The same results are also known for associative algebras, [1], and for groups, [12, 9, 16]. One can obtain similar

implications with the properties right-FP_∞ and bi-FP_∞ , defined with resolutions by right modules and bimodules, respectively.

In the homotopical direction, the condition FDT_3 has been refined into FDT_4 , a property about the existence of a finite presentation with a finite homotopy basis, itself satisfying a homotopical finiteness property, [32]. The condition FDT_4 is also necessary for a monoid to admit a finite convergent presentation and it is sufficient, but not necessary, for the conditions $\text{left/right/bi-FP}_4$.

1.2. Organisation and main results of the article

1.2.1. Polygraphic resolutions. In Section 2, we introduce a notion of homotopical resolution that generalises Squier’s complex, in order to define the homotopical finiteness conditions FDT_n and FDT_∞ . Squier’s complex appears as the first two dimensions of a free $(\infty, 1)$ -category, *i.e.*, a free ∞ -category whose cells of dimension 2 and higher are invertible. Then, homotopy bases and higher homotopy bases generate the higher dimensions of this $(\infty, 1)$ -category, in such a way that the latter is homotopically equivalent to the starting monoid. Moreover, these resolutions further generalise from monoids to \mathfrak{p} -categories, yielding free (∞, \mathfrak{p}) -categories.

More explicitly, let (Σ_1, Σ_2) be a presentation of a monoid \mathbf{M} . Such a presentation of a monoid, or more generally of a (small) category, is called a $(2, 1)$ -polygraph in this article. The terminology comes from Burroni’s polygraphs, [11], also known as Street’s computads, [43, 44]. From the $(2, 1)$ -polygraph (Σ_1, Σ_2) , we generate a free $(2, 1)$ -category Σ_2^\top by taking all the formal composites of 2-cells, modulo exchange relations that correspond exactly to the 2-cells of Squier’s complex associated to (Σ_1, Σ_2) . Informally, the $(2, 1)$ -category Σ_2^\top is homotopically equivalent to Squier’s complex: this allows us to see a homotopy basis as a set Σ_3 of 3-cells, attached to Σ_2^\top , such that the quotient $(2, 1)$ -category Σ_2^\top/Σ_3 is homotopically equivalent to the original monoid or, equivalently, such that any parallel 2-cells of Σ_2^\top are identified in the quotient by Σ_3 .

Then, one considers the free $(3, 1)$ -category Σ_3^\top generated by the $(3, 1)$ -polygraph $(\Sigma_1, \Sigma_2, \Sigma_3)$. One defines a homotopy basis of Σ_3^\top as a set of 4-cells over Σ_3^\top that relate every parallel 3-cells. The same idea is used to define homotopy bases in every dimension, yielding our notion of *polygraphic resolution* of the monoid \mathbf{M} : this is an *acyclic* $(\infty, 1)$ -polygraph $\Sigma = (\Sigma_n)_{n \geq 1}$ such that (Σ_1, Σ_2) is a presentation of \mathbf{M} , where *acyclic* means that each Σ_{n+1} is a homotopy basis of the free $(n, 1)$ -category Σ_n^\top , for $n \geq 3$.

The notion we get is close to Métayer’s polygraphic resolutions, introduced in [36]: these are ∞ -polygraphs that produce cofibrant approximations (free objects that are homotopically equivalent to the original one) in the canonical model structure on ∞ -categories, described in [25]. Our resolutions, called (∞, \mathfrak{p}) -polygraphs, have the same good homotopical properties, with respect to the canonical model structure on (∞, \mathfrak{p}) -categories, obtained by Ara and Métayer, [2].

Theorem 2.3.4. *Let Σ be a polygraphic resolution of a \mathfrak{p} -category \mathcal{C} . The canonical projection $\Sigma^\top \rightarrow \mathcal{C}$ is a cofibrant approximation of \mathcal{C} in the canonical model structure on (∞, \mathfrak{p}) -categories.*

We say that a monoid and, more generally, a \mathfrak{p} -category is of *finite ∞ -derivation type* (FDT_∞) when it admits a polygraphic resolution with finitely many cells in every dimension. This generalises to higher categories and in every dimension the two previously known homotopical finiteness conditions, FDT_3 introduced by Squier for monoids and its refinement FDT_4 .

1. Introduction

1.2.2. Normalisation strategies for polygraphs. In Section 3, we give a constructive characterisation of the acyclicity of an $(\infty, 1)$ -polygraph. Let Σ be a polygraphic resolution of a monoid \mathbf{M} . In particular, if $\pi : \Sigma_1^* \rightarrow \mathbf{M}$ denotes the canonical projection, one can choose a (non-functorial) section ι of π . Moreover, the 2-cells of Σ_2 are generating relations for \mathbf{M} : for every element u of Σ_1^* , one can choose a 2-cell $\sigma_u : u \Rightarrow \iota\pi(u)$ in Σ_2^\top . Then, we use that Σ_3 is a homotopy basis: for every 2-cell $f : u \Rightarrow v$ of Σ_2^\top , one can choose a 3-cell σ_f in Σ_3^\top with shape

$$\begin{array}{ccc}
 & & f \\
 & \curvearrowright & \\
 u & & v \\
 & \curvearrowleft & \\
 & & \sigma_f \\
 & \Downarrow & \\
 & & \iota\pi(u) = \iota\pi(v) \\
 & \Uparrow & \\
 & & \sigma_u \quad \sigma_v
 \end{array}$$

From the acyclicity of Σ , we deduce that similar choices can be made in every dimension, in a coherent way: if $\pi : \Sigma^\top \rightarrow \mathbf{M}$ and $\iota : \mathbf{M} \rightarrow \Sigma^\top$ are now seen as (weak) ∞ -functors, then σ is a (weak) natural isomorphism from the identity of Σ^\top to the composite (weak) ∞ -functor $\iota\pi$.

We call σ a *normalisation strategy*. It generalises to every dimension the notion of strategy appearing in rewriting theory: a canonical way, among all the computations generated by the directed relations, to reduce a word into a normal form. An $(\infty, 1)$ -polygraph with a normalisation strategy is *normalising*.

Theorem 3.3.6. *An $(n, 1)$ -polygraph is acyclic if, and only if, it is normalising.*

Moreover, a normalisation strategy can always be assumed to commute with the monoid product in a sensible way: it can always “reduce” a word by starting on the left or on the right. This leads to the *left-normalising* and *right-normalising* properties for $(\infty, 1)$ -polygraphs, which are also equivalent to acyclicity.

1.2.3. Polygraphic resolutions from convergent presentations. In Section 4, we use normalisation strategies to build, by induction on the dimension, an explicit polygraphic resolution from a convergent presentation. Given a convergent $(2, 1)$ -polygraph Σ , the first dimensions of the polygraphic resolution $c_\infty(\Sigma)$ we get are similar to the ones of Squier’s complex: generators in dimension 1, generating relations in dimension 2, critical branchings in dimension 3. Then, we build the 4-cells from the critical triple branchings, *i.e.*, the minimal overlappings of three 2-cells on the same 1-cell: for such a (f, g, h) , we use a normalisation strategy σ to build the corresponding 4-cell

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & v & \\
 f \nearrow & & \sigma_v \searrow \\
 u & \xrightarrow{g} w & \xrightarrow{\sigma_w} \hat{u} \\
 h \searrow & & \sigma_x \nearrow \\
 & x & \\
 & \Downarrow A & \\
 & \Downarrow B & \\
 & & \hat{u}
 \end{array}
 & \Longrightarrow &
 \begin{array}{ccc}
 & v & \\
 f \nearrow & & \sigma_v \searrow \\
 u & & \hat{u} \\
 h \searrow & & \sigma_x \nearrow \\
 & x & \\
 & \Downarrow C & \\
 & & \hat{u}
 \end{array}
 \end{array}$$

where A , B and C are 3-cells built by using the critical branchings and the normalisation strategy σ . In higher dimensions, we proceed similarly to build the $(n + 1)$ -cells of the resolution from the critical n -fold branchings.

Theorem 4.5.3. *If Σ is a convergent presentation of a category \mathbf{C} , then the $(\infty, 1)$ -polygraph $c_\infty(\Sigma)$ is a polygraphic resolution of \mathbf{C} .*

Since a finite convergent $(2, 1)$ -polygraph has finitely many n -fold critical branchings, a category with a finite convergent presentation is FDT_∞ .

1.2.4. Abelianisation of polygraphic resolutions. In Section 5, we relate the homotopical finiteness condition FDT_∞ to a new homological finiteness condition, called FP_∞ . For that, from a polygraphic resolution Σ of a category \mathbf{C} , we deduce a free Abelian resolution

$$\dots \xrightarrow{\delta_{n+1}} F_{\mathbf{C}}[\Sigma_n] \xrightarrow{\delta_n} F_{\mathbf{C}}[\Sigma_{n-1}] \xrightarrow{\delta_{n-1}} \dots \xrightarrow{\delta_2} F_{\mathbf{C}}[\Sigma_1] \xrightarrow{\delta_1} F_{\mathbf{C}}[\Sigma_0] \xrightarrow{\varepsilon} \mathbb{Z} \longrightarrow 0$$

in the category of *natural systems on \mathbf{C}* , a generalisation of bimodules due to Baues, see [5]. This complex, denoted by $F_{\mathbf{C}}[\Sigma]$, is called the *Reidemeister-Fox-Squier complex of Σ* and extends Squier's exact sequence (2). The acyclicity of $F_{\mathbf{C}}[\Sigma]$ is proved by using a contracting homotopy induced by a normalisation strategy.

Theorem 5.4.3. *If Σ is a polygraphic resolution of a category \mathbf{C} , then the Reidemeister-Fox-Squier complex $F_{\mathbf{C}}[\Sigma]$ is a free resolution of the constant natural system \mathbb{Z} on \mathbf{C} .*

We define the homological properties FP_n , for any n , and FP_∞ as the refined versions of left/right/bi- FP_n and left/right/bi- FP_∞ with natural systems instead of left/right/bi-modules. We get that, for categories, the property FDT_n implies FP_n and, as a consequence, that a category with a finite convergent presentation is of homological type FP_∞ .

Finally, we relate the homological 2-syzygies of a presentation Σ to its *identities among relations*, defined by the authors in [19].

Theorem 5.6.5. *For every 2-polygraph Σ , the natural systems of homological 2-syzygies and of identities among relations of Σ are isomorphic.*

As a consequence, for finitely presented categories, the homological finiteness condition FP_3 is equivalent to the homotopical finiteness condition FDT_{ab} , characterising the existence of a finite homotopy basis of an Abelianised version of a presentation of the category, see Theorem 5.7.3.

1.2.5. Examples. Throughout this article, we apply our constructions to the example of the *reduced standard presentation* of a category, yielding, at the end, an Abelian resolution that is similar to the bar construction. In Section 6, we give two more examples: the monoid with one non-unit and idempotent element and the subcategory of the simplicial category whose morphisms are the monotone surjections only. They give rise to resolutions where the higher-dimensional cells have the shapes of associahedra and permutohedra, respectively.

1.3. Acknowledgements

The authors wish to thank François Métayer, Timothy Porter and the anonymous referee for their comments and suggestions that helped to produce an improved version of this work.

2. POLYGRAPHIC RESOLUTIONS

Throughout this section, we denote by n either a natural number or ∞ .

2.1. Higher-dimensional categories

If \mathcal{C} is an n -category (we always consider strict, globular n -categories), we denote by \mathcal{C}_k the set (and the k -category) of k -cells of \mathcal{C} . If f is a k -cell of \mathcal{C} , then $s_i(f)$ and $t_i(f)$ respectively denote the i -source and i -target of f ; we drop the suffix i when $i = k - 1$. The source and target maps satisfy the *globular relations*:

$$s_i \circ s_{i+1} = s_i \circ t_{i+1} \quad \text{and} \quad t_i \circ s_{i+1} = t_i \circ t_{i+1}.$$

We respectively denote by $f : u \rightarrow v$, $f : u \Rightarrow v$ or $f : u \Rrightarrow v$ a 1-cell, a 2-cell or a 3-cell f with source u and target v .

If f and g are i -composable k -cells, that is when $t_i(f) = s_i(g)$, we denote by $f \star_i g$ their i -composite; we simply use fg when $i = 0$. The compositions satisfy the *exchange relations* given, for every $i \neq j$ and every possible cells f, g, h and k , by:

$$(f \star_i g) \star_j (h \star_i k) = (f \star_j h) \star_i (g \star_j k).$$

If f is a k -cell, we denote by 1_f its identity $(k + 1)$ -cell. When 1_f is composed with cells of dimension $k + 1$ or higher, we simply denote it by f .

2.1.1. (n, p) -categories. A k -cell f of an n -category \mathcal{C} , with i -source u and i -target v , is *i -invertible* when there exists a (necessarily unique) k -cell g in \mathcal{C} , with i -source v and i -target u in \mathcal{C} , called the *i -inverse of f* , that satisfies

$$f \star_i g = 1_u \quad \text{and} \quad g \star_i f = 1_v.$$

When $i = k - 1$, we just say that f is *invertible* and we denote by f^- its *inverse*. As in higher-dimensional groupoids, if a k -cell f is invertible and if its i -source u and i -target v are invertible, then f is $(i - 1)$ -invertible, with $(i - 1)$ -inverse given by

$$v^- \star_{i-1} f^- \star_{i-1} u^-.$$

For a natural number $p \leq n$, or for $p = n = \infty$, an (n, p) -category is an n -category whose k -cells are invertible for every $k > p$. When $n < \infty$, this is a p -category enriched in $(n - p)$ -groupoids and, when $n = \infty$, a p -category enriched in ∞ -groupoids. In particular, an (n, n) -category is an n -category, an $(n, 0)$ -category is an n -groupoid and, when $n < \infty$, an $(n, n - 1)$ -category is a track $(n - 1)$ -category, as defined in [18] after Baues, [5, 6]. If $n < \infty$, any (n, p) -category can be seen as an (∞, p) -category with only identity k -cells for $k > n$.

2.1.2. Spheres and asphericity. Let \mathcal{C} be an n -category. A *0-sphere of \mathcal{C}* is a pair $\gamma = (f, g)$ of 0-cells of \mathcal{C} and, for $1 \leq k \leq n$, a *k -sphere of \mathcal{C}* is a pair $\gamma = (f, g)$ of parallel k -cells of \mathcal{C} , *i.e.*, with $s(f) = s(g)$ and $t(f) = t(g)$; we call f the *source* of γ and g its *target*. If f is a k -cell of \mathcal{C} , for $1 \leq k \leq n$, the *boundary of f* is the $(k - 1)$ -sphere $(s(f), t(f))$. If $n < \infty$, the n -category \mathcal{C} is *aspherical* when the source and the target of each n -sphere of \mathcal{C} coincide, *i.e.*, when every n -sphere of \mathcal{C} has shape (f, f) for some $(n - 1)$ -cell f of \mathcal{C} .

2.1.3. The canonical model structure on (∞, p) -categories. In [2], Ara and Métayer have proved that the canonical model structure on ∞ -categories from [25] transfers to (∞, p) -categories through the adjunction

$$\begin{array}{ccc} & \mathbb{U} & \\ & \curvearrowright & \\ (\infty, p)\text{Cat} & \top & \infty\text{Cat} \\ & \curvearrowleft & \\ & (\cdot)^\top & \end{array}$$

where \mathbb{U} is the forgetful functor and its left adjoint adds to an ∞ -category all the missing inverses for cells of dimension $p + 1$ and above. The proof in [2] is detailed for the specific case $p = 0$, *i.e.*, for ∞ -groupoids, but it works equally well in the general case. Here we are interested in cofibrant replacements in the model structure on (∞, p) -categories, so let us examine the classes of weak equivalences and cofibrations.

From [25], we recall that an ∞ -functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is a *weak equivalence* in the model structure on ∞ -categories if, and only if, it satisfies the following properties:

- for every 0-cell x of \mathcal{D} , there exists a 0-cell \hat{x} of \mathcal{C} such that $F(\hat{x})$ is ω -equivalent to x ,
- for every 0-cells x and y of \mathcal{C} and every 1-cell $u : F(x) \rightarrow F(y)$ of \mathcal{D} , there exists a 1-cell $\hat{u} : x \rightarrow y$ in \mathcal{C} such that $F(\hat{u})$ is ω -equivalent to u ,
- for every parallel n -cells u and v of \mathcal{C} , with $n \geq 1$, and every $(n + 1)$ -cell $f : u \rightarrow v$ of \mathcal{D} , there exists an $(n + 1)$ -cell $\hat{f} : u \rightarrow v$ of \mathcal{C} such that $F(\hat{f})$ is ω -equivalent to f .

The ω -equivalence relation is defined together with the notion of reversible cells, by mutual coinduction:

- two n -cells u and v of an ω -category \mathcal{C} are *ω -equivalent* when there exists a reversible $(n + 1)$ -cell $f : u \rightarrow v$ in \mathcal{C} ,
- an $(n + 1)$ -cell $f : u \rightarrow v$ of an ω -category \mathcal{C} is *reversible* when there exists an $(n + 1)$ -cell $g : v \rightarrow u$ in \mathcal{C} such that $g \star_n f$ and $f \star_n g$ are ω -equivalent to 1_x and 1_y , respectively.

From the result of [2], the weak equivalences for (∞, p) -categories are the images through the forgetful functor \mathbb{U} of the weak equivalences for ∞ -categories, *i.e.*, the ∞ -functors between (∞, p) -categories that are weak equivalences for ∞ -categories.

In the canonical model structure on ∞ -categories, the cofibrations are the retracts of transfinite compositions of pushouts of the ∞ -functors

$$i_n : \partial\mathcal{E}_n \rightarrow \mathcal{E}_n,$$

for $n \geq 0$, where \mathcal{E}_n is the n -globe and $\partial\mathcal{E}_n$ its boundary:

- the n -globe \mathcal{E}_n is the n -category with exactly one n -cell together with its distinct k -source and k -target for every $0 \leq k < n$,
- the boundary $\partial\mathcal{E}_n$ of the n -globe is the same n -category as \mathcal{E}_n but with the n -cell removed.

2. Polygraphic resolutions

By the result of [2], we get that the cofibrations for (∞, p) -categories are the retracts of transfinite compositions of pushouts of the ∞ -functors

$$i_n^\top : \partial \mathcal{E}_n^\top \rightarrow \mathcal{E}_n^\top,$$

for $n \geq 0$, where \mathcal{E}_n^\top and $\partial \mathcal{E}_n^\top$ are obtained from \mathcal{E}_n and $\partial \mathcal{E}_n$ by formal adjunction of inverses for every k -cell, with $1 < k \leq n$.

2.2. Polygraphs

2.2.1. Cellular extensions. Let us assume that $n < \infty$ and let \mathcal{C} be an n -category. A *cellular extension* of \mathcal{C} is a set Γ equipped with a map ∂ from Γ to the set of n -spheres of \mathcal{C} . By considering all the formal compositions of elements of Γ , seen as $(n+1)$ -cells with source and target in \mathcal{C} , one builds the *free $(n+1)$ -category generated by Γ over \mathcal{C}* , denoted by $\mathcal{C}[\Gamma]$. The *size* of an $(n+1)$ -cell f of $\mathcal{C}[\Gamma]$ is the number of $(n+1)$ -cells of Γ it contains.

The *quotient of \mathcal{C} by Γ* , denoted by \mathcal{C}/Γ , is the n -category one gets from \mathcal{C} by identification of the n -cells $s(\gamma)$ and $t(\gamma)$, for every n -sphere γ of Γ . If \mathcal{C} is an (n, p) -category and Γ is a cellular extension of \mathcal{C} , then the *free $(n+1, p)$ -category generated by Γ over \mathcal{C}* is denoted by $\mathcal{C}(\Gamma)$ and defined as follows:

$$\mathcal{C}(\Gamma) = \mathcal{C}[\Gamma, \Gamma^-] / \text{Inv}(\Gamma)$$

where Γ^- contains the same $(n+1)$ -cells as Γ , with source and target reversed, and $\text{Inv}(\Gamma)$ is the cellular extension made of two $(n+2)$ -cells

$$\gamma \star_{n+1} \gamma^- \rightarrow 1_f \quad \text{and} \quad \gamma^- \star_{n+1} \gamma \rightarrow 1_g$$

for each $(n+1)$ -cell γ from f to g in Γ .

2.2.2. Homotopy bases. Let \mathcal{C} be an (n, p) -category, for $p < n < \infty$. A *homotopy basis* of \mathcal{C} is a cellular extension Γ of \mathcal{C} such that the (n, p) -category \mathcal{C}/Γ is aspherical, *i.e.*, such that, for every n -sphere γ of \mathcal{C} , there exists an $(n+1)$ -cell with boundary γ in the $(n+1, p)$ -category $\mathcal{C}(\Gamma)$. For example, the n -spheres of \mathcal{C} form a cellular extension which is a homotopy basis of \mathcal{C} .

2.2.3. (n, p) -polygraphs. An *n -polygraph* is a data Σ made of a set Σ_0 and, for every $0 \leq k < n$, a cellular extension Σ_{k+1} of the free k -category

$$\Sigma_k^* = \Sigma_0[\Sigma_1] \cdots [\Sigma_k].$$

For $p \leq n$, an (n, p) -polygraph is a data Σ made of:


- a p -polygraph $(\Sigma_0, \dots, \Sigma_p)$,
- for every $p \leq k < n$, a cellular extension Σ_{k+1} of the free (k, p) -category

$$\Sigma_k^\top = \Sigma_p^*(\Sigma_{p+1}) \cdots (\Sigma_k).$$

Note that (n, n) -polygraphs coincide with n -polygraphs, so that any notion defined on (n, p) -polygraphs also covers the case of n -polygraphs.

For an (n, p) -polygraph Σ , an element of Σ_k is called *k-cell* of Σ and Σ is *finite* when it has finitely many cells in every dimension. An (n, p) -polygraph Σ is *aspherical* when the free (n, p) -category Σ^\top is aspherical. An (n, p) -polygraph Σ is *acyclic* when, for every $p < k < n$, the cellular extension Σ_{k+1} is a homotopy basis of the (k, p) -category Σ_k^\top .

Remark. An (n, p) -polygraph yields a diagram which is similar to the one given in the original definition of n -polygraphs by Burroni, [11], drawn for the case $n < \infty$ as follows:



This diagram contains the source and target attachment maps of generating $(k + 1)$ -cells on composite k -cells, their extension to composite $(k + 1)$ -cells and the inclusion of generating k -cells into composite k -cells.

2.2.4. Proposition. *Every free (∞, p) -category on an (∞, p) -polygraph is a cofibrant object for the canonical model structure on (∞, p) -categories.*

Proof. Let Σ be an (∞, p) -polygraph. The unique ∞ -functor from the initial (∞, p) -category \emptyset to Σ^\top is obtained as the following countable composition of inclusions:

$$\emptyset \twoheadrightarrow \Sigma_0 \twoheadrightarrow \Sigma_1^* \twoheadrightarrow \dots \twoheadrightarrow \Sigma_p^* \twoheadrightarrow \Sigma_{p+1}^\top \twoheadrightarrow \dots \twoheadrightarrow \Sigma_n^\top \twoheadrightarrow \dots$$

The generating cofibration i_0^\top is the inclusion of $\partial \mathcal{E}_0^\top = \emptyset$ into the singleton \mathcal{E}_0^\top . Thus, for any set X , seen as a 0-polygraph, the inclusion $\emptyset \twoheadrightarrow X$ is equal to

$$\emptyset \simeq \bigsqcup_{x \in X} \partial \mathcal{E}_0^\top \xrightarrow{\bigsqcup_{x \in X} i_0^\top} \bigsqcup_{x \in X} \mathcal{E}_0^\top \simeq X.$$

Then, for $0 < k \leq p$, the inclusion of Σ_{k-1}^* into Σ_k^* is a particular case of an inclusion $\iota : \mathcal{C} \twoheadrightarrow \mathcal{C}[\Gamma]$ for \mathcal{C} a $(k - 1)$ -category and Γ a cellular extension of \mathcal{C} . By seeing each $(k - 1)$ -sphere γ of Γ as an ∞ -functor from $\partial \mathcal{E}_k^\top$ to \mathcal{C} , the inclusion ι is given by the following pushout:

$$\begin{array}{ccc} \bigsqcup_{\gamma \in \Gamma} \partial \mathcal{E}_k^\top & \xrightarrow{\bigsqcup_{\gamma \in \Gamma} i_k^\top} & \bigsqcup_{\gamma \in \Gamma} \mathcal{E}_k^\top \\ \Gamma \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{\iota} & \mathcal{C}[\Gamma]. \end{array}$$

2. Polygraphic resolutions

Finally, the inclusion of Σ_p^* into Σ_{p+1}^\top and, for $n > p$, the inclusion of Σ_n^\top into Σ_{n+1}^\top are particular cases of an inclusion $\iota : \mathcal{C} \hookrightarrow \mathcal{C}(\Gamma)$, for \mathcal{C} an (n, p) -category and Γ a cellular extension of \mathcal{C} . By seeing each n -sphere γ of Γ as an ∞ -functor from $\partial\mathcal{E}_{n+1}^\top$ to \mathcal{C} , the inclusion ι is given by the following pushout:

$$\begin{array}{ccc} \coprod_{\gamma \in \Gamma} \partial\mathcal{E}_{n+1}^\top & \xrightarrow{\coprod_{\gamma \in \Gamma} i_{n+1}^\top} & \coprod_{\gamma \in \Gamma} \mathcal{E}_{n+1}^\top \\ \Gamma \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{\iota} & \mathcal{C}(\Gamma). \end{array}$$

As a conclusion, we get that the inclusion $\emptyset \hookrightarrow \Sigma^\top$ is a countable composition of pushouts of the generating cofibrations $(i_n^\top)_{n \geq 0}$ and, as such, it is a cofibration. \square

2.3. Resolutions by (n, p) -polygraphs

2.3.1. Polygraphic presentations. If $p < n$, given an (n, p) -polygraph Σ , the p -category $\bar{\Sigma}$ presented by Σ is defined by

$$\bar{\Sigma} = \Sigma_p^* / \Sigma_{p+1}.$$

We usually denote by \bar{f} the image of a p -cell of Σ_p^* through the canonical projection $\Sigma_p^* \rightarrow \bar{\Sigma}$. If f is a k -cell of Σ^\top , with $p < k \leq n$, we also denote by \bar{f} the common image in $\bar{\Sigma}$ of the p -cells $s_p(f)$ and $t_p(f)$ by the canonical projection. An (m, p) -polygraph Σ and an (n, p) -polygraph Υ are *Tietze-equivalent* when the p -categories $\bar{\Sigma}$ and $\bar{\Upsilon}$ they present are isomorphic.


2.3.2. Example. Every category \mathbf{C} admits a presentation, called the *standard presentation of \mathbf{C}* , defined as the 2-polygraph whose cells are the following ones:

- one 0-cell for each 0-cell of \mathbf{C} ,
- one 1-cell $\hat{u} : x \rightarrow y$ for every 1-cell $u : x \rightarrow y$ of \mathbf{C} ,
- one 2-cell $\mu_{u,v} : \hat{u}\hat{v} \Rightarrow \widehat{uv}$ for every 1-cells $u : x \rightarrow y$ and $v : y \rightarrow z$ of \mathbf{C} and one 2-cell $\eta_x : 1_x \Rightarrow \hat{1}_x$ for every 0-cell x of \mathbf{C} :

$$\begin{array}{ccc} & y & \\ \hat{u} \nearrow & & \searrow \hat{v} \\ x & & z \\ & \Downarrow \mu_{u,v} & \\ & \widehat{uv} & \end{array} \qquad \begin{array}{ccc} & 1_x & \\ x & \xrightarrow{\quad} & x \\ & \Downarrow \eta_x & \\ & \hat{1}_x & \end{array}$$


In the free category generated by the 1-cells of the standard presentation of \mathbf{C} , we get, for every 0-cell x , the identity 1_x of x and the generating 1-cell $\hat{1}_x$ associated to the identity of x in \mathbf{C} . By removing this last superfluous generator, together with the corresponding 2-cell η_x , we get another presentation of \mathbf{C} , namely the 2-polygraph called the *reduced standard presentation of \mathbf{C}* , with the following cells:

- one 0-cell for each 0-cell of \mathbf{C} ,
- one 1-cell $\widehat{u} : x \rightarrow y$ for every non-identity 1-cell $u : x \rightarrow y$ of \mathbf{C} ,
- one 2-cell



for every non-identity 1-cells $u : x \rightarrow y$ and $v : y \rightarrow z$ of \mathbf{C} such that uv is not an identity,

- one 2-cell



for every non-identity 1-cells $u : x \rightarrow y$ and $v : y \rightarrow x$ of \mathbf{C} such that $uv = 1_x$.

2.3.3. Polygraphic resolutions. Let \mathcal{C} be a p -category. A *polygraphic resolution* of \mathcal{C} is an acyclic (∞, p) -polygraph Σ such that the p -category $\overline{\Sigma}$ is isomorphic to \mathcal{C} . If $p < n < \infty$, a *partial polygraphic resolution of length n* of \mathcal{C} is an acyclic (n, p) -polygraph Σ such that $\overline{\Sigma}$ is isomorphic to \mathcal{C} . Explicitly, the first dimensions of a polygraphic resolution Σ of \mathcal{C} are given as follows:

- For $k < p$, the k -cells of Σ are the ones of \mathcal{C} . In particular, polygraphic resolutions concern the cofibrant p -categories only, *i.e.*, the p -categories that are free up to dimension $p - 1$, which is always the case for $p = 1$.
- The p -cells of Σ are generators for the ones of \mathcal{C} , *i.e.*, the p -category \mathcal{C} is a quotient of the free p -category Σ_p^* .
- The $(p + 1)$ -cells of Σ are relations, *i.e.*, the $(p + 1)$ -polygraph Σ_{p+1} is a presentation of \mathcal{C} .
- The $(p + 2)$ -cells of Σ form a homotopy basis of Σ_{p+1}^\top , *i.e.*, they are generators of the relations between relations of the presentation Σ_{p+1} of \mathcal{C} .

As previously mentioned, Métayer introduced a notion of polygraphic resolution of a p -category \mathcal{C} , with $0 \leq p \leq \infty$, in [36]: this is an ∞ -polygraph Σ such that the free ∞ -category Σ^* is a cofibrant replacement of \mathcal{C} in the canonical model structure on ∞ -categories. The notion we use here is similar, using (∞, p) -polygraphs to produce cofibrant approximations of p -categories in the canonical model structure on (∞, p) -categories:

2.3.4. Theorem. *Let Σ be a polygraphic resolution of a p -category \mathcal{C} . The canonical projection $\Sigma^\top \twoheadrightarrow \mathcal{C}$ is a cofibrant approximation of \mathcal{C} in the canonical model structure on (∞, p) -categories.*

2. Polygraphic resolutions

Proof. From Proposition 2.2.4, we already know that Σ^\top is cofibrant. There remains to check that the canonical projection $\Sigma^\top \rightarrow \mathcal{C}$ is a weak equivalence. Since, by hypothesis, the p -categories \mathcal{C} and $\bar{\Sigma}$ are isomorphic, it is sufficient to prove that the canonical projection $\Sigma^\top \rightarrow \bar{\Sigma}$ is a weak equivalence. First, we note that the ω -equivalence relation is reflexive: hence, proving that two k -cells of $\bar{\Sigma}$ are equal implies that they are ω -equivalent.

By definition, the (∞, p) -categories Σ^\top and $\bar{\Sigma}$ have the same cells up to dimension $p - 1$. Thus, if x is a 0 -cell of $\bar{\Sigma}$, we take $\hat{x} = x$. Moreover, if x and y are parallel k -cells of Σ^\top , for $0 < k < p - 1$, and if $u : x \rightarrow y$ is a $(k + 1)$ -cell of $\bar{\Sigma}$, then we take $\hat{u} = u$.

Now, let x and y be parallel $(p - 1)$ -cells of Σ^\top and let $u : x \rightarrow y$ be a p -cell of $\bar{\Sigma}$. By definition, the p -category $\bar{\Sigma}$ is a quotient of the p -category underlying Σ^\top . Hence, there exists a p -cell \hat{u} in Σ^\top sent to u by the canonical projection.

Then, let u and v be parallel p -cells of Σ^\top and let $f : \bar{u} \rightarrow \bar{v}$ be a $(p + 1)$ -cell of $\bar{\Sigma}$. Since $\bar{\Sigma}$ is a p -category, we must have $\bar{u} = \bar{v}$ and $f = 1_{\bar{u}}$. By definition of $\bar{\Sigma}$, there exists a $(p + 1)$ -cell \hat{f} in Σ^\top from u to v , which is sent to $1_{\bar{u}}$ by the canonical projection.

Finally, let f and g be parallel n -cells of Σ^\top , for $n > p$. Both f and g must be sent to the same n -cell of $\bar{\Sigma}$, so that the only possible $(n + 1)$ -cell of $\bar{\Sigma}$ between their images is an identity. Since Σ is acyclic, there exists an $(n + 1)$ -cell from f to g in Σ^\top , and this $(n + 1)$ -cell must be sent to this same identity $(n + 1)$ -cell of $\bar{\Sigma}$. \square

2.3.5. Polygraphic dimension. Let us note that every cofibrant p -category \mathcal{C} admits a presentation, *i.e.*, a partial polygraphic resolution of length $p + 1$. Indeed, we can take:

- for $k < p$, any choice of generating k -cells of \mathcal{C} ,
- the same p -cells as \mathcal{C} ,
- one $(p + 1)$ -cell from u to v , when u and v are p -cells of the free p -category \mathcal{C}^* that are identified by the projection $\mathcal{C}^* \rightarrow \mathcal{C}$, *i.e.*, that yield the same p -cell of \mathcal{C} by composition.

Furthermore, every partial polygraphic resolution Σ of length n of a p -category \mathcal{C} can be extended into a partial polygraphic resolution of length $n + 1$ of \mathcal{C} by adjunction of the homotopy basis of the n -spheres of Σ^\top . By iterating this process, we can extend any partial polygraphic resolution into a polygraphic resolution of the same p -category. Applied to the generic presentation of a cofibrant p -category, we get that any cofibrant p -category admits a polygraphic resolution.

If \mathcal{C} is a cofibrant p -category, the *polygraphic dimension* of \mathcal{C} is the element $d_{\text{pol}}(\mathcal{C})$ of $\mathbb{N} \amalg \{\infty\}$ defined as follows: if there exists a natural number n such that \mathcal{C} admits an aspherical partial polygraphic resolution of length n , then $d_{\text{pol}}(\mathcal{C})$ is the smallest of those natural numbers; otherwise $d_{\text{pol}}(\mathcal{C}) = \infty$.

2.3.6. Higher-dimensional finite derivation type. For $n \geq p$, a p -category is of *finite n -derivation type* (FDT_n) when it admits a finite partial polygraphic resolution of length n . A p -category is of *finite ∞ -derivation type* (FDT_∞) when it admits a finite polygraphic resolution, *i.e.*, when it is FDT_n for every $n \geq p$. By extension, for $n < p$, a p -category is of *finite n -derivation type* when it admits finitely many generating n -cells.

In particular, a p -category is FDT_p when it is finitely generated, it is FDT_{p+1} when it is finitely presented and it is FDT_{p+2} when it has finite derivation type, a condition introduced by the authors in [18]. When $p = 1$ and for monoids, seen as categories with one 0 -cell, the property FDT_3 corresponds

to the finite derivation type condition originally defined by Squier, [41], while the property FDT_4 was introduced in [32].

Let us note that FDT_{n+1} is harder to fulfil than FDT_n , because of the finiteness condition on $(n + 1)$ -cells, leading to the following chain of implications:


$$FDT_\infty \Rightarrow (\dots) \Rightarrow FDT_{p+2} \Rightarrow FDT_{p+1} \Rightarrow FDT_p \Rightarrow (\dots) \Rightarrow FDT_0.$$

3. NORMALISATION STRATEGIES FOR POLYGRAPHS


3.1. Strategies in rewriting theory

A rewriting system specifies a set of rules that describe valid replacements of subformulas by other ones, [46, 38]. On some formulas, the rewriting rules may produce conflicts, when two or more rules can be applied. For this reason, to transform a rewriting system into a genuine computation algorithm, one specifies a way to apply the rules in a deterministic way by a *strategy*.

For example, in a word rewriting system, formulas are elements of a free monoid. There are two canonical strategies to reduce words where several rewriting rules apply: the leftmost one and the rightmost one, using first the rewriting rule that can be applied on the leftmost or the rightmost subformula:



In term rewriting, formulas are morphisms with target the terminal object in a free Lawvere algebraic theory, [26]. Formulas can be represented by trees and rewriting rules replace subtrees by other subtrees. There exist many possible strategies for term rewriting systems. Among them, outermost and innermost strategies are families of strategies that first use rules that apply closer to the root or closer to the leaves of the term, respectively:



In programming languages based on rewriting mechanisms, such as Caml, [27], and Haskell, [31], strategies are implicitly used by compilers to transform rewriting systems into deterministic algorithms. In that setting, innermost strategies include the call-by-value evaluation, while outermost strategies contain the call-by-need evaluation. Some programming languages, like Tom, [4], include a dedicated grammar to explicitly construct user-defined strategies.

Several models have been introduced to study the computational properties of strategies. In abstract rewriting, a strategy is defined as a subgraph of the ambient abstract rewriting system. This definition allows the introduction of some properties: for example, a normalisation strategy is a strategy that reaches

3. Normalisation strategies for polygraphs

normal forms, [45]. Strategies in functional programming languages are usually classified by corresponding notions of strategies in the λ -calculus, [29]. This has led to the axiomatic setting of standardisation theory, where strategies are seen as standardisation systems of rewriting paths, [33].

In this work, we introduce a notion of normalisation strategy for higher-dimensional rewriting systems that, in turn, induces a notion of normal forms in every dimension, together with a homotopically coherent reduction of every cell to its normal form.

3.2. Normalisation strategies

Before a formal definition of normalisation strategy, let us give the idea underlying this notion. If Σ is an (∞, p) -polygraph, the p -category $\bar{\Sigma}$ it presents can be seen as an (∞, p) -category with identity cells only in dimensions $p + 1$ and higher. This way, the canonical projection $\pi : \Sigma_p^* \twoheadrightarrow \bar{\Sigma}$ can be extended into an (∞, p) -functor $\pi : \Sigma^\top \twoheadrightarrow \bar{\Sigma}$. Given a (non-functorial) section $\iota : \bar{\Sigma} \rightarrow \Sigma_p^*$ of the canonical projection, a normalisation strategy corresponds to an extension of this section into a (∞, p) -functor $\iota : \bar{\Sigma} \rightarrow \Sigma^\top$, in a suitably weak sense, that satisfies $\pi\iota = \text{Id}_{\bar{\Sigma}}$ and $\iota\pi \simeq \text{Id}_{\Sigma^\top}$, with an explicitly chosen natural isomorphism witnessing this last fact: it follows that Σ is a polygraphic resolution of $\bar{\Sigma}$. Let us fix n and p with $0 \leq p \leq n \leq \infty$.

3.2.1. Sections. Let Σ be an (n, p) -polygraph. A *section* of Σ is a choice of a representative p -cell $\hat{u} : x \rightarrow y$ in Σ^\top , for every p -cell $u : x \rightarrow y$ of $\bar{\Sigma}$, such that

$$\hat{1}_x = 1_x$$

holds for every $(p - 1)$ -cell x of $\bar{\Sigma}$. Such an assignment $u \mapsto \hat{u}$ is not assumed to be functorial with respect to the compositions: in general, such a property can only be required for a (p, p) -polygraph, *i.e.*, when $\bar{\Sigma}$ is a free p -category.

Since, by hypothesis, the assignment $u \mapsto \hat{u}$ is compatible with the quotient map, it extends to a mapping of each p -cell u in Σ^* to a parallel p -cell in Σ^* , still denoted by \hat{u} , in such a way that the equality $\bar{u} = \bar{v}$ holds in $\bar{\Sigma}$ if, and only if, we have $\hat{u} = \hat{v}$ in Σ^* . Thereafter, we assume that every (n, p) -polygraph comes with an implicitly chosen section.

3.2.2. Normalisation strategies. Let Σ be an (n, p) -polygraph. A *normalisation strategy* for Σ is a mapping σ of every k -cell f of Σ^\top , with $p \leq k < n$, to a $(k + 1)$ -cell

$$f \xrightarrow{\sigma_f} \hat{f}$$

where, for $k > p$, the notation \hat{f} stands for the k -cell $\hat{f} = \sigma_{s(f)} \star_{k-1} \sigma_{t(f)}^-$, such that the following properties are satisfied:

- for every k -cell f , with $p \leq k < n$,

$$\sigma_{\hat{f}} = 1_{\hat{f}}$$

- for every pair (f, g) of i -composable k -cells, with $p \leq i < k < n$,

$$\sigma_{f \star_i g} = \sigma_f \star_i \sigma_g.$$

An (n, p) -polygraph is *normalising* when it admits a normalisation strategy. This property is independent of the chosen section. Indeed, let us consider an (n, p) -polygraph Σ with two sections $f \mapsto \widehat{f}$ and $f \mapsto \widetilde{f}$ of Σ and let us assume that σ is a normalisation strategy for Σ , equipped with the section $f \mapsto \widehat{f}$. Then, one checks that we get a normalisation strategy τ for the other section by defining τ_f as the following composite:

$$f \xrightarrow{\sigma_f} \widehat{f} \xrightarrow{(\sigma_f^-)^-} \widetilde{f}.$$

3.2.3. Lemma. *Let Σ be an (n, p) -polygraph and let σ be a normalisation strategy for Σ .*

i) *For every k -cell f , with $p - 1 \leq k < n - 1$, we have*

$$\sigma_{1_f} = 1_{1_f}.$$

ii) *For every k -cell f , with $p \leq k < n - 1$, we have*

$$\sigma_{\sigma_f} = 1_{\sigma_f}.$$

iii) *For every k -cell f , with $p < k < n$, we have*

$$\sigma_{f^-} = f^- \star_{k-1} \sigma_f^- \star_{k-1} \widehat{f}^-.$$

Proof. For **i)**, if x is a $(p - 1)$ -cell, we have $\widehat{1_x} = 1_x$ by definition. If f is a k -cell, with $p \leq k < n - 1$, then we have, by definition of $\widehat{1_f}$:

$$\widehat{1_f} = \sigma_{s(1_f)} \star_k \sigma_{t(1_f)}^- = \sigma_f \star_k \sigma_f^- = 1_f.$$

In either case, if f is a k -cell, with $p - 1 \leq k \leq n$, we get $\sigma_{1_f} = 1_{1_f}$ by definition of σ . For **ii)**, if f is a k -cell, with $p \leq k < n - 1$, then the definition of $\widehat{\sigma_f}$ gives:

$$\widehat{\sigma_f} = \sigma_{s(\sigma_f)} \star_k \sigma_{t(\sigma_f)}^- = \sigma_f \star_k \sigma_f^- = \sigma_f \star_k 1_f^- = \sigma_f.$$

As a consequence, we get $\sigma_{\sigma_f} = 1_{\sigma_f}$. Finally, for **iii)**, if f is a k -cell, with $p < k < n$, we have:

$$\sigma_f \star_{k-1} \sigma_{f^-} = \sigma_{f \star_{k-1} f^-} = \sigma_{1_{s(f)}} = 1_{1_{s(f)}}.$$

Thus, σ_{f^-} is the $(k - 1)$ -inverse of σ_f , yielding:

$$\sigma_{f^-} = s(\sigma_f)^- \star_{k-1} \sigma_f^- \star_{k-1} t(\sigma_f)^- = f^- \star_{k-1} \sigma_f^- \star_{k-1} \widehat{f}^-.$$

□

3.3. The case of $(n, 1)$ -polygraphs

Let Σ be an $(n, 1)$ -polygraph. In the lower dimensions, a normalisation strategy σ for Σ specifies the following assignments:


- For every 1-cell u of Σ^\top , a 2-cell

$$u \xrightarrow{\sigma_u} \widehat{u}$$

of Σ^\top that satisfies $\sigma_{\widehat{u}} = 1_{\widehat{u}}$ and thus, in particular, $\sigma_{1_x} = 1_{1_x}$ for every 0-cell x of Σ .


3. Normalisation strategies for polygraphs

- For every 2-cell $f : u \Rightarrow v$ of Σ^\top , a 3-cell




of Σ^\top that satisfies $\sigma_{\hat{f}} = 1_{\hat{f}}$ and the following relations:


- If u is a 1-cell of Σ^\top , then $\sigma_{1_u} = 1_{1_u}$:




- If $f : u \Rightarrow v$ and $g : v \Rightarrow w$ are 2-cells in Σ^\top , then $\sigma_{f \star_1 g} = \sigma_f \star_1 \sigma_g$:



- If $f : u \Rightarrow v$ is a 2-cell in Σ^\top , then $\hat{f}^- = \sigma_v \star_1 \sigma_u^-$ and $\sigma_{f^-} = f^- \star_1 \sigma_f^- \star_1 \hat{f}^-$:




- For every 3-cell $A : f \Rightarrow g : u \Rightarrow v$ of Σ^\top , a 4-cell




of Σ^\top with $\sigma_{\hat{A}} = 1_{\hat{A}}$ and such that the following relations hold:


– If f is a 2-cell of Σ^\top , then $\sigma_{1_f} = 1_{1_f}$:




– If $A : f \Rightarrow f' : u \Rightarrow v$ and $B : g \Rightarrow g' : v \Rightarrow w$ are 3-cells of Σ^\top , then $\sigma_{A \star_1 B} = \sigma_A \star_1 \sigma_B$:



– If $A : f \Rightarrow g : u \Rightarrow v$ and $B : g \Rightarrow h : u \Rightarrow v$ are 3-cells of Σ^\top , then $\sigma_{A \star_2 B} = \sigma_A \star_2 \sigma_B$:



– If $A : f \Rightarrow g : u \Rightarrow v$ is a 3-cell of Σ^\top , then $\hat{A} = \sigma_f \star_2 \sigma_g^-$ and $\sigma_{A^-} = A^- \star_2 \sigma_A^- \star_2 \hat{A}^-$:



3.3.1. Lemma. Let Σ be an $(n, 1)$ -polygraph. Normalisation strategies for Σ are in bijective correspondence with data made of:

- a family with one 2-cell

$$\sigma_u : u \Rightarrow \hat{u}$$

for every 1-cell u of Σ^\top such that $\hat{u} \neq u$,

3. Normalisation strategies for polygraphs

- a family with one $(k + 1)$ -cell

$$\sigma_{u\varphi v} : u\varphi v \rightarrow \widehat{u\varphi v}$$

for every $1 < k < n$, every k -cell φ of Σ and every pair (u, v) of 1-cells of Σ^\top such that the composite k -cell $u\varphi v$ is defined.

Proof. We proceed by induction on the size of cells of Σ^\top . We already know that a normalisation strategy σ has fixed values on normal forms, identities, inverses and i -composites for $i \geq 1$. As a consequence, using the exchange relations, we get that the values of σ are entirely and uniquely determined by its values on 1-cells that are not normal forms and, for every $k \geq 2$, on k -cells with shape $u\varphi v$, where φ is a k -cell of Σ and u and v are 1-cells of Σ^\top . \square

3.3.2. From normalisation strategies to natural transformations. Let σ be a normalisation strategy for an $(n, 1)$ -polygraph Σ . We define, for every 1-cell u of Σ^\top , the 1-cell u^* as u and, by induction on the dimension, for every k -cell f in Σ^\top , with $1 < k \leq n$, the k -cell f^* in Σ^\top is given by

$$f^* = ((f \star_1 \sigma_{t_1(f)^*}) \star_2 \cdots) \star_{k-1} \sigma_{t_{k-1}(f)^*} \cdot$$

For example, for a 2-cell $f : u \Rightarrow v$, the 2-cell f^* is

$$u \xrightarrow{f} v \xrightarrow{\sigma_v} \widehat{u}$$

and, for a 3-cell $A : f \Rightarrow g : u \Rightarrow v$, the 3-cell A^* is

One checks that, for any k -cell f , with $k > 1$, the k -cell f^* has source $s(f)^*$ and target $t(f)^*$. Moreover, we have $(\widehat{f})^* = f^*$, which implies $\sigma_{f^*} = \sigma_f^*$.

Since every k -cell of Σ^\top is invertible for $k \geq 2$, one can recover σ from σ^* , in a unique way, so that the normalisation strategy σ is uniquely and entirely determined by the values

$$\sigma_u^* = \sigma_u : u \Rightarrow \widehat{u}$$

for every 1-cell u with $u \neq \widehat{u}$ and


$$\sigma_{u\varphi v}^* : (u\varphi v)^* \rightarrow \widehat{u\varphi v}^*$$

for every $1 < k < n$, every k -cell φ of Σ and every 1-cells u and v of Σ^\top such that the k -cell $u\varphi v$ is defined. In the lowest dimensions, the natural transformation form σ^* of the strategy σ consists of the following data:


- For every 1-cell u of Σ^\top , a 2-cell $\sigma_u^* = \sigma_u$ from u to \widehat{u} .

3.3. The case of $(n, 1)$ -polygraphs

- For every 2-cell $f : u \Rightarrow v$ of Σ^\top , a 3-cell σ_f^* of Σ^\top corresponding to a (weak) naturality condition:




To simplify subsequent diagrams, we draw the 3-cell σ_f^* in a more compact shape, as follows:




This 3-cell must satisfy the following relations:


- if u is a 1-cell of Σ^\top , then $\sigma_{1_u}^* = 1_{\sigma_u}$ holds:




- if $f : u \Rightarrow v$ and $g : v \Rightarrow w$ are 2-cells of Σ^\top , then $\sigma_{f \star_1 g}^* = (f \star_1 \sigma_g^*) \star_2 \sigma_f^*$ holds:



- if $f : u \Rightarrow v$ is a 2-cell of Σ^\top , then $\sigma_{f^-}^* = f^- \star_1 (\sigma_f^*)^-$ holds:




- For every 3-cell $A : f \Rrightarrow g : u \Rightarrow v$ of Σ^\top , a 4-cell




3. Normalisation strategies for polygraphs

of Σ^\top such that the following relations are satisfied:


- if f is a 2-cell of Σ^\top , then $\sigma_{1_f}^* = 1_{\sigma_f^*}$ holds:




- if $A : f \Rightarrow f' : u \Rightarrow v$ and $B : g \Rightarrow g' : v \Rightarrow w$ are 3-cells of Σ^\top , then $\sigma_{A \star_1 B}^* = (f \star_1 \sigma_B^*) \star_2 \sigma_A^*$ holds:



- if $A : f \Rightarrow g : u \Rightarrow v$ and $B : g \Rightarrow h : u \Rightarrow v$ are 3-cells of Σ^\top , then $\sigma_{A \star_2 B}^* = ((A \star_1 \sigma_v) \star_2 \sigma_B^*) \star_3 \sigma_A^*$ holds:




- if $A : f \Rightarrow g : u \Rightarrow v$ is a 3-cell of Σ^\top , then $\sigma_{A^-}^* = (A^- \star_1 \sigma_v^*) \star_2 (\sigma_A^*)^-$ holds:



3.3.3. Left and right normalisation strategies. Let Σ be an $(n, 1)$ -polygraph. A normalisation strategy σ for Σ is a *left* one when it satisfies the following properties:


- For every pair (u, v) of 0-composable 1-cells of Σ^\top , we have $\sigma_{uv} = \sigma_{uv} \star_1 \sigma_{\widehat{uv}}$, *i.e.*,



- For every pair (f, g) of 0-composable k -cells of Σ^\top , with $2 \leq k \leq p$, $t_1(f) = u'$ and $s_1(g) = v$, we have

$$\sigma_{fg} = \sigma_{fv} \star_1 \sigma_{u'g}.$$

In particular, when $f : u \Rightarrow u'$ and $g : v \Rightarrow v'$ are 0-composable 2-cells of Σ^\top :



In a symmetric way, a normalisation strategy σ is a *right* one when it satisfies:

$$\sigma_{uv} = u\sigma_v \star_1 \sigma_{u\widehat{v}} \quad \text{and} \quad \sigma_{fg} = u\sigma_g \star_1 \sigma_{fv'}.$$

An $(n, 1)$ -polygraph is *left* (resp. *right*) *normalising* when it admits a left (resp. right) normalisation strategy.

3.3.4. Lemma. Let Σ be an $(n, 1)$ -polygraph, let f be a k -cell of Σ^\top , for $1 < k < n$, with 1-source u and 1-target v , and let w and w' be 1-cells of Σ^\top such that wfw' is defined. Then, if σ is a left normalisation strategy for Σ , we have:

$$\sigma_{wfw'} = \sigma_w u w' \star_1 \sigma_{\widehat{w}f} w' \star_1 \sigma_w^- v w' \quad \text{and} \quad \sigma_{wfw'}^* = \sigma_w u w' \star_1 \sigma_{\widehat{w}f}^* w' \star_1 \sigma_{w\widehat{u}w'}.$$

Symmetrically, if σ is a right normalisation strategy, then we have:

$$\sigma_{wfw'} = w u \sigma_{w'} \star_1 w \sigma_{f\widehat{w}'} \star_1 w v \sigma_{w'}^-, \quad \text{and} \quad \sigma_{wfw'}^* = w u \sigma_{w'} \star_1 w \sigma_{f\widehat{w}'}^* \star_1 \sigma_{w\widehat{u}w'}.$$

Proof. In the case of a left normalisation strategy, the proof for right normalisation strategies being symmetric, we have:

$$\sigma_{fw'} = \sigma_f w' \star_1 1_{1_{wvw'}} = \sigma_f w' \star_1 1_{1_{wvw'}} = \sigma_f w'.$$

3. Normalisation strategies for polygraphs

Then, we use the exchange relation to get:

$$\sigma_{\sigma_w f} = \sigma_{w f \star_1 \sigma_w v} = \sigma_{w f} \star_1 \sigma_{\sigma_w v} = \sigma_{w f} \star_1 1_{\sigma_w v} = \sigma_{w f} \star_1 \sigma_w v.$$

Moreover, the definition of left normalisation strategy implies:

$$\sigma_{\sigma_w f} = \sigma_{\sigma_w u} \star_1 \sigma_{\widehat{w} f} = \sigma_w u \star_1 \sigma_{\widehat{w} f}.$$

From the last two computations, we deduce:

$$\sigma_{w f} = \sigma_w u \star_1 \sigma_{\widehat{w} f} \star_1 \sigma_w^{-1} v.$$

Combining all the results, we get the required equality:

$$\sigma_{w f w'} = \sigma_{w f} w' = \sigma_w u w' \star_1 \sigma_{\widehat{w} f} w' \star_1 \sigma_w^{-1} v w'.$$

For $\sigma_{w f w'}^*$, we proceed as follows:

$$\begin{aligned} \sigma_{w f w'}^* &= \sigma_{(w f w')^*} \\ &= \sigma_{w f \star_1 w'} \star_1 \sigma_{w v w'} \\ &= \sigma_w u w' \star_1 \sigma_{\widehat{w} f \star_1 w'} \star_1 \sigma_w^{-1} v w' \star_1 \sigma_{w v w'} \\ &= \sigma_w u w' \star_1 \sigma_{\widehat{w} f \star_1 w'} \star_1 \sigma_{\widehat{w} v} w' \star_1 \sigma_{\widehat{w} u w'} \\ &= \sigma_w^* u w' \star_1 \sigma_{\widehat{w} f w'}^* \star_1 \sigma_{\widehat{w} u w'}^*. \end{aligned} \quad \square$$

3.3.5. Corollary. *Let Σ be an $(n, 1)$ -polygraph. Left (resp. right) normalisation strategies on Σ are in bijective correspondence with the families*

$$\sigma_{\widehat{u}\varphi} : \widehat{u}\varphi \rightarrow \widehat{u}\widehat{\varphi} \quad (\text{resp. } \sigma_{\varphi\widehat{u}} : \varphi\widehat{u} \rightarrow \widehat{\varphi}\widehat{u})$$

and with the families

$$\sigma_{\widehat{u}\varphi}^* : (\widehat{u}\varphi)^* \rightarrow \widehat{\varphi}\widehat{u}^* \quad (\text{resp. } \sigma_{\varphi\widehat{u}}^* : (\varphi\widehat{u})^* \rightarrow \widehat{\varphi}\widehat{u}^*)$$

of $(k+1)$ -cells, indexed by k -cells φ of Σ , for $1 \leq k < n$, and by 1-cells u of $\overline{\Sigma}$ such that the composite k -cell $\widehat{u}\varphi$ (resp. $\varphi\widehat{u}$) exists.

Proof. Let us assume, for example, that σ is a left normalisation strategy. The property satisfied by σ on 1-cells of Σ^\top gives, by induction on the size of 1-cells, that the values of σ on 1-cells of Σ^\top are determined by the 2-cells $\sigma_{\widehat{u}\chi}$, for χ a 1-cell of Σ and u a 1-cell of $\overline{\Sigma}$ such that $\widehat{u}\chi$ is defined. Then, Lemma 3.3.1 tells us that the values of σ on higher-dimensional cells of Σ^\top are determined by the values of σ or of σ^* on k -cells $u\varphi v$ of Σ^\top , where φ is a k -cell of Σ and u and v are 1-cells of Σ^\top . We use Lemma 3.3.4 to conclude. \square

3.3.6. Theorem. *Let Σ be an $(n, 1)$ -polygraph. The following assertions are equivalent:*

- i) Σ is acyclic,
- ii) Σ is normalising,

4. Polygraphic resolutions from convergent presentations

iii) Σ is left normalising,

iv) Σ is right normalising.

Proof. Let us assume that there exists a normalisation strategy σ for Σ . We consider a k -cell f in Σ^\top , for some $1 < k < n$. By definition of a normalisation strategy, the $(k+1)$ -cell σ_f has source f and target \widehat{f} . Thus, if g is a k -cell which is parallel to f , the $(k+1)$ -cell $\sigma_f \star_k \sigma_g^-$ of Σ^\top has source f and target g , proving that Σ_{k+1} forms a homotopy basis of Σ_k^\top . Hence Σ is acyclic.

Conversely, let us assume that Σ is acyclic and let us define a right normalisation strategy σ , the case of a left one being symmetric. By definition of the category $\bar{\Sigma}$, we can choose a 2-cell

$$\sigma_{x\widehat{u}} : x\widehat{u} \Rightarrow \widehat{xu}$$

for every 1-cell x in Σ and every 1-cell u in $\bar{\Sigma}$ such that $x\widehat{u}$ is defined. Then, for $1 < k < n$, we use the fact that Σ_{k+1} is a homotopy basis of Σ_k^\top to choose an arbitrary $(k+1)$ -cell

$$\sigma_{\varphi\widehat{u}} : \varphi\widehat{u} \longrightarrow \widehat{\varphi u}$$

for every k -cell φ in Σ and every 1-cell u in $\bar{\Sigma}$ with $\varphi\widehat{u}$ is defined. We use Corollary 3.3.5 to conclude. \square

3.3.7. Corollary. *Let \mathbf{C} be a category and let n be a non-zero natural number. Then \mathbf{C} is FDT_n if, and only if, there exists a finite, (left, right) normalising $(n, 1)$ -polygraph presenting \mathbf{C} .*

4. POLYGRAPHIC RESOLUTIONS FROM CONVERGENT PRESENTATIONS

4.1. Convergent 2-polygraphs

Let us recall notions and results from rewriting theory for 2-polygraphs, [17, 18, 34]. Let Σ be a fixed 2-polygraph.

4.1.1. Rewriting and normal forms. A *rewriting step* of Σ is a 2-cell of the free 2-category Σ^* with shape

$$y \xrightarrow{w} x \begin{array}{c} \xrightarrow{u} \\ \Downarrow \varphi \\ \xrightarrow{v} \end{array} x' \xrightarrow{w'} y'$$

where $\varphi : u \Rightarrow v$ is a 2-cell of Σ and w and w' are 1-cells of Σ^* . A *rewriting sequence* of Σ is a finite or infinite sequence


$$u_1 \xrightarrow{f_1} u_2 \xrightarrow{f_2} (\dots) \xrightarrow{f_{n-1}} u_n \xrightarrow{f_n} (\dots)$$

of rewriting steps. If Σ has a non-empty rewriting sequence from u to v , we say that u *rewrites into* v . Let us note that every 2-cell f of Σ^* decomposes into a finite rewriting sequence of Σ , this decomposition being unique up to exchange relations.

A 1-cell u of Σ^* is a *normal form* when Σ has no rewriting step with source u . A *normal form* of u is a 1-cell v that is a normal form and such that u rewrites into v . A 1-cell is *reducible* if it is not a normal form.

4. Polygraphic resolutions from convergent presentations


4.1.2. Branchings. A *branching* of Σ is a pair (f, g) of 2-cells of Σ^* with a common source, as in the diagram



The 1-cell u is the *source* of this branching and the pair (v, w) is its *target*, written $(f, g) : u \Rightarrow (v, w)$. We do not distinguish the branchings (f, g) and (g, f) .


A branching (f, g) is *local* when f and g are rewriting steps. Local branchings belong to one of the three following families:

- *aspherical* branchings have shape



with $f : u \Rightarrow v$ a rewriting step of Σ ,

- *Peiffer* branchings have shape



with $f : u \Rightarrow v$ and $g : u' \Rightarrow v'$ rewriting steps of Σ ,

- *overlapping* branchings are the remaining local branchings.


The terms “aspherical” and “Peiffer” come from the corresponding notions for spherical diagrams in Cayley complexes associated to presentations of groups, [30], while the term “critical” comes from rewriting theory, [8, 3].

Local branchings are compared by the order \preceq generated by the relations

$$(f, g) \preceq (ufv, ugv)$$

given for any local branching (f, g) and any 1-cells u and v of Σ^* such that ufv exists (and, thus, so does ugv). An overlapping local branching that is minimal for the order \preceq is called a *critical branching*.

A branching (f, g) is *confluent* when there exists a pair (f', g') of 2-cells of Σ^* with the following shape:



4.1.3. Termination, confluence and convergence. We say that Σ *terminates* when it has no infinite rewriting sequence. In that case, every 1-cell has at least one normal form. Moreover, *Noetherian induction* allows definitions and proofs of properties of 1-cells by induction on the maximum size of the 2-cells leading to normal forms.

We say that Σ is *confluent* (resp. *locally confluent*) when all of its branchings (resp. local branching) are confluent. In a confluent 2-polygraph, every 1-cell has at most one normal form. A fundamental result of rewriting theory is that local confluence is equivalent to confluence of critical branchings. For terminating 2-polygraphs, Newman's Lemma ensures that local confluence and confluence are equivalent properties, [38].

We say that Σ is *convergent* when it terminates and it is confluent. In that case, every 1-cell u has a unique normal form. Such a Σ is called a *convergent presentation of $\bar{\Sigma}$* and has a canonical section sending u to its normal form \hat{u} , so that $\hat{u} = \hat{v}$ holds in Σ^* if, and only if, we have $\bar{u} = \bar{v}$ in $\bar{\Sigma}$. As a consequence, a finite and convergent 2-polygraph Σ yields generators for the 1-cells of the category $\bar{\Sigma}$, together with a decision procedure for the corresponding word problem (finiteness is used to effectively check that a given 1-cell is a normal form).

4.1.4. Reduced 2-polygraphs. A 2-polygraph Σ is *reduced* when, for every 2-cell $\varphi : u \Rightarrow v$ in Σ , the 1-cell u is a normal form for $\Sigma_2 \setminus \{\varphi\}$ and v is a normal form for Σ_2 . Let us note that, in that case, for every 1-cell u of Σ^* , there exist finitely many rewriting steps with source u in Σ^* : indeed, we have exactly one such 2-cell for every decomposition $u = vwv'$ such that w is the source of a 2-cell of Σ and the number of decompositions $u = vwv'$ is finite in a free category.

4.1.5. Lemma. *For every (finite) convergent 2-polygraph, there exists a (finite) Tietze-equivalent, reduced and convergent 2-polygraph.*

Proof. Let Σ be a (finite) convergent 2-polygraph Σ . We successively transform Σ as follows. First, we replace every 2-cell $\varphi : u \Rightarrow v$ in Σ with $\varphi' : u \Rightarrow \hat{u}$. Then, if there exist several 2-cells in Σ with the same source, we drop all of them but one. Finally, we drop all the remaining 2-cells whose source is reducible by another 2-cell. After each step, we check that the (finite) 2-polygraph we get is convergent and that it is Tietze-equivalent to the former one. Moreover, by construction, the result is a reduced 2-polygraph. \square

Remark. This result was proved by Métivier for term rewriting systems, [37], and by Squier for word rewriting systems, [41]. The proof works for any type of rewriting systems, including n -polygraphs for any n .

4.1.6. Example. Let \mathbf{C} be a category and let $N\mathbf{C}$ be the reduced standard polygraphic presentation of \mathbf{C} , i.e., the 2-polygraph with the following cells:

- one 0-cell for each 0-cell of \mathbf{C} ,
- one 1-cell $\hat{u} : x \rightarrow y$ for every non-identity 1-cell $u : x \rightarrow y$ of \mathbf{C} ,
- one 2-cell $\mu_{u,v} : \hat{u}\hat{v} \Rightarrow \widehat{uv}$ for every non-identity 1-cells $u : x \rightarrow y$ and $v : y \rightarrow z$ of \mathbf{C} such that uv is not an identity,
- one 2-cell $\mu_{u,v} : \hat{u}\hat{v} \Rightarrow 1_x$ for every non-identity 1-cells $u : x \rightarrow y$ and $v : y \rightarrow x$ of \mathbf{C} such that $uv = 1_x$.

4. Polygraphic resolutions from convergent presentations


The 2-polygraph \mathbf{NC} is reduced. Let us prove that it is convergent. For termination, one checks that each 2-cell $\mu_{u,v}$ of \mathbf{NC} has source of size 2 and target of size 1 or 0. As a consequence, for every non-identity 2-cell $f : u \Rightarrow v$ of the free 2-category \mathbf{NC}^* , the size of u is strictly greater than the size of v .

For confluence, we check that \mathbf{NC} has one critical branching for every triple (u, v, w) of non-identity composable 1-cells in \mathbf{C} :


$$(\mu_{u,v}\widehat{w}, \widehat{u}\mu_{v,w}).$$

Each of these critical branchings is confluent, with four possible cases, depending on whether uv or vw is an identity or not:


- if neither uv nor vw is an identity:




- if uv is an identity, but not vw :



- if uv is not an identity, but vw is:



- if uv and vw are identities, and thus $u = uvw = w$:



As a conclusion, the reduced standard presentation \mathbf{NC} of the category \mathbf{C} is a reduced convergent presentation of \mathbf{C} .

4.2. Normalisation strategies for convergent 2-polygraphs

4.2.1. The order relation on branchings. Let Σ be a reduced 2-polygraph and let u be a 1-cell in Σ^* . We define the relation \preceq on rewriting steps with source u as follows. If φ and ψ are 2-cells of Σ and if $f = v\varphi v'$ and $g = w\psi w'$ have source u , then we write $f \preceq g$ when v is smaller than w , *i.e.*, informally, when the part of u on which f acts is more at the left than the part on which g acts. By convention, we denote branchings of Σ in increasing order, *i.e.*, (f, g) when $f \preceq g$, which is always possible thanks to the following result.

4.2.2. Lemma. *Let Σ be a reduced 2-polygraph and u be a 1-cell of Σ^* . Then the relation \preceq induces a total ordering on the rewriting steps of Σ with source u .*

Proof. From its definition, we already know that the relation \preceq is reflexive, transitive and total. For antisymmetry, we assume that $f = v\varphi v'$ and $g = w\psi w'$ are rewriting steps with source u , such that $f \preceq g$ and $g \preceq f$, *i.e.*, such that v and w have the same size. Then, using the fact that Σ_1^* is free, we have $v = w$ and either $s(\varphi) = s(\psi)$ or $s(\varphi) = s(\psi)\alpha$ or $s(\varphi)\alpha = s(\psi)$: the latter two cases cannot occur, because Σ is reduced and, from that same hypothesis we get, in the first case, that $\varphi = \psi$, hence that $f = g$. \square

4.2.3. The leftmost and rightmost normalisation strategies. Let Σ be a reduced 2-polygraph and let u be a reducible 1-cell of Σ^* that is not a normal form. The *leftmost* and the *rightmost* rewriting steps on u are denoted by λ_u and ρ_u and defined as the minimum and the maximum elements for \preceq of the (finite, non-empty) set of rewriting steps of Σ with source u . We note that, if u and v are reducible and composable 1-cells of Σ^* , then we have:

$$\lambda_{uv} = \lambda_u v \quad \text{and} \quad \rho_{uv} = u \rho_v.$$

When Σ terminates, the *leftmost normalisation strategy* of Σ is the normalisation strategy σ defined by Noetherian induction as follows. On normal forms, it is given by

$$\sigma_{\hat{u}} = 1_{\hat{u}}$$

and, on reducible 1-cells, by

$$\sigma_u = \lambda_u \star_1 \sigma_{t(\lambda_u)}.$$

One defines the *rightmost normal form* of Σ in a similar way by replacing the leftmost rewriting step by the rightmost one in the case of reducible 1-cells.

4.2.4. Lemma. *The leftmost (resp. rightmost) normalisation strategy σ is a left (resp. right) normalisation strategy for Σ , seen as a $(2, 1)$ -polygraph, with the property that, for every 1-cell u , the 2-cell σ_u lives in $\Sigma^* \subseteq \Sigma^\top$.*

Proof. Let us assume that σ is the leftmost normalisation strategy, the proof in the rightmost case being symmetric. We must prove that, for every composable 1-cells u and v of Σ^* , the following relation holds:

$$\sigma_{uv} = \sigma_u v \star_1 \sigma_{\hat{uv}}.$$

4. Polygraphic resolutions from convergent presentations

We proceed by Noetherian induction on the 1-cell u . If u is a normal form, then $\sigma_u = 1_u$ and $\sigma_{\widehat{uv}} = \sigma_{uv}$, so that the relation is satisfied. Otherwise, we have, using the definition of σ and the properties of λ :

$$\sigma_{uv} = \lambda_{uv} \star_1 \sigma_{t(\lambda_{uv})} = \lambda_{uv} \star_1 \sigma_{t(\lambda_u)v}.$$

We apply the induction hypothesis to $t(\lambda_u)v$ to get:

$$\sigma_{uv} = \lambda_{uv} \star_1 \sigma_{t(\lambda_u)v} \star_1 \sigma_{\widehat{uv}} = \sigma_u v \star_1 \sigma_{\widehat{uv}}.$$

The fact that σ_u is in Σ^* is also proved by Noetherian induction on u , using the definition of σ and the facts that both $1_{\widehat{u}}$ and λ_u are 2-cells of Σ^* . \square

Remark. A reduced and terminating 2-polygraph can have several left or right strategies, beside the leftmost and the rightmost ones. Indeed, let us consider the reduced and terminating 2-polygraph Σ with one 0-cell, three 1-cells a , b and c and the following three 2-cells:

$$aac \xrightarrow{\alpha} a \quad bb \xrightarrow{\beta} cc \quad acc \xrightarrow{\gamma} c.$$

Let us prove that Σ admits at least two different left normalisation strategies. For that, we examine the 1-cell $aabb$ and all the 2-cells of Σ^* from $aabb$ to its normal form ac :

$$aabb \xrightarrow{aa\beta} aacc \begin{array}{c} \xrightarrow{\alpha c} \\ \xleftarrow{a\gamma} \end{array} ac.$$

Thus, if σ is a normalisation strategy, the 2-cell σ_{aabb} can be either $aa\beta \star_1 \alpha c$ or $aa\beta \star_1 a\gamma$. Since the 1-cells a , aa and aab are normal forms, assuming that σ is a left strategy still leaves us with the same choice. Hence, we can define a left normalisation strategy σ for Σ as the leftmost normalisation strategy on every 1-cell of Σ^* , except for $aabb$ where it is given by

$$\sigma_{aabb} = aa\beta \star_1 a\gamma.$$

Thus, we have a left normalisation strategy for Σ , distinct from the leftmost normalisation strategy: indeed, the latter would send $aabb$ to $aa\beta \star_1 \alpha c$.

Let us note that this phenomenon does not come from the fact that Σ is not confluent, since we can add the 2-cell $\delta : bcc \Rightarrow ccb$ to Σ to get a reduced, convergent 2-polygraph which still has at least two different left normalisation strategies. From Σ , we can build a symmetric (for \star_0) 2-polygraph that admits at least two different right normalisation strategies.

However, we can ensure that, if σ is a left (resp. right) normalisation strategy for a reduced and terminating 2-polygraph Σ with the property that, for every 1-cell u of Σ^* , the 2-cell σ_u is in Σ^* , then this same 2-cell admits a decomposition

$$\sigma_u = \lambda_u \star_1 g_u \quad (\text{resp. } \sigma_u = \rho_u \star_1 g_u)$$

with g_u a 2-cell of Σ^* . Indeed, if σ is a left strategy, we consider the decomposition $\lambda_u = v\varphi w$. By definition of λ_u , the 1-cell $vs(\varphi)$ is the source of only one rewriting step of Σ , namely $v\varphi$. Hence, since $\sigma_{vs(\varphi)}$ is a 2-cell of Σ^* with source $vs(\varphi)$, it admits a decomposition

$$\sigma_{vs(\varphi)} = \lambda_{vs(\varphi)} \star_1 h_u$$

4.3. The acyclic (3, 1)-polygraph of generating confluences

with h_u a 2-cell of Σ^* . The 2-cell g_u of Σ^* is given by

$$g_u = h_u \mathcal{W} \star_1 \sigma_{\widehat{vs(\varphi)\mathcal{W}}}$$

and use the hypothesis on σ to get:

$$\sigma_u = \sigma_{vs(\varphi)\mathcal{W}} \star_1 \sigma_{\widehat{vs(\varphi)\mathcal{W}}} = \lambda_{vs(\varphi)\mathcal{W}} \star_1 g_u = \lambda_u \star_1 g_u.$$

The case of a right normalisation strategy is symmetric.

4.2.5. Example. Let \mathbf{C} be a category and let $N\mathbf{C}$ be its reduced standard presentation. A generic 1-cell of $N\mathbf{C}$ is a composite $\widehat{u}_1 \cdots \widehat{u}_n$ of non-identity 1-cells of \mathbf{C} . In the case where no partial composition $u_i u_{i+1} \cdots u_j$ is an identity in \mathbf{C} , the leftmost reduction strategy σ of \mathbf{C} is given on $\widehat{u}_1 \cdots \widehat{u}_n$ by:

$$\sigma_{\widehat{u}_1 \cdots \widehat{u}_n} = \mu_{u_1, u_2} \widehat{u}_3 \cdots \widehat{u}_n \star_1 \mu_{u_1 u_2, u_3} \widehat{u}_4 \cdots \widehat{u}_n \star_1 \cdots \star_1 \mu_{u_1 \cdots u_{n-1}, u_n}.$$

If some partial composition is an identity in \mathbf{C} , we get $\sigma_{\widehat{u}_1 \cdots \widehat{u}_n}$ by removing the corresponding 2-cell μ . For example, in the case $n = 4$, $u_1 u_2 = 1$ and $u_3 u_4 \neq 1$, we have:

$$\sigma_{\widehat{u}_1 \widehat{u}_2 \widehat{u}_3 \widehat{u}_4} = \mu_{u_1, u_2} \widehat{u}_3 \widehat{u}_4 \star_1 \mu_{u_3, u_4} = \mu_{u_1, u_2} \mu_{u_3, u_4}.$$

The rightmost normalisation strategy of $N\mathbf{C}$ is given in a symmetric way.

4.3. The acyclic (3, 1)-polygraph of generating confluences

We fix a reduced convergent 2-polygraph Σ equipped with its rightmost normalisation strategy σ .

4.3.1. Critical branchings of 2-polygraphs. By case analysis on the source of critical branchings of Σ , we can conclude that they must have one of the following two shapes



where φ, ψ are 2-cells of Σ .

We note that, if Σ is finite, then it has finitely many critical branchings: indeed, in that case there are finitely many pairs of 2-cells of Σ and, for φ and ψ fixed, there are finitely many ways to make their sources (which are 1-cells of a free category) overlap as in one of the two diagrams.

In fact, the 2-polygraph Σ being reduced, the first case cannot occur since, otherwise, the source of φ would be reducible by ψ . Thus, every critical branching of Σ must have shape $(\varphi v, u_1 \psi)$. We write the branching in that order since, by definition of \preceq , we have $\varphi v \preceq u_1 \psi$.

We also note that the 1-cells u_1, u_2 and v are normal forms and cannot be identities. Indeed, they are normal forms since, otherwise, at least one of the sources of φ and of ψ would be reducible by another 2-cell, preventing Σ from being reduced. If u_2 was an identity, then the branching would be

4. Polygraphic resolutions from convergent presentations

Peiffer. Moreover, if u_1 (resp. v) was an identity, then the source of ψ (resp. φ) would be reducible by φ (resp. ψ).

Finally, if we write $u = u_1 u_2$, the definitions of λ_{uv} and of ρ_{uv} imply that we have:

$$\lambda_{uv} = \varphi v \quad \text{and} \quad \rho_{uv} = u_1 \psi.$$

From all those observations, we conclude that every critical branching b of Σ must have shape

$$b = (\varphi \widehat{v}, \rho_{u\widehat{v}})$$

where u and v are composable 1-cells of Σ^* and where φ is a 2-cell of Σ with source u .

4.3.2. The basis of generating confluences. The *basis of generating confluences* of Σ is the cellular extension $c_2(\Sigma)$ of Σ^\top made of one 3-cell

$$\begin{array}{ccc} & \widehat{u\widehat{v}} & \\ \varphi \widehat{v} \nearrow & & \searrow \sigma_{\widehat{u\widehat{v}}} \\ u\widehat{v} & \Downarrow \omega_b & \widehat{u\widehat{v}} \\ \sigma_{u\widehat{v}} \searrow & & \nearrow \widehat{u\widehat{v}} \end{array}$$

for every critical branching $b = (\varphi \widehat{v}, \rho_{u\widehat{v}})$ of Σ . Alternatively, since, for a 2-cell f , we have defined f^* as the 2-cell $f \star_1 \sigma_{t(f)}$, the 3-cell ω_b can be pictured in the following, more compact way:

$$\begin{array}{ccc} & (\varphi \widehat{v})^* & \\ \widehat{u\widehat{v}} \nearrow & & \searrow \widehat{u\widehat{v}} \\ u\widehat{v} & \Downarrow \omega_b & \widehat{u\widehat{v}} \\ \widehat{\varphi \widehat{v}} \searrow & & \nearrow \widehat{u\widehat{v}} \end{array}$$

4.3.3. Lemma. *The rightmost normalisation strategy of Σ extends to a right normalisation strategy of $c_2(\Sigma)$.*

Proof. From Corollary 3.3.5, we know that it is sufficient to define a 3-cell

$$\sigma_{\varphi \widehat{w}}^* : (\varphi \widehat{w})^* \Rightarrow \widehat{\varphi w}^*$$

of $c_2(\Sigma)^\top$ for every 2-cell $\varphi : v \Rightarrow \widehat{v}$ of Σ and every 1-cell w in $\bar{\Sigma}$ such that $u = v\widehat{w}$ exists. We note that, by definition, we have:


$$(\varphi \widehat{w})^* = \varphi \widehat{w} \star_1 \sigma_{\widehat{v\widehat{w}}} \quad \text{and} \quad \widehat{\varphi w}^* = \sigma_u = \rho_u \star_1 \sigma_{t(\rho_u)}.$$

Let us proceed by case analysis on the type of the local branching $b = (\varphi \widehat{w}, \rho_u)$.

- If b is aspherical, then $\rho_u = \varphi \widehat{w}$. In that case, we define $\sigma_{\varphi \widehat{w}}^* = 1_{(\varphi \widehat{w})^*}$.
- By hypothesis, the branching b cannot be a Peiffer branching. Indeed, the source of ρ_u cannot be entirely contained in the normal form \widehat{w} .

4.3. The acyclic $(3, 1)$ -polygraph of generating confluences

- Otherwise, we have $\widehat{w} = \widehat{w}_1\widehat{w}_2$ and $b_1 = (\varphi_{\widehat{w}_1}, \rho_{v\widehat{w}_1})$ is a critical branching of Σ . In that case, we define $\sigma_{\varphi_{\widehat{w}}}$ as the composite 3-cell



of $c_2(\Sigma)^\top$. □

4.3.4. Proposition. *The $(3, 1)$ -polygraph $c_2(\Sigma)$ is acyclic.*


Remark. This result is already contained in [18], with a different proof and in a more general form: the generating confluences of a convergent n -polygraph Σ form a homotopy basis of the $(n, n - 1)$ -category Σ^\top .

4.3.5. Corollary. *A category with a finite convergent presentation is FDT_3 .*


In particular, we recover Squier's result: a monoid with a finite convergent presentation has finite derivation type, [42].

4.3.6. Example. Let \mathbf{C} be a category. We consider the reduced standard presentation $N\mathbf{C}$ of \mathbf{C} , equipped with the rightmost normalisation strategy. The basis of generating confluences of $N\mathbf{C}$ has one 3-cell $\mu_{u,v,w}$ for every composable non-identity 1-cells u, v and w of \mathbf{C} . The shape of this 3-cell depends on whether uv or vw is an identity or not:

- if neither uv nor vw is an identity:




- if uv is an identity, but not vw :




4. Polygraphic resolutions from convergent presentations

- if uv is not an identity, but vw is:



- if uv and vw are identities, and thus $u = uvw = w$:



The basis of generating confluences in the case of the leftmost normalisation strategy has the same 3-cells, with source and target reversed.


4.4. The acyclic $(4, 1)$ -polygraph of generating triple confluences

Let Σ be a reduced and convergent 2-polygraph.

4.4.1. Triple branchings of 2-polygraphs. A *triple branching* of Σ is a triple (f, g, h) of 2-cells of Σ^* with the same source and such that $f \preceq g \preceq h$. The triple branching (f, g, h) is *local* when f, g and h are rewriting steps. A local triple branching (f, g, h) is:

- *aspherical* when either (f, g) or (g, h) is aspherical,
- *Peiffer* when either (f, g) or (g, h) is Peiffer,
- *overlapping*, otherwise.

Triple branchings are ordered by inclusion, similarly to branchings. A *critical* triple branching is a minimal overlapping triple branching. Such a triple branching can have two different shapes, where φ, ψ and χ are generating 2-cells :



Those two shapes of critical triple branchings are sufficient for a reduced 2-polygraph but, in a general situation, the other possible type of critical branchings (with an inclusion of one source into the other one) generates several other possibilities. Either way, if Σ is finite, then it has a finite number of critical triple branchings.


4.4. The acyclic $(4, 1)$ -polygraph of generating triple confluences

For both possible shapes, the corresponding critical triple branching b can be written


$$b = (c\hat{v}, \rho_{u\hat{v}}) = (f\hat{v}, \rho_u\hat{v}, \rho_{u\hat{v}})$$

where $c = (f, \rho_u)$ is a critical branching of Σ with source $u = u_1u_2u_3u_4$ and where $\rho_u = u_1\psi$. Indeed, we note that v must be a normal form for Σ to be reduced. Moreover, in the first case, we have $f = \varphi u_4$ and $\rho_{u\hat{v}} = u_1u_2\chi$ and, in the second case, we have $f = \varphi u_3u_4$ and $\rho_{u\hat{v}} = u_1u_2u_3\chi$.

4.4.2. The basis of generating triple confluences. The basis of generating triple confluences of Σ is the cellular extension $c_3(\Sigma)$ of $c_2(\Sigma)^\top$ made of one 4-cell



for every critical triple branching $b = (f\hat{v}, \rho_u\hat{v}, \rho_{u\hat{v}})$ of Σ , where $c = (f, \rho_u)$ is a critical branching of Σ . Using the notations $(\cdot)^*$ and $\hat{\cdot}$ for 2-cells and 3-cells, the 4-cell ω_b can also be written



4.4.3. Lemma. The rightmost normalisation strategy of Σ extends to a right normalisation strategy of $c_3(\Sigma)$.

Proof. Let us define a 4-cell


$$\sigma_{\omega_c\hat{w}}^* : (\omega_c\hat{w})^* \rightrightarrows \widehat{\omega_c\hat{w}}^*$$

of $c_3(\Sigma)^\top$ for every 3-cell ω_c of $c_2(\Sigma)$ and every 1-cell w in $\bar{\Sigma}$ such that $\omega_c\hat{w}$ exists. Let us denote by v the source of the critical branching c of Σ and by f the rewriting step of Σ with source v such that the critical branching c is (f, ρ_v) . We proceed by case analysis on the type of the local triple branching $b = (f\hat{w}, \rho_v\hat{w}, \rho_{v\hat{w}})$.

- If b is aspherical, then $\rho_{v\hat{w}} = \rho_v\hat{w}$. In that case, we define $\sigma_{\omega_c\hat{w}}^* = 1_{(\omega_c\hat{w})^*}$.
- By hypothesis, the triple branching b cannot be a Peiffer one.
- Otherwise, we have $\hat{w} = \hat{w}_1\hat{w}_2$ and $b_1 = (f\hat{w}_1, \rho_v\hat{w}_1, \rho_{v\hat{w}_1})$ is a critical triple branching of Σ .

4. Polygraphic resolutions from convergent presentations

We define the 4-cell $\sigma_{\omega_c \hat{w}}^*$ as the following composite in $c_3(\Sigma)^\top$:




We apply Corollary 3.3.5 to extend the family of 4-cells we have defined to a right normalisation strategy of $c_3(\Sigma)$. \square

4.4.4. Proposition. *The $(4, 1)$ -polygraph $c_3(\Sigma)$ is acyclic.*

4.4.5. Corollary. *A category with a finite convergent presentation is FDT_4 .*

4.4.6. Example. In the case of the reduced standard presentation NC of a category C , the basis of generating triple confluences (for the rightmost normalisation strategy) has one 4-cell $\mu_{u,v,w,x}$ for every composable non-identity 1-cells u, v, w and x of C . The shape of the 4-cell $\mu_{u,v,w,x}$ depends on whether uv, vw, wx, uvw and vwx are identities or not. In the case where neither of these 1-cells is an identity, the corresponding 4-cell is the following one:



4.5. The polygraphic resolution generated by a convergent 2-polygraph

Let Σ be a reduced and convergent 2-polygraph and let us extend it into an acyclic $(\infty, 1)$ -polygraph, *i.e.*, a polygraphic resolution of the category $\bar{\Sigma}$. This $(\infty, 1)$ -polygraph is denoted by $c_\infty(\Sigma)$ and its generating $(n + 1)$ -cells, for $n \geq 2$, are (indexed by) the n -fold critical branchings of Σ . We proceed by induction on n , having already seen the base cases, for $n = 2$ and $n = 3$. The induction case follows the construction of the acyclic $(4, 1)$ -polygraph $c_3(\Sigma)$, so we go faster here.

4.5.1. Higher branchings of 2-polygraphs. An n -fold branching of Σ is a family (f_1, \dots, f_n) of rewriting steps of Σ with the same source and such that $f_1 \preceq \dots \preceq f_n$. We define local, aspherical, Peiffer, overlapping, minimal and critical n -fold branchings in a similar way to the cases $n = 2$ and $n = 3$. As

4.5. The polygraphic resolution generated by a convergent 2-polygraph

before, we study the possible shapes of an n -fold critical branching b of Σ and we conclude that it must have shape

$$b = (c\widehat{v}, \rho_{u\widehat{v}})$$

where c is a critical $(n-1)$ -fold branching of Σ with source u . Moreover, if Σ is finite, then it has finitely many n -fold critical branchings.

4.5.2. The basis of generating n -fold confluences. The *basis of generating n -fold confluences of Σ* is the cellular extension $c_n(\Sigma)$ of $c_{n-1}(\Sigma)^\top$ made of one $(n+1)$ -cell

$$\omega_b : (\omega_c\widehat{v})^* \longrightarrow \widehat{\omega_c v}^*$$

for every critical n -fold branching $b = (c\widehat{v}, \rho_{u\widehat{v}})$ of Σ .


The extension of the right normalisation strategy to $c_n(\Sigma)$ is made in the same way as in the case $n = 3$. It relies on a Corollary 3.3.5 and a case analysis, whose main point is to define an $(n+1)$ -cell

$$\sigma_{\omega_c\widehat{w}}^* : (\omega_c\widehat{w})^* \longrightarrow \widehat{\omega_c w}^*$$

in $c_n(\Sigma)^\top$ for every local n -fold branching

$$b = (c\widehat{w}, \rho_{v\widehat{w}})$$

of Σ such that $\widehat{w} = \widehat{w}_1\widehat{w}_2$ and such that $b_1 = (c\widehat{w}_1, \rho_{v\widehat{w}_1})$ is a critical n -fold branching of Σ . As in the case $n = 3$, we define the $(n+1)$ -cell $\sigma_{\omega_c\widehat{w}}^*$ as the following composite, where f is the first 2-cell of the critical n -fold branching c :



As a conclusion of this construction, we get that the $(n+1, 1)$ -polygraph $c_n(\Sigma)$ is acyclic.

4.5.3. Theorem. *Every convergent 2-polygraph Σ extends to a Tietze-equivalent, acyclic $(\infty, 1)$ -polygraph $c_\infty(\Sigma)$, whose generating n -cells, for every $n \geq 3$, are (indexed by) the critical $(n-1)$ -fold branchings of Σ .*

As a consequence, we get:

4.5.4. Corollary. *A category with a finite convergent presentation is FDT_∞ . Moreover, if \mathbf{C} is a category with a convergent presentation with no critical n -fold branching, for some $n \geq 2$, then $d_{\text{pol}}(\mathbf{C}) \leq n$.*

4.5.5. Example. If $N\mathbf{C}$ is the reduced standard presentation of a category \mathbf{C} , we have already built the dimensions 3 and 4 of the $(\infty, 1)$ -polygraph $c_\infty(N\mathbf{C})$, called the *reduced standard polygraphic resolution of \mathbf{C}* .

5. Abelianisation of polygraphic resolutions

More generally, the $(\infty, 1)$ -polygraph $c_\infty(\mathbf{NC})$ has, for every natural number $n \geq 2$, one n -cell μ_{u_1, \dots, u_n} for every family (u_1, \dots, u_n) of non-identity composable 1-cells of \mathbf{C} . The shape of this cell depends on the partial compositions of the 1-cells u_1, \dots, u_n .

In the case where no such partial composition is an identity, we get an n -cell with the same shape as an n -simplex, representing all the possible ways to transform $\widehat{u}_1 \cdots \widehat{u}_n$ into $\widehat{u_1 \cdots u_n}$, all the homotopies between these different ways, all homotopies between these homotopies, and so on. Indeed, we have seen that each 2-cell $\mu_{u,v} : \widehat{u}v \Rightarrow \widehat{uv}$ is represented by a triangle, each 3-cell $\mu_{u,v,w}$ is pictured as a tetrahedron and each 4-cell $\mu_{u,v,w,x}$ as a 4-simplex. More generally, the source and the target of this n -cell are $(n-1)$ -composites of the following $(n-1)$ -cells:

$$d_i(u_1, \dots, u_n) = \begin{cases} \widehat{u}_1 \mu_{u_2, \dots, u_n} & \text{if } i = 0 \\ \mu_{u_1, \dots, u_i u_{i+1}, \dots, u_n} & \text{if } 1 \leq i \leq n-1 \\ \mu_{u_1, \dots, u_{n-1}} \widehat{u}_n & \text{if } i = n. \end{cases}$$

More precisely, the n -cell μ_{u_1, \dots, u_n} has the same shape as an n -oriental, the higher-categorical equivalent of an n -simplex, see [44].

5. ABELIANISATION OF POLYGRAPHIC RESOLUTIONS

Let us fix a category \mathbf{C} .

5.1. Resolutions of finite type

5.1.1. Modules over a category, [35]. A \mathbf{C} -module is a functor from \mathbf{C} to the category of Abelian groups \mathbf{Ab} . The \mathbf{C} -modules and natural transformations between them form an Abelian category with enough projectives, denoted by $\mathbf{Mod}(\mathbf{C})$. Equivalently, $\mathbf{Mod}(\mathbf{C})$ can be described as the category of additive functors from $\mathbb{Z}\mathbf{C}$ to \mathbf{Ab} , where $\mathbb{Z}\mathbf{C}$ is the free \mathbb{Z} -category over \mathbf{C} : its 0-cells are the ones of \mathbf{C} and each hom-set $\mathbb{Z}\mathbf{C}(x, y)$ is the free Abelian group generated by $\mathbf{C}(x, y)$.

A \mathbf{C} -module M is *free* when it is a coproduct of representable functors $\mathbb{Z}\mathbf{C}(x, -)$ and it is *finitely generated* if there exists an epimorphism of \mathbf{C} -modules $F \rightarrow M$, with F free.

Let M be a \mathbf{C}^0 -module and N be a \mathbf{C} -module. The *tensor product of M and N over \mathbf{C}* is the Abelian group $M \otimes_{\mathbf{C}} N$ defined as the coend

$$M \otimes_{\mathbf{C}} N = \int^{x \in \mathbf{C}_0} M(x) \otimes_{\mathbb{Z}} N(x).$$

In a more explicit way, we have:

$$M \otimes_{\mathbf{C}} N = \left(\bigoplus_{x \in \mathbf{C}_0} M(x) \otimes_{\mathbb{Z}} N(x) \right) / Q$$

where Q is the subgroup of $\bigoplus_{x \in \mathbf{C}_0} M(x) \otimes_{\mathbb{Z}} N(x)$ generated by the elements

$$M(u)(a) \otimes b - a \otimes N(u)(b),$$

for any possible 1-cell $u : x \rightarrow y$ of \mathbf{C} and any elements a of $M(y)$ and b of $N(x)$.

5.1.2. Modules of finite homological type. A \mathbf{C} -module M is of *homological type* FP_n , for a natural number n , when there exists a partial resolution of M of length n by projective, finitely generated \mathbf{C} -modules:

$$P_n \longrightarrow P_{n-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow M \longrightarrow 0.$$

A \mathbf{C} -module M is of *homological type* FP_∞ when there exists a resolution of M by projective, finitely generated \mathbf{C} -modules:

$$\cdots \longrightarrow P_n \longrightarrow P_{n-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow M \longrightarrow 0.$$

As a generalisation of Schanuel's lemma, we have, given two exact sequences

$$0 \longrightarrow P_{n+1} \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_0 \longrightarrow M \longrightarrow 0$$

and

$$0 \longrightarrow P'_{n+1} \longrightarrow P'_n \longrightarrow \cdots \longrightarrow P'_0 \longrightarrow M \longrightarrow 0$$

of projective \mathbf{C} -modules, with P_i and P'_i finitely generated for every $0 \leq i \leq n$, then P_n is finitely generated if and only if P'_n is finitely generated. This yields the following characterisation of the property FP_n :

5.1.3. Lemma. *Let M be a \mathbf{C} -module and let n be a natural number. The following assertions are equivalent:*

- i) *The \mathbf{C} -module M is of homological type FP_n .*
- ii) *There exists a free, finitely generated partial resolution of M of length n :*

$$F_n \longrightarrow F_{n-1} \longrightarrow \cdots \longrightarrow F_0 \longrightarrow M \longrightarrow 0.$$

- iii) *The \mathbf{C} -module M is finitely generated and, for every $0 \leq k < n$ and every projective, finitely generated partial resolution of M of length k*

$$P_k \xrightarrow{d_k} P_{k-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow M \longrightarrow 0,$$

the \mathbf{C} -module $\text{Ker } d_k$ is finitely generated.

5.1.4. Lemma. *Let \mathbf{D} be a category, let $F : \mathbf{C} \rightarrow \mathbf{D}$ be a functor and let $\text{Lan}_F : \text{Mod}(\mathbf{C}) \rightarrow \text{Mod}(\mathbf{D})$ be the additive left Kan extension along F . If M is a \mathbf{C} -module of homological type FP_n then $\text{Lan}_F(M)$ is a \mathbf{D} -module of homological type FP_n .*

Proof. Let us assume that M is a \mathbf{C} -module of type FP_n . Then there exists a projective, finitely generated partial resolution $P_* \rightarrow M$ of length n . If x is a 0-cell in \mathbf{D} , then we have:

$$\text{Lan}_F(M)(x) = \mathbb{Z}\mathbf{D}(F, x) \otimes_{\mathbf{C}} M.$$

Since each \mathbf{C} -module P_i is finitely generated and projective, then so is the \mathbf{D} -module $\text{Lan}_F(P_i)$. Moreover, the functor Lan_F is right-exact: it follows that $\text{Lan}_F(P_*) \rightarrow \text{Lan}_F(M)$ is a projective, finitely generated partial resolution of length n . This proves that $\text{Lan}_F(M)$ is of type FP_n . \square

5. Abelianisation of polygraphic resolutions

5.2. Categories of finite homological type

5.2.1. Natural systems of Abelian groups. The *category of factorisations of \mathbf{C}* is the category, denoted by \mathbf{FC} , whose 0-cells are the 1-cells of \mathbf{C} and whose 1-cells from w to w' are pairs (u, v) of 1-cells of \mathbf{C} such that the following diagram commutes in \mathbf{C} :

$$\begin{array}{ccc} & \cdot & \xrightarrow{w} \cdot \\ & \nearrow u & \searrow v \\ \cdot & & \cdot \\ & \xrightarrow{w'} & \cdot \end{array}$$

In such a situation, the triple (u, w, v) is called a *factorisation of w'* . Composition in \mathbf{FC} is defined by pasting, *i.e.*, if $(u, v) : w \rightarrow w'$ and $(u', v') : w' \rightarrow w''$ are 1-cells of \mathbf{FC} , then the composite $(u, v)(u', v')$ is $(u'u, vv')$:

$$\begin{array}{ccc} & \cdot & \xrightarrow{w} \cdot \\ & \nearrow u & \searrow v \\ \cdot & & \cdot \\ & \xrightarrow{w'} & \cdot \\ & \nearrow u' & \searrow v' \\ \cdot & & \cdot \\ & \xrightarrow{w''} & \cdot \end{array}$$

The identity of w is $(1_{s(w)}, 1_{t(w)})$:

$$\begin{array}{ccc} & \cdot & \xrightarrow{w} \cdot \\ & \nearrow 1_{s(w)} & \searrow 1_{t(w)} \\ \cdot & & \cdot \\ & \xrightarrow{w'} & \cdot \end{array}$$

A *natural system (of Abelian groups) on \mathbf{C}* is an \mathbf{FC} -module D , *i.e.*, a functor $D : \mathbf{FC} \rightarrow \mathbf{Ab}$. As in [7], we denote by D_w the Abelian group which is the image of w by D . If there is no confusion, we denote by uav the image of $a \in D_w$ through the morphism of groups $D(u, v) : D_w \rightarrow D_{w'}$. The category of natural systems on \mathbf{C} is denoted by $\mathbf{Nat}(\mathbf{C})$.

5.2.2. Free natural systems. Given a family X of 1-cells of \mathbf{C} , we denote by $\mathbf{F}_{\mathbf{C}}[X]$ the *free natural system on \mathbf{C} generated by X* , which is defined by

$$\mathbf{F}_{\mathbf{C}}[X] = \bigoplus_{x \in X} \mathbf{FC}(x, -).$$

In particular, if Σ is an $(n, 1)$ -polygraph such that $\bar{\Sigma} \simeq \mathbf{C}$, we consider:

- The free natural system $\mathbf{F}_{\mathbf{C}}[\Sigma_0]$ generated by the 1-cells 1_x , for $x \in \Sigma_0$: if w is a 1-cell in \mathbf{C} , then $\mathbf{F}_{\mathbf{C}}[\Sigma_0]_w$ is the free Abelian group generated by the pairs (u, v) of 1-cells of \mathbf{C} such that $uv = w$.
- For every $1 \leq k \leq p$, the free natural system $\mathbf{F}_{\mathbf{C}}[\Sigma_k]$ generated by a copy of the 1-cell $\bar{\varphi}$ for each k -cell φ of Σ_k : if w is a 1-cell in \mathbf{C} , then $\mathbf{F}_{\mathbf{C}}[\Sigma_k]_w$ is the free Abelian group generated by the triples (u, φ, v) , thereafter denoted by $u[\varphi]v$, made of a k -cell φ of Σ_k and 1-cells u, v of \mathbf{C} such that $u\bar{\varphi}v = w$.

5.2.3. Categories of finite homological type. The property for a category \mathbf{C} to be of homological type FP_n is defined according to a category of modules over one of the categories in the following diagram

$$\begin{array}{ccccc} & & \mathbf{C}^0 & & \\ & & \nearrow & & \searrow \\ \mathbf{FC} & \xrightarrow{\pi} & \mathbf{C}^0 \times \mathbf{C} & & \mathbf{C}^\top \\ & & \searrow & & \nearrow \\ & & \mathbf{C} & & \end{array}$$

where \mathbf{C}^\top is the groupoid generated by \mathbf{C} , π is the projection $u \mapsto (s(u), t(u))$, p_1 and p_2 are the projections of the cartesian product, q_1 and q_2 are the injections $u^0 \mapsto u^-$ and $u \mapsto u$.

Let us denote by \mathbb{Z} the constant natural system on \mathbf{C} given, for any 1-cell u of \mathbf{C} , by

$$\mathbb{Z}_u = \mathbb{Z} \quad \text{and} \quad \mathbb{Z}(u, 1) = \mathbb{Z}(1, u) = \text{Id}_{\mathbb{Z}}.$$

Let $\mathbb{Z}\mathbf{C}$ denote the $\mathbf{C}^0 \times \mathbf{C}$ -module $\mathbb{Z}\mathbf{C}$ whose component at (x, y) is the free \mathbb{Z} -module $\mathbb{Z}\mathbf{C}(x, y)$ generated by $\mathbf{C}(x, y)$.

A category \mathbf{C} is of *homological type*

- i) FP_n when the constant natural system \mathbb{Z} is of type FP_n ,
- ii) *bi*- FP_n when the $\mathbf{C}^0 \times \mathbf{C}$ -module $\mathbb{Z}\mathbf{C}$ is of type FP_n ,
- iii) *left*- FP_n when the constant \mathbf{C} -module \mathbb{Z} is of type FP_n ,
- iv) *right*- FP_n when the constant \mathbf{C}^0 -module \mathbb{Z} is of type FP_n ,
- v) *top*- FP_n when the constant \mathbf{C}^\top -module \mathbb{Z} is of type FP_n .

5.2.4. Proposition. *For categories, we have the following implications:*

$$\begin{array}{ccccc} & & \text{right-}\text{FP}_n & & \\ & & \nearrow & & \searrow \\ \text{FP}_n & \implies & \text{bi-}\text{FP}_n & & \text{top-}\text{FP}_n \\ & & \searrow & & \nearrow \\ & & \text{left-}\text{FP}_n & & \end{array}$$

Proof. We have the following left Kan extensions:

$$\text{Lan}_\pi(\mathbb{Z}) = \mathbb{Z}\mathbf{C} \quad \text{Lan}_{p_i}(\mathbb{Z}) = \mathbb{Z} \quad \text{Lan}_{q_i}(\mathbb{Z}) = \mathbb{Z}.$$

Hence the implications are consequences of Lemma 5.1.4. □

5.2.5. Converse implications. The converse of the implications $\text{bi-}\text{FP}_n \Rightarrow \text{left/right-}\text{FP}_n$ and $\text{left/right-}\text{FP}_n \Rightarrow \text{top-}\text{FP}_n$ of Proposition 5.2.4 does not hold in general. Indeed, Cohen constructed a $\text{right-}\text{FP}_\infty$ monoid which is not $\text{left-}\text{FP}_1$: thus, the properties $\text{top-}\text{FP}_n$, $\text{left-}\text{FP}_n$ and $\text{right-}\text{FP}_n$ are not equivalent in general, [12]. Moreover, monoids with a finite convergent presentation are of types $\text{left-}\text{FP}_\infty$ and $\text{right-}\text{FP}_\infty$, [41, 1, 21], but there exists a finitely presented monoid, of types $\text{left-}\text{FP}_\infty$ and $\text{right-}\text{FP}_\infty$, which

5. Abelianisation of polygraphic resolutions

does not satisfy the homological finiteness condition FHT, introduced by Pride and Wang, [22]; since the properties FHT and bi-FP_3 are equivalent, [23], it follows that the properties left-FP_n and right-FP_n do not imply the property bi-FP_n in general. We conjecture that the converse of the first implication is not true either, but this is still an open problem. However, in the special case of groupoids, all the implications are equivalences.

5.2.6. Proposition. *For groupoids, the properties FP_n , bi-FP_n , left-FP_n , right-FP_n and top-FP_n are equivalent.*

Proof. If \mathbf{G} is a groupoid, then the groupoid \mathbf{G}^\top generated by \mathbf{G} is \mathbf{G} itself. Hence, the properties left-FP_n and top-FP_n are equivalent. As a consequence, it is sufficient to prove that left-FP_n implies FP_n to conclude the proof.

Given a \mathbf{G} -module M , we denote by \widetilde{M} the natural system of \mathbf{G} defined, on a 1-cell $w : x \rightarrow y$ of \mathbf{G} , by $\widetilde{M}_w = M(y)$ and, on a factorisation $w' = uwv$ with $w : x \rightarrow y$ and $w' : x' \rightarrow y'$ in \mathbf{G} , by $\widetilde{M}(u, v) = M(v) : M(y) \rightarrow M(y')$. Let us assume that M is a projective \mathbf{G} -module, *i.e.*, there exists a family X of 1-cells of \mathbf{G} and a surjective morphism of groups

$$\bigoplus_{x \in X} \mathbb{Z}\mathbf{G}(x, y) \xrightarrow{\pi_y} M(y)$$

that is natural in y . Let us denote by L the natural system on \mathbf{G} defined, on a 1-cell u with target y , by

$$L_u = \bigoplus_{x \in X} \mathbb{Z}\mathbf{F}\mathbf{G}(1_x, 1_y).$$

For fixed 0-cells x and y of \mathbf{G} , the set $\mathbf{F}\mathbf{G}(1_x, 1_y)$ is the one of pairs (u, v) of 1-cells of \mathbf{G} such that $u1_x v = 1_y$. Because \mathbf{G} is a groupoid, this is exactly the set of pairs (v^-, v) where $v : x \rightarrow y$ is a 1-cell of \mathbf{G} . Hence, we have a bijection between $\mathbf{G}(x, y)$ and $\mathbf{F}\mathbf{G}(1_x, 1_y)$ that is natural in y . This gives, for any 1-cell u with target y in \mathbf{G} , a surjective morphism of groups

$$L_u = \bigoplus_{x \in X} \mathbb{Z}\mathbf{G}(x, y) \xrightarrow{\pi_{t(u)}} M(y) = \widetilde{M}(u).$$

Moreover, one checks that this is natural in u , yielding an epimorphism $\pi_{t(-)} : L \rightarrow \widetilde{M}$ of natural systems on \mathbf{G} , proving that M is projective. Moreover, by construction, if M is finitely generated, then we can choose the family X to be finite, so that \widetilde{M} is finitely generated too.

As a conclusion, from a partial resolution $P_* \rightarrow \mathbb{Z}$ of length n by projective and finitely generated \mathbf{G} -modules, we can build a partial resolution $\widetilde{P}_* \rightarrow \mathbb{Z}$ of length n by projective and finitely generated natural systems on \mathbf{G} . Thus, the property left-FP_n implies the property FP_n for groupoids. \square

5.2.7. Finite homological type and homology. The cohomology of categories with values in natural systems was defined in [47] and [7]. One can also define the homology of a category \mathbf{C} with values in a *contravariant natural system* D on \mathbf{C} , that is an $(\mathbf{F}\mathbf{C})^0$ -module, and relate its finiteness properties to the fact that \mathbf{C} is FP_n . Note that this is independent of the rest of this article.

5.3. The Reidemeister-Fox-Squier complex

We consider the nerve $N_*(\mathbf{C})$ of \mathbf{C} , with boundary maps $d_i : N_n(\mathbf{C}) \rightarrow N_{n-1}(\mathbf{C})$, for $0 \leq i \leq n$. For $s = (u_1, \dots, u_n)$ in $N_n(\mathbf{C})$, we denote by \bar{s} the composite 1-cell $u_1 \cdots u_n$ of \mathbf{C} . For every natural number n , the n -th chain group $C_n(\mathbf{C}, D)$ is defined as the Abelian group

$$C_n(\mathbf{C}, D) = \bigoplus_{s \in N_n(\mathbf{C})} D_{\bar{s}}.$$

We denote by ι_s the embedding of $D_{\bar{s}}$ into $C_n(\mathbf{C}, D)$. The boundary map $d : C_n(\mathbf{C}, D) \rightarrow C_{n-1}(\mathbf{C}, D)$ is defined, on the component $D_{\bar{s}}$ of $C_n(\mathbf{C}, D)$, by:

$$d\iota_s = \iota_{d_0(s)}u_{1*} + \sum_{i=1}^{n-1} (-1)^i \iota_{d_i(s)} + (-1)^n \iota_{d_n(s)}u_n^*,$$

with $s = (u_1, \dots, u_n)$ and where u_{1*} and u_n^* respectively denote $D(u_1, 1)$ and $D(1, u_n)$. The homology of \mathbf{C} with coefficients in D is defined as the homology of the complex $(C_*(\mathbf{C}, D), d_*)$:

$$H_*(\mathbf{C}, D) = H_*(C_*(\mathbf{C}, D), d_*).$$

We denote by $\text{Tor}_*^{\text{FC}}(D, -)$ the left derived functor from the functor $D \otimes_{\text{FC}} -$. One proves that there is an isomorphism which is natural in D :

$$H_*(\mathbf{C}, D) \simeq \text{Tor}_*^{\text{FC}}(D, \mathbb{Z}).$$

As a consequence, using Lemma 5.1.3, we get:

5.2.8. Proposition. *If a category \mathbf{C} is of homological type FP_n , for a natural number n , then the Abelian group $H_k(\mathbf{C}, \mathbb{Z})$ is finitely generated for every $0 \leq k \leq n$.*

5.3. The Reidemeister-Fox-Squier complex

Let Σ be an $(n, 1)$ -polygraph. The mapping of every 1-cell x of Σ to the element $[x]$ of $F_{\bar{\Sigma}}[\Sigma_1]_{\bar{x}}$ is extended to associate to every 1-cell u of Σ_1^* the element $[u]$ of $F_{\bar{\Sigma}}[\Sigma_1]_{\bar{u}}$, defined by induction on the size of u as follows:

$$[1_x] = 0 \quad \text{and} \quad [uv] = [u]\bar{v} + \bar{u}[v].$$

This is well-defined since the given relations are compatible with the associativity and identity relations of Σ_1^* . Note that the mapping $[\cdot]$ is a special case of the notion of derivation of the category Σ_1^* into the natural system $F_{\bar{\Sigma}}[\Sigma_1]$ on $\bar{\Sigma}$, seen as a natural system on Σ_1^* by composition with the canonical projection $\Sigma_1^* \rightarrow \bar{\Sigma}$, see [7].

Then, for $1 < k \leq n$, the mapping of every k -cell α of Σ to the element $[\alpha]$ of $F_{\bar{\Sigma}}[\Sigma_k]_{\bar{\alpha}}$ is extended to associate to every k -cell f of Σ_k^\top the element $[f]$ of $F_{\bar{\Sigma}}[\Sigma_k]_{\bar{f}}$, defined by induction on the size of f as follows:

$$[1_u] = 0 \quad [f^-] = -[f] \quad [f \star_i g] = \begin{cases} [f]\bar{g} + \bar{g}[f] & \text{if } i = 0 \\ [f] + [g] & \text{otherwise.} \end{cases}$$

Here, to check that this is well-defined, we also have to show that this definition is compatible with exchange relations, for every $0 \leq i < j \leq k$:

$$[(f \star_i g) \star_j (h \star_i k)] = [(f \star_j h) \star_i (g \star_j k)] = \begin{cases} [f]\bar{g} + \bar{f}[g] + [h]\bar{k} + \bar{h}[k] & \text{if } i = 0 \\ [f] + [g] + [h] + [k] & \text{otherwise.} \end{cases}$$

5. Abelianisation of polygraphic resolutions

5.3.1. The Reidemeister-Fox-Squier complex. Let Σ be an $(n, 1)$ -polygraph. For $1 \leq k \leq n$, the k -th Reidemeister-Fox-Squier boundary map of Σ is the morphism of natural systems

$$\delta_k : F_{\bar{\Sigma}}[\Sigma_k] \longrightarrow F_{\bar{\Sigma}}[\Sigma_{k-1}]$$

defined, on the generator $[\alpha]$ corresponding to a k -cell α of Σ , by:

$$\delta_k[\alpha] = \begin{cases} (\bar{\alpha}, 1) - (1, \bar{\alpha}) & \text{if } k = 1 \\ [s(\alpha)] - [t(\alpha)] & \text{otherwise.} \end{cases}$$

The *augmentation map* of Σ is the morphism of natural systems $\varepsilon : F_{\bar{\Sigma}}[\Sigma_0] \rightarrow \mathbb{Z}$ defined, for every pair (u, v) of composable 1-cells of $\bar{\Sigma}$, by:

$$\varepsilon(u, v) = 1.$$

By induction on the size of cells of Σ^\top , one proves that, for every k -cell f in Σ^\top , with $k \geq 1$, the following holds:

$$\delta_k[f] = \begin{cases} (\bar{f}, 1) - (1, \bar{f}) & \text{if } k = 1 \\ [s(f)] - [t(f)] & \text{otherwise.} \end{cases}$$

As a consequence, we have $\varepsilon\delta_1 = 0$ and $\delta_k\delta_{k+1} = 0$, for every $1 \leq k < n$. Thus, we get the following chain complex of natural systems on $\bar{\Sigma}$, called the *Reidemeister-Fox-Squier complex* of Σ and denoted by $F_{\bar{\Sigma}}[\Sigma]$:

$$F_{\bar{\Sigma}}[\Sigma_n] \xrightarrow{\delta_n} F_{\bar{\Sigma}}[\Sigma_{n-1}] \xrightarrow{\delta_{n-1}} \cdots \xrightarrow{\delta_2} F_{\bar{\Sigma}}[\Sigma_1] \xrightarrow{\delta_1} F_{\bar{\Sigma}}[\Sigma_0] \xrightarrow{\varepsilon} \mathbb{Z} \longrightarrow 0.$$

The terminology is due to the fact that this complex is inspired by constructions by Reidemeister, Fox and Squier, see [40, 15, 41]. In particular, the *Fox Jacobian* used by Squier is the boundary map δ_2 , sending every 2-cell $f : u \Rightarrow v$, *i.e.*, every relation $u = v$ of the presentation Σ_2 , to the element $[u] - [v]$ of $F_{\bar{\Sigma}}[\Sigma_1]_{\bar{u}}$.

5.4. Abelianisation of polygraphic resolutions

Let us fix a partial polygraphic resolution Σ of length $n \geq 1$ of the category \mathbf{C} .

5.4.1. Contracting homotopies. Since Σ is acyclic, it admits a left normalisation strategy σ . We denote by σ_k , for $-1 \leq k \leq n$, the following families of morphisms of groups, indexed by a 1-cell w of \mathbf{C} :

$$\begin{array}{ccc} (\sigma_{-1})_w : \mathbb{Z} & \longrightarrow & F_{\mathbf{C}}[\Sigma_0]_w \\ 1 & \longmapsto & (1, w) \end{array} \qquad \begin{array}{ccc} (\sigma_0)_w : F_{\mathbf{C}}[\Sigma_0]_w & \longrightarrow & F_{\mathbf{C}}[\Sigma_1]_w \\ (u, v) & \longmapsto & [\hat{u}]_v \end{array}$$

$$\begin{array}{ccc} (\sigma_k)_w : F_{\mathbf{C}}[\Sigma_k]_w & \longrightarrow & F_{\mathbf{C}}[\Sigma_{k+1}]_w \\ u[x]_v & \longmapsto & [\sigma_{\hat{u}x}]_v \end{array}$$

5.4.2. Lemma. For every $k \in \{1, \dots, n-1\}$, every k -cell f of Σ^\top and every 1-cells u and v of \mathbf{C} such that $u\bar{f}v$ exists, we have:

$$\sigma_k(u[f]v) = [\sigma_{\hat{u}f}]_v.$$

Proof. We proceed by induction on the size of f . If $f = 1_w$, for some $(k-1)$ -cell w of Σ^\top , then we have, on the one hand,

$$\sigma_k(\mathbf{u}[1_w]\mathbf{v}) = \sigma_k(0) = 0$$

and, on the other hand,

$$[\sigma_{1_{\widehat{u}w}}]\mathbf{v} = [1_{1_{\widehat{u}w}}]\mathbf{v} = 0.$$

If f has size 1, then the result holds by definition of σ_k . Let us assume that $f = gh$, where g and h are non-identity k -cells of Σ^\top . Then we use the induction hypothesis on g and h to get, on the one hand:

$$\sigma_k(\mathbf{u}[gh]\mathbf{v}) = \sigma_k(\mathbf{u}[g]\bar{h}\mathbf{v}) + \sigma_k(\mathbf{u}\bar{g}[h]\mathbf{v}) = [\sigma_{\widehat{u}g}]\bar{h}\mathbf{v} + [\sigma_{\widehat{u}gh}]\mathbf{v}.$$

On the other hand, since σ is a left normalisation strategy, we have:

$$[\sigma_{\widehat{u}gh}]\mathbf{v} = [\sigma_{\widehat{u}g}s_1(h) \star_1 \sigma_{\widehat{u}gh}]\mathbf{v} = [\sigma_{\widehat{u}g}]\bar{h}\mathbf{v} + [\sigma_{\widehat{u}gh}]\mathbf{v}.$$

Finally, let us assume that $f = g \star_i h$, where g and h are non-identity k -cells of Σ^\top and $i \geq 1$. Then we get:

$$\sigma_k(\mathbf{u}[g \star_i h]\mathbf{v}) = \sigma_k(\mathbf{u}[g]\mathbf{v}) + \sigma_k(\mathbf{u}[h]\mathbf{v}) = [\sigma_{\widehat{u}g}]\mathbf{v} + [\sigma_{\widehat{u}h}]\mathbf{v}.$$

And we also have:

$$[\sigma_{\widehat{u}(g \star_i h)}]\mathbf{v} = [\sigma_{\widehat{u}g \star_i \widehat{u}h}]\mathbf{v} = [\sigma_{\widehat{u}g} \star_i \sigma_{\widehat{u}h}]\mathbf{v} = [\sigma_{\widehat{u}g}]\mathbf{v} + [\sigma_{\widehat{u}h}]\mathbf{v}. \quad \square$$

5.4.3. Theorem. *If Σ is a (partial) polygraphic resolution (of length n) of a category \mathbf{C} , then the Reidemeister-Fox-Squier complex $F_{\mathbf{C}}[\Sigma]$ is a free (partial) resolution (of length n) of the constant natural system \mathbb{Z} on \mathbf{C} .*

Proof. Let us prove that σ_* is a contracting homotopy. Each $(\sigma_{-1})_w$ is a section of ε_w , hence ε is an epimorphism of natural systems. By linearity of the boundary and section maps, it is sufficient to check the relation

$$\delta_1 \sigma_0(\mathbf{u}, \mathbf{v}) = (\mathbf{u}, \mathbf{v}) - (1, \mathbf{u}\mathbf{v}) = (\mathbf{u}, \mathbf{v}) - \sigma_{-1}\varepsilon(\mathbf{u}, \mathbf{v}),$$

on a generator (\mathbf{u}, \mathbf{v}) of $F_{\mathbf{C}}[\Sigma_0]_w$ to get the exactness of $F_{\mathbf{C}}[\Sigma]$ at $F_{\mathbf{C}}[\Sigma_0]$. Then, on a generator $\mathbf{u}[x]\mathbf{v}$ of $F_{\mathbf{C}}[\Sigma_1]_w$, we compute

$$\delta_2 \sigma_1(\mathbf{u}[x]\mathbf{v}) = \delta_2([\sigma_{\widehat{u}x}])\mathbf{v} = [\widehat{u}x]\mathbf{v} - [\widehat{u}x]\mathbf{v} = [\widehat{u}]\bar{x}\mathbf{v} + \mathbf{u}[x]\mathbf{v} - [\widehat{u}x]\mathbf{v}.$$

and

$$\sigma_0 \delta_1(\mathbf{u}[x]\mathbf{v}) = \sigma_0(\mathbf{u}\bar{x}, \mathbf{v}) - \sigma_0(\mathbf{u}, \bar{x}\mathbf{v}) = [\widehat{u}x]\mathbf{v} - [\widehat{u}]\bar{x}\mathbf{v}.$$

Hence, using the linearity of the boundary and section maps, we get $\delta_2 \sigma_1 + \sigma_0 \delta_1 = 1_{F_{\mathbf{C}}[\Sigma_1]}$, proving exactness at $F_{\mathbf{C}}[\Sigma_1]$. Finally, for $k \in \{2, \dots, p-1\}$ and a generator $\mathbf{u}[\alpha]\mathbf{v}$ of $F_{\mathbf{C}}[\Sigma_k]_w$, we have:

$$\begin{aligned} \delta_{k+1} \sigma_k(\mathbf{u}[\alpha]\mathbf{v}) &= \delta_{k+1}[\sigma_{\widehat{u}\alpha}]\mathbf{v} \\ &= [\widehat{u}\alpha]\mathbf{v} - [\sigma_{\widehat{u}s(\alpha)} \star_{k-1} \sigma_{\widehat{u}t(\alpha)}^-]\mathbf{v} \\ &= \mathbf{u}[\alpha]\mathbf{v} - [\sigma_{\widehat{u}s(\alpha)}]\mathbf{v} + [\sigma_{\widehat{u}t(\alpha)}]\mathbf{v} \\ &= \mathbf{u}[\alpha]\mathbf{v} - \sigma_{k-1}(\mathbf{u}[s\alpha]\mathbf{v}) + \sigma_{k-1}(\mathbf{u}[t\alpha]\mathbf{v}) \\ &= \mathbf{u}[\alpha]\mathbf{v} - \sigma_{k-1} \delta_k(\mathbf{u}[\alpha]\mathbf{v}). \end{aligned}$$

Thus, by linearity of the boundary and section maps, we get $\delta_{k+1} \sigma_k + \sigma_{k-1} \delta_k = 1_{F_{\mathbf{C}}[\Sigma_k]}$, proving exactness at $F_{\mathbf{C}}[\Sigma_k]$ and concluding the proof. \square

5. Abelianisation of polygraphic resolutions

By construction, if Σ is a finite $(n, 1)$ -polygraph, then each $F_{\bar{\Sigma}}[\Sigma_k]$ is finitely generated. In particular, every category with a finite number of 0-cells is of homological type FP_0 and every category that is finitely generated (resp. presented) is of homological type FP_1 (resp. FP_2). More generally, we have the following result, generalising the fact that a finite derivation type monoid is of homological type FP_3 , see [13, 39]:

5.4.4. Corollary. *For categories, the property FDT_n implies the property FP_n , for every $0 \leq n \leq \infty$.*

Finally, as a consequence of Theorem 4.5.3, we get:

5.4.5. Corollary. *If a category admits a finite and convergent presentation, then it is of homological type FP_∞ .*

5.4.6. Example. Let \mathbf{C} be a category. The Reidemeister-Fox-Squier complex corresponding to the reduced standard polygraphic resolution $c_\infty(N\mathbf{C})$ of \mathbf{C} is (isomorphic to) the following one

$$\dots \xrightarrow{\delta_{n+1}} F_{\mathbf{C}}[\mathbf{C}_n] \xrightarrow{\delta_n} F_{\mathbf{C}}[\mathbf{C}_{n-1}] \xrightarrow{\delta_{n-1}} \dots \xrightarrow{\delta_2} F_{\mathbf{C}}[\mathbf{C}_1] \xrightarrow{\delta_1} F_{\mathbf{C}}[\mathbf{C}_0] \xrightarrow{\varepsilon} \mathbb{Z} \longrightarrow 0$$

where \mathbf{C}_n is the set of composable non-identity 1-cells u_1, \dots, u_n of \mathbf{C} . The differential map is given, on a generator $[u_1, \dots, u_n]$, by:

$$\begin{aligned} \delta[u_1, \dots, u_n] &= \sum_{i=0}^n (-1)^{n-i} [d_i(u_1, \dots, u_n)] \\ &= (-1)^n u_1 [u_2, \dots, u_n] + \sum_{i=1}^{n-1} (-1)^{n-i} [u_1, \dots, u_i u_{i+1}, \dots, u_n] + [u_1, \dots, u_{n-1}] u_n. \end{aligned}$$

5.5. Homological syzygies and cohomological dimension

5.5.1. Homological syzygies. For every k in $\{1, \dots, p+1\}$, the kernel of δ_k is denoted by $h_k(\Sigma)$ and called *the natural system of homological k -syzygies of Σ* . The kernel of ε is denoted by $h_0(\bar{\Sigma})$ and called *the augmentation ideal of $\bar{\Sigma}$* .

When Σ_0 is finite, then the natural system $h_0(\bar{\Sigma})$ is finitely generated if, and only if, the category $\bar{\Sigma}$ has homological type FP_1 . If Σ is a generating 1-polygraph for a category \mathbf{C} , one checks that $h_0(\mathbf{C})$ is generated by the set $\{(x, 1) - (1, x) \mid x \in \Sigma_1\}$. Thus a category has homological type FP_1 if, and only if, it is finitely generated.

From Theorem 5.4.3, we get a characterisation of the homological properties FP_n in terms of polygraphic resolutions:

5.5.2. Proposition. *Let \mathbf{C} be a category and Σ be a partial finite polygraphic resolution of \mathbf{C} of length n . If the natural system $h_n(\Sigma)$ of homological n -syzygies of Σ is finitely generated, then \mathbf{C} is of homological type FP_{n+1} .*

Theorem 5.4.3 also gives a description of homological n -syzygies in terms of critical n -fold branchings of a convergent presentation:

5.6. Homological syzygies and identities among relations

5.5.3. Proposition. *Let \mathbf{C} be a category with a convergent presentation Σ . Then, for every $n \geq 2$, the natural system $h_n(\Sigma)$ of homological n -syzygies of Σ is generated by the elements*

$$\delta_{n+1}[\omega_b] = [(\omega_c \widehat{v})^*] - [\widehat{\omega_c v}^*]$$

where $b = (c\widehat{v}, \rho_{u\widehat{v}})$ ranges over the critical n -fold branchings of Σ .

5.5.4. Cohomological dimension. Finally, Theorem 5.4.3 gives the following bounds for the cohomological dimension of a category. We recall that the *cohomological dimension* of a category \mathbf{C} , is defined, when it exists as the lowest $0 \leq n \leq \infty$ such that the constant natural system \mathbb{Z} on \mathbf{C} admits a projective resolution

$$0 \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow \mathbb{Z} \longrightarrow 0.$$

In that case, the cohomological dimension of \mathbf{C} is denoted by $\text{cd}(\mathbf{C})$. In particular, when \mathbf{C} is free, then we have $\text{cd}(\mathbf{F}) \leq 1$, see [7].

5.5.5. Proposition. *The cohomological dimension of a category \mathbf{C} admits the following upper bounds:*

- i) *The inequality $\text{cd}(\mathbf{C}) \leq d_{\text{pol}}(\mathbf{C})$ holds.*
- ii) *If \mathbf{C} admits an aspherical partial polygraphic resolution of length n , then $\text{cd}(\mathbf{C}) \leq n$.*
- iii) *If \mathbf{C} admits a convergent presentation with no critical n -fold branching, then $\text{cd}(\mathbf{C}) \leq n$.*

5.6. Homological syzygies and identities among relations

In [19], the authors have introduced the natural system of identities among relations of an n -polygraph Σ . If Σ is a convergent 2-polygraph, this natural system on $\bar{\Sigma}$ is generated by the critical branchings of Σ . In Proposition 5.5.3, we have seen that this is also the case of the natural system of homological 2-syzygies of Σ . In this section, we prove that, more generally, the natural systems of homological 2-syzygies and of identities among relations of any 2-polygraph are isomorphic.

5.6.1. Natural systems on n -categories. We recall from [18], that a *context* of an n -category \mathcal{C} is an n -cell C of the n -category $\mathcal{C}[x]$, where x is an $(n-1)$ -sphere of \mathcal{C} , and such that C contains exactly one occurrence of the n -cell x of $\mathcal{C}[x]$. Such a context admits a (generally non-unique) decomposition

$$C = f_n \star_{n-1} (f_{n-1} \star_{n-2} (\cdots \star_1 f_1 \times g_1 \star_1 \cdots) \star_{n-2} g_{n-1}) \star_{n-1} g_n,$$

where, for every k in $\{1, \dots, n\}$, f_k and g_k are k -cells of \mathcal{C} . The context C is a *whisker of \mathcal{C}* if the n -cells f_n and g_n are identities. If f is an n -cell of \mathcal{C} with boundary x , one denotes by $C[f]$ the n -cell of \mathcal{C} obtained by replacing x with f in C .

If Γ is a cellular extension of \mathcal{C} , then every non-identity $(n+1)$ -cell f of $\mathcal{C}[\Gamma]$ has a decomposition

$$f = C_1[\varphi_1] \star_n \cdots \star_n C_k[\varphi_k],$$

with $k \geq 1$ and, for every i in $\{1, \dots, k\}$, φ_i in Γ and C_i a whisker of $\mathcal{C}[\Gamma]$.

The *category of contexts of \mathcal{C}* is denoted by $\text{Ct}(\mathcal{C})$, its objects are the n -cells of \mathcal{C} and its morphisms from f to g are the contexts C of \mathcal{C} such that $C[f] = g$ holds. When \mathbf{C} is a category, the category $\text{Ct}(\mathbf{C})$ of contexts of \mathbf{C} is isomorphic to the category FC of factorisations of \mathbf{C} . A *natural system on \mathcal{C}* is a $\text{Ct}(\mathcal{C})$ -module, *i.e.*, a functor D from $\text{Ct}(\mathcal{C})$ to the category Ab of Abelian groups; we denote by D_u and by D_C the images of an n -cell u and of a context C of \mathcal{C} through the functor D .

5. Abelianisation of polygraphic resolutions

5.6.2. Identities among relations. Let Σ be an n -polygraph, seen as an $(n, n - 1)$ -polygraph. An n -cell f of Σ^\top is *closed* when its source and target coincide. The natural system $\Pi(\Sigma)$ on $\bar{\Sigma}$ of *identities among relations of Σ* is defined as follows:

- If u is an $(n - 1)$ -cell of $\bar{\Sigma}$, the Abelian group $\Pi(\Sigma)_u$ is generated by one element $[f]$, for each n -cell $f : v \Rightarrow v$ of Σ^\top such that $\bar{v} = u$, submitted to the following relations:

- if $f : v \rightarrow v$ and $g : v \rightarrow v$ are n -cells of Σ^\top , with $\bar{v} = u$, then

$$[f \star_{n-1} g] = [f] + [g];$$

- if $f : v \rightarrow w$ and $g : w \rightarrow v$ are n -cells of Σ^\top , with $\bar{v} = \bar{w} = u$, then

$$[f \star_{n-1} g] = [g \star_{n-1} f].$$

- If $g = C[f]$ is a factorisation in $\bar{\Sigma}$, then the morphism $\Pi(\Sigma)_C : \Pi(\Sigma)_f \rightarrow \Pi(\Sigma)_g$ of groups is defined by

$$\Pi(\Sigma)_C([f]) = [\widehat{C}[f]],$$

where \widehat{C} is any representative context for C in Σ^* . We recall from [19] that the value of $\Pi(\Sigma)$ does not depend on the choice of \widehat{C} , proving that $\Pi(\Sigma)$ is a natural system on $\bar{\Sigma}$ and allowing one to denote this element of $\Pi(\Sigma)_g$ by $C[f]$.

As consequences of the defining relations of each group $\Pi(\Sigma)_u$, we get the following equalities:

$$[1_u] = 0 \quad [f^-] = -[f] \quad [g \star_{n-1} f \star_{n-1} g^-] = [f]$$

for every $n - 1$ -cell u and every n -cells $f : u \rightarrow u$ and $g : v \rightarrow u$ of Σ^\top .

5.6.3. Lemma. *Let Σ be a 2-polygraph. For every closed 2-cell f of Σ^\top , we have $[f] = 0$ in $F_C[\Sigma_2]$ if, and only if, $[f] = 0$ holds in $\Pi(\Sigma)$.*

Proof. To prove that $[f] = 0$ implies $[f] = 0$, we check that the relations defining $\Pi(\Sigma)$ are also satisfied in $F_C[\Sigma_2]$. The first relation is given by the definition of the map $[\cdot]$. The second relation is checked as follows:

$$[f \star_1 g] = [f] + [g] = [g] + [f] = [g \star_1 f].$$

Conversely, let us consider a 2-cell $f : w \Rightarrow w$ in Σ^\top such that $[f] = 0$ holds. We decompose f into:

$$f = u_1 \varphi_1^{\varepsilon_1} v_1 \star_1 \cdots \star_1 u_p \varphi_p^{\varepsilon_p} v_p$$

where φ_i is a 2-cell of Σ , u_i and v_i are 1-cells of Σ^\top , ε_i is an element of $\{-, +\}$. Then we get:

$$0 = [f] = \sum_{i=1}^p \varepsilon_i \bar{u}_i [\varphi_i] \bar{v}_i$$

Since $F_C[\Sigma_2]$ is freely generated, as an FC -module, by the elements $[\varphi]$ of $F_C[\Sigma_2]_{\bar{\varphi}}$, for φ a 2-cell of Σ , this implies the existence of a self-inverse permutation τ of $\{1, \dots, p\}$ such that the following relations are satisfied:

$$\varphi_i = \varphi_{\tau(i)} \quad \bar{u}_i = \bar{u}_{\tau(i)} \quad \bar{v}_i = \bar{v}_{\tau(i)} \quad \varepsilon_i = -\varepsilon_{\tau(i)}.$$

5.6. Homological syzygies and identities among relations

Let us denote, for every $1 \leq i \leq p$, the source and target of $\varphi_i^{\varepsilon_i}$ by w_i and w'_i respectively. They satisfy $\bar{w}_i = \bar{w}'_i$. We also fix a section $\hat{\cdot}$ and a left strategy σ for the 2-polygraph Σ . In particular, the section satisfies $\hat{u} = \hat{v}$ for every 1-cells u and v such that $\bar{u} = \bar{v}$.

For every $1 \leq i \leq p$, we denote by f_i the following 2-cell of Σ^\top :

$$f_i = \sigma_{u_i w_i v_i}^- \star_1 u_i \varphi_i^{\varepsilon_i} v_i \star_1 \sigma_{u_i w'_i v_i}.$$

Using the facts that w is equal to both $u_1 w_1 v_1$ and $u_p w'_p v_p$ and that $u_i w'_i v_i$ is equal to $u_{i+1} w_{i+1} v_{i+1}$ for every $1 \leq i < p$, we can write the 2-cell f of the $(2, 1)$ -category Σ^\top as the following composite:

$$f = \sigma_w \star_1 f_1 \star_1 f_2 \star_1 \cdots \star_1 f_p \star_1 \sigma_w^-.$$

As a consequence, we get:

$$[f] = [\sigma_w^- \star_1 f \star_1 \sigma_w] = \sum_{i=1}^p [f_i].$$

In order to conclude this proof, it is sufficient to check that, for every $1 \leq i \leq p$, we have the equality $[f_{\tau(i)}] = -[f_i]$. Let us fix an i in $\{1, \dots, p\}$ and let us compute $[f_i]$. Since σ is a left normalisation strategy, we have

$$\sigma_{u_i w_i v_i} = \sigma_{u_i w_i v_i} \star_1 \sigma_{\hat{u}_i w_i v_i} \star_1 \sigma_{\widehat{u}_i w_i v_i}$$

and, using the fact that $\widehat{u}_i w_i = \widehat{u}_i w'_i$,

$$\sigma_{u_i w'_i v_i} = \sigma_{u_i w'_i v_i} \star_1 \sigma_{\widehat{u}_i w'_i v_i} \star_1 \sigma_{\widehat{u}_i w_i v_i}.$$

This gives:

$$[f_i] = [\sigma_{\widehat{u}_i w_i v_i}^- \star_1 \sigma_{\widehat{u}_i w_i v_i}^- \star_1 \sigma_{\widehat{u}_i w_i v_i}^- \star_1 u_i \varphi_i^{\varepsilon_i} v_i \star_1 \sigma_{u_i w'_i v_i} \star_1 \sigma_{\widehat{u}_i w'_i v_i} \star_1 \sigma_{\widehat{u}_i w_i v_i}].$$

We can remove the final 2-cell $\sigma_{\widehat{u}_i w_i v_i}$ and its inverse at the beginning of the composition, to get:

$$[f_i] = [\sigma_{\widehat{u}_i w_i v_i}^- \star_1 \sigma_{\widehat{u}_i w_i v_i}^- \star_1 u_i \varphi_i^{\varepsilon_i} v_i \star_1 \sigma_{u_i w'_i v_i} \star_1 \sigma_{\widehat{u}_i w'_i v_i}].$$

Then, we use exchange relations to get:

$$[f_i] = [\sigma_{\widehat{u}_i w_i v_i}^- \star_1 \widehat{u}_i \varphi_i^{\varepsilon_i} \star_1 \sigma_{\widehat{u}_i w'_i v_i}] \bar{v}_i.$$

Now, let us compute $[f_{\tau(i)}]$. We already know that $\varphi_{\tau(i)} = \varphi_i$ and $\varepsilon_{\tau(i)} = -\varepsilon_i$. As a consequence, we get $w_{\tau(i)} = w'_i$ and $w'_{\tau(i)} = w_i$. Moreover, we have $\widehat{u}_{\tau(i)} = \widehat{u}_i$, so that we have:

$$[f_{\tau(i)}] = [\sigma_{\widehat{u}_i w_i v_{\tau(i)}}^- \star_1 \sigma_{\widehat{u}_i w_i v_{\tau(i)}}^- \star_1 \sigma_{\widehat{u}_i w'_i v_{\tau(i)}} \star_1 u_{\tau(i)} \varphi_i^{-\varepsilon_i} v_{\tau(i)} \star_1 \sigma_{u_i w_i v_{\tau(i)}} \star_1 \sigma_{\widehat{u}_i w_i v_{\tau(i)}} \star_1 \sigma_{\widehat{u}_i w_i v_{\tau(i)}}].$$

We remove the 2-cell $\sigma_{\widehat{u}_i w_i v_{\tau(i)}}$ and its inverse and, then, we use exchange relations and $\bar{v}_{\tau(i)} = \bar{v}_i$, in order to get:

$$[f_{\tau(i)}] = [\sigma_{\widehat{u}_i w'_i v_i}^- \star_1 \widehat{u}_i \varphi_i^{-\varepsilon_i} \star_1 \sigma_{\widehat{u}_i w_i v_i}] \bar{v}_i = -[f_i].$$

This implies $[f] = 0$, thus concluding the proof. \square

5. Abelianisation of polygraphic resolutions

5.6.4. Lemma. *Let Σ be a 2-polygraph. For every element α in $\mathfrak{h}_2(\Sigma)$, there exists a closed 2-cell f in Σ^\top such that $\alpha = [f]$ holds.*

Proof. Let w be the 1-cell of $\bar{\Sigma}$ such that α belongs to $F_{\bar{\Sigma}}[\Sigma_2]_w$ and let Σ_3 be a homotopy basis of the $(2, 1)$ -category Σ^\top . Since $\delta_2(\alpha) = 0$, Theorem 5.4.3 implies the existence of an element b in $F_{\bar{\Sigma}}[\Sigma_3]_w$ such that $\alpha = \delta_3(b)$ holds. By definition of $F_{\bar{\Sigma}}[\Sigma_3]_w$, we can write

$$b = \sum_{i=1}^p \varepsilon_i u_i [\alpha_i] v_i$$

with, for every $1 \leq i \leq p$, α_i in Σ_3 , u_i and v_i in $\bar{\Sigma}$ and ε_i in $\{-, +\}$ such that $u_i \bar{\alpha}_i v_i = w$ holds. We fix a section $\hat{\cdot}$ of Σ and we choose 2-cells

$$g_i : \hat{w} \Rightarrow \hat{u}_i s_1(\alpha_i^{\varepsilon_i}) \hat{v}_i \quad \text{and} \quad h_i : \hat{u}_i t_1(\alpha_i^{\varepsilon_i}) \hat{v}_i \Rightarrow \hat{w}.$$

Let A be the following 3-cell of Σ_3^\top :

$$A = (g_1 \star_1 \hat{u}_1 \alpha_1^{\varepsilon_1} \hat{v}_1 \star_1 h_1) \star_1 \cdots \star_1 (g_p \star_1 \hat{u}_p \alpha_p^{\varepsilon_p} \hat{v}_p \star_1 h_p).$$

By definition of $[\cdot]$ on 3-cells, we have

$$[A] = \sum_{i=1}^p [g_i \star_1 \hat{u}_i \alpha_i^{\varepsilon_i} \hat{v}_i \star_1 h_i] = \sum_{i=1}^p ([1_{g_i}] + \varepsilon_i u_i [\alpha_i] v_i + [1_{h_i}]) = b.$$

Finally, we get:

$$\alpha = \delta_3[A] = [s(A)] - [t(A)] = [s(A) \star_1 t(A)^-].$$

Hence $f = s(A) \star_1 t(A)^-$ is a closed 2-cell of Σ^\top that satisfies $\alpha = [f]$. \square

5.6.5. Theorem. *For every 2-polygraph Σ , the natural systems of homological 2-szygies and of identities among relations of Σ are isomorphic.*

Proof. We define a morphism of natural systems $\Phi : \Pi(\Sigma) \rightarrow \mathfrak{h}_2(\Sigma)$ by

$$\Phi([f]) = [f].$$

This definition is correct, since the defining relations of $\Pi(\Sigma)$ also hold in $F_{\mathbb{C}}[\Sigma_2]$, hence in $\mathfrak{h}_2(\Sigma)$. Let us check that Φ is a morphism of natural systems. Indeed, we have

$$\Phi(u[f]v) = \Phi([\hat{u}f\hat{v}]) = [\hat{u}f\hat{v}] = u[f]v = u\Phi([f])v$$

for every 2-cell $f : w \Rightarrow w$ in Σ^\top and 1-cells u, v in $\bar{\Sigma}$ such that $\hat{u}f\hat{v}$ is defined.

Now, let us define a morphism of natural systems $\Psi : \mathfrak{h}_2(\Sigma) \rightarrow \Pi(\Sigma)$. Let α be an element of $\mathfrak{h}_2(\Sigma)_w$. Then there exists a closed 2-cell $f : u \Rightarrow u$ such that $\alpha = [f]$ and $w = \bar{u}$. We define

$$\Psi(\alpha) = [f].$$

This definition does not depend on the choice of f . Indeed, let us assume that $g : v \Rightarrow v$ is a closed 2-cell such that $\alpha = [g]$ holds. It follows that $\bar{v} = w = \bar{u}$. Hence, we can choose a 2-cell $h : u \Rightarrow v$ in Σ^\top . Then we have:

$$\alpha = [f] = [g] = [h \star_1 g \star_1 h^-].$$

As a consequence, we get:

$$[f \star_1 h^- \star_1 g^- \star_1 h] = [f] - [h \star_1 g \star_1 h^-] = 0.$$

Thus

$$0 = [f \star_1 h^- \star_1 g^- \star_1 h] = [f] - [h \star_1 g \star_1 h^-] = [f] - [g].$$

The relations $\Psi\Phi = 1_{\Pi(\Sigma)}$ and $\Phi\Psi = 1_{h_2(\Sigma)}$ are direct consequences of the definitions of Φ and Ψ . \square

5.7. Abelian finite derivation type

An $(n, n-1)$ -category \mathcal{C} is *Abelian* if, for every $(n-1)$ -cell u of \mathcal{C} , the group $\text{Aut}_u^{\mathcal{C}}$ of closed n -cells of \mathcal{C} with source u is Abelian. The Abelian $(n, n-1)$ -category generated by \mathcal{C} is the $(n, n-1)$ -category, denoted by \mathcal{C}_{ab} and defined as the quotient of \mathcal{C} by the cellular extension that contains one n -sphere

$$f \star_{n-1} g \longrightarrow g \star_{n-1} f$$

for every closed n -cells f and g of \mathcal{C} with the same source.

One says that an n -polygraph Σ is of *Abelian finite derivation type*, FDT_{ab} for short, when the Abelian $(n, n-1)$ -category Σ_{ab}^\top admits a finite homotopy basis. In this section, we prove that an n -polygraph is FDT_{ab} if, and only if, the natural system $\Pi(\Sigma)$ of identities among relations of Σ is finitely generated.

In [19], it is proved that, given an n -polygraph Σ , there exists an isomorphism of natural systems on the free $(n-1)$ -category Σ_{n-1}^* :

$$\Pi(\Sigma)_{\bar{u}} \simeq \text{Aut}_{\bar{u}}^{\Sigma_{\text{ab}}^\top}.$$

In fact, this property characterises the natural system $\Pi(\Sigma)$ on the $(n-1)$ -category $\bar{\Sigma}$ up to isomorphism. In [19], we also proved the following result.

5.7.1. Lemma ([19]). *Let \mathcal{C} be an $(n, n-1)$ -category and let \mathcal{B} be a family of closed n -cells of \mathcal{C} . The following assertions are equivalent:*

- i) *The cellular extension $\tilde{\mathcal{B}} = \left\{ \tilde{\beta} : \beta \rightarrow 1_{s\beta}, \beta \in \mathcal{B} \right\}$ of \mathcal{C} is a homotopy basis.*
- ii) *Every closed n -cell f in \mathcal{C} has a decomposition*

$$f = (g_1 \star_{n-1} C_1 [\beta_1^{\varepsilon_1}] \star_{n-1} g_1^-) \star_{n-1} \cdots \star_{n-1} (g_p \star_{n-1} C_p [\beta_p^{\varepsilon_p}] \star_{n-1} g_p^-) \quad (3)$$

with, for every $1 \leq i \leq p$, β_i in \mathcal{B} , ε_i in $\{-, +\}$, C_i a whisker of \mathcal{C} and g_i an n -cell of \mathcal{C} .

5.7.2. Proposition. *An n -polygraph Σ is FDT_{ab} if, and only if, the natural system $\Pi(\Sigma)$ on $\bar{\Sigma}$ is finitely generated.*

5. Abelianisation of polygraphic resolutions

Proof. Let us assume that the n -polygraph Σ is FDT_{ab} . Then the Abelian $(n, n-1)$ -category $\Sigma_{\text{ab}}^{\top}$ has a finite homotopy basis \mathcal{B} . Let $\partial\mathcal{B}$ be the set of closed n -cells of $\Sigma_{\text{ab}}^{\top}$ defined by:

$$\partial\mathcal{B} = \{ \partial\beta = s(\beta) \star_{n-1} t(\beta)^{-}, \beta \in \mathcal{B} \}.$$

By Lemma 5.7.1, any closed n -cell f in $\Sigma_{\text{ab}}^{\top}$ can be written

$$f = (g_1 \star_{n-1} C_1[\partial\beta_1^{\varepsilon_1}] \star_{n-1} g_1^{-}) \star_{n-1} \cdots \star_{n-1} (g_p \star_{n-1} C_p[\partial\beta_p^{\varepsilon_p}] \star_{n-1} g_p^{-}),$$

with, for every $1 \leq i \leq p$, β_i in \mathcal{B} , ε_i in $\{-, +\}$, C_i a whisker of Σ^{\top} and g_i an n -cell of Σ^{\top} . As a consequence, for any identity among relations $[f]$ in $\Pi(\Sigma)$, we have:

$$[f] = \sum_{i=1}^k \varepsilon_i [g_i \star_{n-1} C_i[\partial\beta_i] \star_{n-1} g_i^{-}] = \sum_{i=1}^k \varepsilon_i C_i[\partial\beta_i].$$

Thus, the elements of $[\partial\mathcal{B}]$ form a generating set for the natural system $\Pi(\Sigma)$. Hence $\Pi(\Sigma)$ is finitely generated.

Conversely, suppose that the natural system $\Pi(\Sigma)$ is finitely generated. This means that there exists a finite set \mathcal{B} of closed n -cells of $\Sigma_{\text{ab}}^{\top}$ such that the following property is satisfied: for every $(n-1)$ -cell u of $\bar{\Sigma}$ and every closed n -cell f with source v of $\Sigma_{\text{ab}}^{\top}$ such that $\hat{v} = u$, one can write

$$[f] = \sum_{i=1}^p \varepsilon_i C_i[\beta_i]$$

with, for every $1 \leq i \leq p$, β_i in \mathcal{B} , C_i a whisker of $\bar{\Sigma}$ and ε_i an integer; moreover, each whisker C_i is such that, for every representative \hat{C}_i of C_i in $\Sigma_{\text{ab}}^{\top}$, $\hat{C}_i[\beta_i]$ is a closed n -cell of $\Sigma_{\text{ab}}^{\top}$ whose source v_i satisfies $\bar{v}_i = v$. We fix, for every i , an n -cell $g_i : \hat{v} \Rightarrow v_i$ in Σ^{\top} . Then, the properties of $\Pi(\Sigma)$ imply:

$$\begin{aligned} [f] &= \sum_{i=1}^p [g_i \star_{n-1} \hat{C}_i[\beta_i^{\varepsilon_i}] \star_{n-1} g_i^{-}] \\ &= [(g_1 \star_{n-1} \hat{C}_1[\beta_1^{\varepsilon_1}] \star_{n-1} g_1^{-}) \star_{n-1} \cdots \star_{n-1} (g_p \star_{n-1} \hat{C}_p[\beta_p^{\varepsilon_p}] \star_{n-1} g_p^{-})]. \end{aligned}$$

We use the isomorphism between $\Pi(\Sigma)_u$ and $\text{Aut}_v^{\Sigma_{\text{ab}}^{\top}}$ and Lemma 5.7.1 to deduce that the cellular extension $\tilde{\mathcal{B}} = \{ \tilde{\beta} : \beta \rightarrow 1_{s\beta}, \beta \in \mathcal{B} \}$ of $\Sigma_{\text{ab}}^{\top}$ is a homotopy basis. Thus Σ is FDT_{ab} . \square

In [19], the authors have proved that the property to be finitely generated for $\Pi(\Sigma)$ is Tietze-invariant for finite polygraphs: if Σ and Υ are two Tietze-equivalent finite n -polygraphs, then the natural system $\Pi(\Sigma)$ is finitely generated if, and only if, the natural system $\Pi(\Upsilon)$ is finitely generated.

By Proposition 5.7.2, we deduce that the property FDT_{ab} is Tietze-invariant for finite polygraphs, so that one can say that an n -category is FDT_{ab} when it admits a presentation by a finite $(n+1)$ -polygraph which is FDT_{ab} . In this way, the following result relates the homological property FP_3 and the homotopical property FDT_{ab} .

5.7.3. Theorem. *Let \mathbf{C} be a category with a finite presentation Σ . The following conditions are equivalent:*

- i) the category \mathbf{C} is of homological type FP_3 ,
- ii) the natural system $h_2(\Sigma)$ on \mathbf{C} is finitely generated,
- iii) the natural system $\Pi(\Sigma)$ on \mathbf{C} is finitely generated,
- iv) the category \mathbf{C} is FDT_{ab} .

Proof. The equivalence between **i)** and **ii)** comes from the definition of the property FP_3 . The equivalence between **ii)** and **iii)** is a consequence of Theorem 5.6.5. The last equivalence is given by Proposition 5.7.2. \square

In Corollary 5.4.4, we have seen that FDT_3 implies FP_3 . We expect that the reverse implication is false in general, which is equivalent to proving that FDT_{ab} does not imply FDT_3 , since FP_3 is equivalent to FDT_{ab} for finitely presented categories.

6. EXAMPLES

6.1. A concrete example of reduced standard polygraphic resolution

Let us denote by A the monoid with one non-unit element, a , and with product given by $aa = a$. The standard presentation of A , seen as a category, is the reduced and convergent 2-polygraph, denoted by As , with one 0-cell (for 1), one 1-cell (for a) and one 2-cell $aa \Rightarrow a$. Here we use diagrammatic notations, where a is denoted by a vertical string $|$ and the 2-cell $aa \Rightarrow a$ is pictured as \blacktriangledown . The 2-polygraph As has one critical branching:

$$(\blacktriangledown|, | \blacktriangledown).$$

The corresponding generating confluence (for the rightmost normalisation strategy) is the 3-cell:

By extending As with that 3-cell, one gets a finite, acyclic $(3, 1)$ -polygraph, still denoted by As and which is a partial polygraphic resolution of A of length 3. We conclude that the monoid A has the property FDT_3 and, thus, is of homological type FP_3 . In particular, the natural system $h_2(As)$ of homological 2-szygies of As is generated by the following element:

$$\delta_3[\blacktriangledown] = [\blacktriangledown|] - [|\blacktriangledown] = [\blacktriangledown|] + [|\blacktriangledown] - [|\blacktriangledown] - [\blacktriangledown|] = [\blacktriangledown|]a - a[\blacktriangledown|].$$

The 2-polygraph As has exactly one critical triple branching:

$$b = (\blacktriangledown||, | \blacktriangledown|, || \blacktriangledown).$$

This triple critical branching b has shape $(f\hat{a}, g\hat{a}, \rho_{u\hat{a}})$ with


$$f = \blacktriangledown| \quad \text{and} \quad g = | \blacktriangledown.$$

6. Examples


We compute the 3-cells $\omega_{f,g}$, $\sigma_{f\hat{a}}^*$ and $\sigma_{g\hat{a}}^*$, using their definitions and the properties of the rightmost normalisation strategy σ , to get:

$$\omega_{f,g}\hat{a} = \text{[Diagram: a triangle with a downward arrow on the right side]} \quad \sigma_{f\hat{a}}^* = \text{[Diagram: two triangles pointing down, one to the left of the other, with a star symbol between them]} \quad \sigma_{g\hat{a}}^* = \text{[Diagram: two triangles pointing down, one to the right of the other, with a star symbol between them]} .$$

We fill the diagram defining $\omega_b = \text{[Diagram: a triangle with a downward arrow on the right side]}$, obtaining:



Contracting one dimension, we see that the 4-cell $\text{[Diagram: a triangle with a downward arrow on the right side]}$ is, in fact, Mac Lane's pentagon, or Stasheff's polytope K_4 :



We get a finite, acyclic $(4, 1)$ -polygraph $c_3(As)$ which is a partial polygraphic resolution of A of length 4, proving that A has the property FDT_4 and, as a consequence, that it is of homological type FP_4 . In particular, the natural system $h_3(As)$ of homological 3-syzygies of As is generated by the following element:

$$\begin{aligned} \delta_4[\text{[Diagram: a triangle with a downward arrow on the right side]}] &= [\text{[Diagram: a triangle with a downward arrow on the right side]}] + [\text{[Diagram: a triangle with a downward arrow on the right side]}] + [\text{[Diagram: a triangle with a downward arrow on the right side]}] - [\text{[Diagram: a triangle with a downward arrow on the right side]}] - [\text{[Diagram: a triangle with a downward arrow on the right side]}] \\ &= [\text{[Diagram: a triangle with a downward arrow on the right side]}] + [\text{[Diagram: a triangle with a downward arrow on the right side]}] + [\text{[Diagram: a triangle with a downward arrow on the right side]}] - [\text{[Diagram: a triangle with a downward arrow on the right side]}] - [\text{[Diagram: a triangle with a downward arrow on the right side]}] \\ &= a [\text{[Diagram: a triangle with a downward arrow on the right side]}] - [\text{[Diagram: a triangle with a downward arrow on the right side]}] + [\text{[Diagram: a triangle with a downward arrow on the right side]}] a. \end{aligned}$$

Iterating the process, we get a resolution of A by an acyclic $(\infty, 1)$ -polygraph $c_\infty(As)$. For every natural number, $c_\infty(As)$ has exactly one n -cell, whose shape is Stasheff's polytope K_n . For example, in dimension 5, the generating 5-cell ω_b is associated to the following critical quadruple branching:

$$b = (\text{[Diagram: a triangle with a downward arrow on the right side]} \text{[Diagram: a triangle with a downward arrow on the right side]} \text{[Diagram: a triangle with a downward arrow on the right side]} \text{[Diagram: a triangle with a downward arrow on the right side]} \text{[Diagram: a triangle with a downward arrow on the right side]}).$$


To compute the source and target of the corresponding 4-cell ω_b , we use the inductive construction of the rightmost strategy σ . Alternatively, one can also start from the 2-dimensional source and target of ω_b ,

6.1. A concrete example of reduced standard polygraphic resolution


which are obtained as the 2-cells associated to the source $aaaaa$ of the critical quadruple branching b by the leftmost and the rightmost strategies, respectively:

$$s_2(\omega_b) = \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \quad \text{and} \quad t_2(\omega_b) = \begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} .$$

Then one computes all the possible 3-cells from $s_2(\omega_b)$ to $t_2(\omega_b)$ and one fills all the 3-dimensional spheres with 4-cells built from the generating 4-cell $\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array}$. Either way, we obtain the following composite 4-cell as the source of ω_b :



And the following composite 4-cell is the target of ω_b :



6. Examples

The corresponding generator of the natural system $h_4(As)$ of homological 4-syzygies of As is:

$$\begin{aligned} \delta_5[\omega_b] &= \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] + \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] + \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] - \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] - \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] - \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] \\ &= \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right] a - a \left[\begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right]. \end{aligned}$$

6.2. The category \mathbf{Epi}

We denote by \mathbf{Epi} the subcategory of the simplicial category whose 0-cells are the natural numbers and whose morphisms from n to p are the monotone surjections from $\{0, \dots, n\}$ to $\{0, \dots, p\}$. This category is studied in [28], where it is denoted by Δ^{epi} .

The category \mathbf{Epi} admits a presentation by the (infinite) 2-polygraph Σ with the natural numbers as 0-cells, with 1-cells

$$n + 1 \xrightarrow{s_i} n \quad 0 \leq i \leq n,$$

where s_i represents the map

$$s_i(j) = \begin{cases} j & \text{if } 0 \leq j \leq i, \\ j - 1 & \text{if } i + 1 \leq j \leq n + 1, \end{cases}$$

and with 2-cells

$$\begin{array}{ccc} & s_i \rightarrow n + 1 & s_j \rightarrow n \\ n + 2 & \Downarrow s_{i,j} & n \\ & s_{j+1} \rightarrow n + 1 & s_i \rightarrow n \end{array} \quad 0 \leq i \leq j \leq n + 1.$$

Let us prove that this 2-polygraph is convergent. For termination, given a 1-cell $u = s_{i_1} \dots s_{i_k}$ of Σ^* , we define the natural number $\nu(u)$ as the number of pairs (i_p, i_q) such that $i_p \leq i_q$, with $1 \leq p < q \leq k$. In particular, we have $\nu(s_i s_j) = 1$ and $\nu(s_{j+1} s_i) = 0$ when $i \leq j$, *i.e.*, when $s_i s_j$ is the source and $s_{j+1} s_i$ is the target of a 2-cell of Σ . Moreover, we have $\nu(w u w') > \nu(w v w')$ when $\nu(u) > \nu(v)$ holds. Thus, for every non-identity 2-cell $f : u \Rightarrow v$ in Σ^* , the strict inequality $\nu(u) > \nu(v)$ is satisfied, yielding termination of Σ .

The 2-polygraph Σ has one critical branching $(s_{i,j} s_k, s_i s_{j,k})$ for every possible $0 \leq i \leq j \leq k \leq n + 2$ and it is confluent, so that we get a partial resolution of \mathbf{Epi} by the acyclic $(3, 1)$ -polygraph $c_2(\Sigma)$ obtained by extending Σ with all the 3-cells filling the confluence diagrams associated to the critical branchings:

$$\begin{array}{ccccc} & & s_{j+1} s_i s_k & \xrightarrow{s_{j+1} s_{i,k}} & s_{j+1} s_{k+1} s_i & \xrightarrow{s_{j+1, k+1} s_i} & s_{k+2} s_{j+1} s_i \\ & s_{i,j} s_k \nearrow & & & & & \\ s_i s_j s_k & & & & & & \\ & s_i s_{j,k} \searrow & & & & & \\ & & s_i s_{k+1} s_j & \xrightarrow{s_{i, k+1} s_j} & s_{k+2} s_i s_j & \xrightarrow{s_{k+2} s_{i,j}} & s_{k+2} s_{j+1} s_i \\ & & & & & & \end{array}$$

To simplify notations, we draw each 1-cell s_i as a vertical string $|_i$, each 2-cell $s_{i,j}$ as $\bowtie_{i,j}$, so that each 3-cell $s_{i,j,k}$ has the same shape as the Yang-Baxter relation, or permutohedron of order 3:

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \end{array} \xrightarrow{\text{Diagram 4}} \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \\ \text{Diagram 7} \end{array}$$

Using those notations, the natural system $h_2(\Sigma)$ of homological 2-syzygies of Σ is generated by the elements

$$\delta_3 [\bowtie_{i,j,k}] = \left[\text{Diagram 1} \right] - \left[\text{Diagram 2} \right] = \begin{cases} ([\bowtie_{i,j}]s_k - s_{k+2}[\bowtie_{i,j}]) \\ + (s_{j+1}[\bowtie_{i,k}] - [\bowtie_{i,k+1}]s_j) \\ + ([\bowtie_{j+1,k+1}]s_i - s_i[\bowtie_{j,k}]). \end{cases}$$


The 2-polygraph Σ has one critical triple branching

$$(s_{i,j}s_k s_l, s_i s_{j,k} s_l, s_i s_j s_{k,l})$$

for every possible $0 \leq i \leq j \leq k \leq l \leq n + 3$. This yields a partial resolution of **Epi** by the acyclic $(4, 1)$ -polygraph $c_3(\Sigma)$ with one 4-cell $s_{i,j,k,l}$ for every possible $0 \leq i \leq j \leq k \leq l \leq n + 3$. In string diagrams, omitting the subscripts, each critical triple branching is written

$$(\bowtie ||, | \bowtie |, || \bowtie).$$

With the same conventions, the corresponding 4-cell has the shape of the permutohedron of order 4:



As usual, the elements $\delta_4 [\bowtie_{i,j,k,l}]$, for $0 \leq i \leq j \leq k \leq l \leq n + 3$, form a generating set for the natural system $h_3(\Sigma)$ of homological 3-syzygies of Σ .

More generally, this construction extends to a polygraphic resolution of **Epi** where, for every natural number n , the generating n -cell has the shape of a permutohedron of order n .

REFERENCES

- [1] David J. Anick, *On the homology of associative algebras*, Transactions of the American Mathematical Society **296** (1986), no. 2, 641–659.
- [2] Dimitri Ara and François Métayer, *The Brown-Golasinski model structure on strict ∞ -groupoids revisited*, Homology Homotopy Appl. **13** (2011), no. 1, 121–142.
- [3] Franz Baader and Tobias Nipkow, *Term rewriting and all that*, Cambridge University Press, 1998.
- [4] Jean-Christophe Bach, Émilie Balland, Paul Brauner, Radu Kopetz, Pierre-Étienne Moreau, and Antoine Reilles, *TOM manual*, 2009.
- [5] Hans-Joachim Baues, *Combinatorial homotopy and 4-dimensional complexes*, de Gruyter Expositions in Mathematics, vol. 2, Walter de Gruyter & Co., Berlin, 1991.
- [6] Hans-Joachim Baues and Elias Gabriel Minian, *Track extensions of categories and cohomology*, K-Theory **23** (2001), no. 1, 1–13.
- [7] Hans-Joachim Baues and Günther Wirsching, *Cohomology of small categories*, J. Pure Appl. Algebra **38** (1985), no. 2-3, 187–211.
- [8] Ronald V. Book and Friedrich Otto, *String-rewriting systems*, Texts and Monographs in Computer Science, Springer-Verlag, 1993.
- [9] Kenneth S. Brown, *The geometry of rewriting systems: a proof of the Anick-Groves-Squier theorem*, Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989), Math. Sci. Res. Inst. Publ., vol. 23, Springer, New York, 1992, pp. 137–163.
- [10] Ronald Brown and Johannes Huebschmann, *Identities among relations*, Low-dimensional topology (Bangor, 1979), London Math. Soc. Lecture Note Ser., vol. 48, Cambridge Univ. Press, Cambridge, 1982, pp. 153–202.
- [11] Albert Burroni, *Higher-dimensional word problems with applications to equational logic*, Theoretical Computer Science **115** (1993), no. 1, 43–62.
- [12] Daniel E. Cohen, *A monoid which is right-FP $_{\infty}$ but not left-FP $_1$* , Bull. London Math. Soc. **24** (1992), no. 4, 340–342.
- [13] Robert Cremanns and Friedrich Otto, *Finite derivation type implies the homological finiteness condition FP $_3$* , J. Symbolic Comput. **18** (1994), no. 2, 91–112.
- [14] ———, *For groups the property of having finite derivation type is equivalent to the homological finiteness condition FP $_3$* , J. Symbolic Comput. **22** (1996), no. 2, 155–177.
- [15] Ralph H. Fox, *Free differential calculus. I. Derivation in the free group ring*, Ann. of Math. (2) **57** (1953), 547–560.
- [16] John R. J. Groves, *Rewriting systems and homology of groups*, Groups—Canberra 1989, Lecture Notes in Mathematics, vol. 1456, Springer-Verlag, 1990, pp. 114 – 141.
- [17] Yves Guiraud, *Termination orders for 3-dimensional rewriting*, J. Pure and Appl. Algebra **207** (2006), no. 2, 341–371.
- [18] Yves Guiraud and Philippe Malbos, *Higher-dimensional categories with finite derivation type*, Theory and Applications of Categories **22** (2009), no. 18, 420–478.
- [19] ———, *Identities among relations for higher-dimensional rewriting systems*, Séminaires et Congrès **26** (2011), 145–161.

-
- [20] Deepak Kapur and Paliath Narendran, *A finite Thue system with decidable word problem and without equivalent finite canonical system*, Theoret. Comput. Sci. **35** (1985), no. 2-3, 337–344.
- [21] Yuji Kobayashi, *Complete rewriting systems and homology of monoid algebras*, J. Pure Appl. Algebra **65** (1990), no. 3, 263–275.
- [22] Yuji Kobayashi and Friedrich Otto, *On homotopical and homological finiteness conditions for finitely presented monoids*, Internat. J. Algebra Comput. **11** (2001), no. 3, 391–403.
- [23] ———, *For finitely presented monoids the homological finiteness conditions FHT and bi-FP₃ coincide*, J. Algebra **264** (2003), no. 2, 327–341.
- [24] Yves Lafont, *A new finiteness condition for monoids presented by complete rewriting systems (after Craig C. Squier)*, J. Pure Appl. Algebra **98** (1995), no. 3, 229–244.
- [25] Yves Lafont, François Métayer, and Krzysztof Worytkiewicz, *A folk model structure on omega-cat*, Adv. Math. **224** (2010), no. 3, 1183–1231.
- [26] Francis William Lawvere, *Functorial semantics of algebraic theories*, Reprints in Theory and Applications of Categories **5** (2004), 1–121.
- [27] Xavier Leroy, *The Objective Caml system*, 2008, (with Damien Doligez, Jacques Garrigue, Didier Rémy and Jérôme Vouillon).
- [28] Muriel Livernet and Birgit Richter, *An interpretation of E_n -homology as functor homology*, Math. Z. **269** (2011), no. 1–2, 193–219.
- [29] Jean-Jacques Lévy, *Réductions correctes et optimales dans le λ -calcul*, 1978, Thèse de doctorat d’État, Université Paris 7.
- [30] Roger C. Lyndon and Paul E. Schupp, *Combinatorial group theory*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1977 edition.
- [31] Simon Marlow (ed.), *Haskell 2010 language report*, 2010.
- [32] Stuart McGlashan, Elton Pasku, and Stephen J. Pride, *Finiteness conditions for rewriting systems*, Internat. J. Algebra Comput. **15** (2005), no. 1, 175–205.
- [33] Paul-André Mellies, *A diagrammatic standardization theorem*, Processes, Terms and Cycles: Steps on the Road to Infinity (Aart Middeldorp, Vincent van Oostrom, Femke van Raamsdonk, and Roel de Vrijer, eds.), vol. 3838, 2002.
- [34] Samuel Mimram, *Computing critical pairs in 2-dimensional rewriting systems*, preprint, 2010.
- [35] Barry Mitchell, *Rings with several objects*, Advances in Mathematics **8** (1972), 1–161.
- [36] François Métayer, *Resolutions by polygraphs*, Theory and Applications of Categories **11** (2003), 148–184.
- [37] Yves Métivier, *About the rewriting systems produced by the knuth-bendix completion algorithm*, Inf. Process. Lett. **16** (1983), no. 1, 31–34.
- [38] Maxwell Herman Alexander Newman, *On theories with a combinatorial definition of "equivalence"*, Annals of Mathematics **43** (1942), no. 2, 223–243.
- [39] Stephen J. Pride, *Low-dimensional homotopy theory for monoids*, Internat. J. Algebra Comput. **5** (1995), no. 6, 631–649.
- [40] Kurt Reidemeister, *Über Identitäten von Relationen*, Abh. Math. Sem. Univ. Hamburg **16** (1949), 114–118.
- [41] Craig C. Squier, *Word problems and a homological finiteness condition for monoids*, J. Pure Appl. Algebra **49** (1987), no. 1-2, 201–217.

REFERENCES

- [42] Craig C. Squier, Friedrich Otto, and Yuji Kobayashi, *A finiteness condition for rewriting systems*, Theoret. Comput. Sci. **131** (1994), no. 2, 271–294.
- [43] Ross Street, *Limits indexed by category-valued 2-functors*, J. Pure Appl. Algebra **8** (1976), no. 2, 149–181.
- [44] ———, *The algebra of oriented simplexes*, J. Pure Appl. Algebra **49** (1987), no. 3, 283–335.
- [45] Terese, *Term rewriting systems*, Cambridge Tracts in Theoretical Computer Science, vol. 55, Cambridge University Press, 2003.
- [46] Axel Thue, *Probleme über Veränderungen von Zeichenreihen nach gegebenen Regeln.*, Kristiania Vidensk. Selsk, Skr. (1914), no. 10, 493–524.
- [47] Charles Wells, *Extension theories for categories*, preliminary report, 1979.