Accepted Manuscript

Title: T3-mediated expression of PGC-1 α via a far upstream located thyroid hormone response element

Authors: Anne Wulf, Angelika Harneit, Meike Kröger, Maxim Kebenko, Marianne G. Wetzel, Joachim M. Weitzel

PII: S0303-7207(08)00027-0
DOI: doi:10.1016/j.mce.2008.01.017
Reference: MCE 6805

To appear in: *Molecular and Cellular Endocrinology*

Received date: 15-11-2007
Revised date: 9-1-2008
Accepted date: 23-1-2008

Please cite this article as: Wulf, A., Harneit, A., Kröger, M., Kebenko, M., Wetzel, M.G., Weitzel, J.M., T3-mediated expression of PGC-1 α via a far upstream located thyroid hormone response element, *Molecular and Cellular Endocrinology* (2007), doi:10.1016/j.mce.2008.01.017

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**T3-mediated expression of PGC-1 α via a far upstream located thyroid hormone
response element**

5

Anne Wulf¹, Angelika Harneit¹, Meike Kröger¹, Maxim Kebenko¹,
Marianne G. Wetzel¹, Joachim M. Weitzel^{1,2,3}

10

¹ Zentrum für Experimentelle Medizin, Institut für Biochemie und Molekularbiologie I,
Universitätsklinikum Hamburg-Eppendorf, 20246 Hamburg, Germany; ² Institut für
Experimentelle Endokrinologie, Charité Universitätsmedizin Berlin, 10117 Berlin, Germany;
³ Department of Reproductive Biology, FBN Dummerstorf, 18196 Dummerstorf, Germany.

15

³ To whom correspondence should be addressed:

20

Department of Reproductive Biology, FBN Dummerstorf,
Wilhelm-Stahl-Allee 2, 18196 Dummerstorf, Germany.

Tel.: +49-38208-68788

Fax: +49-38208-68752

E-mail: weitzel@fbn-dummerstorf.de

25

Running Title: T3 regulation of PGC-1 α

Abstract

30

Thyroid hormone (T3) has a profound influence on normal development, differentiation and metabolism. T3 induces complex gene expression patterns raises the question of how these expression patterns might be regulated. Since the transcriptional coactivator PGC-1 α (peroxisome proliferators-activated receptor γ coactivator-1 α) induces very similar cellular energy metabolic pathways, we investigated the molecular mechanism of T3 regulation of PGC-1 α . PGC-1 α is rapidly regulated by T3, both in vivo and in cell culture. Transient transfection experiments demonstrated binding of the thyroid hormone receptor (TR) to a response element located at -4 kb upstream of the transcriptional start site within the PGC-1 α gene. Introducing of a single copy of the -4 kb TRE in a heterologous promoter context is sufficient to maintain T3 responsiveness. Chromatin immunoprecipitation analysis revealed increased histone acetylation upon stimulation of T3. Finally, TR binds the -4 kb TRE in electrophoretic mobility shift assays, identifying PGC-1 α as a direct target of TR action. Since T3 directly regulates PGC-1 α and PGC-1 α coactivates liganded TR, we suggest an autoregulatory feed-forward loop of PGC-1 α activation upon T3 treatment.

45

Keywords: thyroid hormone, metabolism, transcription factor, nuclear receptor, coactivator.

Introduction

50

Thyroid hormone (T3) triggers mitochondrial biogenesis, adaptive thermogenesis and hepatic gluconeogenesis (Yen, 2001). Thyroid hormone regulates gene expression via binding to thyroid hormone receptors (TRs), which belong to the family of nuclear receptors. TRs can bind to thyroid hormone response elements (TREs) either as a homodimer or as a

55

heterodimer preferentially with retinoid X receptor (RXR). The TR interacts with co-factors, which mediate T3 signalling to the basal transcriptional machinery and activates or represses gene transcription (Zhang and Lazar, 2000). On positively T3-regulated genes unliganded TR suppresses basal gene transcription by interacting with corepressors such as nuclear receptor corepressor (NCoR) and silencing mediator of retinoid and thyroid receptors (SMRT).

60

Corepressors are associated with histone deacetylases (HDACs), which modify chromatin into a compact and transcriptionally silent structure. Binding of T3 to the receptor (liganded TR) leads to conformational changes of the TR, which releases the corepressor and recruits the coactivators such as steroid receptor coactivator 1 (SRC-1) or peroxisome proliferators-activated receptor γ coactivator-1 α (PGC-1 α). Coactivators possess or recruit proteins with

65

histone acetyltransferase activity (HAT), which remodels chromatin into an open structure allowing for transcription (Rosenfeld et al., 2006). In contrast to positively T3-regulated genes the mechanism of negatively T3-regulated gene transcription is less clear (Lazar, 2003).

70

Beside positively and negatively regulated target genes, gene expression analysis indicated different induction kinetics (Miller et al., 2001; Weitzel et al., 2001b; Flores-Morales et al., 2002; Weitzel et al., 2003a). While some target genes were rapidly induced within 6 hours after the administration of T3 ('early' induction), a second subset of T3-target genes were delayed regulated after a lag period of 12-24 hours ('late' induction). Whereas numerous TREs have been identified in early-induced gene promoters (Zilz et al., 1990; Weitzel et al., 2001a), we failed to detect TREs in late-induced gene promoters (Weitzel et al., 2003a).

75

These later genes might be regulated in a different manner, probably via an 'intermediate factor'. As a candidate factor for such an intermediate factor we suggested PGC-1 α (Weitzel et al., 2003b). PGC-1 α regulates adaptive thermogenesis in brown adipose tissue, hepatic gluconeogenesis and mitochondrial biogenesis in skeletal muscle and heart (among others) all of which resembles those effects which are concurrently regulated by thyroid hormone as well; reviewed in (Finck and Kelly, 2006; Handschin and Spiegelman, 2006). PGC-1 α itself is regulated by several environmental and endocrine signals, including T3 (Weitzel et al.,

80

2001b; Irrcher et al., 2003; Weitzel et al., 2003a). Furthermore, PGC-1 α binds and coactivates a great variety of transcription factors implicated in the control of cellular energy metabolic pathways (e.g. TRs, ERRs and NRFs) probably serving as bridging factor for inducing late T3-regulated genes.

The aim of the following study was to investigate the molecular mechanism of T3-mediated activation of PGC-1 α . Our data indicate that PGC-1 α is regulated via a far upstream located TRE within the PGC-1 α gene. PGC-1 α is a direct target of TR and is rapidly regulated by T3. Given that T3 activates PGC-1 α and PGC-1 α coactivates liganded TR, our data indicate an autoregulatory feed-forward loop upon stimulation by T3.

Material and Methods

Animals and treatments

Adult male Wistar rats (Charles River Deutschland GmbH, Sulzfeld, Germany) were housed in the animal care facility of the University Hospital Hamburg-Eppendorf according to the UKE animal care guidelines. Hypothyroidism was provoked by an intraperitoneal injection of Na¹³¹I (Amersham Biosciences, Freiburg, Germany) (250 μ Ci/100 g body weight) and hyperthyroidism was provoked by i.p. injection of T3 (50 μ g/ 100 g body weight) (Weitzel et al., 2001b; Weitzel et al., 2003a). Zero and 6 hours after thyroid hormone injection, rats were killed by decapitation and livers were isolated for RNA preparation. Total RNA was prepared using RNA-Clean (Hybaid-AGS, Heidelberg, Germany) according to the recommendations of the manufacturer and were subject to real-time PCR using the LC-DNA Master SYBR Green kit in a LightCycler (Roche Applied Science, Mannheim, Germany) as described (Weitzel et al., 2001b; Weitzel et al., 2003a; Wulf et al., 2007).

Isolation and characterisation of DNA sequences

The TRE-containing promoter fragment B(-316/+109) of rat mGPDH in pGL2-basic has been described previously (Weitzel et al., 2001a) and was further subcloned into pRL-MA (Promega, Mannheim, Germany). The promoter fragment from -4483 to +78 relative to the transcriptional start site of mouse PGC-1 α in pGL3-basic was a kind gift of James Rhee (Boston, USA). Further deletion fragments thereof were amplified by PCR and subcloned into pGL3-basic or pRL-MA vectors. Point mutations were introduced using the QuickChange

site-directed mutagenesis system (Stratagene, Amsterdam, The Netherlands) according to the manufacturer's recommendations. A sequence portion from -4025 to -4010 (5'-tgcaacttgcttagct-3') was designed as double stranded oligonucleotides and ligated upstream of the thymidine kinase (TK) promoter into pRL-MA, generating the construct TRE-TK-Luc. All clones were confirmed by sequencing. In silico analysis to predict nuclear receptor binding sites were performed using NUBIScan (Podvinec et al., 2002) and MatInspector (Genomatix, München, Germany).

Transient Transfection analyses

Human hepatocarcinoma HepG2 were cultured under standard conditions in Dulbecco's modified Eagle's medium plus Glutamax (Invitrogen, Karlsruhe, Germany) and 10% fetal calf serum (Biochrom, Berlin, Germany) as described (Weitzel et al., 2001a; Weitzel et al., 2003a). Rat pituitary GC cells (a gift of Danielle Gourdji, Paris, France) were cultured in a 1:1 mixture of Dulbecco's modified Eagle's and Ham's F12 medium plus 10% fetal calf serum (final concentration of T3: ~ 180 pM). Cells were treated with 100 nM T3 for various time points before cells were harvested and RNA was prepared. Transient transfection experiments and mammalian one-hybrid assays were performed using a calcium phosphate technique described previously.

Electrophoretic mobility shift assay (EMSA)

In vitro-translated chicken TR α 1 was synthesized by a transcription/translation-coupled reticulocyte lysate system (Promega) according to the manufacturer's instructions. One μ l of in vitro-translated TR was incubated with 40 fmol of radioactively labelled, double-stranded oligonucleotide probe (-4029 to -4006 of the PGC-1 α gene, 5'-ctgctgcacttgcttagcttag-3') in the presence of 1 μ g poly (dA-dT) x (dA-dT) as described previously (Weitzel et al., 2001a). For competition experiments a 100-fold molar excess of double-stranded oligonucleotides 5' mut (5'-ctgcgaattcttgcttagcttag-3'), 3' mut (5'-ctgctgcactttgcgaattcgtag-3') or TRE-mGPDH (5'-accgtggccccgctgacccgggtg-3') were added to the binding reaction. Shifted DNA bands were separated on 5% polyacrylamide gels and visualized by autoradiography on XAR films (Kodak, Rochester, NY).

Chromatin Immunoprecipitation (ChIP)

For ChIP analyses, 1 μ g of luciferase reporter, 0.1 μ g TR and 0.1 μ g RXR expression plasmids were mixed and transfected into HepG2 cells using 4 μ l Metafectene (Biontix,

München, Germany) per 9.6 cm² dish according to the instructions of the manufacturer and stimulated with 100 nM T3 for 24 hours. Transfection efficiencies were ~ 11%. Cells were fixed with 1% formaldehyde for 4 min at 37°C, washed with PBS (136 mM NaCl, 50 mM Na₂HPO₄, pH 7.4) containing 1 mM protease inhibitor AEBSF (Roche Applied Bioscience) and resolved in Lysis buffer (Upstate, Hamburg, Germany). DNA was sheared by ultra sound for 30 sec in a Sonorex ultrasonic bath (Bandelin, Berlin, Germany), centrifuged and the supernatant was diluted in ChIP dilution buffer. Twenty µl of this material was taken away as input control and to the remaining portion a 1: 500 dilution of anti-acetylated histone H3 antibodies (Upstate) or irrelevant antibodies was added and incubated overnight at 4°C. Sixty µl of salmon sperm DNA/protein A agarose slurry was added and incubated at 4°C for 1 hour, centrifuged, washed several times according to the instructions of the manufacturer and finally suspended in elution buffer (1% SDS, 100 mM NaHCO₃). Histone-DNA cross links were reversed by adding NaCl to a final concentration of 200 mM and incubation at 65°C for 4 hours. Immunoprecipitated DNA was purified by phenol/chloroform extraction according to standard procedures, and finally resolved in 20 µl water. PCR amplification reactions were performed using the forward primer WL328 (5'-ctaagcggtacttactgag-3') and the reverse primer WL339 (5'-ctttatgttttggcgcttc-3'). DNA amplification products (~ 300 bp in length) were radioactively labelled due to the incorporation of radioactively labelled nucleotides, separated on 5% polyacrylamide gels and visualized by autoradiography on XAR films (Kodak).

Results

We previously described PGC-1α as a target of thyroid hormone action (Weitzel et al., 2001b; Weitzel et al., 2003a). After administration of T3 to hypothyroid rats, endogenous steady state levels of PGC-1α mRNA rise 13-fold in liver compared to untreated control animals (Figure 1A). This activation is accomplished by increased PGC-1α protein concentrations. A similar induction has also been detected in rat pituitary GC cells (Figure 1B) which mimic thyroid hormone-mediated hepatic gene expression patterns (Miller et al., 2001; Weitzel et al., 2001b; Weitzel et al., 2003a; Wulf et al., 2007). Since this activation occurs fast (within 4 to 6 hours, Figure 1B) we suggested a direct mechanism of activation by thyroid hormone, probably via a thyroid hormone response element.

185 To analyze whether PGC-1 α is directly regulated via a TRE, we investigated a fragment comprising sequences from -4483 to +78 relative to the transcriptional start site of the PGC-1 α gene. Transient transfection experiment of a luciferase reporter construct of this 4.5 kb fragment together with a fusion protein of TR with the viral activation domain VP16 (TR-VP16) demonstrated an activation of gene expression in a mammalian one-hybrid assay (Figure 2). As positive control for this assay we used an mGPDH promoter-luciferase reporter construct which contains a well characterized TRE (Weitzel et al., 2001a) (not shown), whereas as negative control we used a VP16-only construct without a binding site for a TRE. Similar results have been observed using the 4.5 kb reporter by cotransfection of TR and stimulation with T3 in ‘classical’ transfection experiments (supplementary figure 1).

190 Consecutive 5’ deletion of the 4.5 kb sequences (-2500, -1200, -200 fragments, Figure 2) lost the ability to activate TR-mediated gene expression, indicating a TRE within the 2 kb most upstream located sequence portion. In silico analyses indicated the sequence 5’-TGCACTttgcTTAGCT-3’ at position -4025 with the highest score matching a DR+4 thyroid hormone response element. This sequences was excised and integrated upstream of a TK promoter-driven luciferase reporter. As demonstrated in Figure 2, a single copy of this element was sufficient to activate TR-mediated gene expression (construct TRE-TK-Luc). Thus, the TRE-like sequence at -4 kb is functionally active in cell experiments even in a heterologous promoter context.

200 To investigate the molecular consequences of thyroid hormone-mediated gene expression we performed chromatin immunoprecipitations. The 4.5 kb PGC-1 α promoter reporter construct together with TR was transfected into HepG2 cells and stimulated with T3. Chromatin was fixed and immunoprecipitated with an antibody directed against acetylated histone H3, a marker for transcriptionally active chromatin. Immunoprecipitated DNA was amplified by PCR reaction. As shown in Figure 3, withdraw of T3 leads to under-acetylated histone occupancy of DNA. In contrast, administration of T3 leads to higher acetylated histone occupancy of target sequences. These data indicate that thyroid hormone-mediated activation of luciferase reporter activity correlates with a higher acetylated histone occupancy, indicating an open, transcriptionally active chromatin structure upon T3 stimulation.

205 To finally test whether TR is able to physically bind the -4 kb TRE sequence, we performed electrophoretic mobility shift assays. As demonstrated in Figure 4, an in vitro-translated thyroid hormone receptor α 1 is able to bind the -4 kb TRE sequence (Figure 4, arrow). The

interaction could be competed with a 100-fold molar excess of unlabelled fragment (wt) or
 220 with a control TRE from the mGPDH promoter (TRE-mGPDH). In contrast, TR-DNA
 binding could not be blocked by an excess of mutated TRE variants bearing mutations either
 within the 5' or 3' nuclear receptor half sites (5' mut and 3' mut, respectively; for sequences
 see Material and Methods). These data indicate that TR is able to bind the -4 kb TRE in
 EMSA experiments.

Discussion

Gene expression analysis in vivo and cell culture indicated remarkable differences in thyroid
 230 hormone-mediated induction kinetics. Whereas ~ 50% of T3-induced genes are rapidly
 regulated within 6 hours, another ~ 50% of T3 target genes are regulated after a lag period of
 12-24 hours within 48 hours. Surprisingly, these expression patterns are very similar in vivo
 (Weitzel et al., 2001b; Weitzel et al., 2003a; Helbing et al., 2003), primary cell culture
 (Moeller et al., 2005) and permanent cell lines (Miller et al., 2001; Shih et al., 2004),
 235 suggesting common regulation mechanisms. In particular rat pituitary GC cells have been
 shown to be an ideal model system to investigate thyroid hormone-mediated gene expression
 patterns (Miller et al., 2001; Wulf et al., 2007). We hypothesized that different regulation
 mechanisms might be responsible for the different induction kinetics. In line with this
 hypothesis, we suggested that early regulated genes are direct targets of TR. Indeed thyroid
 240 hormone response elements have been described in the regulatory elements of Spot 14 and
 mGPDH (among others), both of which are regulated rapidly by T3 (Zilz et al., 1990; Weitzel
 et al., 2001a). In contrast delayed regulated target genes might be indirectly regulated via an
 intermediate factor. Indeed, we failed to identify TREs in several late induced gene promoters
 (Weitzel et al., 2003a); however, we could not completely rule out the existence of TREs in
 245 far distally located sequences which are difficult to address experimentally. Based on a
 candidate approach, we suggested PGC-1 α to be an intermediate factor of T3 action (Weitzel
 et al., 2003b). PGC-1 α has a profound influence on development, differentiation and
 metabolism. In particular, activation of PGC-1 α in muscle induces the full program of
 mitochondrial biogenesis including an increase in mitochondrial mass and oxygen
 250 consumption capacity, whereas a decrease of PGC-1 α leads to a reduced mitochondrial mass
 and oxygen consumption rates. Furthermore, PGC-1 α activation in liver induces
 gluconeogenesis whereas blunted PGC-1 α expression leads to reduced expression levels of

gluconeogenic key enzymes; reviewed in (Finck and Kelly, 2006; Handschin and Spiegelman, 2006). These and other effects resemble those effects which are regulated by thyroid hormone.

A prerequisite of an intermediate factor would be that its gene expression is rapidly regulated by T3. For PGC-1 α , this has been demonstrated in rat liver in vivo and GC cells, respectively. Endogenous transcript levels rise within 4 – 6 hours after administration of T3 (Figure 1). A second prerequisite would be that PGC-1 α is directly regulated by a TRE. As demonstrated in Figure 2, TR activates PGC-1 α gene expression via a TRE which is located at -4 kb upstream of the transcriptional start site. Integration of a single copy of this TRE sequence is sufficient to activate a luciferase reporter in transfection experiments (Figure 2; construct TRE-TK-Luc). Thyroid hormone-mediated gene expression via its nuclear receptor is accomplished by alterations in histone acetylation rates in the direction toward a higher acetylated ‘open’ chromatin (Figure 3). Finally, TR is able to bind the TRE sequence at -4 kb physically in EMSA experiments. These data indicate that the PGC-1 α gene is a direct target of TR via binding to a response element at -4 kb. Of note, this TRE is not identical to other described nuclear receptor binding sites within the PGC-1 α gene, in particular a PPRE site at -2 kb and a ERR binding site at -600 bp (Hondares et al., 2006; Laganier et al., 2004) which are not used by TR (Figure 2).

Since thyroid hormone regulates PGC-1 α gene expression via thyroid hormone receptor binding (see above) and PGC-1 α coactivates TR (Puigserver et al., 1998; Knutti et al., 2001; Zhang et al., 2004; Sadana et al., 2007) a T3-mediated activation of PGC-1 α leads to an autoregulatory feed-forward loop. Interestingly, a feed-forward mechanism has also been described for PPAR-, ERR- and MEF2-mediated activation of PGC-1 α (Hondares et al., 2006; Laganier et al., 2004; Handschin et al., 2003). However; this raises the question of how this activating signal might be switched off. One possibility would be the RNA processing capacity of PGC-1 α . PGC-1 α has been shown to regulate the splicing ratio of the TR α gene toward the TR α 2 variant which has dominant negative properties (Thijssen-Timmer et al., 2006). Thus, a decreased TR α 1 concentration relative to TR α 2 concentration would lead to dampened thyroid hormone-mediated stimulation. Another mechanism might be interference between the thyroid hormone and the cAMP signal transduction pathway. cAMP activates phosphorylation and transcriptional activity of transcription factors such as CREB which binds to CRE sites within target genes. A CRE site has been described within the PGC-

1 α gene (Herzig et al., 2001) and we and others could detect interference between the thyroid hormone and the cAMP signal pathway (Irrcher et al., 2007) (and not shown) which has also been described for PPAR and cAMP-mediated regulation of the PGC-1 α gene (Hondares et al., 2006). Furthermore, an antagonism of the thyroid hormone and the cAMP signal pathway via protein-protein interaction between CREB and TR has been described (Mendez-Pertuz et al., 2003). Finally, the presence of CRE sites in several thyroid hormone target genes (e.g. PEPCK, a key gluconeogenic gene) further supports this idea.

What might be the role of PGC-1 α in thyroid hormone-mediated gene expression? Previous data from our laboratory indicated that complex thyroid hormone-mediated gene expression patterns are not completely depending on PGC-1 α activation (Wulf et al., 2007). Furthermore, recent data by Holloszy and coworkers indicated that PGC-1 α activation appears to be rather a cause than a reason of exercise-induced mitochondrial biogenesis (Wright et al., 2007). This observation would also fit the phenotype of PGC-1 α knockout mice which show relative subtle alterations in mitochondrial biogenesis and architecture (Lin et al., 2004; Leone et al., 2005; Arany et al., 2005). However, isolated hepatocytes from PGC-1 α knockout mice clearly demonstrated reduced mitochondrial performance suggesting a role of PGC-1 α in fine tuning of mitochondrial biogenesis (Lin et al., 2004; Leone et al., 2005). Our data indicate that PGC-1 α is a direct target of thyroid hormone via its nuclear receptor. Since PGC-1 α coactivates transcription factors such as TR, NRF-1, NRF-2 and ERRs, all of which are involved in regulating mitochondrial biogenesis, a thyroid hormone-mediated activation of PGC-1 α might help this coactivator to exert its function in adaptation to nutritive, endocrine and other environmental signals.

Acknowledgments

We are indebted to James Rhee and Christoph Handschin for kind gifts of plasmid DNA, to Danielle Gourdji for kind gift of rat pituitary GC cells and to Hans J. Seitz and Josef Köhrle for continued support during the project. The work was supported by a grant from the Deutsche Forschungsgemeinschaft (WE2458/3-2) to JMW.

Figure Legends

Figure 1

Thyroid hormone induces PGC-1 α gene expression in rat liver and pituitary GC cells.

(A) Hypothyroid male rats were treated with T3 for 0 and 6 hours. Livers were prepared, RNA isolated and quantified by real-time PCR. Expression levels were normalized to the expression levels of the housekeeping gene ubiquitin. (B) Rat pituitary GC cells were treated with T3 for 0, 2, 4, 6 and 10 hours. Cells were lysed, RNA isolated and quantified by real-time PCR. Expression levels were normalized to the housekeeping gene Gs α . Data are mean $\pm$ SD of at least three independent experiments performed in duplicates and are expressed as fold activation data relative to un-stimulated expression values.

Figure 2

TR activates gene expression via a distal thyroid hormone response element. A chimeric protein of TR α 1 with the viral activator domain VP16 (TR-VP16) or the VP16 domain alone (VP16) plus the indicated promoter fragments of PGC-1 α in pGL3 luciferase reporter vectors were transfected into HepG2 cells. Promoter activities were calculated as fold activation rates of the construct by TR-VP16 relative to VP16-only cotransfection $\pm$ S.D. A TRE-like sequence at -4 kb (see text) was integrated upstream of a TK luciferase reporter, given the construct TRE-TK-Luc.

Figure 3

Acetylated histone H3 is associated with the PGC-1 α gene upon T3 stimulation.

The promoter fragment of PGC-1 α (-4483/+78-Luc) was transfected into HepG2 cells. An expression vector for TR was cotransfected and stimulated with 100 nM T3 for 24 hours. Chromatin was fixed by formaldehyde and preserved as input control or immunoprecipitated with an anti-acetylated histone H3 antibody (IP: acH3) or irrelevant antibody (IgG). Immunoprecipitated DNA was quantified by PCR and visualised on 5% polyacrylamide gels. A graphic representation is shown in the lower part of the figure.

Figure 4

TR binds to the PGC-1 α gene at the -4 kb TRE. In vitro-translated chicken TR α 1 was incubated with a labelled DNA fragment from -4029 to -4006 of the PGC-1 α gene. For competition experiments, a 100-fold molar excess of wild type fragment (wt), mutated

versions of this DNA fragment or the TRE from the mGPDH promoter (TRE-mGPDH) was added. Mutated DNA sequences contain mutations either within the 5' nuclear receptor half-site (5' mut) or the 3' half-site (3' mut) of the suggested TRE (for sequences, see Material and Methods). Specific protein-DNA complexes are indicated by an arrow.

355

Reference List

- Arany,Z., He,H.M., Lin,J.D., Hoyer,K., Handschin,C., Toka,O., Ahmad,F., Matsui,T.,
360 Chin,S., Wu,P.H., Rybkin,I.I., Shelton,J.M., Manieri,M., Cinti,S., Schoen,F.J., Bassel-Duby,R., Rosenzweig,A., Ingwall,J.S., and Spiegelman,B.M. (2005). Transcriptional coactivator PGC-1 alpha controls the energy state and contractile function of cardiac muscle. *Cell Metabolism* 1, 259-271.
- Finck,B.N. and Kelly,D.P. (2006). PGC-1 coactivators: inducible regulators of energy
365 metabolism in health and disease. *Journal of Clinical Investigation* 116, 615-622.
- Flores-Morales,A., Gullberg,H., Fernandez,L., Stahlberg,N., Lee,N.H., Vennstrom,B., and Norstedt,G. (2002). Patterns of liver gene expression governed by TRbeta. *Mol. Endocrinol.* 16, 1257-1268.
- Handschin,C., Rhee,J., Lin,J., Tarr,P.T., and Spiegelman,B.M. (2003). An autoregulatory loop
370 controls peroxisome proliferator-activated receptor gamma coactivator 1alpha expression in muscle. *Proc. Natl. Acad. Sci. U. S. A* 100, 7111-7116.
- Handschin,C. and Spiegelman,B.M. (2006). Peroxisome proliferator-activated receptor gamma coactivator 1 coactivators, energy homeostasis, and metabolism. *Endocr. Rev.* 27, 728-735.
- 375 Helbing,C.C., Werry,K., Crump,D., Domanski,D., Veldhoen,N., and Bailey,C.M. (2003). Expression profiles of novel thyroid hormone-responsive genes and proteins in the tail of *Xenopus laevis* tadpoles undergoing precocious metamorphosis. *Mol Endocrinol* 17, 1395-409.
- Herzig,S., Long,F., Jhala,U.S., Hedrick,S., Quinn,R., Bauer,A., Rudolph,D., Schutz,G.,
380 Yoon,C., Puigserver,P., Spiegelman,B., and Montminy,M. (2001). CREB regulates hepatic gluconeogenesis through the coactivator PGC-1. *Nature* 413, 179-83.
- Hondares,E., Mora,O., Yubero,P., Rodriguez de la,C.M., Iglesias,R., Giralt,M., and Villarroya,F. (2006). Thiazolidinediones and rexinoids induce peroxisome proliferator-activated receptor-coactivator (PGC)-1alpha gene transcription: an autoregulatory loop
385 controls PGC-1alpha expression in adipocytes via peroxisome proliferator-activated receptor-gamma coactivation. *Endocrinology* 147, 2829-2838.
- Irrcher,I., Adhihetty,P.J., Sheehan,T., Joseph,A.M., and Hood,D.A. (2003). PPARgamma coactivator-1alpha expression during thyroid hormone- and contractile activity-induced mitochondrial adaptations. *Am J Physiol Cell Physiol* 284, 1669-77.

- 390 Irrcher,I., Walkinshaw,D.R., Sheehan,T.E., and Hood,D.A. (2007). Thyroid hormone (T3) rapidly activates p38 and AMPK in skeletal muscle in vivo. *J. Appl. Physiol.* [Epub ahead of print]
- Knutti,D., Kressler,D., and Kralli,A. (2001). Regulation of the transcriptional coactivator PGC-1 via MAPK-sensitive interaction with a repressor. *Proc. Natl. Acad. Sci. U. S. A* 98, 9713-9718.
- 395 Laganiere,J., Tremblay,G.B., Dufour,C.R., Giroux,S., Rousseau,F., and Giguere,V. (2004). A polymorphic autoregulatory hormone response element in the human estrogen-related receptor alpha (ERRalpha) promoter dictates peroxisome proliferator-activated receptor gamma coactivator-1alpha control of ERRalpha expression. *J. Biol. Chem.* 279, 18504-18510.
- 400 Lazar,M.A. (2003). Thyroid hormone action: a binding contract. *J. Clin. Invest* 112, 497-499.
- Leone,T.C., Lehman,J.J., Finck,B.N., Schaeffer,P.J., Wende,A.R., Boudina,S., Courtois,M., Wozniak,D.F., Sambandam,N., Bernal-Mizrachi,C., Chen,Z., Holloszy,J.O., Medeiros,D.M., Schmidt,R.E., Saffitz,J.E., Abel,E.D., Semenkovich,C.F., and Kelly,D.P. (2005). PGC-1alpha deficiency causes multi-system energy metabolic derangements: muscle dysfunction, abnormal weight control and hepatic steatosis. *PLoS Biol* 3, e101.
- 405 Lin,J., Wu,P.H., Tarr,P.T., Lindenberg,K.S., St-Pierre,J., Zhang,C.Y., Mootha,V.K., Jager,S., Vianna,C.R., Reznick,R.M., Cui,L., Manieri,M., Donovan,M.X., Wu,Z., Cooper,M.P., Fan,M.C., Rohas,L.M., Zavacki,A.M., Cinti,S., Shulman,G.I., Lowell,B.B., Krainc,D., and Spiegelman,B.M. (2004). Defects in adaptive energy metabolism with CNS-linked hyperactivity in PGC-1alpha null mice. *Cell* 119, 121-35.
- 410 Mendez-Pertuz,M., Sanchez-Pacheco,A., and Aranda,A. (2003). The thyroid hormone receptor antagonizes CREB-mediated transcription. *Embo J* 22, 3102-12.
- Miller,L.D., Park,K.S., Guo,Q.M., Alkharouf,N.W., Malek,R.L., Lee,N.H., Liu,E.T., and Cheng,S.Y. (2001). Silencing of Wnt signaling and activation of multiple metabolic pathways in response to thyroid hormone-stimulated cell proliferation. *Mol. Cell Biol.* 21, 6626-6639.
- 415 Moeller,L.C., Dumitrescu,A.M., Walker,R.L., Meltzer,P.S., and Refetoff,S. (2005). Thyroid hormone responsive genes in cultured human fibroblasts. *J Clin Endocrinol Metab* 90, 936-43.
- Podvinec,M., Kaufmann,M.R., Handschin,C., and Meyer,U.A. (2002). NUBIScan, an in silico approach for prediction of nuclear receptor response elements. *Mol. Endocrinol.* 16, 1269-1279.
- 420 Puigserver,P., Wu,Z., Park,C.W., Graves,R., Wright,M., and Spiegelman,B.M. (1998). A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92, 829-39.
- 425 Rosenfeld,M.G., Lunyak,V.V., and Glass,C.K. (2006). Sensors and signals: a coactivator/corepressor/epigenetic code for integrating signal-dependent programs of transcriptional response. *Genes Dev.* 20, 1405-1428.
- Sadana,P., Zhang,Y., Song,S., Cook,G.A., Elam,M.B., and Park,E.A. (2007). Regulation of carnitine palmitoyltransferase I (CPT-1alpha) gene expression by the peroxisome proliferator activated receptor gamma coactivator (PGC-1) isoforms. *Mol. Cell Endocrinol.* 267, 6-16.
- 430

Shih,C.H., Chen,S.L., Yen,C.C., Huang,Y.H., Chen,C.D., Lee,Y.S., and Lin,K.H. (2004).
Thyroid hormone receptor-dependent transcriptional regulation of fibrinogen and coagulation
proteins. *Endocrinology* 145, 2804-14.

Thijssen-Timmer,D.C., Schiphorst,M.P., Kwakkel,J., Emter,R., Kralli,A., Wiersinga,W.M.,
and Bakker,O. (2006). PGC-1alpha regulates the isoform mRNA ratio of the alternatively
spliced thyroid hormone receptor alpha transcript. *J. Mol. Endocrinol.* 37, 251-257.

Weitzel,J.M., Hamann,S., Jauk,M., Lacey,M., Filbry,A., Radtke,C., Iwen,K.A., Kutz,S.,
Harneit,A., Lizardi,P.M., and Seitz,H.J. (2003a). Hepatic gene expression patterns in thyroid
hormone-treated hypothyroid rats. *J Mol Endocrinol* 31, 291-303.

Weitzel,J.M., Iwen,K.A., and Seitz,H.J. (2003b). Regulation of mitochondrial biogenesis by
thyroid hormone. *Exp Physiol* 88, 121-8.

Weitzel,J.M., Kutz,S., Radtke,C., Grott,S., and Seitz,H.J. (2001a). Hormonal regulation of
multiple promoters of the rat mitochondrial glycerol-3-phosphate dehydrogenase gene:
identification of a complex hormone-response element in the ubiquitous promoter B. *Eur J*
Biochem 268, 4095-103.

Weitzel,J.M., Radtke,C., and Seitz,H.J. (2001b). Two thyroid hormone-mediated gene
expression patterns in vivo identified by cDNA expression arrays in rat. *Nucleic Acids Res*
29, 5148-55.

Wright,D.C., Han,D.H., Garcia-Roves,P.M., Geiger,P.C., Jones,T.E., and Holloszy,J.O.
(2007). Exercise-induced mitochondrial biogenesis begins before the increase in muscle PGC-
1 alpha expression. *Journal of Biological Chemistry* 282, 194-199.

Wulf,A., Harneit,A., and Weitzel,J.M. (2007). T3-mediated gene expression is independent of
PGC-1alpha. *Mol. Cell Endocrinol.* 270, 57-63.

Yen,P.M. (2001). Physiological and molecular basis of thyroid hormone action. *Physiol Rev*
81, 1097-142.

Zhang,J. and Lazar,M.A. (2000). The mechanism of action of thyroid hormones. *Annu. Rev.*
Physiol 62, 439-466.

Zhang,Y., Ma,K., Song,S., Elam,M.B., Cook,G.A., and Park,E.A. (2004). Peroxisomal
proliferator-activated receptor-gamma coactivator-1 alpha (PGC-1 alpha) enhances the
thyroid hormone induction of carnitine palmitoyltransferase I (CPT-I alpha). *J. Biol. Chem.*
279, 53963-53971.

Zilz,N.D., Murray,M.B., and Towle,H.C. (1990). Identification of multiple thyroid hormone
response elements located far upstream from the rat S14 promoter. *J Biol Chem* 265, 8136-43.

Figure 1

Figure 2

Figure 3

Figure 4