Mini-Review: New insights into the evolution of chorionic gonadotrophin

Alexander Henke, Jörg Gromoll

► To cite this version:

Alexander Henke, Jörg Gromoll. Mini-Review: New insights into the evolution of chorionic gonadotrophin. *Molecular and Cellular Endocrinology*, 2008, 291 (1-2), pp.11. 10.1016/j.mce.2008.05.009 . hal-00532027

HAL Id: hal-00532027

<https://hal.science/hal-00532027>

Submitted on 4 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Mini-Review: New insights into the evolution of chorionic gonadotrophin

Authors: Alexander Henke, Jörg Gromoll

PII: S0303-7207(08)00225-6
DOI: doi:10.1016/j.mce.2008.05.009
Reference: MCE 6881

To appear in: *Molecular and Cellular Endocrinology*

Received date: 12-2-2008
Revised date: 17-5-2008
Accepted date: 19-5-2008

Please cite this article as: Henke, A., Gromoll, J., Mini-Review: New insights into the evolution of chorionic gonadotrophin, *Molecular and Cellular Endocrinology* (2007), doi:10.1016/j.mce.2008.05.009

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Revised version of manuscript MCE-D-08-00055

Mini-Review: new insights into the evolution of chorionic gonadotrophin

Alexander Henke & Jörg Gromoll

Corresponding author:

Jörg Gromoll

Joerg.gromoll@ukmuenster.de

Tel.: +49-251-8356447

Fax.: +49-251-8356093

University Clinic Münster

Institute of Reproductive Medicine

Domagkstrasse 11

D-48149 Münster, Germany

Abstract

The glycoprotein hormones luteinizing hormone (LH) and chorionic gonadotrophin (CG) are crucial for reproduction, as LH induces sex hormone production and ovulation, and CG is essential for the establishment of pregnancy and fetal male sexual differentiation. Both consist of two heterodimeric peptides of which the α -subunit is common to both hormones whereas the β -subunit is hormone-specific. The CGB gene was derived from LHB by gene duplication and frame shift mutation that led to a read-through into the formerly 3'-untranslated region, giving rise to the carboxyl-terminal peptide. Owing to nucleotide changes within the 5'-region of CGB, a new transcriptional start site and regulatory region was gained. These changes led to the specific expression of CGB in the placenta and its decrease in the pituitary. Recent findings on gonadotrophins led to an extended model for the sequence of events in the evolution of the CGB gene in primates and its tissue-specific expression.

Gonadotrophin physiology

Vertebrate reproduction is controlled by hormones of the hypothalamic-pituitary-gonadal axis. The adenohypophysis synthesizes and releases the gonadotrophins luteinizing hormone (LH) and follicle stimulating hormone (FSH), which act on the gonads and induce sex steroid synthesis and gamete production and maturation in both sexes. FSH is necessary for follicle maturation in the ovary and the stimulation of testicular Sertoli cells to support spermatogenesis. LH triggers sex steroid production in the ovary, whereas LH peak levels induce ovulation. In the testis, LH induces testosterone biosynthesis in Leydig cells.

The third gonadotrophin, chorionic gonadotrophin (CG), is the first biochemical signal from the embryo to the maternal environment. It is produced by extraembryonic trophoblasts at the beginning of a pregnancy and stimulates the corpus luteum to maintain progesterone synthesis that represses further follicle maturation, thereby ensuring embryo implantation and continuity of the pregnancy (Pierce and Parsons 1981). During pregnancy, CG is produced by the fetal part of the placenta and is also crucial for male fetal sexual differentiation, as CG stimulates the fetal testosterone synthesis in the testicular Leydig cells (Abramovich et al. 1974; Clements et al. 1976; Huhtaniemi et al. 1977; Tapanainen et al. 1981; Gromoll et al. 2000). Further biochemical and physiological details on LH and CG are compiled in Table 1.

Both LH and CG bind to the same receptor, the LH-/CG-receptor (LHR). Reports show that the expression of LHR is not restricted to the gonads, but is also expressed in the endometrium and other reproduction-associated tissues (Ziecik et al. 2007). It was therefore speculated that the special structure of the primate placenta favoured CG function in facilitating embryo implantation (Licht et al. 2007). Reports of CG's promotion of angiogenesis support the hypothesis that the embryo fosters maternal blood vessel growth via CG for better supply of nutrients and oxygen and easier release of CG and other factors (Zygmunt et al. 2002; Herr et al. 2007). Furthermore, it was reported that CG is necessary for the invasion of cytotrophoblasts into the endometrium during embryo implantation (Islami et

al. 2001; Carver et al. 2003). One study demonstrated the association between recurrent miscarriage and low serum CG values in patients; it was concluded that low serum CG hinders embryo implantation (Rull and Laan 2005).

Biochemical studies have revealed that LH, FSH and CG consist of two subunits, of which the α -subunit is identical in all hormones and the β -subunit confers hormone-specificity (Bousfield and Ward 2006). Together with thyroid stimulating hormone (TSH) they form a family of glycoprotein hormones and the resolution of their three-dimensional structure classified them as part of the superfamily of cystine knot growth factors such as PDGF, NGF, VEGF or TGF β (Lapthorn et al. 1994; Wu et al. 1994).

Prepubertal gonadotrophin expression

LH and FSH in serum are not only detectable in adults but also at very low serum concentrations in prepubertal and pubertal children (Apter et al. 1989). In 5-6 year-old girls, a pulsatile and diurnal LH pattern is present and levels rise clearly before the onset of puberty (Dunkel et al. 1990a; Dunkel et al. 1990b; Dunkel et al. 1992; Demir et al. 1995; Clark et al. 1997; Mitamura et al. 2000). As found in man, a rise of LH and testosterone levels from the prepubertal to the adult could be shown in the rhesus macaque (Meeran et al. 2003). Despite these findings, the function, if any, of childhood gonadotrophins remains to be established.

Human LH variants

There are several reports on common variant LH (V-LH), which escaped detection by conventional LH-antibodies (Pettersson et al. 1992). The investigation of the V-LHB gene identified two new SNPs in protein-coding parts that result in two amino acid changes: Trp8 => Arg8 and Ile15 => Thr15 (Furui et al. 1994; Okuda et al. 1994; Pettersson et al. 1994). Eight additional SNPs in the promoter of V-LHB lead to higher promoter activity compared

with the normal LHB promoter (Jiang et al. 1999). A study of a homozygous individual showed a higher activity of V-LH compared with normal LH but also a shorter half-life: 26 min in males and 44 min in females for V-LH versus 48 min in males and 53 min in females for WT-LT (Haavisto et al. 1995).

V-LH is present in nearly all populations worldwide, posing an important genetic LHB variant: among 30 different ethnicities, the carrier frequency had a mean $\pm$ SD of 16.5 ± 10.8 %, ranging from 0 % to 54% (Nilsson et al. 1997; Nilsson et al. 1998; Ramanujam et al. 1998; Elter et al. 1999; Ramanujam et al. 1999; Starka et al. 1999; Lamminen and Huhtaniemi 2001).

Ectopic human CG

Additional expression of CG was also found in the pituitary and at very low levels in the serum of normal individuals, in whom CG levels were slightly elevated in elderly individuals >50 years (Borkowski et al. 1984; Stenman et al. 1987; Hoermann et al. 1990; Odell et al. 1990; Alfthan et al. 1992; Birken et al. 1996; Dirnhofer et al. 1996). Furthermore, it was found that the CG levels were responsive to positive feedback via GnRH *in vivo* and *in vitro* and to negative feedback via progesterone and oestrogen *in vivo* (Stenman et al. 1987; Henke et al. 2007). Despite all these findings, so far no physiological function could be attributed to the CG expression in the pituitary and resulting serum CG in non-pregnant individuals.

A common evolutionary origin of gonadotrophin subunits

The peptide sequences of the β -subunits of the gonadotrophins revealed a homology among them and therefore a common evolutionary origin (Dayhoff 1976). Sequence comparison of the human LHB and CGB mRNAs and genes displays high homology and a duplication of CGB from LHB, confirming the concept of a common evolutionary origin (Talmadge et al. 1984). The LHB and CGB transcripts encompass 1.1 and 1.5 kilobases, respectively, and are

1 therefore relatively small transcripts. The detailed structures and lengths of the LHB/CGB
2 genes and transcripts are given in Figure 1 and Table 2, respectively.

4 **The structure of the LHB and CGB genes**

5 Gene duplication, copy diversification and the acquisition of new function are important
6 mechanisms of gene evolution (Zhang 2003). In the case of primate gonadotrophin subunit
7 genes, LHB underwent a gene duplication, in which one copy was conserved as LHB, while
8 the other copy developed into CGB by a single nucleotide deletion in codon 114; this led to a
9 frame shift mutation, thereby elongating the open reading frame (ORF) into the previously 3'
10 untranslated region (Table 2). In addition, two nucleotides inserted in codon 138 for human
11 CGB led to a further frame-shift in the ORF. These changes resulted in eight different and 24
12 additional codons for CGB in comparison with LHB (Table 2).

14 **Primate LHB/CGB genes are organised in clusters**

15 It was found that human CGB is a multicopy gene, as six CGB gene copies and one LHB
16 gene copy form a cluster (Figure 1A) (Boorstein et al. 1982; Policastro et al. 1983; Talmadge
17 et al. 1983; Fiddes and Talmadge 1984). The cluster is located on chromosome 19 at
18 19q13.32 (Naylor et al. 1983; Brook et al. 1984; Julier et al. 1984). However, the exact map
19 of the cluster itself remained elusive until it was cloned completely as a 58 kb DNA fragment
20 by Policastro *et al.*, (Policastro et al. 1986) and confirmed by others (Jameson and Lindell
21 1988). The single genes are adjacent to each other in tandem and inverted repeats within the
22 gene cluster. A recent re-sequencing of this cluster in humans of different ethnicities revealed
23 a high rate of SNPs and a low linkage disequilibrium due to excessive gene conversions so
24 that this gene cluster represents a hot spot of genomic diversity (Hallast et al. 2005). CGB1
25 and CGB2 underwent further differentiation from canonical CGB owing to a DNA insertion
26 (736 bp for CGB1, 724 bp for CGB2) and a deletion of 52 bp plus 356 bp in the region

serving as 5'UTR in the paralogous CGB3/5/7/8 genes. These rearrangements led to a novel exon 1 (58 bp) and constitutive skipping of canonical exon 1, in which the donor splice site is disrupted. The two known exons 2 and 3 are included in the ORF but are frame-shifted, thereby leading to a hypothetical ORF of 132 codons with a new stop codon and elongated 3'UTR (Figure 1 and Table 2). Despite extensive descriptions of their mRNAs, no CGB1/2 protein product has been demonstrated yet (Bo and Boime 1992; Hollenberg et al. 1994; Dirnhofer et al. 1996; Hallast et al. 2007).

A BLAST search for CGB gene copies in the chimpanzee genome (Chimpanzee Sequencing and Analysis Consortium 2005) resulted in several hits within a contig allocated to chromosome 19 (NW_001228244.1) but the hits were adjacent to gaps and did not allow an allocation of human CGB homologues to the chimpanzee counterparts except for LHB. We believe that a correct contig is not available yet.

A BLAST search in the rhesus macaque genome (Rhesus Macaque Genome Sequencing and Analysis Consortium et al. 2007) resulted in four hits in the BAC CH250-325I15, available under accession number AC202849.6. These four hits were identified as one LHB and three CGB genes, which spanned approximately 30 kb and are shown as schematic cluster in Figure 1 B.

Transcription of human CGB genes

In the human, the placental syncytiotrophoblasts are the major expression site of CGB during the first trimester of pregnancy. Gene copies CGB3/5/8 are expressed strongest whereas CGB1/2/7 genes are expressed at very low levels. CGB7 was also detected in normal breast tissue with an as yet unknown function (Talmadge et al. 1984; Bo and Boime 1992; Rull and Laan 2005). Interestingly, human CGB isoforms 3,5,8 were also detected, albeit at low expression, in the pituitary at the mRNA and protein levels (Hoermann et al. 1990; Odell et al. 1990; Birken et al. 1996; Dirnhofer et al. 1996). It follows that among the different human CGB gene copies there are differences in the strength of their promoters, dependent on the

tissue (Rull et al. 2008). It is also well known that epithelial tumours of the urogenital system, breast tumours and testis tumours can express *CGB* ectopically. Real time PCR of the mRNA isoforms in breast cancer revealed an up-regulation of *CGB* isoforms 3, 5 and 8 but not 1, 2 and 7 (Bellet et al. 1997; Giovangrandi et al. 2001). However, the ultimate reason for ectopic *CGB* expression in some tumours is unknown.

New World monkey LH/CG

A surprising plasticity of the LH/CG system was recently discovered in the common marmoset (*Callithrix jacchus*), a New World monkey. It appears that LHB became a pseudogene and its loss was compensated for by *CGB* (Muller et al. 2004; Scammell et al. 2008). The marmoset LHB and *CGB* genes have exon/intron structures homologous to the human situation, however marmoset *CGB* transcripts of the pituitary have a remarkably short 5'UTR of only 7 bp, which is comparable with other mammalian LHB 5'UTRs that are 9 bp in length (Jameson et al. 1986; Muller et al. 2004; Henke et al. 2007).

Gonadotropins in equid species

In the equine species horse (*Equus ferus caballus*), donkey (*Equus asinus*) and zebra (*Equus burchelli*), a single copy gene gives rise to an LHB/*CGB*-like gonadotrophin peptide that also harbours a CTP (Leigh and Stewart 1990; Sherman et al. 1992; Chopineau et al. 1995; Chopineau et al. 1999). The horse CTP was derived from a 10 bp deletion in exon 3, which was different from the single nucleotide deletion in the primate *CGB* gene. The same promoter region of the horse LHB/*CGB* gene serves as promoter in the pituitary and placenta. The main difference from other mammalian LHB promoters is a small insertion (50 bp) that gave rise to a second TATA-box, leaving the remaining promoter largely unchanged (Sherman et al. 1992). Owing to these characteristics it was concluded that this single copy

gene evolved independently from the primate CGB, occurs only in equine species and therefore demonstrates convergent evolution at the molecular level (Sherman et al. 1992).

This gene is expressed in both the pituitary and placenta of equine species and was confirmed in several studies at the protein level (Bousfield et al. 1987; Sugino et al. 1987).

During gestation, equine CG is synthesised and secreted by the endometrial cups that arise from the placental trophoblast chorionic girdle cells. In serum, CG appears from day 40 of gestation onwards, increases until it reaches its peak level between days 55-70, decreases until day 130 and then becomes undetectable (Moore et al. 1975; Spincemaille et al. 1975; Wooding et al. 2001). In contrast, in humans, CG is detectable earlier in the pregnancy than in horses and drops after a peak to a low, but still detectable, level, until parturition.

There are no reports on whether pregnant mare CG contributes to pregnancy recognition, since in horses the conceptus signals its presence to the maternal environment *via* prostaglandin F₂ α (PGF₂ α) in a paracrine manner. PGF₂ α normally induces luteolysis of the *corpus luteum* and results in pregnancy loss, when embryos secrete too much PGF₂ α . (Allen 2001). This mechanism contrasts not only the primate CG mechanism of pregnancy recognition but also the pregnancy recognition in ruminants via interferon tau (Roberts 2007; Roberts et al. 2008). Unfortunately, there are no studies on the mechanism of pregnancy recognition in prosimians and tarsier primate species that lack the CGB gene; otherwise it might have been possible to track the switch from prostaglandin/interferon-tau to CG-mediated pregnancy recognition.

Glycosylation patterns and the role of the carboxy-terminal peptide (CTP) in CG

The human LHB and CGB peptides are N-glycosylated at Asn13 and Asn 30. In addition, the CTP of CGB prolongs the β -subunit peptide and enables extensive glycosylation of four novel O-glycoslation sites at Ser121, Ser 127, Ser 132 and Ser 138 (Birken and Canfield 1977; Keutmann and Williams 1977). CTP glycosylation leads to a prolonged half-life of 5-6 hours

1 for CG compared with 40-60 minutes for LH and an increased molecular weight from 31 kDa
 2 for LH holohormone to 36 kDa for CG holohormone (Wehmann and Nisula 1981; Storrington
 3 al. 1988; Haavisto et al. 1995; le Cotonnec et al. 1998a; le Cotonnec et al. 1998b; le Cotonnec
 4 et al. 1998c). The glycosylated CTP also results in an altered route of hormone secretion: LH
 5 is secreted baso-laterally from gonadotrophic storage granules after GnRH-stimulation,
 6 whereas CG is released constantly and apically by trophoblasts towards the maternal blood
 7 circulation system (Table 2) (Childs et al. 1983; Handwerger et al. 1987; Lloyd and Childs
 8 1988; Currie and McNeilly 1995; Jablonka-Shariff et al. 2002). *In vitro* experiments with
 9 polarised Madin-Darby canine kidney cells demonstrated that monomeric CGB subunit is
 10 released 30% baso-laterally and 70% apically, whereas the LHB subunit is released 30%
 11 apically and 70% baso-laterally. When the CTP is removed from the CGB peptide, it is
 12 released like LHB and thereby proving that the CTP determines the route of secretion
 13 (Jablonka-Shariff et al. 2002). As in equine species a previously cryptic CTP became
 14 expressed, and Nakav *et al.* (Nakav et al. 2005) investigated whether other species also
 15 possess a cryptic CTP in the 3' region of their LHB genes. *In vitro*, bovine LHB was fused to
 16 either bovine or human CTP. In both cases efficient glycosylation failed and therefore the
 17 protein was secreted baso-laterally as found for LH. The absence of apical routing, which is
 18 necessary for efficient placental release of CG into the maternal blood stream, provides an
 19 explanation of why CGB did not emerge in ruminants (Nakav et al. 2005). Furthermore,
 20 predominantly oviparous vertebrate taxa such as fishes, amphibians, reptiles and birds, do not
 21 show *in silico* potential CTP sequences. Only mammals do so, and even marsupials (e.g.,
 22 kangaroo and opossum) show a putative CTP, although its sequence deviates significantly
 23 from CTP sequences found in the taxon Placentalia. Thus, genuine CTP-like sequences, either
 24 encrypted or expressed, correlate well with the presence of a placenta (Nakav et al. 2005).
 25 However, the glycosylation pattern of a potential CTP would not be beneficial in all species,

1 with respect to the mode of secretion, and we speculate that this is one reason why CG is
2 restricted to primate and equine species.

3 The marmoset differs from the human with regard to the glycosylation pattern of CGB, as
4 marmoset CGB possesses one N-glycosylation and two O-glycosylation sites, and therefore
5 there are differences both with human CGB and LHB (see also Table1). Thus, marmoset CGB
6 represents a peptide like hCGB but with a glycosylation pattern intermediate between hLHB
7 and hCGB (Simula et al. 1995; Amato et al. 1998). These peculiar biochemical properties
8 lead to a disruption of clear CG pulses emanating from the pituitary, as the half-life of
9 marmoset CG appears to be intermediate between that of human CG and human LH
10 (Tannenbaum et al. 2007a; Tannenbaum et al. 2007b). The half-life of marmoset CG has not
11 been investigated yet, nor has its placental glycosylation pattern, which might differ from that
12 in the pituitary, since the exact composition of glycosylation depends upon the tissue-specific
13 enzymes expressed. For instance, there is a hyperglycosylated form of human CG (hCG-H)
14 that has a molecular weight of approximately 40 kDa. This form is produced in the first
15 trimester of pregnancy (O'Connor et al. 1998). Its biological role is the cytotrophoblast
16 invasion into the endometrium. Ectopic expression of hCG-H occurs in pregnancies with
17 Down-syndrome and in some malignant tumours. The more malignant the cells were, the
18 more CG was hyperglycosylated (Valmu et al. 2006; Cole and Khanlian 2007). Although the
19 reason for ectopic CGB expression in tumours is still unclear, it could be demonstrated that
20 CG can drive prostate carcinoma malignancy (Wu and Walker 2006).

22 **An extended model of primate CGB evolution**

23 The recent findings on gonadotrophins led us to propose a revised and extended version of
24 primate CGB gene evolution, covering the sequence of events such as duplication, mutation
25 and acquisition of new functions (Fiddes and Talmadge 1984).

1 **1. Duplication of the LHB gene**

2 Approximately 55-50 million years ago (mya) the split of the tarsiers from the lineage of
 3 Simiiformes occurred. The anthropoid line encompasses New World monkeys (NWM), Old
 4 World monkeys (OWM), small apes, great apes and humans. The latter three groups form the
 5 taxon Hominoidea. As there is no CGB or second LHB gene in prosimians and tarsiers, the
 6 LHB gene duplication must have occurred before the NWM diverged from the OWM,
 7 approximately 55-35 mya (see also Fig. 2) (Bailey et al. 1991; Schrago and Russo 2003;
 8 Opazo et al. 2006; Schrago 2007). Because LHB and CGB have not only highly homologous
 9 coding sequences, but also highly homologous promoters, we assume that, after the LHB gene
 10 duplicated, both LHB copies were functionally expressed in the pituitary.

12 **2. The appearance of CGB**

13 In one LHB gene copy a single nucleotide was deleted in the 114th codon and led to a frame
 14 shift mutation. This slipped the last seven codons out of frame compared with LHB, and led
 15 to a read-through into the previous 3'UTR and an incorporation of this sequence into the
 16 coding region. An insertion of the two nucleotides C and G in the 138th codon further
 17 elongated the ORF, expanding the CGB subunit for 24 additional codons that encode the
 18 CTP. These frame shifts resulted in a new stop codon, TAA, in the previous 3'UTR. It is part
 19 of the polyadenylation signal and yields a very short 3'UTR of only 16 nucleotides (Talmadge
 20 et al. 1984). However, it is not possible to elucidate which event occurred first; the single
 21 nucleotide deletion or the insertion of the two bases.

22 This concept of events for the mechanism of CGB gene evolution by LHB gene duplication
 23 and nucleotide modifications that led to the CTP, has already been proposed (Dayhoff 1976;
 24 Fiddes and Goodman 1980; Fiddes and Talmadge 1984; Talmadge et al. 1984).

1 Since CGB derived from LHB, including the promoter, and because CGB is still expressed at
 2 very low levels in human pituitaries, we speculate that CGB was still expressed in the
 3 pituitary in ancient primates at this stage of evolution.

4 **3. CG gains expression in the placenta**

5 Changes in the promoter region of CGB led to a shift in expression from the pituitary to the
 6 placenta (Hollenberg et al. 1994). The hCGB transcriptional start site relocated to 365 bp
 7 upstream of ATG, so that the 5' UTR of the mRNA comprises the region which serves as core
 8 promoter and transcriptional start site for LHB transcription. Human CGB transcription is
 9 mainly regulated by a TATA-less promoter, which is regulated by the transcription factors
 10 (TF) cAMP element binding protein (CREB), v-ETS erythroblastosis virus E26 oncogene
 11 homolog 2 (ETS2), activating protein 2 (AP2) and selective promoter factor 1 (SP1).
 12 Although each of these factors is expressed in several different tissues, they form together a
 13 trophoblast-specific combinatorial code for CGB gene transactivation (Jameson and
 14 Hollenberg 1993; Hollenberg et al. 1994; Pestell et al. 1994; Johnson and Jameson 1999;
 15 Johnson and Jameson 2000; Ghosh et al. 2003). At the blastula stage, CG is synthesised by
 16 the trophectoderm but not by the embryonic stem cells in the inner cell mass. The
 17 pluripotency marker Oct3/4 is expressed by the embryonic stem cells and acts as a
 18 transcription factor. Oct3/4 is able to bind to the promoters of the glycoprotein hormone alpha
 19 gene (GPHA) and CGB and thereby repress their transcription (Liu and Roberts 1996; Liu et
 20 al. 1997). Thus, in the placenta, CGB expression is controlled tissue-specifically by a
 21 completely different regulatory region and set of TFs compared with pituitary-expressed
 22 LHB.

23 The glycoprotein α -subunit, which is necessary for full functional and biological activity of
 24 the gonadotrophins, also gained expression in the placenta, otherwise no functional CG
 25 holohormone could have formed. However, it remains to be clarified when exactly this
 26 happened and which of the two subunit genes was expressed first in the placenta.

4. Differences of CG expression in pituitaries depend on taxonomic group

The LHB core promoter for mouse, rat, cattle and man comprises two transcription factor binding sites for steroidogenic factor 1 (SF1), two for early response growth protein 1 (EGR1), one for pituitary homeobox factor 1 (PITX1) and a TATA box. SF1 is involved in steroid synthesis in a wide spectrum of tissues (e.g. gonads and adrenals) and EGR1 is expressed mainly during embryogenesis. PITX1 is a pituitary-specific transcription factor and all three factors together form a gonadotrophe-specific, tripartite regulatory complex for LHB activation (Kaiser et al. 2000; Jorgensen et al. 2004).

It was demonstrated that in human CGB only three nucleotide changes in transcription factor binding sites (TFBS) were sufficient to cause CGB down-regulation in the pituitary and reduced GnRH-responsiveness in comparison with LHB (Henke et al. 2007). In one TFBS for EGR1, a C-T transition and a G-C transversion occurred at positions -119 and -112, respectively, in relation to the start codon ATG. An A-G transition at -106 in the TFBS for PITX1 decreased its protein binding ability, and all three nucleotide changes together demonstrated synergistic effects in pituitary CGB down-regulation (Henke et al. 2007).

5. CGB gene duplications in OWM and Hominoidea and a new LH/CG system in NWM

In the lineages of OWM and Hominoidea, the CGB gene itself underwent several duplications, leading to different copy numbers varying from three to six, as indicated in Table 3 and Fig. 2. This was reported first for humans and later investigated systematically for the order of primates (Boorstein et al. 1982; Talmadge et al. 1983; Maston and Ruvolo 2002). A recent study demonstrated that CGB1 is the most recent gene copy, as it evolved in the last common ancestor of African great apes (gorilla, chimpanzee and human), after the split from gibbons and orang utan, approximately 15 to 8 mya (Hallast et al. 2007).

1 Originally both genes, LHB and CGB, must have been present in NWM. However, for yet
2 unknown reasons, LHB became non-functional in the common marmoset and its loss was
3 compensated for by CGB, which is a single copy gene, expressed in both pituitary and
4 placenta (Muller et al. 2004).

5 This is valid for the complete branch of NWM because of the conserved CGB promoter
6 architecture across all four NWM families (unpublished data from our laboratory), and the
7 lack of LHB and presence of CGB in pituitaries in other NWM species (Scammell et al.
8 2008). Further striking evidence comes from the similar exon architecture of the LHR type 2
9 (Gromoll et al. 2003; Gromoll et al. 2007). The LHR types 1 and 2 are G protein-coupled,
10 seven transmembrane receptors. In OWM and Hominoidea, the LHR type 1 is able to bind
11 and transduce signals from LH and CG equally well. However, in NWM, the LHR type 2
12 lacks exon 10 of 11 exons at the mRNA and protein levels, although it is present at the
13 genomic level (Gromoll et al. 2003; Gromoll et al. 2007). Exon 10 encodes the extra-cellular
14 hinge region of the receptor and its absence in LHR type 2 impairs LH signal transduction but
15 not CG signalling (Muller et al. 2003). Thus, the LHR type 2 of the NWM is able to
16 discriminate between LH and CG and transduces CG signalling only. Therefore Fig. 2
17 illustrates not only the CGB evolution in primates but also includes the LHR type 2 co-
18 evolution and important dates of primate speciation.

20 **Summary and Conclusion**

21 In this review we have discussed the evolutionary events that led to the rise of CGB in
22 primates and were able to approximate the time lines during which each event took place. In
23 brief, we propose the following sequence of events:

24 LHB gene duplication, emergence of the CTP, co-expression of CGB & GPH α in the placenta
25 and gain-of-function of CG in the placenta occurred approximately 55-35 mya, a relatively
26 short time in evolutionary terms.

Revised version of manuscript MCE-D-08-00055

The events specific for NWM, such as emergence of LHR type 2 and LHB's becoming a pseudogene, most likely happened 35-20 mya. In the lineage leading to OWM, small apes, great apes and the human, CGB was duplicated in the period 35-25 mya. A further duplication, which is specific for gorilla, chimpanzee and humans, occurred 15-8 mya.

The CGB gain-of-function, its gene copy number variance in OWM and Hominoidea, and the strong pituitary CG expression in the NWMs demonstrate a remarkable plasticity of the LH/CG hormone system not previously anticipated. Thus, the CGB gene cluster represents a hallmark of primate evolution and is an excellent example of how a few changes in duplicated DNA sequences can lead to a novel gene, characterised by a changed cellular expression, peptide extension as well as a novel physiological role. Therefore this gene cluster gives insight into currently ongoing evolution in primates, including the human.

Acknowledgment

The authors thank Dr. J. Wistuba and Dr. T.G. Cooper for useful comments on the manuscript and language editing. Financial support was received from the German Research council (DFG), grant Gr 1541-8-1 and the Innovative Medical Research Fund of the University of Münster, grant SI 520601.

1 **Table 1: Main differences between human LH and CG.**

Characteristic	human LH	human CG
Major site of expression	pituitary	placenta
Time of physiological relevant expression	starts at adolescence	starts at pregnancy
Major function	ovulation & synthesis of sex steroids and INSL3	establishment of pregnancy in ♀ Fetal differentiation in ♂
Initial half-life in blood	40-60 min	5-6 h
Peptide length and of glycosylation pattern of mature β -subunit	121 amino acids 2 x N-glycan no O-glycan	145 amino acids 2x N-glycan 4x O-glycan
Secretion pattern	storage in vesicles; pulsatile release in baso-lateral direction	constitutive secretion in apical direction

2 The data are based on Pierce and Parsons 1981; Wehmann and Nisula 1981; Storrington et al.
3 1988; le Cotonnec et al. 1998a, b, c ; Jablonka-Shariff et al. 2002; Bousfield and Ward 2006.
4 INSL3 = insulin-like growth factor 3.

5

Table 2: Comparison of human gonadotrophin β -subunit transcripts.

	LHB [bp]	CGB3/5/7/8 [bp]	CGB1/2 [bp]
5'UTR	9	365	174
Exon 1*	15	15	58
Intron 1	352	352 (CGB 3/5/7) 353 (CGB 8)	399
Exon 2	168	168	168
Intron 2	235	235	235
Exon 3*	240	312	170
3'UTR	91	20	162
Total	1110	1467	1366

* length of UTR sequence is given separately and therefore excluded from given exon size

1 **Table 3: CGB copy numbers in different primate species.**

Primate group	Species	CGB copy numbers
Hominoidea	Human (<i>Homo sapiens</i>)	6
	Chimpanzee (<i>Pan troglodytes</i>)	5
	Orang utan (<i>Pongo pygmaeus</i>)	4
Cercopithecoidea (Old World monkeys)	Rhesus macaque (<i>Macaca mulatta</i>)	3
	Guereza monkey (<i>Colobus guereza</i>)	4
	Dusky leaf monkey (<i>Presbytis obscura</i>)	5
Platyrrhini (New World monkeys)	Owl monkey (<i>Aotus trivirgatus</i>)	1
	Dusk titi monkey (<i>Callicebus moloch</i>)	1
Prosimii	Ring-tailed lemur (<i>Lemur catta</i>)	0

2

3 Data are based on (Maston and Ruvolo 2002; Hallast et al. 2007).

1 **Figure 1: The structure of LH/CGB gene clusters and their genes.**

2 A schematic overview of the (A) organisation of the human LHB/CGB gene cluster on
3 chromosome 19.3 with indications of the orientation of the genes and the intergenic
4 distances, (B) the organisation of the rhesus macaque's LHB/CGB gene cluster and
5 (C) details of the human genes themselves. The numbering of rhesus macaque CGB
6 genes is in numerical order and does not imply special homology to the accordingly
7 numbered human CGB genes. Black boxes indicate UTR, white boxes the exons
8 (Ex) and solid vertical lines introns (Int) and promoter regions. The original CGB
9 structure for CGB1/2 before deletions and insertions is indicated by grey shading.

Figure 2: Schematic evolution of CGB within the primates.

This is a schematic overview of CGB evolution in primates. Prosimians and tarsiers possess only LHB but not CGB. Solid arrows indicate when an event happened, based on published time points. Dashed arrows indicate events for which the point of time is unknown. The period in which the event most likely happened is explained in the text. When more than one CGB copy number is present in a lineage, it is indicated by (n+2) or (n+3). Mya, million years ago; NWM, New World monkeys; OWM, Old World monkeys. Branching time points are according to Bailey et al. 1991; Kumar and Hedges 1998; Schrago and Russo 2003; Poux et al. 2006; Rhesus Macaque Genome Sequencing and Analysis Consortium et al. 2007 and Schrago 2007.

References

- Abramovich D. R., T. G. Baker, and P. Neal. 1974. Effect of human chorionic gonadotrophin on testosterone secretion by the foetal human testis in organ culture. *J. Endocrinol.* 60:179-185.
- Alfthan H., C. Haglund, J. Dabek, and U. Stenman. 1992. Concentrations of human choriogonadotropin, its beta-subunit, and the core fragment of the beta-subunit in serum and urine of men and nonpregnant women. *Clin Chem* 38:1981-1987.
- Allen W. 2001. Fetomaternal interactions and influences during equine pregnancy. *Reproduction* 121:513-527.
- Amato F., A. P. Simula, L. J. Gameau, and R. J. Norman. 1998. Expression, characterisation and immunoassay of recombinant marmoset chorionic gonadotrophin dimer and beta-subunit. *J. Endocrinol.* 159:141-151.
- Apter D., B. Cacciatore, H. Alfthan, and U. H. Stenman. 1989. Serum luteinizing hormone concentrations increase 100-fold in females from 7 years to adulthood, as measured by time-resolved immunofluorometric assay. *J. Clin. Endocrinol. Metab.* 68:53-57.
- Bailey W. J., D. H. Fitch, D. A. Tagle, J. Czelusniak, J. L. Slightom, and M. Goodman. 1991. Molecular evolution of the psi eta-globin gene locus: gibbon phylogeny and the hominoid slowdown. *Mol. Biol. Evol.* 8:155-184.
- Bellet D., V. Lazar, I. Bieche, V. Paradis, Y. Giovangrandi, P. Paterlini, R. Lidereau, P. Bedossa, J. M. Bidart, and M. Vidaud. 1997. Malignant transformation of nontrophoblastic cells is associated with the expression of chorionic gonadotropin beta genes normally transcribed in trophoblastic cells. *Cancer Res.* 57:516-523.
- Birken S., and R. E. Canfield. 1977. Isolation and amino acid sequence of COOH-terminal fragments from the beta subunit of human choriogonadotropin. *J. Biol. Chem.* 252:5386-5392.
- Birken S., Y. Maydelman, M. A. Gawinowicz, A. Pound, Y. Liu, and A. S. Hartree. 1996. Isolation and characterization of human pituitary chorionic gonadotropin. *Endocrinology* 137:1402-1411.
- Bo M., and I. Boime. 1992. Identification of the transcriptionally active genes of the chorionic gonadotropin beta gene cluster in vivo. *J. Biol. Chem.* 267:3179-3184.
- Boorstein W. R., N. C. Vamvakopoulos, and J. C. Fiddes. 1982. Human chorionic gonadotropin beta-subunit is encoded by at least eight genes arranged in tandem and inverted pairs. *Nature* 300:419-422.
- Borkowski A., V. Puttaert, M. Gyling, C. Muquardt, and J. J. Body. 1984. Human chorionic gonadotropin-like substance in plasma of normal nonpregnant subjects and women with breast cancer. *J. Clin. Endocrinol. Metab.* 58:1171-1178.
- Bousfield G. R., W. K. Liu, H. Sugino, and D. N. Ward. 1987. Structural studies on equine glycoprotein hormones. Amino acid sequence of equine lutropin beta-subunit. *J. Biol. Chem.* 262:8610-8620.

- 1 Bousfield G. R., and D. N. Ward. 2006. Gonadotropins: Chemistry and Biosynthesis. Pp.
2 1581-1634 in J. D. Neill, ed. Knobil and Neill's Physiology of Reproduction. Elsevier
3 Academic Press, St Louis, MO, USA.
- 4 Brook J. D., D. J. Shaw, L. Meredith, G. A. Bruns, and P. S. Harper. 1984. Localisation of
5 genetic markers and orientation of the linkage group on chromosome 19. Hum. Genet.
6 68:282-285.
- 7 Carver J., K. Martin, I. Spyropoulou, D. Barlow, I. Sargent, and H. Mardon. 2003. An in-vitro
8 model for stromal invasion during implantation of the human blastocyst. Hum. Reprod.
9 18:283-290.
- 10 Childs G. V., C. Hyde, Z. Naor, and K. Catt. 1983. Heterogeneous luteinizing hormone and
11 follicle-stimulating hormone storage patterns in subtypes of gonadotropes separated by
12 centrifugal elutriation. Endocrinology 113:2120-2128.
- 13 Chimpanzee Sequencing and Analysis Consortium. 2005. Initial sequence of the chimpanzee
14 genome and comparison with the human genome. Nature 437:69-87.
- 15 Chopineau M., F. Stewart, and W. R. Allen. 1995. Cloning and analysis of the cDNA
16 encoding the horse and donkey luteinizing hormone beta-subunits. Gene 160:253-256.
- 17 Chopineau M., N. Martinat, C. Pourchet, F. Stewart, Y. Combarnous, and F. Guillou. 1999.
18 Cloning, sequencing and functional expression of zebra (*Equus burchelli*) LH. J. Reprod.
19 Fertil. 115:159-166.
- 20 Clark P. A., A. Iranmanesh, J. D. Veldhuis, and A. D. Rogol. 1997. Comparison of Pulsatile
21 Luteinizing Hormone Secretion Between Prepubertal Children and Young Adults:
22 Evidence for a Mass/Amplitude-Dependent Difference Without Gender or Day/Night
23 Contrasts. J Clin Endocrinol Metab 82:2950-2955.
- 24 Clements J. A., F. I. Reyes, J. S. Winter, and C. Faïman. 1976. Studies on human sexual
25 development. III. Fetal pituitary and serum, and amniotic fluid concentrations of LH, CG,
26 and FSH. J. Clin. Endocrinol. Metab. 42:9-19.
- 27 Cole L. A., and S. A. Khanlian. 2007. Hyperglycosylated hCG: A variant with separate
28 biological functions to regular hCG. Molecular and Cellular Endocrinology, 260-
29 262:228-236.
- 30 Currie R. J., and A. S. McNeilly. 1995. Mobilization of LH secretory granules in
31 gonadotrophs in relation to gene expression, synthesis and secretion of LH during the
32 preovulatory phase of the sheep oestrous cycle. J. Endocrinol. 147:259-270.
- 33 Dayhoff M. O. 1976. Atlas of Protein Sequence and Structure Vol. 5 Supplement 2.
34 Washington, D.C., USA.
- 35 Demir A., L. Dunkel, U. Stenman, and R. Voutilainen. 1995. Age-related course of urinary
36 gonadotropins in children. J Clin Endocrinol Metab 80:1457-1460.
- 37 Dirnhofer S., M. Hermann, A. Hittmair, R. Hoermann, K. Kapelari, and P. Berger. 1996.
38 Expression of the human chorionic gonadotropin-beta gene cluster in human pituitaries
39 and alternate use of exon 1. J. Clin. Endocrinol. Metab. 81:4212-4217.

- 1 Dunkel L., H. Alfthan, U. H. Stenman, and J. Perheentupa. 1990a. Gonadal control of
2 pulsatile secretion of luteinizing hormone and follicle-stimulating hormone in prepubertal
3 boys evaluated by ultrasensitive time-resolved immunofluorometric assays. *J. Clin.*
4 *Endocrinol. Metab.* 70:107-114.
- 5 Dunkel L., H. Alfthan, U. H. Stenman, P. Tapanainen, and J. Perheentupa. 1990b. Pulsatile
6 secretion of LH and FSH in prepubertal and early pubertal boys revealed by ultrasensitive
7 time-resolved immunofluorometric assays. *Pediatr. Res.* 27:215-219.
- 8 Dunkel L., H. Alfthan, U. Stenman, G. Selstam, S. Rosberg, and K. Albertsson-Wikland.
9 1992. Developmental changes in 24-hour profiles of luteinizing hormone and follicle-
10 stimulating hormone from prepuberty to midstages of puberty in boys. *J Clin Endocrinol*
11 *Metab* 74:890-897.
- 12 Elter K., C. T. Erel, N. Cine, U. Ozbek, B. Hacıhanefioglu, and E. Ertunçalp. 1999. Role of
13 the mutations Trp8 => Arg and Ile15 => Thr of the human luteinizing hormone beta-
14 subunit in women with polycystic ovary syndrome. *Fertil. Steril.* 71:425-430.
- 15 Fiddes J. C., and K. Talmadge. 1984. Structure, expression, and evolution of the genes for the
16 human glycoprotein hormones. *Recent Prog. Horm. Res.* 40:43-78.
- 17 Fiddes J. C., and H. M. Goodman. 1980. The cDNA for the beta-subunit of human chorionic
18 gonadotropin suggests evolution of a gene by readthrough into the 3'-untranslated region.
19 *Nature* 286:684-687.
- 20 Furui K., N. Suganuma, S. Tsukahara, Y. Asada, F. Kikkawa, M. Tanaka, T. Ozawa, and Y.
21 Tomoda. 1994. Identification of two point mutations in the gene coding luteinizing
22 hormone (LH) beta-subunit, associated with immunologically anomalous LH variants. *J.*
23 *Clin. Endocrinol. Metab.* 78:107-113.
- 24 Ghosh D., T. Ezashi, M. C. Ostrowski, and R. M. Roberts. 2003. A central role for Ets-2 in
25 the transcriptional regulation and cyclic adenosine 5'-monophosphate responsiveness of
26 the human chorionic gonadotropin-beta subunit gene. *Mol. Endocrinol.* 17:11-26.
- 27 Giovannardi Y., B. Parfait, M. Asheuer, M. Olivi, R. Lidereau, M. Vidaud, and I. Bieche.
28 2001. Analysis of the human CGB/LHB gene cluster in breast tumors by real-time
29 quantitative RT-PCR assays. *Cancer Lett.* 168:93-100.
- 30 Gromoll J., U. Eiholzer, E. Nieschlag, and M. Simoni. 2000. Male hypogonadism caused by
31 homozygous deletion of exon 10 of the luteinizing hormone (LH) receptor: differential
32 action of human chorionic gonadotropin and LH. *J. Clin. Endocrinol. Metab.* 85:2281-
33 2286.
- 34 Gromoll J., J. Wistuba, N. Terwort, M. Godmann, T. Muller, and M. Simoni. 2003. A new
35 subclass of the luteinizing hormone/chorionic gonadotropin receptor lacking exon 10
36 messenger RNA in the New World monkey (*Platyrrhini*) lineage. *Biol. Reprod.* 69:75-80.
- 37 Gromoll J., L. Lahrmann, M. Godmann, T. Muller, C. Michel, S. Stamm, and M. Simoni.
38 2007. Genomic Checkpoints for Exon 10 Usage in the Luteinizing Hormone Receptor
39 Type 1 and Type 2. *Mol. Endocrinol.* 21:1984-1996.

- 1 Haavisto A. M., K. Pettersson, M. Bergendahl, A. Virkamaki, and I. Huhtaniemi. 1995.
2 Occurrence and biological properties of a common genetic variant of luteinizing
3 hormone. *J. Clin. Endocrinol. Metab.* 80:1257-1263.
- 4 Hallast P., K. Rull, and M. Laan. 2007. The evolution and genomic landscape of CGB1 and
5 CGB2 genes. *Mol. Cell. Endocrinol.* 260-262:2-11.
- 6 Hallast P., L. Nagirnaja, T. Margus, and M. Laan. 2005. Segmental duplications and gene
7 conversion: Human luteinizing hormone/chorionic gonadotropin beta gene cluster.
8 *Genome Res.* 15:1535-1546.
- 9 Handwerger S., S. P. Wilson, L. Tyrey, and P. M. Conn. 1987. Biochemical evidence that
10 human placental lactogen and human chorionic gonadotropin are not stored in
11 cytoplasmic secretion granules. *Biol. Reprod.* 37:28-32.
- 12 Henke A., C. M. Luetjens, M. Simoni, and J. Gromoll. 2007. Chorionic gonadotropin beta
13 subunit gene expression in the marmoset pituitary is controlled by steroidogenic factor 1
14 (SF1), early growth response protein 1 (Egr1) and pituitary homeobox factor 1 (Pitx1).
15 *Endocrinology* .
- 16 Herr F., N. Baal, K. Reisinger, A. Lorenz, T. McKinnon, K. T. Preissner, and M. Zygmunt.
17 2007. HCG in the regulation of placental angiogenesis. Results of an in vitro study.
18 *Placenta* 28 Suppl A:S85-93.
- 19 Hoermann R., G. Spoettl, R. Moncayo, and K. Mann. 1990. Evidence for the presence of
20 human chorionic gonadotropin (hCG) and free beta-subunit of hCG in the human
21 pituitary. *J. Clin. Endocrinol. Metab.* 71:179-186.
- 22 Hollenberg A. N., R. G. Pestell, C. Albanese, M. E. Boers, and J. L. Jameson. 1994. Multiple
23 promoter elements in the human chorionic gonadotropin beta subunit genes distinguish
24 their expression from the luteinizing hormone beta gene. *Mol. Cell. Endocrinol.* 106:111-
25 119.
- 26 Huhtaniemi I. T., C. C. Korenbrot, and R. B. Jaffe. 1977. HCG binding and stimulation of
27 testosterone biosynthesis in the human fetal testis. *J. Clin. Endocrinol. Metab.* 44:963-
28 967.
- 29 Islami D., P. Mock, and P. Bischof. 2001. Effects of human chorionic gonadotropin on
30 trophoblast invasion. *Semin. Reprod. Med.* 19:49-53.
- 31 Jablonka-Shariff A., V. Garcia-Campayo, and I. Boime. 2002. Evolution of lutropin to
32 chorionic gonadotropin generates a specific routing signal for apical release in vivo. *J.*
33 *Biol. Chem.* 277:879-882.
- 34 Jameson J. L., C. M. Lindell, and J. F. Habener. 1986. Evolution of different transcriptional
35 start sites in the human luteinizing hormone and chorionic gonadotropin beta-subunit
36 genes. *DNA* 5:227-234.
- 37 Jameson J. L., and A. N. Hollenberg. 1993. Regulation of chorionic gonadotropin gene
38 expression. *Endocr. Rev.* 14:203-221.

- 1 Jameson J. L., and C. M. Lindell. 1988. Isolation and characterization of the human chorionic
2 gonadotropin beta subunit (CG beta) gene cluster: regulation of transcriptionally active
3 CG beta gene by cyclic AMP. *Mol. Cell. Biol.* 8:5100-5107.
- 4 Jiang M., P. Pakarinen, F. P. Zhang, T. El-Hefnawy, P. Koskimies, K. Pettersson, and I.
5 Huhtaniemi. 1999. A common polymorphic allele of the human luteinizing hormone
6 beta-subunit gene: additional mutations and differential function of the promoter
7 sequence. *Hum. Mol. Genet.* 8:2037-2046.
- 8 Johnson W., and J. L. Jameson. 2000. Role of Ets2 in cyclic AMP regulation of the human
9 chorionic gonadotropin beta promoter. *Mol. Cell. Endocrinol.* 165:17-24.
- 10 Johnson W., and J. L. Jameson. 1999. AP-2 (activating protein 2) and Sp1 (selective promoter
11 factor 1) regulatory elements play distinct roles in the control of basal activity and cyclic
12 adenosine 3',5'-monophosphate responsiveness of the human chorionic gonadotropin-beta
13 promoter. *Mol. Endocrinol.* 13:1963-1975.
- 14 Jorgensen J. S., C. C. Quirk, and J. H. Nilson. 2004. Multiple and overlapping combinatorial
15 codes orchestrate hormonal responsiveness and dictate cell-specific expression of the
16 genes encoding luteinizing hormone. *Endocr. Rev.* 25:521-542.
- 17 Julier C., D. Weil, P. Couillin, J. C. Cote, V. C. Nguyen, C. Foubert, A. Boue, J. P. Thirion, J.
18 C. Kaplan, and C. Junien. 1984. The beta chorionic gonadotropin-beta luteinizing gene
19 cluster maps to human chromosome 19. *Hum. Genet.* 67:174-177.
- 20 Kaiser U. B., L. M. Halvorson, and M. T. Chen. 2000. Sp1, steroidogenic factor 1 (SF-1), and
21 early growth response protein 1 (egr-1) binding sites form a tripartite gonadotropin-
22 releasing hormone response element in the rat luteinizing hormone-beta gene promoter:
23 an integral role for SF-1. *Mol. Endocrinol.* 14:1235-1245.
- 24 Keutmann H. T., and R. M. Williams. 1977. Human chorionic gonadotropin. Amino acid
25 sequence of the hormone-specific COOH-terminal region. *J. Biol. Chem.* 252:5393-5397.
- 26 Kumar S., and S. B. Hedges. 1998. A molecular timescale for vertebrate evolution. *Nature*
27 392:917-920.
- 28 Lamminen T., and I. Huhtaniemi. 2001. A common genetic variant of luteinizing hormone;
29 relation to normal and aberrant pituitary-gonadal function. *European Journal of*
30 *Pharmacology*, 414:1-7.
- 31 Laphorn A. J., D. C. Harris, A. Littlejohn, J. W. Lustbader, R. E. Canfield, K. J. Machin, F. J.
32 Morgan, and N. W. Isaacs. 1994. Crystal structure of human chorionic gonadotropin.
33 *Nature* 369:455-461.
- 34 le Cotonnec J. Y., H. C. Porchet, V. Beltrami, and A. Munafo. 1998a. Clinical pharmacology
35 of recombinant human luteinizing hormone: Part II. Bioavailability of recombinant
36 human luteinizing hormone assessed with an immunoassay and an in vitro bioassay.
37 *Fertil. Steril.* 69:195-200.
- 38 le Cotonnec J. Y., H. C. Porchet, V. Beltrami, and A. Munafo. 1998b. Clinical pharmacology
39 of recombinant human luteinizing hormone: Part I. Pharmacokinetics after intravenous
40 administration to healthy female volunteers and comparison with urinary human
41 luteinizing hormone. *Fertil. Steril.* 69:189-194.

- 1 le Cotonnec J. Y., E. Loumaye, H. C. Porchet, V. Beltrami, and A. Munafo. 1998c.
2 Pharmacokinetic and pharmacodynamic interactions between recombinant human
3 luteinizing hormone and recombinant human follicle-stimulating hormone. *Fertil. Steril.*
4 69:201-209.
- 5 Leigh S. E., and F. Stewart. 1990. Partial cDNA sequence for the donkey chorionic
6 gonadotrophin-beta subunit suggests evolution from an ancestral LH-beta gene. *J. Mol.*
7 *Endocrinol.* 4:143-150.
- 8 Licht P., H. Fluhr, J. Neuwinger, D. Wallwiener, and L. Wildt. 2007. Is human chorionic
9 gonadotropin directly involved in the regulation of human implantation? *Mol. Cell.*
10 *Endocrinol.* 269:85-92.
- 11 Liu L., and R. M. Roberts. 1996. Silencing of the gene for the beta subunit of human
12 chorionic gonadotropin by the embryonic transcription factor Oct-3/4. *J. Biol. Chem.*
13 271:16683-16689.
- 14 Liu L., D. Leaman, M. Villalta, and R. M. Roberts. 1997. Silencing of the gene for the alpha-
15 subunit of human chorionic gonadotropin by the embryonic transcription factor Oct-3/4.
16 *Mol. Endocrinol.* 11:1651-1658.
- 17 Lloyd J. M., and G. V. Childs. 1988. Differential storage and release of luteinizing hormone
18 and follicle-releasing hormone from individual gonadotropes separated by centrifugal
19 elutriation. *Endocrinology* 122:1282-1290.
- 20 Maston G. A., and M. Ruvolo. 2002. Chorionic gonadotropin has a recent origin within
21 primates and an evolutionary history of selection. *Mol. Biol. Evol.* 19:320-335.
- 22 Meeran D., H. F. Urbanski, S. J. Gregory, J. Townsend, and D. J. Tortonese. 2003.
23 Developmental Changes in the Hormonal Identity of Gonadotroph Cells in the Rhesus
24 Monkey Pituitary Gland. *J Clin Endocrinol Metab* 88:2934-2942.
- 25 Mitamura R., K. Yano, N. Suzuki, Y. Ito, Y. Makita, and A. Okuno. 2000. Diurnal Rhythms
26 of Luteinizing Hormone, Follicle-Stimulating Hormone, Testosterone, and Estradiol
27 Secretion before the Onset of Female Puberty in Short Children. *J Clin Endocrinol Metab*
28 85:1074-1080.
- 29 Moore R. M., W. R. Allen, and D. W. Hamilton. 1975. Origin and histogenesis of equine
30 endometrial cups. *J Reprod Fert* 23:Supplement 391-396.
- 31 Muller T., J. Gromoll, and M. Simoni. 2003. Absence of exon 10 of the human luteinizing
32 hormone (LH) receptor impairs LH, but not human chorionic gonadotropin action. *J.*
33 *Clin. Endocrinol. Metab.* 88:2242-2249.
- 34 Muller T., M. Simoni, E. Pekel, C. M. Luetjens, R. Chandolia, F. Amato, R. J. Norman, and J.
35 Gromoll. 2004. Chorionic gonadotrophin beta subunit mRNA but not luteinising
36 hormone beta subunit mRNA is expressed in the pituitary of the common marmoset
37 (*Callithrix jacchus*). *J. Mol. Endocrinol.* 32:115-128.
- 38 Nakav S., A. Jablonka-Shariff, S. Kaner, P. Chadna-Mohanty, H. E. Grotjan, and D. Ben-
39 Menahem. 2005. The LHbeta gene of several mammals embeds a carboxyl-terminal
40 peptide-like sequence revealing a critical role for mucin oligosaccharides in the evolution

- 1 of lutropin to chorionic gonadotropin in the animal phyla. *J. Biol. Chem.* 280:16676-
2 16684.
- 3 Naylor S. L., W. W. Chin, H. M. Goodman, P. A. Lalley, K. H. Grzeschik, and A. Y.
4 Sakaguchi. 1983. Chromosome assignment of genes encoding the alpha and beta subunits
5 of glycoprotein hormones in man and mouse. *Somatic Cell Genet.* 9:757-770.
- 6 Nilsson C., K. Pettersson, R. P. Millar, K. A. Coerver, M. M. Matzuk, and I. T. Huhtaniemi.
7 1997. Worldwide frequency of a common genetic variant of luteinizing hormone: an
8 international collaborative research. International Collaborative Research Group. *Fertil.*
9 *Steril.* 67:998-1004.
- 10 Nilsson C., M. Jiang, K. Pettersson, A. Iitia, M. Makela, H. Simonsen, S. Easteal, R. J.
11 Herrera, and I. Huhtaniemi. 1998. Determination of a common genetic variant of
12 luteinizing hormone using DNA hybridization and immunoassays. *Clin. Endocrinol.*
13 *(Oxf)* 49:369-376.
- 14 O'Connor J. F., N. Elish, T. Kakuma, J. Schlatterer, and G. Kovalevskaya. 1998. Differential
15 urinary gonadotrophin profiles in early pregnancy and early pregnancy loss. *Prenat.*
16 *Diagn.* 18:1232-1240.
- 17 Odell W. D., J. Griffin, H. M. Bashey, and P. J. Snyder. 1990. Secretion of chorionic
18 gonadotropin by cultured human pituitary cells. *J. Clin. Endocrinol. Metab.* 71:1318-
19 1321.
- 20 Okuda K., T. Yamada, H. Imoto, H. Komatsubara, and O. Sugimoto. 1994. Antigenic
21 alteration of an anomalous human luteinizing hormone caused by two chorionic
22 gonadotropin-type amino-acid substitutions. *Biochem. Biophys. Res. Commun.* 200:584-
23 590.
- 24 Opazo J. C., D. E. Wildman, T. Prychitko, R. M. Johnson, and M. Goodman. 2006.
25 Phylogenetic relationships and divergence times among New World monkeys
26 (Platyrrhini, Primates). *Molecular Phylogenetics and Evolution*, 40:274-280.
- 27 Pestell R. G., A. N. Hollenberg, C. Albanese, and J. L. Jameson. 1994. c-Jun represses
28 transcription of the human chorionic gonadotropin alpha and beta genes through distinct
29 types of CREs. *J. Biol. Chem.* 269:31090-31096.
- 30 Pettersson K., M. M. Mäkelä, P. Dahlén, T. Lamminen, K. Huoponen, and I. Huhtaniemi.
31 1994. Gene polymorphism found in the LH beta gene of an immunologically anomalous
32 variant of human luteinizing hormone. *Eur. J. Endocrinol.* 130:Suppl. 2 Abstract S17.03.
- 33 Pettersson K., Y. Q. Ding, and I. Huhtaniemi. 1992. An immunologically anomalous
34 luteinizing hormone variant in a healthy woman. *J. Clin. Endocrinol. Metab.* 74:164-171.
- 35 Pierce J. G., and T. F. Parsons. 1981. Glycoprotein hormones: structure and function. *Annu.*
36 *Rev. Biochem.* 50:465-495.
- 37 Policastro P., C. Ovitt, M. Hoshina, H. Fukuoka, M. Boothby, and I. Boime. 1983. The beta
38 subunit of human chorionic gonadotropin is encoded by multiple genes. *J. Biol. Chem.*
39 258:11492-11499.

- 1 Policastro P., S. Daniels-McQueen, G. Carle, and I. Boime. 1986. A map of the hCG beta-LH
2 beta gene cluster. *J. Biol. Chem.* 261:5907-5916.
- 3 Poux C., P. Chevret, D. Huchon, W. W. de Jong, and E. J. Douzery. 2006. Arrival and
4 diversification of caviomorph rodents and platyrrhine primates in South America. *Syst.*
5 *Biol.* 55:228-244.
- 6 Ramanujam L., W. X. Liao, A. C. Roy, S. C. Ng, and S. S. Ratnam. 1998. Molecular variants
7 of luteinizing hormone in three populations of Southeast Asia. *Hum. Hered.* 48:232-234.
- 8 Ramanujam L. N., W. X. Liao, A. C. Roy, A. Loganath, H. H. Goh, and S. C. Ng. 1999.
9 Association of molecular variants of luteinizing hormone with menstrual disorders. *Clin.*
10 *Endocrinol. (Oxf)* 51:243-246.
- 11 Rhesus Macaque Genome Sequencing and Analysis Consortium, R. A. Gibbs, J. Rogers, M.
12 G. Katze, R. Bumgarner, G. M. Weinstock, E. R. Mardis, K. A. Remington, R. L.
13 Strausberg, J. C. Venter, R. K. Wilson, M. A. Batzer, C. D. Bustamante, E. E. Eichler, M.
14 W. Hahn, R. C. Hardison, K. D. Makova, W. Miller, A. Milosavljevic, R. E. Palermo, A.
15 Siepel, J. M. Sikela, T. Attaway, S. Bell, K. E. Bernard, C. J. Buhay, M. N. Chandrabose,
16 M. Dao, C. Davis, K. D. Delehaunty, Y. Ding, H. H. Dinh, S. Dugan-Rocha, L. A.
17 Fulton, R. A. Gabisi, T. T. Garner, J. Godfrey, A. C. Hawes, J. Hernandez, S. Hines, M.
18 Holder, J. Hume, S. N. Jhangiani, V. Joshi, Z. M. Khan, E. F. Kirkness, A. Cree, R. G.
19 Fowler, S. Lee, L. R. Lewis, Z. Li, Y. S. Liu, S. M. Moore, D. Muzny, L. V. Nazareth, D.
20 N. Ngo, G. O. Okwuonu, G. Pai, D. Parker, H. A. Paul, C. Pfannkoch, C. S. Pohl, Y. H.
21 Rogers, S. J. Ruiz, A. Sabo, J. Santibanez, B. W. Schneider, S. M. Smith, E. Sodergren,
22 A. F. Svatek, T. R. Utterback, S. Vattathil, W. Warren, C. S. White, A. T. Chinwalla, Y.
23 Feng, A. L. Halpern, L. W. Hillier, X. Huang, P. Minx, J. O. Nelson, K. H. Pepin, X. Qin,
24 G. G. Sutton, E. Venter, B. P. Walenz, J. W. Wallis, K. C. Worley, S. P. Yang, S. M.
25 Jones, M. A. Marra, M. Rocchi, J. E. Schein, R. Baertsch, L. Clarke, M. Csuros, J.
26 Glasscock, R. A. Harris, P. Havlak, A. R. Jackson, H. Jiang, Y. Liu, D. N. Messina, Y.
27 Shen, H. X. Song, T. Wylie, L. Zhang, E. Birney, K. Han, M. K. Konkel, J. Lee, A. F.
28 Smit, B. Ullmer, H. Wang, J. Xing, R. Burhans, Z. Cheng, J. E. Karro, J. Ma, B. Raney,
29 X. She, M. J. Cox, J. P. Demuth, L. J. Dumas, S. G. Han, J. Hopkins, A. Karimpour-Fard,
30 Y. H. Kim, J. R. Pollack, T. Vinar, C. Addo-Quaye, J. Degenhardt, A. Denby, M. J.
31 Hubisz, A. Indap, C. Kosiol, B. T. Lahn, H. A. Lawson, A. Marklein, R. Nielsen, E. J.
32 Vallender, A. G. Clark, B. Ferguson, R. D. Hernandez, K. Hirani, H. Kehrer-Sawatzki, J.
33 Kolb, S. Patil, L. L. Pu, Y. Ren, D. G. Smith, D. A. Wheeler, I. Schenck, E. V. Ball, R.
34 Chen, D. N. Cooper, B. Giardine, F. Hsu, W. J. Kent, A. Lesk, D. L. Nelson, W. E.
35 O'brien, K. Prufer, P. D. Stenson, J. C. Wallace, H. Ke, X. M. Liu, P. Wang, A. P. Xiang,
36 F. Yang, G. P. Barber, D. Haussler, D. Karolchik, A. D. Kern, R. M. Kuhn, K. E. Smith,
37 and A. S. Zweig. 2007. Evolutionary and biomedical insights from the rhesus macaque
38 genome. *Science* 316:222-234.
- 39 Roberts R. M. 2007. Interferon-tau, a Type 1 interferon involved in maternal recognition of
40 pregnancy. *Cytokine Growth Factor Rev.* 18:403-408.
- 41 Roberts R. M., Y. Chen, T. Ezashi, and A. M. Walker. 2008. Interferons and the maternal-
42 conceptus dialog in mammals. *Semin. Cell Dev. Biol.* 19:170-177.
- 43 Rull K., and M. Laan. 2005. Expression of beta-subunit of HCG genes during normal and
44 failed pregnancy. *Hum. Reprod.* 20:3360-3368.

- 1 Rull K., P. Hallast, L. Uuskula, J. Jackson, M. Punab, A. Salumets, R. K. Campbell, and M.
2 Laan. 2008. Fine-scale quantification of HCG beta gene transcription in human
3 trophoblastic and non-malignant non-trophoblastic tissues. *Mol. Hum. Reprod.* 14:23-31.
- 4 Scammell J. G., J. D. Funkhouser, F. S. Moyer, S. V. Gibson, and D. L. Willis. 2008.
5 Molecular cloning of pituitary glycoprotein α -subunit and follicle stimulating hormone
6 and chorionic gonadotropin β -subunits from New World squirrel monkey and owl
7 monkey. *Gen. Comp. Endocrinol.* 155:534-541.
- 8 Schrago C. G. 2007. On the time scale of New World primate diversification. *Am. J. Phys.*
9 *Anthropol.* 132:344-354.
- 10 Schrago C. G., and C. A. M. Russo. 2003. Timing the Origin of New World Monkeys. *Mol*
11 *Biol Evol* 20:1620-1625.
- 12 Sherman G. B., M. W. Wolfe, T. A. Farmerie, C. M. Clay, D. S. Threadgill, D. C. Sharp, and
13 J. H. Nilson. 1992. A single gene encodes the beta-subunits of equine luteinizing
14 hormone and chorionic gonadotropin. *Mol. Endocrinol.* 6:951-959.
- 15 Simula A. P., F. Amato, R. Faast, A. Lopata, J. Berka, and R. J. Norman. 1995. Luteinizing
16 hormone/chorionic gonadotropin bioactivity in the common marmoset (*Callithrix*
17 *jacchus*) is due to a chorionic gonadotropin molecule with a structure intermediate
18 between human chorionic gonadotropin and human luteinizing hormone. *Biol. Reprod.*
19 53:380-389.
- 20 Spincemaille J., R. Bouters, M. Vandeplassche, and P. Bonte. 1975. Some aspects of
21 endometrial cup formation and PMSG production. *J Reprod Fert* 23, Supplement 415-
22 418.
- 23 Starka L., M. Hill, R. Hampl, and I. T. Huhtaniemi. 1999. Genetic variant of luteinizing
24 hormone in Czech Republic. *Endocr. Regul.* 33:103-108.
- 25 Stenman U. H., H. Alfthan, T. Ranta, E. Vartiainen, J. Jalkanen, and M. Seppala. 1987. Serum
26 levels of human chorionic gonadotropin in nonpregnant women and men are modulated
27 by gonadotropin-releasing hormone and sex steroids. *J. Clin. Endocrinol. Metab.* 64:730-
28 736.
- 29 Storrington P. L., S. A. Khan, Y. G. Mistry, and E. Diczfalussy. 1988. Plasma clearance in the rat
30 of the LH bioactivity of two human LH standards of differing molecular composition. *J.*
31 *Endocrinol.* 119:327-324.
- 32 Sugino H., G. R. Bousfield, W. T. Moore Jr, and D. N. Ward. 1987. Structural studies on
33 equine glycoprotein hormones. Amino acid sequence of equine chorionic gonadotropin
34 beta-subunit. *J. Biol. Chem.* 262:8603-8609.
- 35 Talmadge K., N. C. Vamvakopoulos, and J. C. Fiddes. 1984. Evolution of the genes for the
36 beta subunits of human chorionic gonadotropin and luteinizing hormone. *Nature* 307:37-
37 40.
- 38 Talmadge K., W. R. Boorstein, and J. C. Fiddes. 1983. The human genome contains seven
39 genes for the beta-subunit of chorionic gonadotropin but only one gene for the beta-
40 subunit of luteinizing hormone. *DNA* 2:281-289.

- 1 Talmadge K., W. R. Boorstein, N. C. Vamvakopoulos, M. J. Gething, and J. C. Fiddes. 1984.
2 Only three of the seven human chorionic gonadotropin beta subunit genes can be
3 expressed in the placenta. *Nucleic Acids Res.* 12:8415-8436.
- 4 Tannenbaum P. L., N. J. Schultz-Darken, M. J. Woller, and D. H. Abbott. 2007a.
5 Gonadotrophin-Releasing Hormone (GnRH) Release In Marmosets II: Pulsatile Release
6 of GnRH and Pituitary Gonadotrophin in Adult Females. *J. Neuroendocrinol.* 19:354-
7 363.
- 8 Tannenbaum P. L., N. J. Schultz-Darken, W. Saltzman, E. Terasawa, M. J. Woller, and D. H.
9 Abbott. 2007b. Gonadotrophin-releasing hormone (GnRH) release in marmosets I: in
10 vivo measurement in ovary-intact and ovariectomised females. *J. Neuroendocrinol.*
11 19:342-353.
- 12 Tapanainen J., M. Koivisto, R. Vihko, and I. Huhtaniemi. 1981. Enhanced activity of the
13 pituitary-gonadal axis in premature human infants. *J. Clin. Endocrinol. Metab.* 52:235-
14 238.
- 15 Valmu L., H. Alfthan, K. Hotakainen, S. Birken, and U. H. Stenman. 2006. Site-specific
16 glycan analysis of human chorionic gonadotropin beta-subunit from malignancies and
17 pregnancy by liquid chromatography-electrospray mass spectrometry. *Glycobiology*
18 16:1207-1218.
- 19 Wehmann R. E., and B. C. Nisula. 1981. Metabolic and renal clearance rates of purified
20 human chorionic gonadotropin. *J. Clin. Invest.* 68:184-194.
- 21 Wooding F. B., G. Morgan, A. L. Fowden, and W. R. Allen. 2001. A structural and
22 immunological study of chorionic gonadotrophin production by equine trophoblast girdle
23 and cup cells. *Placenta* 22:749-767.
- 24 Wu H., J. W. Lustbader, Y. Liu, R. E. Canfield, and W. A. Hendrickson. 1994. Structure of
25 human chorionic gonadotropin at 2.6 Å resolution from MAD analysis of the
26 selenomethionyl protein. *Structure* 2:545-558.
- 27 Wu W., and A. M. Walker. 2006. Human chorionic gonadotropin beta (HCGbeta) down-
28 regulates E-cadherin and promotes human prostate carcinoma cell migration and
29 invasion. *Cancer* 106:68-78.
- 30 Zhang J. 2003. Evolution by gene duplication: an update. *Trends in Ecology & Evolution*,
31 18:292-298.
- 32 Ziecik A. J., M. M. Kaczmarek, A. Blitek, A. E. Kowalczyk, X. Li, and N. A. Rahman. 2007.
33 Novel biological and possible applicable roles of LH/hCG receptor. *Mol. Cell.*
34 *Endocrinol.* 269:51-60.
- 35 Zygmunt M., F. Herr, S. Keller-Schoenwetter, K. Kunzi-Rapp, K. Munstedt, C. V. Rao, U.
36 Lang, and K. T. Preissner. 2002. Characterization of human chorionic gonadotropin as a
37 novel angiogenic factor. *J. Clin. Endocrinol. Metab.* 87:5290-5296.

38
39