Molecular biomarkers of endocrine disruption in small model fish

Stefan Scholz, Ian Mayer

► To cite this version:

Stefan Scholz, Ian Mayer. Molecular biomarkers of endocrine disruption in small model fish. Molecular and Cellular Endocrinology, 2008, 293 (1-2), pp.57. 10.1016/j.mce.2008.06.008 . hal-00532039

HAL Id: hal-00532039

<https://hal.science/hal-00532039>

Submitted on 4 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Molecular biomarkers of endocrine disruption in small model fish

Authors: Stefan Scholz, Ian Mayer

PII: S0303-7207(08)00268-2
DOI: doi:10.1016/j.mce.2008.06.008
Reference: MCE 6897

To appear in: *Molecular and Cellular Endocrinology*

Received date: 11-2-2008
Revised date: 5-5-2008
Accepted date: 11-6-2008

Please cite this article as: Scholz, S., Mayer, I., Molecular biomarkers of endocrine disruption in small model fish, *Molecular and Cellular Endocrinology* (2007), doi:10.1016/j.mce.2008.06.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title

Molecular biomarkers of endocrine disruption in small model fish

Authors

Stefan Scholz^{1c}, Ian Mayer²

¹ UFZ - Helmholtz Centre for Environmental Research

Department of Cell Toxicology

Permoserstr. 15

04318 Leipzig

Germany

Email: stefan.scholz@ufz.de

Phone: +49 341 235 1217

Fax: +49 341 235 2324

² Department of Biology

University of Bergen

Thormøhlensgt. 55

5020 Bergen

Norway

Email: ian.mayer@bio.uib.no

Phone: +47 55 584442

^c Corresponding author: Stefan Scholz, UFZ - Helmholtz Centre for Environmental Research, Department of Cell Toxicology, Permoserstr. 15, 04318 Leipzig, Germany, Email: stefan.scholz@ufz.de, phone: +49 341 235 1217, fax: +49 341 235 2324

1 Fax: +47 55 584450

3 **Summary**

4 A wide range of environmental contaminants can interfere with the hormonal regulation in
5 vertebrates. These endocrine disrupting chemicals (EDCs) are of high relevance for human
6 and wildlife health, since endocrine signalling controls many essential physiological
7 processes which impact on the individual's health, such as growth and development, stress
8 response, and ultimately reproduction and population development. Small fish represent a
9 cost-effective model for testing potential EDCs allowing the possibility to integrate from
10 molecular to phenotypic and functional effects. We have comprehensively reviewed
11 exposure-effect data from four different small model fish: zebrafish, medaka, fathead
12 minnow, and the three-spined stickleback. The majority of available data refer to EDCs
13 interfering with reproductive hormones. However, we have also included interactions with
14 other hormone systems, particularly the thyroid hormones. We demonstrate that the available
15 data clearly indicates the predictive potential of molecular biomarkers, supporting the
16 development and regulatory application of simple molecular-based screening assays using
17 small model fish for EDC testing.

21 **Keywords**

22 Endocrine disruption, biomarker, hormone receptors, vitellogenin, spiggin, (teleost)

24 **1. Introduction**

25 There is an increasing public concern over the adverse effects of anthropogenic
26 contaminants on the health and well-being of both humans and wildlife (WHO, 2002). To

1 date, a wide variety of both developmental and reproductive disorders observed in wildlife
2 species have clearly been linked to the exposure to environmental contaminants which act as
3 endocrine disrupting chemicals (EDCs, Edwards et al., 2006; Tyler et al., 1998; WHO, 2002),
4 that is “exogenous agents that interfere with the production, release, transport, metabolism,
5 binding, action or elimination of natural hormones in the body responsible for the
6 maintenance of homeostasis and the regulation of developmental processes” (Kavlock et al.
7 1996). EDCs are of high environmental relevance, since many essential physiological
8 processes which impact on the individual’s health, such as growth and development, stress
9 response, and ultimately reproduction and population development are controlled by
10 hormones. Regardless of the initial type of interaction, the effects of EDCs with different
11 modes of action are often very similar and resemble changes in endogenous hormone levels.

12 In vertebrates, different types of hormone systems could principally be targets of
13 environmental chemicals: reproductive hormones such as estrogens and androgens, thyroidal
14 hormones, corticosteroids, growth hormone and their associated hypothalamus/pituitary
15 releasing and stimulating hormones. However, research has to date focused nearly entirely on
16 the interference with the steroidal reproductive hormones for obvious reasons. Firstly,
17 reproductive hormones control one of the most important endpoints in the risk assessment of
18 environmental chemicals, that is all aspects of reproductive development from sex
19 differentiation to puberty, aspects of which are critical for population viability. Secondly,
20 reproductive hormones are of vital importance during the early critical stages of embryonic
21 development and sex differentiation. As a result, even though the binding affinity of most
22 EDCs to the ER/AR is low compared to endogenous steroid hormones, exposure to even low
23 concentrations of xenoestrogens/xenoandrogens can disrupt normal
24 developmental/reproductive processes. This is evident for exposure to micropollutants, that is,
25 anthropogenic chemicals continuously detected at low concentrations in the environment. A
26 well known example is ethinylestradiol (EE₂), which is discharged into the aquatic

environment via municipal effluents due to its use in contraceptive pills. EE₂ is found in the lower ng/L-range in aquatic surface water, a concentration that leads to sustainable impacts on reproduction and population development in fish (Larsson et al., 1999).

Fish are considered as one of the primary risk organisms for EDCs, especially those interacting with the reproductive hormones. Firstly, sex determination in teleost fishes is characterised as being very labile, and can be disturbed or even functionally reversed by the external application of hormones at critical developmental stages (Francis, 1992). Secondly, as the aquatic environment represents the ultimate sink for most anthropogenic contaminants, fish populations are directly exposed to a wide variety of EDCs, originating from industrial, agricultural or municipal effluents. There is now unequivocal evidence showing that EDCs can have long-term effects on reproduction and subsequent population development in natural fish populations (Kidd et al., 2007). However, investigations based on the evaluation of reproductive endpoints are costly, time-consuming and laborious. This has generally resulted in these endpoints not being routinely integrated into most chemical safety assessments. However, since endocrine disruption can be clearly linked to molecular interactions, expression of appropriate biomarkers could be used as predictors for reproductive or other effects.

1.1. The relevance of using small fish models for endocrine disruption research

Small fish represent a cost-effective model for large scale chemical testing. They are easy to maintain in small laboratory aquaria, have a short life-cycle, and readily produce many young. In contrast to targeted *in vitro* cellular models such as the yeast screen assay for estrogenic compounds or primary cell cultures, small fish models allow the possibility to integrate alternative modes of action, such as compounds that modify endogenous hormone levels. Furthermore, uptake, metabolism and excretion as important constraints of bioavailability are considered. The attractiveness of using small fish models in chemical

testing programmes is further enhanced by recent advances in molecular approaches. DNA markers for molecular sex determination are now available for some small fish models (medaka and stickleback, Griffiths et al., 2000; Matsuda et al., 2002). Moreover, the genome has now been sequenced for three of the species highlighted in this review (medaka, zebrafish and stickleback), resulting in the potential development of new molecular tools.

For this review we have selected 4 different model fish that are ideally suited to establish links between molecular biomarkers and phenotypic and functional effects of EDCs: zebrafish, medaka, fathead minnow, and the three-spined stickleback. Each of these organisms is well established for the testing of environmental chemicals providing different advantages and possibilities such as detailed genetic characterisation, sex specific markers, and/or use as an experimental and biomonitoring model. Due to their short life-cycle, the selected species allow for the analysis of reproductive performance, which is one of the most important endpoints for long-term effects with implications on population development. A brief description of the model small fish species is given in Table 1 (detailed information on the different models can be obtained from the review of Ankley and Johnson, 2004 and Katsiadaki et al. 2007).

1.2. Scope of the review

In this review, we summarise the available data on endocrine disruption in fish highlighting the links between molecular, phenotypic and physiological effects (in particular reproduction). We focus nearly exclusively on small fish models, since they provide the experimental basis to establish appropriate correlations between the different effect levels including reproductive and transgenerational effects. For the first time, data are presented in a comprehensive manner, providing an overview of the effect levels of different molecular and phenotypic endpoints in the selected model species. The majority of available data refer to EDCs interfering with reproductive hormones. However, we have also reviewed interactions

with other hormone systems, indicating further potential applications for small fish models. We show that the available data clearly indicate the predictive potential of molecular biomarkers supporting the development of screening assays for endocrine disruption such as initiated by the OECD (OECD, 2006).

2. Data sources and graphical presentation

The Web of Science (<http://portal.isiknowledge.com/>) and references have been searched for publications (up to April 2008) on endocrine disrupting effects in the following four small fish models: fathead minnow (*Pimephales promelas*), zebrafish (*Danio rerio*), medaka (*Oryzias latipes*), and the three-spined stickleback (*Gasterosteus aculeatus*). Combinations of the terms “endocrine”, “disruption” and partial or entire names of species and hormones following a manual selection of relevant articles have been used for the literature survey. The review has been prepared independently and in parallel to an USEPA report (USEPA, 2007) summarising endocrine disrupting effects on the fathead minnow. Data reported for the fathead minnow largely overlap with the USEPA report, which contains additional data without graphical representation but is partially more detailed.

With respect to concentration response levels of experimental studies we established a quality control. Only those studies that used water-borne exposure, tested more than one concentration of a potential EDC (in case several studies with the same endpoint have been available) and provided statistical analysis to extract a LOEC (lowest observed effect concentration) were considered. *Ex vivo* studies (i.e. incubation of oocytes or sperm) have been excluded. In case the lowest tested concentrations caused an effect this concentration was included as the LOEC as well. In the supplemented table S1 these values were marked by “ $\leq$ ”. Furthermore, studies with conflicting results (contradictive effects in repeated experiments) were excluded. Occasionally, effects were reported at one low concentration but not at higher levels. If no mechanistic basis could be ascribed to these findings, these studies

were also excluded. Based on these criteria, the data from 147 studies performed with one of the selected small model species were selected for this review. Concentration levels reflect measured or nominal concentrations. Not all studies included nominal concentrations validated by chemical analysis. We decided to represent effect concentration in $\mu\text{g/L}$ or mg/L , since for the majority of studies concentrations are not given as molar concentration.

In order to summarise the large data set on reproductive steroid hormones we chose a graphical representation. Although such a representation cannot provide all the information that might be of interest it is ideal to give a comprehensive and easy to grasp overview of a large number of data. However, it was not possible to include information on exposure conditions into the graphs. Therefore, for the interested reader we have provided tables with detailed data as supplementary information (Table S1). Data were summarised in different categories: effects on molecular biomarkers, secondary sex characteristics (if available for the species), histopathology (gonad histology, sex reversal, gonado-somatic index, thyroid histology) and reproduction (spawning, fecundity, sperm quality and population development). For thyroid interfering compounds, information on hormone levels, teratogenic effects and toxicity have been included as well. The molecular markers were subdivided as vitellogenin expression/level or others (aromatase expression/activity, gene expression of other genes, steroid levels).

3. Reproductive hormones

In common with all vertebrates, reproduction in teleost fish is under the control of the brain-pituitary-gonadal (BPG) axis. In seasonally spawning fish external factors such as light and water temperature control the timing of gonad recrudescence and maturation via the brain. Signals from the brain control the hypothalamic secretion of gonadotropin-releasing hormones which stimulate the release of gonadotropins (luteinizing hormone [LH] and follicle-stimulating hormone [FSH]) from the pituitary. The gonadotropins in turn control the

production and secretion of steroid hormones from the gonads. These hormones (17β -estradiol and testosterone in females, and testosterone and predominantly 11-ketotestosterone in males, as well as maturation-inducing hormones in both sexes) initiate changes in secondary sex characteristics, behaviour as well as development and maturation of gametes. Via negative or positive feedback mechanisms, depending on the physiological stage, reproductive steroids regulate the secretion of pituitary gonadotropins (Peter & Yu 1997, Schulz & Goos 1999).

The BPG axis provides several potential modes of interaction for EDCs ranging from interference with hormone synthesis and secretion, binding to hormone receptors, or altering the availability of endogenous hormones. Eventually, endocrine disrupting effects are mediated via the available level of steroid hormones or its analogs. Thus, biomarkers directly responsive to steroid hormone levels are the most widely used molecular indicators of endocrine disruption. A popular example is the synthesis and secretion of the yolk protein vitellogenin which is induced in the liver by the circulating female steroid hormone, 17β -estradiol and has been specifically applied to identify xenoestrogens (Tyler et al. 1996).

3.1. Estrogenic and anti-estrogenic effects

A wide variety of structurally diverse anthropogenic compounds have been shown to exhibit estrogenic or anti-estrogenic effects (Jobling et al., 1995; Kime, 1998). Estrogenic effects refer to compounds mimicking the action of estrogens via binding to the estrogen receptor (ER) and activation of responsive genes. In teleosts, two receptor isoforms (α and β and subtypes of these) with differential tissue distribution patterns and binding affinities have been identified (e.g., Hawkins & Thomas 2004, Menuet et al. 2002). Furthermore, estrogenic effects may also be mediated by elevation of endogenous free plasma estrogen levels caused by competitive binding of xenobiotics to sex steroid binding globulins (e.g., Danzo 1997, Tollefsen 2002). Anti-estrogenic effects refer to compounds with weak or no estrogenic

activity but the ability to competitively bind to the ER. Furthermore, anti-estrogenic action can be mediated by affecting the expression and activity of the cytochrome P450A19 enzyme (aromatase). A number of clinical compounds and pesticides are known to inhibit aromatase, which catalyses the final step in the synthesis of estrogens, i.e. the introduction of an aromatic ring in the androgen hormone precursor. Inhibition of this step leads to a reduction of 17β -estradiol plasma concentration resulting in reduced vitellogenin levels in female fish (see supplemented Table S1). Aromatase inhibitors could also be classified as androgenic compounds due to their associated masculinising effects (e.g. Kwon et al., 2000; Mak et al., 1999). These masculinising effects are presumably provoked by accumulation of androgens since fish treated with the aromatase inhibitor fadrozole show elevated 11-ketotestosterone levels (e.g., Ankley et al. 2002). However, due to their primary involvement in estrogen-synthesis inhibition, we decided to summarise aromatase inhibitors along with (anti)estrogenic and estrogenic compounds. Effect levels from 112 references - in which an effect of an endocrine disruptor using one of the model teleosts has been described for waterborne exposures - were extracted (Fig. 1). The test compounds comprised of known and putative estrogens (17α -ethinylestradiol, 17β -estradiol, estrone, bisphenol A, 4-aminobenzoate, 3-benzylidene camphor, benzophenone, diethylstilbestrol, β -HCH, o,p-DDT, pp-DDE, letrozole, octyl methoxycinnamate, 4-nonylphenol, 4-tert-octylphenol, 4-pentylphenol), anti-estrogens (equol, genistein, tamoxifen, ZM 189,156), aromatase inhibitors (fadrozole, tributyltin) or potential xenoestrogens of unknown mechanisms (atrazine, benzo(a)pyrene, butyl benzyl phthalate, cyproconazole, p-dichlorobenzene, o-dichlorobenzene, m-dichlorobenzene, nianserin, methoxychlor, propylparaben). Some compounds are difficult to classify, since they interfere with the synthesis of steroids and may affect both female and male hormonal regulation (e.g. ketoconazole). Aromatase inhibitors that also inhibit androgen synthesis and bind to the androgen receptor, such as prochloraz and fenimarol, are described in section 3.2 (anti-androgenic and androgenic effects). The majority

1 of studies (43) were performed with the fathead minnow and zebrafish (42), followed by
2 medaka (24) and stickleback (11). Effect levels for individual compounds vary by up to four
3 orders of magnitude (or 5 orders of magnitude if the androgenic marker spiggin is included).
4 However, particularly for the zebrafish and medaka, this can be attributed to the different
5 exposure regimes using either different stages or exposure durations. Most of the studies
6 performed with fathead minnow and stickleback followed OECD recommendations (21 days
7 of exposure of adult fish; OECD, 2006). In these studies, inter-laboratory variation and use of
8 either nominal or measured concentrations to describe the LOEC may represent the major
9 source of variation. Many studies may also have underestimated the effect levels, since the
10 appropriate effects were already detected in the lowest tested concentrations. On the other
11 hand, the ranking of the (anti)estrogenic activity of the different compounds is for most of the
12 compounds highly conserved among the species with average differences in sensitivity below
13 one order of magnitude. Furthermore, despite the variability in experimental parameters, a
14 large overlap in effect concentrations for the endpoint vitellogenin (induction or inhibition),
15 various molecular endpoints (aromatase inhibition, hormone levels, expression of various
16 genes), secondary sex characteristics (tubercles, fins, nuptial coloration), histopathologic
17 effects (gonadal structure, gonadosomatic indices, incidence of intersex and gonadal sex
18 reversal) and reproduction (spawning, fecundity, fertilisation rate in offspring etc.) is apparent
19 (see also supplement Table S1). For those compounds, for which several endpoints have been
20 analysed, the molecular endpoints (VTG levels and others) generally appear to be among the
21 most sensitive endpoints with particular overlap to environmentally relevant endpoints, i.e.
22 reproduction. Exposure periods of 2-3 weeks generally appear to be sensitive enough to affect
23 these molecular endpoints in the same concentration that cause reproductive effects. It may
24 thus be concluded that for screening of chemicals for their (anti)estrogenic capacity, analysis
25 of molecular endpoints in relatively simple screening assays may be sufficient to predict
26 potential adverse effects for higher tier level testing. The inhibition of the androgen marker

spiggin (see below) has also been analysed in male sticklebacks exposed to estrogens. Notably, exposure of sticklebacks to relatively high concentrations of estrogens can result in a decrease in spiggin levels probably caused by indirect effects of estrogens on this androgenic marker (Fig. 1 D, Jolly, 2007, Katsiadaki et al., unpublished data). This indicates the need for molecular markers closely linked to the mode-of-action.

3.2. Androgenic and anti-androgenic effects

Androgenic effects are of similar environmental relevance to estrogenic effects, and in fact have historically been detected in wildlife as early as the first reports of estrogenic effects (Howell et al., 1980; Jafri and Ensor, 1979). In these early studies, before the concept of endocrine disruption had been fully recognised, female mosquitofish (*Gambusia affinis*) sampled downstream of a paper mill were found to have developed a gonopodium, a male specific secondary sexual character. Various subsequent studies on fish have confirmed the presence of androgenic ('masculinising') compounds in paper and pulp mill effluent (Cody and Bortone, 1997; Larsson et al., 2000). Anti(androgenic) effects are mediated via direct binding of a xenobiotic to the androgen receptor (AR), but may also be provoked by inhibition of estradiol synthesis and accumulation of endogenous androgen levels (see 3.2). Two AR subtypes were reported for rainbow trout (*Oncorhynchus mykiss*), eel kelp bass (*Paralabrax clathratus*) and Atlantic croaker (*Micropogonias undulatus*), termed either AR1 and 2 or AR α and β with distinctive binding properties or transactivation activity for androgens including testosterone and 11-ketotestosterone (Sperry & Thomas 1999, Takeo & Yamashita 2000). For other species only a single AR with homology to either a teleost α or β subtype could be identified so far (Hossain et al. 2007)

A number of contaminants interfering with androgen regulation have been identified, such as the pesticides linuron and fenitrothion, and the fungicides vinclozolin and prochloraz (Makynen et al., 2000; Katsiadaki et al., 2006; Vinggaard et al., 2006).

Furthermore, there is growing concern over the environmental impact of animal feedlot effluents due to the use of anabolic growth promoters commonly used on livestock in the USA, such as the highly potent synthetic androgen trenbolone (Lange et al., 2002). Two isomers of this androgen, 17 α -trenbolone and 17 β -trenbolone, have recently been detected in animal feedlot effluent in Ohio (Durhan et al., 2006) both of which have been shown to be strongly androgenic in the fathead minnow (Ankley et al., 2003).

Detection of (anti)androgenic effects has been hampered by the unavailability of androgen regulated marker genes in fish which may explain the greater focus on (anti)estrogenic effects. However, since the detection, characterisation and cloning of spiggin - a glue protein synthesized in the kidney of male sticklebacks for the purpose of nest building - an appropriate and quantifiable biomarker for screening of androgens both *in vivo* and *in vitro* is available in teleosts as well (Jolly et al., 2006; Jones et al., 2001; Nagae et al., 2007).

Furthermore, the antiandrogenic properties of test compounds can also be quantitatively evaluated by assessing their ability to suppress the androgen-stimulated induction of spiggin release in male sticklebacks. However, due to the large variability and high levels of spiggin in males the use of females co-exposed with androgens have been preferred (Jolly, 2007; Katsiadaki et al., 2006).

Despite the advantages of spiggin measurement, the majority of studies investigating (anti) androgenic effects have been performed using the fathead minnow (19) and zebrafish (6) and only three reports – based on the above listed criteria - are available for the medaka (Figure 2, supplement Table S1). Molecular effects of (anti)androgens were detected indirectly by analysis of vitellogenin induction or suppression, by analysis of various genes/proteins (LH, FSH, aromatase, estrogen receptor a.o.) or testosterone levels. Although less quantitative, (anti)androgenic properties of test compounds could also be identified from secondary sex characters (tubercles), histopathology (gonadal structure, GSI, gonadal sex ratio) and reproductive impairment (spawning, fecundity, fertilisation rate in offspring). Nine

1 publications report (anti)androgens in the stickleback in waterborne exposure experiments.
2 The only endpoints reported were spiggin levels and kidney hypertrophy (kidney epithelial
3 cells are enlarged in male stickleback due to spiggin production, Figure 2, supplement Table
4 S1).

5 In common with the (anti)estrogen studies considered above, data on (anti)androgens
6 exhibited a variability in effect levels by up to four orders of magnitude for the different
7 reproductive endpoints, and up to two orders of magnitude for the same endpoint. The same
8 reasons as discussed for (anti)estrogenic effects may similarly apply for this variability.
9 However, a comparison of those compounds (vinclozolin, flutamide, trenbolone and
10 methyltestosterone) that have been tested on at least three of the selected model species
11 indicate a highly conserved ranking of anti(androgenic) effect levels. Due to the lower
12 number of available studies in general and particularly those analysing reproductive
13 endpoints, it is difficult to judge whether molecular endpoints may be used as a predictor of
14 adverse reproductive effects in screening assays. Although vitellogenin is among the most
15 sensitive endpoints, a molecular marker that is directly linked to the mode of action may be
16 more desirable and may provide a closer link to higher tier effect levels. However,
17 impairment of reproduction has not been investigated in parallel to the androgenic marker
18 spiggin.

19 Identification of androgen-sensitive genes may help to establish sensitive markers for
20 fathead minnow, zebrafish and medaka. A current microarray study by Leon et al. (2008) in
21 medaka exposed to the teleost-specific androgen 11-ketotestosterone, the antiandrogen
22 flutamide, and the antiandrogenic fungicide vinclozolin identified 578 androgen-sensitive
23 genes, of which 50 were able to discriminate between the test compounds. These genes
24 include gonadotropin-releasing hormones and their receptors, and genes involved in
25 steroidogenesis. Detailed analysis of the affected genes may reveal marker genes directly

regulated by androgens and would allow comparing higher tier levels with molecular responses in medaka, fathead minnow and zebrafish as well.

3.3. Other hormones of the BPG-axis

The analysis of endocrine disrupting effects has to date focussed mostly on estrogens and androgens and their target organs, the gonads or liver. However, the hormonal control of reproduction via the BPG-axis provides a number of additional potential targets for EDCs such as the synthesis and binding of releasing and trophic hormones or the control of final maturation by "maturation-inducing steroids" (C21-progestins). Interferences with releasing or trophic hormones are likely to be reflected in alterations in steroid hormones levels, and cause effects similar to direct interference (receptor binding or modulation of synthesis) with estrogens or androgens (Katti and Sathyanesan, 1986). Vice versa, changes in expression or circulating levels of gonadotropin by environmental contaminants may originate from exposure to (anti)estrogens or (anti)androgens and subsequent feedback adaptations via the hypothalamus and pituitary (Singh et al., 1994; Villeneuve et al., 2007b). Thus, *in vivo*, any type of interference with hypothalamus or pituitary hormones may be detected by the same endpoints demonstrating interactions with reproductive steroids, such as vitellogenin expression. Cloning and sequencing of genes for releasing and stimulating hormones in small model teleosts allows for a detailed analysis of interactions between hormones of the BPG axis (Gopinath et al., 2004; Hellqvist et al., 2004; Kwok et al., 2005; Okubo et al., 2001; Villeneuve et al., 2007b). A good example of this mode of action was demonstrated by Villeneuve et al. (2007b), who showed that exposure to the chemical ketoconazole, an inhibitor of steroid synthesis, led to an elevation of FSH β mRNA expression in male fathead minnows. No effect on LH and FSH expression could be observed in zebrafish exposed to benzo(a)pyren, albeit both aromatase and VTG expression were induced (Hoffmann and Oris, 2006).

Another important target of EDCs may involve maturation of oocytes and sperm, which is triggered by gonadotropins. Specifically, maturation of germ cells is triggered by gonadotropins but mediated by specific testicular and ovarian C21-steroid hormones (maturation-inducing hormone, MIH; $17\alpha,20\beta$ -dihydroxy-4-pregnen-3-one, $17\alpha,20\beta$ -DP in salmonid fishes; $17\alpha,20\beta,21$ -trihydroxy-4-pregnen-3-one, 20β -S in sciaenid fishes) synthesised from the ovarian follicular layer or sperm (Nagahama, 1994; Nagahama et al., 1993). Effects on maturation by EDCs have been studied for oocytes *in vivo* and *in vitro* in zebrafish. Dithylstilbestrol (DES) induced the maturation of oocytes, while pentachlorophenol caused an inhibition (Hachisuka et al., 2005; Tokumoto et al., 2004). The time and change in germinal vesicle breakdown and synthesis of cyclin B indicated that there was no distinguishable difference between exposure to either DES or the natural maturation-inducing hormone $17\alpha,20\beta$ -DP. Binding of DES to the membrane progesterin receptor- α suggests a specific interaction with oocyte maturation via a non-genomic pathway as well (Tokumoto et al., 2007). The analysis of target genes or proteins of the maturation hormones would allow the quantification of the strength of interaction, and to estimate the specificity of EDC interference with these hormones. However, these target genes – with the exception of the receptor itself - have yet not been identified for teleosts.

The previous reported examples of EDC action on the BPG-axis demonstrate the strength of *in vivo* testing systems in identifying potential adverse effects. On the other hand, it may be difficult to detect the underlying mode of action. This may be facilitated by the analysis of molecular targets used in combination with simple *in vitro* systems, in particular those allowing for the determination of binding to steroid hormone receptors (Coldham et al., 1997).

4. Interaction with thyroid hormones

1 The thyroid hormones play a crucial role in the maintenance of a normal physiological
 2 state in all vertebrates, notably in the control of development, growth and behaviour. In
 3 addition, in teleost fishes, the thyroid hormones also play a pivotal role in the regulation of
 4 metamorphosis, either the transition from larval to juvenile stages in marine flat fishes and
 5 eels, the transformation from adult yellow to silver eels, or smoltification in salmonids
 6 (Brown et al., 2004; Power et al., 2001). Seasonal variations of thyroid hormones are
 7 associated with the reproductive status in teleost fishes. These changes indicate a potential
 8 function in the control of both sexual differentiation and reproductive development (Cyr and
 9 Eales, 1996). The latter is also supported by a recent report on the masculinisation of
 10 sticklebacks by perchlorate, an inhibitor of thyroid hormone synthesis (Bernhardt et al.,
 11 2006). Furthermore, thyroids are also involved in the regulation of the immune system, as
 12 indicated by the suppression of immune function by hyperthyroidism (Yada and Nakanishi,
 13 2002).

14 In common with the reproductive steroid hormones, the synthesis and release of the
 15 thyroid hormones is under the control of a central brain-pituitary-thyroid (BPT) axis.
 16 Releasing and stimulating hormones trigger the production and release of the thyroid
 17 hormones. Thyroid-disrupting compounds can interact at the various levels of this regulatory
 18 pathway, in particular with the synthesis (via thyroglobulin iodination, or deiodination of
 19 thyroxine [T_4] to triiodothyronine [T_3]) and metabolism (conjugation) of thyroid hormones,
 20 interference with the plasma binding proteins, or by binding to the thyroid receptor (TR,
 21 Brown et al., 2004). Upon binding to its ligand, corepressors are replaced by activator
 22 proteins and TR homodimers or heterodimers with the retinoic X receptors induce the
 23 expression of target genes containing thyroid responsive elements - such as the
 24 phosphoenolpyruvate carboxykinase (Park et al., 1995) or 5'diodinase (Jakobs et al., 1997).
 25 Similar to what has been observed in mammals, two TR isoforms (α and β) and splicing
 26 variants of these have been identified for many teleosts (Marchand et al. 2001). All the

subtypes tested so far have exhibited transactivation activity for thyroid hormones.
(Kawakami et al. 2006, Marchand et al. 2001)

Environmental chemicals mimicking the thyroid hormone have not been described nor have TR-binding or transactivation properties been included in current screening strategies. In contrast, a wide variety of structurally diverse chemicals has been shown to interfere with thyroid hormone levels in fish (Brown et al., 2004). The majority of these effects, however, are mediated apparently indirectly through a general and/or complex interference of the contaminant with the physiology of the fish rather than specific interaction with thyroid regulation. Examples of environmental compounds that have been shown to specifically disrupt thyroid hormone synthesis include perchlorate, PCBs (Kime, 1998) and arsenic (Davey et al., 2008). In order to identify thyroid-disrupting compounds which can interfere with the synthesis of thyroid hormones or act as agonists/antagonists of the thyroid hormones, it is necessary to define the typical effect patterns of model compounds. This has been performed in zebrafish, fathead minnow and stickleback for inhibition of the thyroid action using compounds that interfere with thyroid hormone synthesis via thyroidal iodine uptake, thyroperoxidase and/or 5-deiodinase (Table 2). These model compounds derive from medical applications in the treatment of thyroidal disease. One of these model compounds, perchlorate, is also an environmentally relevant contaminant due to its application in solid fuel rockets and in the manufacture of flares, fireworks and automobile airbags (Greer et al., 2002). Typical effects observed in fish are lowering of endogenous T4 and/or T3 levels and thyroidal hypertrophy (Table 2). Frequently, a reduction of growth and developmental retardations has also been reported. Maternal effects on oocyte size, hatching and – in one case reported for the stickleback – sexual differentiation and spawning also suggest an interference with reproduction (Bernhardt et al., 2006). However, the effects occur generally in the mg/L range close to concentrations that elicit morphologically visible toxic effects. Interestingly, the effective concentrations overlap with those for compounds known to inhibit

metamorphosis in amphibians (Opitz et al., 2005). The potential long-term effects of lower dose treatments have not been studied in detail. Relevant long-term effects could include reduced reproductive performance, immune suppression (e.g. Lam et al., 2005) and growth effects. These types of effects would indicate a reduced individual fitness and performance with implications at the population level. It is obvious that such subtle effects may be difficult and laborious to detect. Therefore, molecular markers that are clearly linked to adverse effects could serve as indicators or early warning signals in short-term experiments. Genes (or their corresponding gene products) containing TRE elements such as phosphoenolpyruvate carboxy kinase (Park et al., 1995) could be – similar to estrogen regulated genes such as vitellogenin – potential sensitive markers for the interference with the thyroid axis. However, the differential regulation of fish genes with TRE elements, e.g. growth hormone (Farchi-Pisanty et al., 1997), has not yet been studied in response to potential endocrine disrupting chemicals.

5. Corticosteroids

Corticosteroids (glucocorticoids and mineralcorticoids) are involved in the regulation of various important physiological processes of environment adaptation and the stress response (changes in salinity, low oxygen concentrations, capture, presence of predators, pollutants etc., Barton, 2002). Like thyroid and reproductive steroid hormones, they are controlled by hypothalamus and pituitary hormones and exert their action through binding to intracellular receptors. Some species of fish possess two receptor isoforms of glucocorticoids resulting from a genome duplication (Schaaf et al., 2008, Alsop & Vijayan 2008, Bury et al. 2003). Although the mineral corticoid receptor has been detected, its function is not clear, since teleosts lack the capacity to synthesise the mineralcorticoid aldosterone. However, corticosteroids with high binding affinity that are not inactivated in MR-specific tissues, such

1 as 11-deoxycorticosterone, have been identified. It has therefore been suggested that these
 2 glucocorticoids control osmoregulatory functions associated with mineralcorticosteroids in
 3 higher vertebrates (Bury & Sturm 2007, Prunet et al. 2006, Sturm et al. 2005). Piscine
 4 corticosteroids are synthesised in the interrenal cells of the head kidney (Chester Jones &
 5 Mosley 1980) in response to various stress or environmental stimuli, and mediated via the
 6 hypothalamus-pituitary-interrenal (HPI) axis. Corticosteroids can have many diverse
 7 physiologic roles, such as, for instance, osmoregulatory adaptations in migrating euryhaline
 8 species or the redirection of energy to the central nervous system and suppression of immune
 9 function during a stress response (McCormick 2001, Barton, 2002, Charmandari et al., 2005).
 10 Thus, corticosteroid hormone-disrupting compounds could have severe impacts on adaptive
 11 responses with obvious implications for individual health and population development even
 12 though these effects may be subtle and difficult to detect.

13 Various anthropogenic compounds, such as PCBs, can suppress corticosteroid levels
 14 and/or attenuate the corticosteroid response (Pottinger, 2003; Quabius et al., 1997). However,
 15 an elevation of cortisol levels has also been reported in fishes following exposure to
 16 estrogenic compounds such as estradiol and 4-nonylphenol (Lerner et al., 2007; Pottinger,
 17 2003). While suppression of the corticosteroid response could impair the organism's ability to
 18 adapt to changing environments or stress, a chronic or frequent hyperactivation may more
 19 seriously lead to growth suppression, reproductive dysfunction, and immunosuppression
 20 (Pottinger, 2003). Similar to that reported for thyroid-disrupting compounds, many of the
 21 compounds interfering with corticosteroid regulation may act via an unspecific, complex
 22 interaction. These compounds may also interfere with the early steps of steroid synthesis and
 23 thus effect reproductive steroid hormones as well (Aluru and Vijayan, 2006). Finally, there
 24 are strong links between corticosteroids and reproductive hormones explaining the suppressed
 25 immune response during sexual maturation (Yada and Nakanishi, 2002) and by elevated
 26 levels of estradiol and 11-ketotestosterone (Kurtz et al., 2007; Pottinger, 2003).

Molecular markers could assist in both identifying mechanisms of action, and in determining the specificity of compounds interfering with corticosteroid regulation. The genes directly under control of the corticosteroid receptors are largely unknown, but early studies in mammals have revealed various genes with glucocorticoid receptor promoter elements that might be useful markers to study the effects of contaminants on glucocorticoid response also in fish as well (Phuc Le et al., 2005; Wang et al., 2004). An example of differential gene regulation in fish has already been reported, i.e. the upregulation of the glucocorticoid receptor and interleukin-1-beta by PCBs in rainbow trout anterior kidney cells (Quabius et al., 2005). Current knowledge on the stress response – and particularly its interference by environmental contaminants - of the species covered in this review is scarce. Cortisol has been shown to be involved in the adaptation to crowding stress, salinity and acidity in zebrafish and medaka (Ramsay et al., 2006; Sakamoto et al., 2001; Yada and Ito, 1999). Thus, the stress response in these model species is likely to resemble those of other fish species. With regards to their extensive use as laboratory fish (including endocrine disruption research) and the extensive knowledge on their molecular genetics, these species would also be ideally suited for a molecular-based model for screening the ability of environmental compounds to interfere with the glucocorticoid stress response.

6. Other hormones

Studies on endocrine disruption in fish have focused almost exclusively on the reproductive steroids, thyroidal hormones and glucocorticoids. However, these hormones regulate only a portion of the vital physiological processes under endocrine control. In principle, a number of different hormones and their corresponding physiological processes could be disrupted by environmental contaminants, provoking adverse effects. Potential targets could be those processes regulated by growth hormones, somatostatin, vasotocin, isotocin, melanin-concentrating hormones, endorphin, somatolactin, prolactin, atrial natriuretic peptide and

1 parathyroid hormones among others. In a few cases, potential target genes have been
2 identified in the zebrafish (Hoshijima and Hirose, 2007). However, it will be difficult to
3 include this array of potential endocrine targets into present testing schemes. Hence, model
4 species with extensive genetic characterisation could be used for a comprehensive approach
5 using modern techniques for screening whole genomes or proteoms. This type of screening
6 may indicate relevant hormone regulated processes and pathways which could be integrated
7 in targeted testing approaches.

10 **7. Conclusions**

11 A diverse array of important physiological functions are controlled by secreted hormones. As
12 these processes are susceptible to interference by EDCs, the identification of these chemicals
13 is highly relevant to the health and well-being of both wildlife and humans. Therefore, the
14 inclusion of endocrine interference in the regulation of chemical use has been demanded
15 (Lyons, 2006). In fish – which are directly exposed to discharged industrial, agricultural and
16 municipal effluents - endocrine disrupting effects have been reported for a diverse array of
17 anthropogenic chemicals which can mimic/interfere with reproductive steroids, thyroidal
18 hormones and glucocorticoids. Molecular biomarkers for EDC effects have to date only been
19 investigated for (anti)estrogenic and (anti)androgenic effects. For these EDCs, the existing
20 data in small model fish indicate a high correlation between the upregulation of endocrine
21 target genes and long-term adverse effects such as reproduction. This finding is not elusive,
22 since many compounds – modifying endogenous hormone levels via binding to nuclear
23 receptors – directly interfere with the transcription of target genes. The potential of molecular
24 screening approaches has already been recognised as a valuable tool in regulatory
25 applications, and has led to the development of fish screening assays for compounds
26 interfering with reproductive hormones (Hutchinson et al., 2006; OECD, 2006; USEPA,

2007) The strong correlation between xenoestrogen exposure and reproductive impairment allows for the establishment of simple but integrative screening assays for regulatory purposes. Such screening assays may be established additionally for other hormones and corresponding physiologic processes, once appropriate target genes/proteins have been identified. Thus, screening assays could contain a battery of molecular targets allowing a more comprehensive approach in the identification of endocrine disrupting compounds in fish and vertebrates in general.

Acknowledgement

We acknowledge Karl Fent (Fent et al., 2006), who inspired us to summarise effect levels graphically.

References

- Allen, Y.T., Katsiadaki, I., Pottinger, T.G., Jolly, C., Matthiessen, P., Mayer, I., Smith, A., Scott, A.P., Eccles, P., Sanders, M.B., Pulman, K.G.T. and Feist, S. 2008. Intercalibration exercise using a stickleback endocrine disrupter screening assay. *Environ. Toxicol. Chem.* **27**, 404-412.
- Alsop, D., Vijayan, M.M., 2008. Development of the corticosteroid stress axis and receptor expression in zebrafish. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* 294, R711-R719
- Aluru, N. and Vijayan, M.M., 2006. Aryl hydrocarbon receptor activation impairs cortisol response to stress in rainbow trout by disrupting the rate-limiting steps in steroidogenesis. *Endocrinology* 147, 1895-1903.

- 1 Andersen, L., Goto-Kazeto, R., Trant, J.M., Nash, J.P., Korsgaard, B. and Bjerregaard, P.,
2 2006. Short-term exposure to low concentrations of the synthetic androgen
3 methyltestosterone affects vitellogenin and steroid levels in adult male zebrafish (*Danio*
4 *rerio*). *Aquat. Toxicol.* 76, 343-352.
- 5 Andersson, C., Katsiadaki, I., Lundstedt-Enkel, K. and Örberg, J., 2007. Effects of 17 alpha-
6 ethynylestradiol on EROD activity, spiggin and vitellogenin in three-spined stickleback
7 (*Gasterosteus aculeatus*). *Aquat. Toxicol.* 83, 33-42.
- 8 Ankley, G.T., Jensen, K.M., Durhan, E.J., Makynen, E.A., Butterworth, B.C., Kahl, M.D.,
9 Villeneuve, D.L., Linnam, A., Gray, L.E., Cardon, M. and Wilson, V.S., 2005. Effects of
10 two fungicides with multiple modes of action on reproductive endocrine function in the
11 fathead minnow (*Pimephales promelas*). *Toxicol. Sci.* 86, 300-308.
- 12 Ankley, G.T., Jensen, K.M., Kahl, M.D., Korte, J.J. and Makynen, E.A., 2001. Description
13 and evaluation of a short-term reproduction test with the fathead minnow (*Pimephales*
14 *promelas*). *Environ. Toxicol. Chem.* 20, 1276-1290.
- 15 Ankley, G.T., Jensen, K.M., Kahl, M.D., Makynen, E.A., Blake, L.S., Greene, K.J., Johnson,
16 R.D. and Villeneuve, D.L., 2007. Ketoconazole in the fathead minnow (*Pimephales*
17 *promelas*): Reproductive toxicity and biological compensation. *Environ. Toxicol. Chem.* 26,
18 1214-1223.
- 19 Ankley, G.T., Jensen, K.M., Makynen, E.A., Kahl, M.D., Korte, J.J., Hornung, M.W., Henry,
20 T.R., Denny, J.S., Leino, R.L., Wilson, V.S., Cardon, M.C., Hartig, P.C. and Gray, L.E.,
21 2003. Effects of the androgenic growth promoter 17-beta-trenbolone on fecundity and
22 reproductive endocrinology of the fathead minnow. *Environ. Toxicol. Chem.*, 1350-1360.
- 23 Ankley, G.T. and Johnson, R.D., 2004. Small fish models for identifying and assessing the
24 effects of endocrine-disrupting chemicals. *ILAR Journal* 45, 469-483.
- 25 Ankley, G.T., Kahl, M.D., Jensen, K.M., Hornung, M.W., Korte, J.J., Makynen, E.A. and
26 Leino, R.L., 2002. Evaluation of the aromatase inhibitor fadrozole in a short-term

- 1 reproduction assay with the fathead minnow (*Pimephales promelas*). Toxicol. Sci. 67, 121-
2 130.
- 3 Ankley, G.T. and Villeneuve, D.L., 2006. The fathead minnow in aquatic toxicology: Past,
4 present and future. Aquat. Toxicol. 78, 91-102.
- 5 Barton, B.A., 2002. Stress in fishes: A diversity of responses with particular reference to
6 changes in circulating corticosteroids. Integr. Comp. Biol. 42, 517-525.
- 7 Bell, A.M., 2001. Effects of an endocrine disrupter on courtship and aggressive behaviour of
8 male three-spined stickleback, *Gasterosteus aculeatus*. Anim. Behav. 62, 775-780.
- 9 Bell, A.M., 2004. An endocrine disrupter increases growth and risky behavior in threespined
10 stickleback (*Gasterosteus aculeatus*). Horm. Behav. 45, 108-114.
- 11 Bernhardt, R.R., von Hippel, F.A. and Cresko, W.A., 2006. Perchlorate induces
12 hermaphroditism in threespine sticklebacks. Environ. Toxicol. Chem. 25, 2087-2096.
- 13 Biales, A.D., Bencic, D.C., Flick, R.W., Lazorchak, J. and Lattier, D.L., 2007a.
14 Quantification and associated variability of induced vitellogenin gene transcripts in fathead
15 minnow (*Pimephales promelas*) by quantitative real-time polymerase chain reaction assay.
16 Environ. Toxicol. Chem. 26, 287-296.
- 17 Biales, A.D., Bencic, D.C., Lazorchak, J.L. and Lattier, D.L., 2007b. A quantitative real-time
18 polymerase chain reaction method for the analysis of vitellogenin transcripts in model and
19 nonmodel fish species. Environ. Toxicol. Chem. 26, 2679-2686.
- 20 Bistodeau, T.J., Barber, L.B., Bartell, S.E., Cediell, R.A., Grove, K.J., Klaustermeier, J.,
21 Woodard, J.C., Lee, K.E. and Schoenfuss, H.L., 2006. Larval exposure to environmentally
22 relevant mixtures of alkylphenolethoxylates reduces reproductive competence in male
23 fathead minnows. Aquat. Toxicol. 79, 268-277.
- 24 Bogers, R., De Vries-Buitenweg, S., Van Gils, M., Baltussen, E., Hargreaves, A., van de
25 Waart, B., De Roode, D., Legler, J. and Murk, A., 2006a. Development of chronic tests for

- 1 endocrine active chemicals - Part 2: An extended fish early-life stage test with an androgenic
- 2 chemical in the fathead minnow (*Pimephales promelas*). *Aquat. Toxicol.* 80, 119-130.
- 3 Bogers, R., Mutsaerds, E., De Roode, D.F., Druke, J., Legler, J., Murk, A.J. and Van Der
- 4 Burg, B., 2006b. Estrogenic endpoints in fish early life-stage tests: Luciferase and
- 5 vitellogenin induction in estrogen-responsive transgenic zebrafish. *Environ. Toxicol. Chem.*
- 6 25, 241-247.
- 7 Brian, J.V., Augley, J. and Braithwaite, V.A., 2006. Endocrine disrupting effects on the
- 8 nesting behaviour of male three-spined stickleback *Gasterosteus aculeatus* L. *J. Fish Biol.*
- 9 68, 1883-1890.
- 10 Bringolf, R.B., Belden, J.B. and Summerfelt, R.C., 2004. Effects of atrazine on fathead
- 11 minnow in a short-term reproduction assay. *Environ. Toxicol. Chem.* 23, 1019-1025.
- 12 Brion, F., Tyler, C.R., Palazzi, X., Laillet, B., Porcher, J.M., Garric, J. and Flammarion, P.,
- 13 2004. Impacts of 17 beta-estradiol, including environmentally relevant concentrations, on
- 14 reproduction after exposure during embryo-larval-, juvenile- and adult-life stages in
- 15 zebrafish (*Danio rerio*). *Aquat. Toxicol.* 68, 193-217.
- 16 Brown, S.B., Adams, B.A., Cyr, D.G. and Eales, J.G., 2004. Contaminant effects on the
- 17 teleost fish thyroid. *Environ. Toxicol. Chem.* 23, 1680-1701.
- 18 Bury, N., Sturm, A., Le Rouzic, P., Lethimonier, C., Ducouret, B., Guiguen, Y., Robinson-
- 19 Rechavi, M., Laudet, V., Rafestin-Oblin, M., Prunet, P., 2003: Evidence for two distinct
- 20 functional glucocorticoid receptors in teleost fish. *J. Mol. Endocrinol.* 31, 141-156
- 21 Bury, N.R., Sturm, A., 2007: Evolution of the corticosteroid receptor signalling pathway in
- 22 fish. *Gen. Comp. Endocrinol.* 153, 47-56.
- 23 Charmandari, E., Tsigos, C. and Chrousos, G., 2005. Endocrinology of the stress response. In:
- 24 *Annu. Rev. Physiol.*, pp. 259-284.

- 1 Chester Jones, I., Mosley, W., 1980.: The interrenal gland in Pisces. In: Jones IC , Henderson
2 IW (Editors), General, Comparative and Clinical Endocrinology of the Adrenal Cortex.
3 Academic Press, London, New York, pp. 395-523.
- 4 Cody, R.P. and Bortone, S.A., 1997. Masculinization of mosquitofish as an indication of
5 exposure to kraft mill effluent. Bull. Environ. Contam. Toxicol. 58, 429-436.
- 6 Coldham, N.G., Dave, M., Sivapathasundaram, S., McDonnell, D.P., Connor, C. and Sauer,
7 M.J., 1997. Evaluation of a recombinant yeast cell estrogen screening assay. Environ. Health
8 Perspect. 105, 734-742.
- 9 Coverdale, L.E., Lean, D. and Martin, C.C., 2004. Not just a fishing trip - Environmental
10 genomics using zebrafish. Current Genomics 5, 395-407.
- 11 Crane, H.M., Pickford, D.B., Hutchinson, T.H. and Brown, J.A., 2004. Developmental
12 changes of thyroid hormones in the fathead minnow, *Pimephales promelas*. Gen. Comp.
13 Endocrinol. 139, 55-60.
- 14 Crane, H.M., Pickford, D.B., Hutchinson, T.H. and Brown, J.A., 2005. Effects of ammonium
15 perchlorate on thyroid function in developing fathead minnows, *Pimephales promelas*.
16 Environ. Health Perspect. 113, 396-401.
- 17 Cyr, D.G. and Eales, J.G., 1996. Interrelationships between thyroidal and reproductive
18 endocrine systems in fish. Rev. in Fish Biol. Fisheries 6, 165-200.
- 19 Danzo, B.J., 1997: Environmental xenobiotics may disrupt normal endocrine function by
20 interfering with the binding of physiological ligands to steroid receptors and binding proteins.
21 Environ. Health. Perspect. 105, 294-301.
- 22 Davey, J., Nomikos, A., Wungjiranirun, M., Sherman, J., Ingram, L., Batki, C., Lariviere, J.
23 and Hamilton, J., 2008. Arsenic as an endocrine disruptor: arsenic disrupts retinoic acid
24 receptor- and thyroid hormone receptor-mediated gene regulation and thyroid hormone-
25 mediated amphibian tail metamorphosis. Environ. Health Perspect. 116, 165-172.

- 1 Durhan, E.J., Lambright, C.S., Makynen, E.A., Lazorchak, J., Hartig, P.C., Wilson, V.S.,
2 Gray, L.E. and Ankley, G.T., 2006. Identification of metabolites of trenbolone acetate in
3 androgenic runoff from a beef feedlot. *Environ. Health Perspect.* 114, 65-68.
- 4 Eaton, R.C. and Farley, R.D., 1974. Spawning cycle and egg production of zebrafish,
5 *Brachydanio rerio*, in the laboratory. *Copeia* 1, 195-209.
- 6 Edwards, T.M., Moore, B.C. and Guillette, L.J., 2006. Reproductive dysgenesis in wildlife: a
7 comparative view. *Int. J. Androl.* 29, 109-121.
- 8 Elsalini, O.A. and Rohr, K.B., 2003. Phenylthiourea disrupts thyroid function in developing
9 zebrafish. *Dev. Genes Evol.* 212, 593-598.
- 10 Farchi-Pisanty, O., Sternberg, H. and Moav, B., 1997. Transcriptional regulation of fish
11 growth hormone gene. *Fish Physiol. Biochem.* 17, 237-246.
- 12 Fenske, M., van Aerle, R., Brack, S., Tyler, C.R. and Segner, H., 2001. Development and
13 validation of a homologous zebrafish (*Danio rerio* Hamilton-Buchanan) vitellogenin
14 enzyme-linked immunosorbent assay (ELISA) and its application for studies on estrogenic
15 chemicals. *Comp. Biochem. Physiol. C* 129, 217-232.
- 16 Fent, K., Weston, A.A. and Caminada, D., 2006. Ecotoxicology of human pharmaceuticals.
17 *Aquat. Toxicol.* 76, 122-159.
- 18 Filby, A.L., Thorpe, K.L., Maack, G. and Tyler, C.R., 2007. Gene expression profiles
19 revealing the mechanisms of anti-androgen- and estrogen-induced feminization in fish.
20 *Aquat. Toxicol.* 81, 219-231.
- 21 Filby, A.L., Thorpe, K.L. and Tyler, C.R., 2006. Multiple molecular effect pathways of an
22 environmental oestrogen in fish. *J. Mol. Endocrinol.* 37, 121-134.
- 23 Francis, R.C., 1992. Sexual lability in teleosts: developmental factors. *The Quarterly Review*
24 *of Biology* 67, 1-18.
- 25 Giesy, J.P., Pierens, S.L., Snyder, E.M., Miles-Richardson, S., Kramer, V.J., Snyder, S.A.,
26 Nichols, K.M. and Villeneuve, D.A., 2000. Effects of 4-nonylphenol on fecundity and

- 1 biomarkers of estrogenicity in fathead minnows (*Pimephales promelas*). Environ. Toxicol.
- 2 Chem. 19, 1368-1377.
- 3 Goldsmith, P., 2004. Zebrafish as a pharmacological tool: the how, why and when. Curr.
- 4 Opin. Pharmacol. 4, 504-512.
- 5 Gopinath, A., Tseng, L.A. and Whitlock, K.E., 2004. Temporal and spatial expression of
- 6 gonadotropin releasing hormone (GnRH) in the brain of developing zebrafish (*Danio rerio*).
- 7 Gene Expression Patterns 4, 65-70.
- 8 Gray, M.A. and Metcalfe, C.D., 1997. Induction of testis-ova in Japanese medaka (*Oryzias*
- 9 *latipes*) exposed to p-nonylphenol. Environ. Toxicol. Chem. 16, 1082-1086.
- 10 Gray, M.A., Niimi, A.J. and Metcalfe, C.D., 1999. Factors effecting the development of
- 11 testis-ova in medaka, *Oryzias latipes*, exposed to octylphenol. Environ. Toxicol. Chem. 18,
- 12 1835-1842.
- 13 Greer, M.A., Goodman, G., Pleus, R.C. and Greer, S.E., 2002. Health effects assessment for
- 14 environmental perchlorate contamination: the dose response for inhibition of thyroidal
- 15 radioiodine uptake in humans. Environ. Health Perspect. 110, 927-37.
- 16 Griffiths, R., Orr, K.L., Adam, A. and Barber, I., 2000. DNA sex identification in the three-
- 17 spined stickleback. J. Fish Biol. 57, 1331-1334.
- 18 Grist, E.P.M., Wells, N.C., Whitehouse, P., Brighty, G. and Crane, M., 2003. Estimating the
- 19 effects of 17 alpha-ethinylestradiol on populations of the fathead minnow *Pimephales*
- 20 *promelas*: Are conventional toxicological endpoints adequate? Environ. Sci. Technol. 37,
- 21 1609-1616.
- 22 Gronen, S., Denslow, N., Manning, S., Barnes, S., Barnes, D. and Brouwer, M., 1999. Serum
- 23 vitellogenin levels and reproductive impairment of male Japanese medaka (*Oryzias latipes*)
- 24 exposed to 4-tert-octylphenol. Environ. Health Perspect. 107, 385-390.

- 1 Hachisuka, K., Ii, S., Yamaguti, T., Tokumoto, M. and Tokumoto, T., 2005. In vivo analysis
2 of effect of endocrine-disrupting chemicals on induction of oocyte maturation in zebrafish.
3 *Zoolog. Sci.* 22, 1452-1452.
- 4 Hahlbeck, E., Griffiths, R. and Bengtsson, B.-E., 2004a. The juvenile three-spined stickleback
5 (*Gasterosteus aculeatus* L.) as a model organism for endocrine disruption: I. Sexual
6 differentiation. *Aquat. Toxicol.* 70, 287.
- 7 Hahlbeck, E., Katsiadaki, I., Mayer, I., Adolfsson-Erici, M., James, J. and Bengtsson, B.-E.,
8 2004b. The juvenile three-spined stickleback (*Gasterosteus aculeatus* L.) as a model
9 organism for endocrine disruption II--kidney hypertrophy, vitellogenin and spiggin
10 induction. *Aquat. Toxicol.* 70, 311.
- 11 Halm, S., Pounds, N., Maddix, S., Rand-Weaver, M., Sumpter, J.P., Hutchinson, T.H. and
12 Tyler, C.R., 2002. Exposure to exogenous 17 beta-oestradiol disrupts P450aromB mRNA
13 expression in the brain and gonad of adult fathead minnows (*Pimephales promelas*). *Aquat.*
14 *Toxicol.* 60, 285-299.
- 15 Harries, J.E., Runnalls, T., Hill, E., Harris, C.A., Maddix, S., Sumpter, J.P. and Tyler, C.R.,
16 2000. Development of a reproductive performance test for endocrine disrupting chemicals
17 using pair-breeding fathead minnows (*Pimephales promelas*). *Environ. Sci. Technol.* 34,
18 3003-3011.
- 19 Hawkins, M.B., Thomas, P., 2004. The Unusual Binding Properties of the Third Distinct
20 Teleost Estrogen Receptor Subtype ER{beta}a Are Accompanied by Highly Conserved
21 Amino Acid Changes in the Ligand Binding Domain. *Endocrinology* 145, 2968-2977.
- 22 Hellqvist, A., Bornestaf, C., Borg, B. and Schmitz, M., 2004. Cloning and sequencing of the
23 FSH-beta and LH beta-subunit in the three-spined stickleback, *Gasterosteus aculeatus*, and
24 effects of photoperiod and temperature on LH-beta and FSH-beta mRNA expression. *Gen.*
25 *Comp. Endocrinol.* 135, 167-174.

- 1 Hill, A.J., Teraoka, H., Heideman, W. and Peterson, R.E., 2005. Zebrafish as a model
2 vertebrate for investigating chemical toxicity. *Toxicol Sci* 86, 6-19.
- 3 Hill, J.R.L. and Janz, D.M., 2003. Developmental estrogenic exposure in zebrafish (*Danio*
4 *rerio*): I. Effects on sex ratio and breeding success. *Aquat. Toxicol.* 63, 417-429.
- 5 Hirai, N., Nanba, A., Koshio, M., Kondo, T., Morita, M. and Tatarazako, N., 2006.
6 Feminization of Japanese medaka (*Oryzias latipes*) exposed to 17[beta]-estradiol: Formation
7 of testis-ova and sex-transformation during early-ontogeny. *Aquat. Toxicol.* 77, 78-86.
- 8 Hoffmann, J.L. and Oris, J.T., 2006. Altered gene expression: A mechanism for reproductive
9 toxicity in zebrafish exposed to benzo[a]pyrene. *Aquat. Toxicol.* 78, 332-340.
- 10 Hossain MS, Larsson A, Scherbak N, Olsson P-E, Orban L (2007): Zebrafish Androgen
11 Receptor: Isolation, Molecular, and Biochemical Characterization.
12 10.1095/biolreprod.107.062018.
- 13 Hornung, M.W., Jensen, K.M., Korte, J.J., Kahl, M.D., Durhan, E.J., Denny, J.S., Henry, T.R.
14 and Ankley, G.T., 2004. Mechanistic basis for estrogenic effects in fathead minnow
15 (*Pimephales promelas*) following exposure to the androgen 17 alpha-methyltestosterone:
16 conversion of 17 alpha-methyltestosterone to 17 alpha-methylestradiol. *Aquat. Toxicol.* 66,
17 15-23.
- 18 Hoshijima, K. and Hirose, S., 2007. Expression of endocrine genes in zebrafish larvae in
19 response to environmental salinity. *J. Endocrinol.* 193, 481-491.
- 20 Howell, W.M., Black, D. and Bortone, S.A., 1980. Abnormal expression of secondary sex
21 characters in a population of mosquitofish, *Gambusia affinis holbrooki*: evidence for
22 environmentally-induced masculinization. *Copeia* 1980, 676-681.
- 23 Hutchinson, T.H., Ankley, G.T., Segner, H. and Tyler, C.R., 2006. Screening and testing for
24 endocrine disruption in fish-biomarkers as "signposts," not "traffic lights," in risk
25 assessment. *Environ. Health Perspect.* 114 Suppl 1, 106-14.

- 1 Inui, M., Adachi, T., Takenaka, S., Inui, H., Nakazawa, M., Ueda, M., Watanabe, H., Mori,
2 C., Iguchi, T. and Miyatake, K., 2003. Effect of UV screens and preservatives on
3 vitellogenin and choriogenin production in male medaka (*Oryzias latipes*). *Toxicology* 194,
4 43-50.
- 5 Islinger, M., Willimski, D., Volkl, A. and Braunbeck, T., 2003. Effects of 17 α -ethinylestradiol
6 on the expression of three estrogen-responsive genes and cellular ultrastructure of liver and
7 testes in male zebrafish. *Aquat. Toxicol.* 62, 85-103.
- 8 Islinger, M., Yuan, H., Voelkl, A. and Braunbeck, T., 2002. Measurement of vitellogenin
9 gene expression by RT-PCR as a tool to identify endocrine disruption in Japanese medaka
10 (*Oryzias latipes*). *Biomarkers* 7, 80-93.
- 11 Jafri, S.I.H. and Ensor, D.M., 1979. Occurrence of an intersex condition in the roach *Rutilus*
12 *rutilus* (L.). *J. Fish Biol.* 14, 547-549.
- 13 Jakobs, T.C., Schmutzler, C., Meissner, J. and Kohrle, J., 1997. The promoter of the human
14 type I 5'-deiodinase gene--mapping of the transcription start site and identification of a
15 DR+4 thyroid-hormone-responsive element. *Eur. J. Biochem.* 247, 288-97.
- 16 Jensen, K.M. and Ankley, G.T., 2006. Evaluation of a commercial kit for measuring
17 vitellogenin in the fathead minnow (*Pimephales promelas*). *Ecotoxicol. Environ. Saf.* 64,
18 101-105.
- 19 Jensen, K.M., Kahl, M.D., Makynen, E.A., Korte, J.J., Leino, R.L., Butterworth, B.C. and
20 Ankley, G.T., 2004. Characterization of responses to the antiandrogen flutamide in a short-
21 term reproduction assay with the fathead minnow. *Aquat. Toxicol.* 70, 99-110.
- 22 Jobling, S., Casey, D., Rodgers-Gray, T., Oehlmann, J., Schulte-Oehlmann, U., Pawlowski,
23 S., Baunbeck, T., Turner, A.P. and Tyler, C.R., 2003. Comparative responses of molluscs
24 and fish to environmental estrogens and an estrogenic effluent. *Aquat. Toxicol.* 65, 205-220.

- 1 Jobling, S., Reynolds, T., White, R., Parker, M.G. and Sumpter, J.P., 1995. A variety of
2 environmental persistent chemicals, including some phthalate plasticizers, are weakly
3 estrogenic. Environ. Health Perspect. 503, 582-587.
- 4 Jolly, C., 2007. The three-spined stickleback (*Gasterosteus aculeatus*) as a bioindicator
5 species for the identification of (anti-)androgenic compounds. Dissertation. University of
6 Bergen, Norway, Bergen.
- 7 Jolly, C., Katsiadaki, I., Le Belle, N., Mayer, I. and Dufour, S., 2006. Development of a
8 stickleback kidney cell culture assay for the screening of androgenic and anti-androgenic
9 endocrine disrupters. Aquat. Toxicol. 79, 158-166.
- 10 Jones, I., Lindberg, C., Jakobsson, S., Hellqvist, A., Hellman, U., Borg, B. and Olsson, P.-E.,
11 2001. Molecular Cloning and Characterization of Spiggin. An androgen-regulated
12 extraorganismal adhesive with structural similarities to Von Willebrand factor-related
13 proteins. J. Biol. Chem. 276, 17857-17863.
- 14 Jurcikova, J., Mikula, P., Dobsikova, R., Nemethova, D. and Svobodova, Z., 2007. Effects of
15 metazachlor on vitellogenin induction in zebrafish (*Danio rerio*). Acta Veterinaria Brno 76,
16 S61-S66.
- 17 Kang, I., Yokota, H., Oshima, Y., Tsuruda, Y., Yamaguchi, T., Maeda, M., Imada, N.,
18 Tadokoro, H. and Honjo, T., 2002. Effect of 17beta-estradiol on the reproduction of
19 Japanese medaka (*Oryzias latipes*). Chemosphere 47, 71-80.
- 20 Katsiadaki, I., Sanders, M., Sebire, M., Nagae, M., Soyano, K. and Scott, A.P. 2007. Three-
21 spined stickleback: an emerging model in environmental endocrine disruption.
22 Environmental Sciences 14, 263-283.
- 23 Katsiadaki, I., Morris, S., Squires, C., Hurst, M., James, J. and Scott, A., 2006. Use of the
24 three-spined stickleback (*Gasterosteus aculeatus*) as a sensitive in vivo test for detection of
25 environmental antiandrogens. Environ. Health Perspect. 114, 115-121.

- 1 Katsiadaki, I., Scott, A.P., Hurst, M.R., Matthiessen, P. and Mayer, I., 2002a. Detection of
2 environmental androgens: a novel method based on enzyme-linked immunosorbent assay of
3 spiggin, the stickleback (*Gasterosteus aculeatus*) glue protein. Environ. Toxicol. Chem. 21,
4 1946-1954.
- 5 Katsiadaki, I., Scott, A.P. and Mayer, I., 2002b. The potential of the three-spined stickleback
6 (*Gasterosteus aculeatus* L.) as a combined biomarker for oestrogens and androgens in
7 European waters. Mar. Environ. Res. 54, 725-728.
- 8 Katti, S.R. and Sathyanesan, A.G., 1986. Changes in the hypothalamoneurohypophyseal
9 complex of lead treated teleostean fish *Clarias batrachus* (L). Z. Mikrosk. Anat. Forsch.
10 100, 347-352.
- 11 Kausch, U., Alberti, M., Haindl, S., Budczies, J. and Hock, B., 2008. Biomarkers for exposure
12 to estrogenic compounds: Gene expression analysis in zebrafish (*Danio rerio*). Environ.
13 Toxicol. 23, 15-24.
- 14 Kavlock, R.J., Daston, G.P., DeRosa, C., Fenner-Crisp, P., Gray, L.E., Kaattari, S., Lucier,
15 G., Luster, M., Mac, M.J., Maczka, C., Miller, R., Moore, J., Rolland, R., Scott, G.,
16 Sheehan, D.M., Sinks, T., Tilson, H.A., 1996. Research needs for the risk assessment of
17 health and environmental effects of endocrine disruptors: a report of the U.S. EPA-
18 sponsored workshop. Environ. Health Perspect. 104 Suppl 4, 715-40.
- 19 Kawakami, Y., Shin, D.H., Kitano, T., Adachi, S., Yamauchi, K., Ohta, H., 2006.
20 Transactivation activity of thyroid hormone receptors in fish (*Conger myriaster*) in response
21 to thyroid hormones. Comparative Biochemistry and Physiology B-Biochemistry &
22 Molecular Biology 144, 503-509.
- 23 Kazeto, Y., Goto-Kazeto, R., Place, A.R. and Trant, J.M., 2003. Aromatase expression in
24 zebrafish and channel catfish brains: changes in transcript abundance associated with the
25 reproductive cycle and exposure to endocrine disrupting chemicals. Fish Physiol. Biochem.
26 28, 29-32.

- 1 Kazeto, Y., Place, A.R. and Trant, J.M., 2004. Effects of endocrine disrupting chemicals on
2 the expression of CYP19 genes in zebrafish (*Danio rerio*) juveniles. *Aquat. Toxicol.* 69, 25-
3 34.
- 4 Kidd, K.A., Blanchfield, P.J., Mills, K.H., Palace, V.P., Evans, R.E., Lazorchak, J.M. and
5 Flick, R.W., 2007. Collapse of a fish population after exposure to a synthetic estrogen. *Proc.*
6 *Natl. Acad. Sci. U. S. A.* 104, 8897-8901.
- 7 Kime, D.E., 1998. *Endocrine disruption in fish*, pp. 396, Kluwer Academic Publishers,
8 Dordrecht, NL.
- 9 Kinnberg, K., Holbech, H., Petersen, G.I. and Bjerregaard, P., 2007. Effects of the fungicide
10 prochloraz on the sexual development of zebrafish (*Danio rerio*). *Comp. Biochem. Physiol.*
11 *C* 145, 165-170.
- 12 Kiparissis, Y., Balch, G.C., Metcalfe, T.L. and Metcalfe, C.D., 2003a. Effects of the
13 isoflavones genistein and equol on the gonadal development of Japanese medaka *Oryzias*
14 *latipes*. *Environ. Health Perspect.* 111, 1158-63.
- 15 Kiparissis, Y., Metcalfe, T.L., Balch, G.C. and Metcalfe, C.D., 2003b. Effects of the
16 antiandrogens, vinclozolin and cyproterone acetate on gonadal development in the Japanese
17 medaka (*Oryzias latipes*). *Aquat. Toxicol.* 63, 391-403.
- 18 Knörr, S. and Braunbeck, T., 2002. Decline in reproductive success, sex reversal, and
19 developmental: alterations in Japanese medaka (*Oryzias latipes*) after continuous exposure
20 to octylphenol. *Ecotoxicol. Environ. Saf.* 51, 187-196.
- 21 Kuhl, A. and Brouwer, M., 2006. Antiestrogens inhibit xenoestrogen-induced brain aromatase
22 activity but do not prevent xenoestrogen-induced feminization in Japanese medaka (*Oryzias*
23 *latipes*). *Environ. Health Perspect.* 114, 500-506.
- 24 Kunz, P.Y. and Fent, K., 2006. Multiple hormonal activities of UV filters and comparison of
25 in vivo and in vitro estrogenic activity of ethyl-4-aminobenzoate in fish. *Aquat. Toxicol.* 79,
26 305-324.

- 1 Kunz, P.Y., Galicia, H.F. and Fent, K., 2006a. Comparison of in vitro and in vivo estrogenic
2 activity of UV filters in fish. *Toxicol. Sci.* 90, 349-361.
- 3 Kunz, P.Y., Gries, T. and Fent, K., 2006b. The ultraviolet filter 3-benzylidene camphor
4 adversely affects reproduction in fathead minnow (*Pimephales promelas*). *Toxicol. Sci.* 93,
5 311-321.
- 6 Kurtz, J., Kalbe, M., Langefors, A., Mayer, I., Milinski, M. and Hasselquist, D., 2007. An
7 experimental test of the immunocompetence handicap hypothesis in a teleost fish: 11-
8 ketotestosterone suppresses innate immunity in three-spined sticklebacks. *American*
9 *Naturalist* 170, 509-519.
- 10 Kwok, H.F., So, W.K., Wang, Y. and Ge, W., 2005. Zebrafish gonadotropins and their
11 receptors: I. Cloning and characterization of zebrafish follicle-stimulating hormone and
12 luteinizing hormone receptors--evidence for their distinct functions in follicle development.
13 *Biol Reprod* 72, 1370-81.
- 14 Kwon, J.Y., Haghpanah, V., Kogson-Hurtado, L.M., McAndrew, B.J. and Penman, D.J.,
15 2000. Masculinization of genetic female Nile tilapia (*Oreochromis niloticus*) by dietary
16 administration of an aromatase inhibitor during sexual differentiation. *J. Exp. Zool.* 287, 46-
17 53.
- 18 Lam, S.H., Sin, Y.M., Gong, Z. and Lam, T.J., 2005. Effects of thyroid hormone on the
19 development of immune system in zebrafish. *Gen. Comp. Endocrinol.* 142, 325-335.
- 20 Lange, I.G., Daxenberger, A., Schiffer, B., Witters, H., Ibarreta, D. and Meyer, H.H.D., 2002.
21 Sex hormones originating from different livestock production systems: fate and potential
22 disrupting activity in the environment. *Anal. Chim. Acta* 473, 27-37.
- 23 Lange, R., Hutchinson, T.H., Croudace, C.P., Siegmund, F., Schweinfurth, H., Hampe, P.,
24 Panter, G.H. and Sumpter, J.P., 2001. Effects of the synthetic estrogen 17 alpha-
25 ethinylestradiol on the life-cycle of the fathead minnow (*Pimephales promelas*). *Environ.*
26 *Toxicol. Chem.* 20, 1216-1227.

- 1 Larsson, D.G.J., Adolfson-Erici, M., Parkkonen, J., Pettersson, M., Berg, A.H., Olsson, P.-E.
2 and Förlin, L., 1999. Ethinylestradiol - an undesired fish contraceptive? *Aquat. Toxicol.* 45,
3 91-97.
- 4 Larsson, D.G.J., Hällman, H. and Förlin, L., 2000. More male fish embryos near a pulp mill.
5 *Environ. Toxicol. Chem.* 19, 2911-2917.
- 6 Lattier, D., Reddy, T., Gordon, D., Lazorchak, J.M., Smith, M., Williams, D., Wiechman, B.,
7 Flick, R., Miracle, A. and Toth, G., 2002. 17alpha-ethynylestradiol induced vitellogenin
8 gene transcription quantified in livers of adult males, larvae, and gills of fathead minnows
9 (*Pimephales promelas*). *Environ. Toxicol. Chem.* 21, 2385-2393.
- 10 Leon, A., Wu, P.-S., Hall, L.C., Johnson, M.L. and Teh, S.J., 2008. Global gene expression
11 profiling of androgen disruption in qurt strain medaka. *Environmental Science and*
12 *Technology* 42, 962-969.
- 13 Lerner, D.T., Bjornsson, B.T. and McCormick, S.D., 2007. Aqueous exposure to 4-
14 nonylphenol and 17 beta-estradiol increases stress sensitivity and disrupts ion regulatory
15 ability of juvenile Atlantic salmon. *Environ. Toxicol. Chem.* 26, 1433-1440.
- 16 Liu, F.J., Jia-Sheng, W. and Theodorakis, C.W., 2006. Thyrotoxicity of sodium arsenate,
17 sodium perchlorate, and their mixture in zebrafish *Danio rerio*. *Environ. Sci. Technol.* 40,
18 3429-3436.
- 19 Liu, Y.W. and Chan, W.K., 2002. Thyroid hormones are important for embryonic to larval
20 transitory phase in zebrafish. *Differentiation* 70, 36-45.
- 21 Lyons, G., 2006. Viewpoint: Policy requirements for protecting wildlife from endocrine
22 disruptors. *Environ. Health Perspect.* 114, 142-146.
- 23 Maack, G. and Segner, H., 2004. Life-stage-dependent sensitivity of zebrafish (*Danio rerio*)
24 to estrogen exposure. *Comp. Biochem. Physiol. C* 139, 47-55.

- 1 Majewski, A.R., Blanchfield, P.J., Palace, V.P. and Wautier, K., 2002. Waterborne 17alpha-
- 2 ethynylestradiol affects aggressive behaviour of male fathead minnows (*Pimephales*
- 3 *promelas*) under artificial spawning conditions. Water Qual. Res. J. Can. 37, 697-710.
- 4 Mak, P., Cruz, F.D. and Chen, S., 1999. A yeast screen system for aromatase inhibitors and
- 5 ligands for androgen receptor: yeast cells transformed with aromatase and androgen
- 6 receptor. Environ. Health Perspect. 107, 855-860.
- 7 Makynen, E.A., Kahl, M.D., Jensen, K.M., Tietge, J.E., Wells, K.L., Van der Kraak, G. and
- 8 Ankley, G.T., 2000. Effects of the mammalian antiandrogen vinclozolin on development
- 9 and reproduction of the fathead minnow (*Pimephales promelas*). Aquat. Toxicol. 48, 461-
- 10 475.
- 11 Marchand O, Safi R, Escriva H, Van Rompaey E, Prunet P, Laudet V, 2001: Molecular
- 12 cloning and characterization of thyroid hormone receptors in teleost fish. J. Mol. Endocrinol.
- 13 26, 51-65.
- 14 Martinovic, D., Blake, L.S., Durhan, E.J., Greene, K.J. and Kahl, M.D., 2008. Reproductive
- 15 toxicity of vinclozolin in the fathead minnow: Confirming an anti-androgenic mode of
- 16 action. Environ. Toxicol. Chem. 27, 478-488.
- 17 Martyniuk, C.J., Gerrie, E.R., Popescu, J.T., Ekker, M. and Trudeau, V.L., 2007. Microarray
- 18 analysis in the zebrafish (*Danio rerio*) liver and telencephalon after exposure to low
- 19 concentration of 17alpha-ethynylestradiol. Aquat. Toxicol. 84, 38-49.
- 20 Matsuda, M., Nagahama, Y., Shinomiya, A., Sato, T., Matsuda, C., Kobayashi, T., Morrey,
- 21 C.E., Shibata, N., Asakawa, S., Shimizu, N., Hori, H., Hamaguchi, S. and Sakaizumi, M.,
- 22 2002. DMY is a Y-specific DM-domain gene required for male development in the medaka
- 23 fish. Nature 417, 559-63.
- 24 Maunder, R.J., Matthiessen, P., Sumpter, J.P. and Pottinger, T.G., 2007. Impaired
- 25 reproduction in three-spined sticklebacks exposed to ethinyl estradiol as juveniles. Biol.
- 26 Reprod., biolreprod.107.062240.

- 1 McCormick SD, 2001. Endocrine Control of Osmoregulation in Teleost Fish. Am. Zool. 41,
2 781-794.
- 3 Menuet, A., Pellegrini, E., Anglade, I., Blaise, O., Laudet, V., Kah, O., Pakdel, F., 2002.
4 Molecular characterization of three estrogen receptor forms in zebrafish: binding
5 characteristics, transactivation properties, and tissue distributions. Biol Reprod 66, 1881-92.
- 6 Miles-Richardson, S.R., Kramer, V.J., Fitzgerald, S.D., Render, J.A., Yamini, B., Barbee, S.J.
7 and Giesy, J.P., 1999. Effects of waterborne exposure of 17 beta-estradiol on secondary sex
8 characteristics and gonads of fathead minnows (*Pimephales promelas*). Aquat. Toxicol. 47,
9 129-145.
- 10 Miracle, A., Ankley, G. and Lattier, D., 2006. Expression of two vitellogenin genes (vg1 and
11 vg3) in fathead minnow (*Pimephales promelas*) liver in response to exposure to steroidal
12 estrogens and androgens. Ecotoxicol. Environ. Saf. 63, 337-342.
- 13 Mukhi, S., Carr, J.A., Anderson, T.A. and Patino, R., 2005. Novel biomarkers of perchlorate
14 exposure in zebrafish. Environ. Toxicol. Chem. 24, 1107-1115.
- 15 Mukhi, S. and Patino, R., 2007. Effects of prolonged exposure to perchlorate on thyroid and
16 reproductive function in zebrafish. Toxicol. Sci. 96, 246-254.
- 17 Muncke, J., Junghans, M. and Eggen, R., 2007. Testing estrogenicity of known and novel
18 (xeno-)estrogens in the MolDarT using developing zebrafish *Danio rerio*. Environ. Toxicol.
19 22, 185-193.
- 20 Nagae, M., Kawasaki, F., Tanaka, Y., Ohkubo, N., Matsubara, T., Soyano, K., Hara, A.,
21 Arizono, K., Scott, A.P. and Katsiadaki, I., 2007. Detection and assessment of androgenic
22 potency of endocrine-disrupting chemicals using three-spined stickleback, *Gasterosteus*
23 *aculeatus*. Environmental Sciences . 14, 255-261.
- 24 Nagahama, Y., 1994. Endocrine regulation of gametogenesis in fish. Int J Dev Biol 38, 217-
25 29.

- 1 Nagahama, Y., Yoshikuni, M., Aamashita, M., Sakai, N. and Tanaka, N., 1993. Molecular
2 endocrinology of oocyte growth and maturation in fish. *Fish Physiol. Biochem.* 11, 3-14.
- 3 Nagel, R., 2002. DarT: The embryotest with the zebrafish *Danio rerio* - a general model in
4 ecotoxicology and toxicology. *ALTEX* 19 Suppl 1/02, 38-48.
- 5 Nanda, I., Kondo, M., Hornung, U., Asakawa, S., Winkler, C., Shimizu, A., Shan, Z., Haaf,
6 T., Shimizu, N., Shima, A., Schmid, M. and Scharf, M., 2002. A duplicated copy of
7 DMRT1 in the sex-determining region of the Y chromosome of the medaka, *Oryzias latipes*.
8 *Proc. Natl. Acad. Sci. U. S. A.* 22, 11778-11783.
- 9 Nash, J.P., Kime, D.E., Van der Ven, L.T.M., Wester, P.W., Brion, F., Maack, G.,
10 Stahlschmidt-Allner, P. and Tyler, C.R., 2004. Long-term exposure to environmental
11 concentrations of the pharmaceutical ethinylestradiol causes reproductive failure in fish.
12 *Environ. Health Perspect.* 112, 1725-1733.
- 13 Nichols, K.M., Snyder, E.M., Snyder, S.A., Pierens, S.L., Miles-Richardson, S.R. and Giesy,
14 J.P., 2001. Effects of nonylphenol ethoxylate exposure on reproductive output and
15 bioindicators of environmental estrogen exposure in fathead minnows, *Pimephales*
16 *promelas*. *Environ. Toxicol. Chem.* 20, 510-522.
- 17 Nimrod, A.C. and Benson, W.H., 1998. Reproduction and development of Japanese medaka
18 following an early life stage exposure to xenoestrogens. *Aquat. Toxicol.* 44, 141-156.
- 19 OECD, 2006. Report of the validation of the 21-day fish screening assay for the detection of
20 endocrine substances (phase 1b), series on testing and assessment, Number 61,
21 ENV/JM/MONO(2006)29.
- 22 Okubo, K., Nagata, S., Ko, R., Kataoka, H., Yoshiura, Y., Mitani, H., Kondo, M., Naruse, K.,
23 Shima, A. and Aida, K., 2001. Identification and characterization of two distinct GnRH
24 receptor subtypes in a teleost, the medaka *Oryzias latipes*. *Endocrinology* 142, 4729-4739.
- 25 Opitz, R., Braunbeck, T., Bögi, C., Pickford, D.B., Nentwig, G., Oehlmann, J., Tooi, O., Lutz,
26 I. and Kloas, W., 2005. Description and initial evaluation of a xenopus metamorphosis assay

- 1 for detection of thyroid system-disrupting activities of environmental compounds. Environ.
- 2 Toxicol. Chem. 24, 653-664.
- 3 Örn, S., Holbech, H., Madsen, T.H., Norrgren, L. and Petersen, G.I., 2003. Gonad
- 4 development and vitellogenin production in zebrafish (*Danio rerio*) exposed to
- 5 ethinylestradiol and methyltestosterone. Aquat. Toxicol. 65, 397-411.
- 6 Örn, S., Yamani, S. and Norrgren, L., 2006. Comparison of vitellogenin induction, sex ratio,
- 7 and gonad morphology between zebrafish and Japanese medaka after exposure to 17 α -
- 8 ethinylestradiol and 17 β -Trenbolone. Arch. Environ. Contam. Toxicol. 51, 237-243.
- 9 Palace, V.P., Evans, R.E., Wautier, K., Baron, C., Vandenbyllardt, L., Vandersteen, W. and
- 10 Kidd, K., 2002. Induction of vitellogenin and histological effects in wild fathead minnows
- 11 from a lake experimentally treated with the synthetic estrogen, ethinylestradiol. Water Qual.
- 12 Res. J. Can. 37, 637-650.
- 13 Panter, G.H., Hutchinson, T.H., Hurd, K.S., Bamforth, J., Stanley, R.D., Duffell, S.,
- 14 Hargreaves, A., Gimeno, S. and Tyler, C.R., 2006. Development of chronic tests for
- 15 endocrine active chemicals Part 1. An extended fish early-life stage test for oestrogenic
- 16 active chemicals in the fathead minnow (*Pimephales promelas*). Aquat Toxicol 77, 279-90.
- 17 Panter, G.H., Hutchinson, T.H., Hurd, K.S., Sherren, A., Stanley, R.D. and Tyler, C.R., 2004.
- 18 Successful detection of (anti-) androgenic and aromatase inhibitors in pre-spawning adult
- 19 fathead minnows (*Pimephales promelas*) using easily measured endpoints of sexual
- 20 development. Aquat. Toxicol. 70, 11-21.
- 21 Panter, G.H., Hutchinson, T.H., Lange, R., Lye, C.M., Sumpter, J.P., Zerulla, M. and Tyler,
- 22 C.R., 2002. Utility of a juvenile fathead minnow screening assay for detecting (anti-
- 23)estrogenic substances. Environ. Toxicol. Chem. 21, 319-326.
- 24 Park, E.A., Jerden, D.C. and Bahouth, S.W., 1995. Regulation of phosphoenolpyruvate
- 25 carboxykinase gene transcription by thyroid hormone involves two distinct binding sites in
- 26 the promoter. Biochem. J. 309 (Pt 3), 913-9.

- 1 Parng, C., Seng, W.L., Semino, C. and McGrath, P., 2002. Zebrafish: A preclinical model for
2 drug screening. *Assay Drug Dev. Technol.* 1, 41-48.
- 3 Parrott, J.L. and Blunt, B.R., 2005. Life-cycle exposure of fathead minnows (*Pimephales*
4 *promelas*) to an ethinylestradiol concentration below 1 ng/L reduces egg fertilization success
5 and demasculinizes males. *Environ. Toxicol.* 20, 131-141.
- 6 Parrott, J.L. and Wood, C.S., 2002. Fathead minnow lifecycle tests for detection of endocrine-
7 disrupting substances in effluents. *Water Qual. Res. J. Can.* 37, 651-667.
- 8 Patino, R., Waincott, M.R., Cruz-Li, E.I., Balakrishnan, S., McMurry, C., Blazer, V.S. and
9 Anderson, T.A., 2003. Effects of ammonium perchlorate on the reproductive performance
10 and thyroid follicle histology of zebrafish. *Environ. Toxicol. Chem.* 22, 1115-1121.
- 11 Pawlowski, S., Sauer, A., Shears, J.A., Tyler, C.R. and Braunbeck, T., 2004a. Androgenic and
12 estrogenic effects of the synthetic androgen 17 alpha-methyltestosterone on sexual
13 development and reproductive performance in the fathead minnow (*Pimephales promelas*)
14 determined using the gonadal recrudescence assay. *Aquat. Toxicol.* 68, 277-291.
- 15 Pawlowski, S., van Aerle, R., Tyler, C.R. and Braunbeck, T., 2004b. Effects of 17 alpha-
16 ethinylestradiol in a fathead minnow (*Pimephales promelas*) gonadal recrudescence assay.
17 *Ecotoxicol. Environ. Saf.* 57, 330-345.
- 18 Peichel, C.L., Ross, J.A., Matson, C.K., Dickson, M., Grimwood, J., Schmutz, J., Myers,
19 R.M., Mori, S., Schluter, D. and Kingsley, D.M., 2004. The master sex-determination locus
20 in threespine sticklebacks is on a nascent Y chromosome. *Curr. Biol.* 14, 1416-1424.
- 21 Peter, R.E., Yu, K.L., 1997. Neuroendocrine regulation of ovulation in fishes: basic and
22 applied aspects. *Rev. in Fish Biol. Fisheries* 7, 173-197.
- 23 Phuc Le, P., Friedman, J.R., Schug, J., Brestelli, J.E., Parker, J.B., Bochkis, I.M. and
24 Kaestner, K.H., 2005. Glucocorticoid receptor-dependent gene regulatory networks. *PLoS*
25 *Genetics* 1, e16 EP -.

- 1 Pottinger, T.G., 2003. Interactions of endocrine-disrupting chemicals with stress responses in
2 wildlife. *Pure Appl. Chem.* 75, 2321-2333.
- 3 Pottinger, T.G., Carrick, T.R. and Yeomans, W.E., 2002. The three-spined stickleback as an
4 environmental sentinel: effects of stressors on whole-body physiological indices. *J. Fish*
5 *Biol.* 61, 207-229.
- 6 Power, D.M., Llewellyn, L., Faustino, M., Nowell, M.A., Bjornsson, B.T., Einarsdottir, I.E.,
7 Canario, A.V.M. and Sweeney, G.E., 2001. Thyroid hormones in growth and development
8 of fish. *Comp. Biochem. Physiol. C* 130, 447-459.
- 9 Prunet, P., Sturm, A. and Milla, S., 2006. Multiple corticosteroid receptors in fish: From old
10 ideas to new concepts. *Gen. Comp. Endocrinol.* 147, 17-23.
- 11 Quabius, E., Krupp, G. and Secombes, C., 2005. Polychlorinated biphenyl 126 affects
12 expression of genes involved in stress-immune interaction in primary cultures of rainbow
13 trout anterior kidney cells. *Environ. Toxicol. Chem.* 24, 3053-60.
- 14 Quabius, E.S., Balm, P.H.M. and Wendelaar Bonga, S.E., 1997. Interrenal stress
15 responsiveness of tilapia (*Oreochromis mossambicus*) is impaired by dietary exposure to
16 PCB 126. *Gen. Comp. Endocrinol.* 108, 472-482.
- 17 Ramsay, J.M., Feist, G.W., Varga, Z.M., Westerfield, M., Kent, M.L. and Schreck, C.B.,
18 2006. Whole-body cortisol is an indicator of crowding stress in adult zebrafish, *Danio rerio*.
19 *Aquaculture* 258, 565-574.
- 20 Rose, J., Holbech, H., Lindholst, C., Norum, U., Povlsen, A., Korsgaard, B. and Bjerregaard,
21 P., 2002. Vitellogenin induction by 17 beta-estradiol and 17 alpha-ethinylestradiol in male
22 zebrafish (*Danio rerio*). *Comp. Biochem. Physiol. C* 131, 531-539.
- 23 Runnalls, T.J., Hala, D.N. and Sumpter, J.P., 2007. Preliminary studies into the effects of the
24 human pharmaceutical clofibric acid on sperm parameters in adult fathead minnow. *Aquat.*
25 *Toxicol.* 84, 111-118.

- 1 Sakamoto, T., Kozaka, T., Takahashi, A., Kawauchi, H. and Ando, M., 2001. Medaka
2 (*Oryzias latipes*) as a model for hypoosmoregulation of euryhaline fishes. *Aquaculture* 193,
3 347-354.
- 4 Sanchez, W., Goin, C., Brion, F., Olsson, P.E., Goksoyr, A. and Porcher, J.M., 2008. A new
5 ELISA for the three-spined stickleback (*Gasterosteus aculeatus* L.) spiggin, using
6 antibodies against synthetic peptide. *Comp. Biochem. Physiol. C* 147, 129-137.
- 7 Schaaf, M.J.M., Champagne, D., van Laanen, I.H.C., van Wijk, D., Meijer, A.H., Meijer,
8 O.C., Spalink, H.P. and Richardson, M.K., 2008. Discovery of a functional glucocorticoid
9 receptor beta-isoform in zebrafish. *Endocrinology* 149, 1591-1599.
- 10 Schafers, C., Teigeler, M., Wenzel, A., Maack, G., Fenske, M. and Segner, H., 2007.
11 Concentration- and time-dependent effects of the synthetic estrogen, 17 alpha-
12 ethinylestradiol, on reproductive capabilities of the zebrafish, *Danio rerio*. *J. Toxicol.*
13 *Environ. Health - Part A* 70, 768-779.
- 14 Schoenfuss, H.L., Bartell, S.E., Bistodeau, T.B., Cediel, R.A., Grove, K.J., Zintek, L., Lee,
15 K.E. and Barber, L.B., 2008. Impairment of the reproductive potential of male fathead
16 minnows by environmentally relevant exposures to 4-nonylphenol. *Aquat. Toxicol.* 86, 91-
17 98.
- 18 Scholz, S. and Gutzeit, H.O., 2000. 17-alpha-ethinylestradiol affects reproduction, sexual
19 differentiation and aromatase gene expression of the medaka (*Oryzias latipes*). *Aquat.*
20 *Toxicol.* 50, 363-373.
- 21 Scholz, S., Kordes, C., Hamann, J. and Gutzeit, H.O., 2004. Induction of vitellogenin in vivo
22 and in vitro in the model teleost medaka (*Oryzias latipes*): comparison of gene expression
23 and protein levels. *Mar. Environ. Res.* 57, 235-244.
- 24 Schulz, R.W., Bogerd, J., Male, R., Ball, J., Fenske, M., Olsen, L.C. and Tyler, C.R., 2007.
25 Estrogen-induced alterations in amh and dmrt1 expression signal for disruption in male
26 sexual development in the zebrafish. *Environ. Sci. Technol.* 41, 6305-10.

- 1 Schulz, R.W. and Goos, H.J.T., 1999. Puberty in male fish: concepts and recent developments
2 with special reference to the African catfish (*Clarias gariepinus*). *Aquaculture* 177, 5-12.
- 3 Segner, H., Navas, J.M., Schafers, C. and Wenzel, A., 2003. Potencies of estrogenic
4 compounds in in vitro screening assays and in life cycle tests with zebrafish in vivo.
5 *Ecotoxicol. Environ. Saf.* 54, 315-322.
- 6 Seki, M., Fujishima, S., Nozaka, T., Maeda, M. and Kobayashi, K., 2006. Comparison of
7 response to 17 beta-estradiol and 17 beta-trenbolone among three small fish species.
8 *Environ. Toxicol. Chem.* 25, 2742-2752.
- 9 Seki, M., Yokota, H., Maeda, M., Tadokoro, H. and Kobayashi, K., 2003. Effects of 4-
10 nonylphenol and 4-tert-octylphenol on sex differentiation and vitellogenin induction in
11 medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.* 22, 1507-1516.
- 12 Seki, M., Yokota, H., Matsubara, H., Tsuruda, Y., Maeda, N., Tadokoro, H. and Kobayashi,
13 K., 2002. Effect of ethinylestradiol on the reproduction and induction of vitellogenin and
14 testis-ova in medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.* 21, 1692-1698.
- 15 Singh, P.B., Kime, D.E., Epler, P. and Chyb, J., 1994. Impact of beta-hexachlorocyclohexane
16 exposure on plasma gonadotropin levels and in vitro stimulation of gonadal steroid
17 production by carp hypophyseal homogenate in *Carassius auratus*. In: *J. Fish Biol.*, pp. 195-
18 204.
- 19 Sperry, T.S. and Thomas, P., 1999. Characterization of Two Nuclear Androgen Receptors in
20 Atlantic Croaker: Comparison of Their Biochemical Properties and Binding Specificities.
21 *Endocrinology* 140, 1602-1611.
- 22 Sun, L.W., Zha, J.M., Spear, P.A. and Wang, Z.J., 2007a. Tamoxifen effects on the early life
23 stages and reproduction of Japanese medaka (*Oryzias latipes*). *Environmental Toxicology*
24 *and Pharmacology* 24, 23-29.

- 1 Sun, L.W., Zha, J.M., Spear, P.A. and Wang, Z.J., 2007b. Toxicity of the aromatase inhibitor
2 letrozole to Japanese medaka (*Oryzias latipes*) eggs, larvae and breeding adults. Comp.
3 Biochem. Physiol. C 145, 533-541.
- 4 Sohoni, P., Tyler, C.R., Hurd, K., Caunter, J., Hetheridge, M., Williams, T., Woods, C.,
5 Evans, M., Toy, R., Gargas, M. and Sumpter, J.P., 2001. Reproductive effects of long-term
6 exposure to bisphenol a in the fathead minnow (*Pimephales promelas*). Environ. Sci.
7 Technol. 35, 2917-2925.
- 8 Sturm, A., Bury, N., Dengreville, L., Fagart, J., Flouriot, G., Rafestin-Oblin, M.E. and Prunet,
9 P., 2005. 11-deoxycorticosterone is a potent agonist of the rainbow trout (*Oncorhynchus*
10 *mykiss*) mineralocorticoid receptor. Endocrinology 146, 47-55.
- 11 Tagawa, M. and Hirano, T., 1991. Effects of thyroid-hormone deficiency in eggs on early
12 development of the medaka, *Oryzias latipes*. J. Exp. Zool. 257, 360-366.
- 13 Takeo, J. and Yamashita, S., 2000. Rainbow trout androgen receptor-alpha fails to distinguish
14 between any of the natural androgens tested in transactivation assay, not just 11-
15 ketotestosterone and testosterone. General and Comparative Endocrinology 117, 200-206.
- 16 Thorpe, K.L., Benstead, R., Hutchinson, T.H. and Tyler, C.R., 2007. Associations between
17 altered vitellogenin concentrations and adverse health effects in fathead minnow
18 (*Pimephales promelas*). Aquat. Toxicol. 85, 176-183.
- 19 Tokumoto, T., Tokumoto, M., Horiguchi, R., Ishikawa, K. and Nagahama, Y., 2004.
20 Diethylstilbestrol induces fish oocyte maturation. Proc. Natl. Acad. Sci. U. S. A. 101, 3686-
21 3690.
- 22 Tokumoto, T., Tokumoto, M. and Thomas, P., 2007. Interactions of diethylstilbestrol (DES)
23 and DES analogs with membrane progestin receptor-alpha and the correlation with their
24 nongenomic progestin activities. Endocrinology 148, 3459-3467.

- 1 Tollefsen, K.E., 2002. Interaction of estrogen mimics, singly and in combination, with plasma
2 sex steroid-binding proteins in rainbow trout (*Oncorhynchus mykiss*). *Aquat. Toxicol.* 56,
3 215-25.
- 4 Tong, Y., Shan, T., Poh, Y.K., Yan, T., Wang, H., Lam, S.H. and Gong, Z.Y., 2004.
5 Molecular cloning of zebrafish and medaka vitellogenin genes and comparison of their
6 expression in response to 17 beta-estradiol. *Gene* 328, 25-36.
- 7 Tyler, C.R., Eerden, v.d., Jobling, S., Panter, G. and Sumpter, J.P., 1996. Measurement of
8 vitellogenin, a biomarker for exposure to oestrogenic chemicals, in a wide variety of
9 cyprinid fish. *Journal of Comparative Physiology B* 166, 418-426.
- 10 Tyler, C.R., Jobling, S. and Sumpter, J.P., 1998. Endocrine disruption in wildlife: a critical
11 review of the evidence. *Crit. Rev. Toxicol.* 28, 319-361.
- 12 Tyler, C.R., van Aerle, R., Hutchinson, T.H., Maddix, S. and Trip, H., 1999. An in vivo
13 testing system for endocrine disruptors in fish early life stages using induction of
14 vitellogenin. *Environ. Toxicol. Chem.* 18, 337-347.
- 15 USEPA, 2007. Validation of the fish short-term reproduction assay: integrated summary
16 report, U.S. Environmental Protection Agency, Endocrine Disruptor Screening Program,
17 Washington, D.C., December 15, 2007.
- 18 Van den Belt, K., Berckmans, P., Vangenechten, C., Verheyen, R. and Witters, H., 2004.
19 Comparative study on the in vitro/in vivo estrogenic potencies of 17beta-estradiol, estrone,
20 17alpha-ethynylestradiol and nonylphenol. *Aquat. Toxicol.* 66, 183-195.
- 21 Van den Belt, K., Verheyen, R. and Witters, H., 2001. Reproductive effects of
22 ethynylestradiol and 4t-octylphenol on the zebrafish (*Danio rerio*). *Arch. Environ. Contam.*
23 *Toxicol.* 41, 458-467.
- 24 Van den Belt, K., Verheyen, R. and Witters, H., 2003a. Comparison of vitellogenin responses
25 in zebrafish and rainbow trout following exposure to environmental estrogens. *Ecotoxicol.*
26 *Environ. Saf.* 56, 271-281.

- 1 Van den Belt, K., Verheyen, R. and Witters, H., 2003b. Effects of 17 alpha-ethynylestradiol
2 in a partial life-cycle test with zebrafish (*Danio rerio*): effects on growth, gonads and female
3 reproductive success. *Sci. Total Environ.* 309, 127-137.
- 4 Van den Belt, K., Wester, P.W., van der Ven, L.T., Verheyen, R. and Witters, H., 2002.
5 Effects of ethynylestradiol on the reproductive physiology in zebrafish (*Danio rerio*): time
6 dependency and reversibility. *Environ Toxicol Chem* 21, 767-75.
- 7 van der Ven, L.T.M., van den Brandhof, E.J., Vos, J.H., Power, D.M. and Wester, P.W.,
8 2006. Effects of the antithyroid agent propylthiouracil in a partial life cycle assay with
9 zebrafish. *Environ. Sci. Technol.* 40, 74-81.
- 10 Versonnen, B.J., Arijs, K., Verslycke, T., Lema, W. and Janssen, C.R., 2003. In vitro and in
11 vivo estrogenicity and toxicity of o-, m-, and p-dichlorobenzene. *Environ. Toxicol. Chem.*
12 22, 329-335.
- 13 Versonnen, B.J. and Janssen, C.R., 2004. Xenoestrogenic effects of ethynylestradiol in
14 zebrafish (*Danio rerio*). *Environ. Toxicol.* 19, 198-206.
- 15 Villeneuve, D.L., Blake, L.S., Brodin, J.D., Greene, K.J., Knoebl, I., Miracle, A.L.,
16 Martinovic, D. and Ankley, G.T., 2007a. Transcription of key genes regulating gonadal
17 steroidogenesis in control and ketoconazole- or vinclozolin-exposed fathead minnows.
18 *Toxicol. Sci.* 98, 395-407.
- 19 Villeneuve, D.L., Knoebl, I., Kahl, M.D., Jensen, K.M., Hammermeister, D.E., Greene, K.J.,
20 Blake, L.S. and Ankley, G.T., 2006. Relationship between brain and ovary aromatase
21 activity and isoform-specific aromatase mRNA expression in the fathead minnow
22 (*Pimephales promelas*). *Aquat. Toxicol.* 76, 353-368.
- 23 Villeneuve, D.L., Miracle, A.L., Jensen, K.M., Degitz, S.J., Kahl, M.D., Korte, J.J., Greene,
24 K.J., Blake, L.S., Linnum, A.L. and Ankley, G.T., 2007b. Development of quantitative real-
25 time PCR assays for fathead minnow (*Pimephales promelas*) gonadotropin beta subunit
26 mRNAs to support endocrine disruptor research. *Comp. Biochem. Physiol. C* 145, 171-183.

- 1 Vinggaard, A.M., Hass, U., Dalgaard, M., Andersen, H.R., Bonefeld-Jorgensen, E.,
2 Christiansen, S., Laier, P. and Poulsen, M.E., 2006. Prochloraz: an imidazole fungicide with
3 multiple mechanisms of action. In: Int. J. Androl., pp. 186-192.
- 4 Wang, J.-C., Derynck, M.K., Nonaka, D.F., Khodabakhsh, D.B., Haqq, C. and Yamamoto,
5 K.R., 2004. Chromatin immunoprecipitation (ChIP) scanning identifies primary
6 glucocorticoid receptor target genes. Proc. Natl. Acad. Sci. U. S. A. 101, 15603-15608.
- 7 Weber, L.P., Hill, J., Robert L. and Janz, D.M., 2003. Developmental estrogenic exposure in
8 zebrafish (*Danio rerio*): II. Histological evaluation of gametogenesis and organ toxicity.
9 Aquat. Toxicol. 63, 431-446.
- 10 Weisbrod, C.J., Kunz, P.Y., Zenker, A.K. and Fent, K., 2007. Effects of the UV filter
11 benzophenone-2 on reproduction in fish. Toxicol. Appl. Pharmacol. 225, 255-266.
- 12 Wester, P., van den Brandhof, E., Vos, J. and van der Ven, L., 2003. Identification of
13 endocrine disruptive effects in the aquatic environment a partial life cycle assay in zebrafish.
14 RIVM report 640920001/2003.
- 15 WHO, 2002. World Health Organization: Global Assessment of the State-of-the-Science of
16 Endocrine Disruptors. (Ed. T. Damstra, S. Barlow, A. Bergman, R. Kavlock and G. Van Der
17 Kraak). Report WHO/PCS/EDC/02.2, 180 p.
- 18 Williams, T.D., Caunter, J.E., Lillicrap, A.D., Hutchinson, T.H., Gillings, E.G. and Duffell,
19 S., 2007. Evaluation of the reproductive effects of tamoxifen citrate in partial and full life-
20 cycle studies using fathead minnows (*Pimephales promelas*). Environ. Toxicol. Chem. 26,
21 695-707.
- 22 Yada, T. and Ito, F., 1999. Sodium-retaining effects of cortisol, prolactin, and estradiol-17
23 beta in medaka *Oryzias latipes* exposed to acid water. Fish. Sci. 65, 405-409.
- 24 Yada, T. and Nakanishi, T., 2002. Interaction between endocrine and immune systems in fish.
25 In: International Review of Cytology - a Survey of Cell Biology, Vol 220, pp. 35-92.

- 1 Yamamoto, T.-O., 1975. Medaka (Killifish): Biology and strains, Keigaku Publishing
2 Company, Tokyo.
- 3 Yang, F.X., Xu, Y. and Hui, Y., 2006. Reproductive effects of prenatal exposure to
4 nonylphenol on zebrafish (*Danio rerio*). Comp. Biochem. Physiol. C 142, 77-84.
- 5 Yokota, H., Tsuruda, Y., Maeda, M., Oshima, Y., Tadokoro, H., Nakazono, A., Honjo, T. and
6 Kobayashi, K., 2000. Effect of bisphenol A on the early life stage in Japanese medaka
7 (*Oryzias latipes*). Environ. Toxicol. Chem. 19, 1925-1930.
- 8 Zerulla, M., Lange, R., Steger-Hartmann, T., Panter, G., Hutchinson, T. and Dietrich, D.R.,
9 2002. Morphological sex reversal upon short-term exposure to endocrine modulators in
10 juvenile fathead minnow (*Pimephales promelas*). Toxicol. Lett. 131, 51-63.
- 11 Zhang, Z.B. and Hu, J.Y., 2008. Effects of p,p '-DDE exposure on gonadal development and
12 gene expression in Japanese medaka (*Oryzias latipes*). Journal of Environmental Sciences-
13 China 20, 347-352.

Figure legends

Table 1: Brief description of the model organisms covered in this review

Table 2: Summary of lowest observed effect concentrations (mg/L) of thyroid disrupting compounds in zebrafish, fathead minnow and stickleback. Perchlorate is an inhibitor of thyroidal iodine uptake. Propylthiouracil, phenylthiourea, and thiourea inhibit thyroperoxidase and 5'-deiodinase. Methimazole is an inhibitor of thyroperoxidase. The mechanism of arsenite interference is not known but is discussed as a general interference with nuclear hormone receptors (Davey et al., 2008).

Fig. 1: Estrogenic and anti-estrogenic chemicals and their effect levels (lowest observed effect concentrations, LOEC) for vitellogenin/choriogenin expression, various molecular endpoints, secondary sex characters, histopathology and reproduction in males and or females of the small model species fathead minnow (A), zebrafish (B), medaka (C) and three-spined stickleback (D). The data were extracted from Allen et al., 2008; Andersen et al., 2004; Andersson et al., 2007a,b; Ankley et al., 2001; Ankley et al., 2002; Ankley et al., 2007; Bell, 2001; Bell, 2004; Biales et al., 2007a,b; Bistodeau et al., 2006; Bogers et al., 2006; Brian et al., 2006; Bringolf et al., 2004; Brion et al., 2004; Fenske et al., 2001; Filby et al., 2006; Giesy et al., 2000; Gray et al., 1997; Gray et al., 1999; Grist et al., 2003; Gronen et al., 1999; Hahlbeck et al., 2004a,b; Halm et al., 2002; Harries et al., 2000; Hill and Janz, 2003; Hirai et al., 2006; Hoffmann and Oris, 2006; Inui et al., 2003; Islinger et al., 2002; Jobling et al., 2003; Jolly, 2007; Katsiadaki, Pottinger and Mayer, unpublished data; Jurcikova et al., 2007; Kang et al., 2002; Kausch et al., 2008; Kazeto et al., 2003; Kazeto et al., 2004; Kidd et al., 2007; Kiparissis et al., 2003; Kishida et al., 2001; Knörr and Braunbeck, 2002; Kuhl and Brouwer, 2006; Kunz and Fent, 2006; Kunz et al., 2006a,b;

1 Lange et al., 2001; Lattier et al., 2002; Legler et al., 2002; Maack and Segner, 2004;
 2 Majewski et al., 2002; Martyniuk et al., 2007; Maunder et al., 2007; McAllister and Kime,
 3 2003; Mikula et al., 2006; Miles-Richardson et al., 1999; Miracle et al., 2006; Muncke et
 4 al., 2007; Nash et al., 2004; Nichols et al., 2001; Nimrod and Benson, 1998; Nimrod and
 5 Benson, 1998; OECD, 2006; Örn et al., 2003; Örn et al., 2006; Palace et al., 2002; Panter
 6 et al., 2002; Panter et al., 2004; Panter et al., 2006; Parrott and Blunt, 2005; Parrott and
 7 Wood, 2002; Pawlowski et al., 2004; Pottinger et al., 2002; Rose et al., 2002; Runnalls et
 8 al., 2007; Schafers et al., 2007; Schoenfuss et al., 2008; Scholz and Gutzeit, 2000; Scholz
 9 and Gutzeit, 2001; Scholz et al., 2004; Schulz et al., 2007; Segner et al., 2003; Seki et al.,
 10 2002; Seki et al., 2003; Seki et al., 2006; Sohoni et al., 2001; Sun et al., 2007a,b; Thorpe et
 11 al., 2007; Tong et al., 2004; Tyler et al., 1999; Van den Belt et al., 2001; Van den Belt et
 12 al., 2002; Van den Belt et al., 2003a,b; Van den Belt et al., 2004; van der Ven et al.,
 13 2006a,b; Versonnen and Janssen, 2004; Versonnen et al., 2003; Versonnen et al., 2004;
 14 Villeneuve et al., 2006; Villeneuve et al., 2007; Weber et al., 2003; Weisbrod et al., 2007;
 15 Wester et al., 2003; Williams et al., 2007; Yang et al., 2006; Yokota et al., 2000; Zerulla et
 16 al., 2002; Zhang et al., 2008. Detailed information on exposure conditions (stage, period) and
 17 description of effects can be found in the supplement available online (Tables S1). Criteria for
 18 the selection of references are outlined in the paragraph 2 (Data sources and graphical
 19 presentation). Please note different scale of the x-axis for medaka (C).
 20
 21

22 **Fig. 2:** Androgenic and anti-androgenic chemicals and their effect levels (lowest observed
 23 effect concentrations) for vitellogenin, spiggin, various molecular endpoints, secondary sex
 24 characters, histopathology and reproduction in males and or females of the small model
 25 species fathead minnow (A), zebrafish (B), medaka and three-spined stickleback (C). The data
 26 were extracted from Allen et al., 2008; Andersen et al., 2006; Ankley et al., 2001; Ankley et

1 al., 2003; Ankley et al., 2005; Bogers et al., 2006; Filby et al., 2007; Hahlbeck et al.,
2 2004a,b; Hornung et al., 2004; Jensen and Ankley, 2006; Jensen et al., 2004; Jolly, 2007;
3 Katsiadaki et al., 2002a,b; Kinnberg et al., 2007; Kiparissis et al., 2003; Makynen et al.,
4 2000; Martinovic et al., 2008; Miracle et al., 2006; OECD, 2006; Örn et al., 2003; Örn et
5 al., 2006; Panter et al., 2004; Parrott and Wood, 2002; Pawlowski et al., 2004; Sanchez et
6 al., 2008; Seki et al., 2006; Thorpe et al., 2007; Villeneuve et al., 2007; Wester et al., 2003;
7 Wester et al., 2003; Zerulla et al., 2002. Detailed information on exposure conditions (stage,
8 period) and description of effects can be found in the supplement available online (Tables
9 S1). Criteria for the selection of references are outlined in the paragraph 2 (Data sources and
10 graphical presentation).

11

12

Tables and figures

Table 1

The **fathead minnow** (*Pimephales promelas*) is a cyprinid fresh water fish with broad distribution across North America. It is used for a number of different regulatory tests for the risk assessment of chemicals. Appropriate testing guidelines for EDCs including reproductive endpoints have been established and genes involved in sexual development have been identified. With respect to endocrine disruption, secondary sex characteristics – androgen-dependent formation of tubercles – are of interests (Ankley and Villeneuve, 2006; Panter et al., 2004) (photo: Matthew Winter, AstraZeneca UK).

The **zebrafish** (*Danio rerio*), a small tropical fish native to the rivers of India and South Asia (Eaton and Farley, 1974), is one of the most popular model organisms in developmental genetics and (eco)toxicology (e.g. Coverdale et al., 2004; Goldsmith, 2004; Hill et al., 2005; Nagel, 2002; Parg et al., 2002). Its rapid development, easy maintenance in the laboratory and large number of offspring provides advantages for experimental studies. The extensive genetic characterisation and corresponding functional data are in particular useful for molecular studies in endocrine disruption (photo: Andre Künzelmann, UFZ).

The **medaka** (*Oryzias latipes*) native to Japan, Korea, China and Taiwan, is the only fish species of which the sex determination locus has been identified. Furthermore, sex specific markers (chromatophores, PCR-marker) and secondary sex characters (shape of the anal and dorsal fin) have lead to increasing attraction of the medaka as a model for endocrine disruption (Matsuda et al., 2002; Nanda et al., 2002; Yamamoto, 1975) (photo: Andre Künzelmann, UFZ).

The **stickleback** (*Gasterosteus aculeatus*) is unique among the small model fish due to the synthesis of an androgen-regulated protein called spiggin, a glue protein produced in the male kidney and used for nest building. Furthermore, the stickleback's quantifiable courtship behaviour (Bell, 2001) and male breeding colour (Maunder et al., 2007) have been used to analyse endocrine disruption. DNA markers for sex determination are available for the stickleback (Griffiths et al., 2000; Peichel et al., 2004). The stickleback is a euryhaline species and the only model indicator species that can be used for biomonitoring in marine and freshwaters in the Northern hemisphere (photo: Ian Mayer, UIB).

Table 2

Compound	Species	Hormone levels, molecular effects	Histology	Teratogenicity	Toxicity	Reproduction	Exposure period	Reference
Perchlorate	ZF		≤ 0.011 colloidal follicular structure, angiogenesis				Adults, 12 w + 12 w recovery	(Mukhi et al., 2005)
	ZF		≤ 6 Follicle structure, angiogenesis			≤ 6 PEV reduced	Adults, 8 w	(Patino et al., 2003)
	ZF	≤ 10 T4 reduced in females and embryos	≤ 10 thyroidal hypertrophy, jar malformation		≤ 10 reduced weight in female	≤ 10 PEV reduced, egg and larval offspring size increased	Adults, 10 w + 6 w recovery	(Mukhi and Patino, 2007)
	ZF	≤ 10 Collidal area, epithelial cell height					Adult, 90 d	(Liu et al., 2006)
	ZF	250 T4 reduced		250 (2500) Delayed hatch (retarded, less motile)	25000 Cell death		6 hpf – 6 dpf	(Elsalini and Rohr, 2003)
	FM		≤ 1 Thyroid hyperplasia	≤ 10 Retardation, reduced pigmentation			0-28 dph	(Crane et al., 2005)
	SB ^a		≤ 30 (100) Male nuptial coloration (intersexual gonads)		≤ 30 Reduced survival	≤ 30 Reduced spawning success and courtship behaviour	Adult, 350 d	(Bernhardt et al., 2006)
Propylthiouracil	ZF	10	≤ 1	100	100	10-100 ^b	Adult, 21 d	(van der Ven

		T3, T4 decreased	thyroid follicle synthetic activity	Body weight F1 generation	Reduced body weight	Increased egg number and egg size	Juvenile, 42 d	et al., 2006)
	ZF	100 T4 reduced		100 Retarded, curved body	1000 Reduced survival		6 hpf – 6 dpf	(Elsalini and Rohr, 2003)
Phenylthiourea	ZF	30 T4 reduced		30 (300) no pigmentation (retarded growth, curved body)	300 Reduced survival		6 hpf – 6 dpf	(Elsalini and Rohr, 2003)
Thiourea	ZF	≤ 300 T3, T4 decreased					0 hpf – 16 dpf	(Tagawa and Hirano, 1991)
Methimazole	ZF	117 T4 reduced		117 (234) Delayed hatch (retarded. less motile, curved body shape)	1170 Reduced survival		6 hpf – 6 dpf	(Elsalini and Rohr, 2003)
	ZF			34 mg/L Various morphometric parameters reduced	≤ 34,000 Reduced length		5 hpf-5dpf	(Liu and Chan, 2002)
	ZF	≤ 34 RagI, IgL-C-expression,			≤ 34 Reduced length		5-32 dpf	(Lam et al., 2005)
	FM	0.032 T3 (28 d inhibition, 56 induction)		0.032 Delayed hatching (not at 0.32 mg/L)			0-48 dpf	(Crane et al., 2004)
Arsenite	ZF		≤ 10 Colloid area and height (not in all concentrations and time points)				Adults, 90 d	(Liu et al., 2006)
Thyroxine (T4)	ZF	≤ 0.008		0.023	0.023		5 hpf-5dpf	(Liu and Chan,

Accepted Manuscript

	TR α , TR β gene expression increased		Deflation of swim bladder	Reduced length		2002)
ZF	≤ 23.3	≤ 23.3			5-32 dpf	(Lam et al., 2005)
	RagI-expression	Thymus structure				

^a Data on statistic significance not presented in the paper.

^b No LOEC calculated, p-value significant for regression analysis

C

D

A

B

C