Accepted Manuscript

Title: Identification of novel DNA fragments and partial sequence of a genomic island specific of *Brucella pinnipedialis*

Authors: Marianne Maquart, Yann Fardini, Michel S. Zygmunt, Axel Cloeckeaert

PII: S0378-1135(08)00150-8
DOI: doi:10.1016/j.vetmic.2008.04.015
Reference: VETMIC 4013

To appear in: *VETMIC*

Received date: 5-3-2008
Revised date: 21-3-2008
Accepted date: 10-4-2008

Please cite this article as: Maquart, M., Fardini, Y., Zygmunt, M.S., Cloeckeaert, A., Identification of novel DNA fragments and partial sequence of a genomic island specific of *Brucella pinnipedialis*, *Veterinary Microbiology* (2007), doi:10.1016/j.vetmic.2008.04.015

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Identification of novel DNA fragments and partial sequence of a genomic island specific
of *Brucella pinnipedialis***

Marianne Maquart , Yann Fardini, Michel S. Zygmunt , Axel Cloeckaert*

INRA, UR1282, Infectiologie Animale et Santé Publique, IASP, Nouzilly, F-37380, France

Corresponding author : Axel Cloeckaert, INRA, UR1282, Infectiologie Animale et Santé
Publique, IASP, Nouzilly, F-37380, France.

Tel.: +33 2 47 42 77 50; Fax: +33 2 47 42 77 74; e-mail: cloeckae@tours.inra.fr

REVISED MANUSCRIPT VETMIC-D-08-2354

29 **Abstract**

30 Since the 1990s, *Brucella* strains have been isolated from a wide variety of marine
31 mammals and were recently recognized as two different species i.e. *B. pinnipedialis* for
32 pinniped isolates and *B. ceti* for cetacean isolates. The aim of this study was to identify
33 specific DNA fragments of marine mammal *Brucella* strains using a previously described
34 Infrequent Restriction Site-PCR (IRS-PCR) method but with three new couples of restriction
35 enzymes applied on a larger panel of marine mammal *Brucella* isolates (n=74) and one human
36 isolate from New Zealand likely from marine mammal origin. This study revealed five DNA
37 fragments specific of *Brucella* strains isolated from marine mammals. Among them two new
38 DNA fragments were specific of *B. pinnipedialis* but were not detected in hooded seal
39 isolates. DNA fragment I identified in the previous IRS-PCR study and fragment VI of this
40 study were located on a cloned and sequenced 6 kb *SacI* fragment. Its nucleotide sequence
41 revealed that it is likely part of a putative genomic island. Sequence analysis showed that it
42 carries four ORFs coding for putative metabolic functions. Although hooded seal isolates are
43 classified within *B. pinnipedialis* it was shown in this study that they do not carry this
44 genomic island and this raises the question about their evolutionary history within *B.*
45 *pinnipedialis*.

1. Introduction

Brucellae are Gram-negative, facultatively, intracellular bacteria that can infect many species of animals and man. Six species were initially recognized within the genus *Brucella*: *B. abortus*, *B. melitensis*, *B. suis*, *B. ovis*, *B. canis*, and *B. neotomae* (Corbel et al., 1984; Moreno et al., 2002). This classification is mainly based on differences in pathogenicity, host preference, and phenotypic characteristics (Alton et al., 1998; Corbel et al., 1984; Moreno et al., 2002). Three additional species have been recently included in the genus *Brucella* i.e. *B. ceti* and *B. pinnipedialis* isolated from marine mammals, with cetaceans (dolphin, porpoise, and whale species) and pinnipeds (various seal species) as preferred host respectively, and *B. microti* isolated from the common vole (Foster et al., 2007; Scholz et al., 2008).

At the molecular level, evidence for two marine mammal *Brucella* species i.e. *B. ceti* and *B. pinnipedialis* has been firstly provided by study of DNA polymorphism at the *omp2* locus (Cloeckaert et al., 2001). This was further confirmed by an infrequent restriction site-PCR (IRS-PCR) method, taking into account the higher number of IS711 elements in the genome of marine mammal isolates compared to terrestrial mammal *Brucella* species (Bricker et al., 2000; Clavareau et al., 1998; Cloeckaert et al., 2003). IRS-PCR revealed four specific DNA fragments useful for the detection and identification of marine mammal *Brucella* isolates (Cloeckaert et al., 2003). Interestingly to date three human cases with *Brucella* infections presumably of marine mammal origin, according to specific molecular markers cited above, have been published and may point towards a zoonotic potential of these marine mammal *Brucella* species (Sohn et al., 2003; McDonald et al., 2006).

In the present study we used IRS-PCR with three new couples of restriction enzymes on a larger panel of marine mammal *Brucella* isolates (n=74) to identify novel DNA fragments specific of these marine mammal isolates.

79 2. Materials and methods

80 2.1. Bacterial strains

81 The marine mammal *Brucella* strains studied are listed in Table 1. Most isolates were
 82 from individual stranded dead animals over the period from 1994 to 2005. Isolates from
 83 hooded seals in Norway were from apparently healthy animals and caught in their natural
 84 habitat (Tryland et al., 2005). Terrestrial mammal *Brucella* strains used in this study were
 85 those of the previous IRS-PCR study (Cloeckeaert et al., 2003). Culture conditions of these
 86 strains were those described previously by Cloeckeaert et al. (2003).

88 2.2. IRS-PCR

89 IRS-PCR was performed as described previously by Cloeckeaert et al. (2003) with
 90 some modification, using the previous couple of restriction enzymes *Pst*I/*Sal*I and three new
 91 couples of restriction enzymes: *Bgl*II/*Sal*I, *Bam*HI/*Sal*I, *Eco*RI/*Sal*I.

92 One microgram of DNA was digested with 10 U of *Pst*I (or *Eco*RI, *Bgl*II, *Bam*HI)
 93 (Promega) and 10 U of *Sal*I (Promega) in 1x buffer H (Promega) for 2 h 30 min at 37 °C in a
 94 volume of 15 µl. Sterile distilled water, 2.5 U of T4 DNA ligase (Promega), 1x ligase buffer
 95 (Promega), the *Pst*I adapter (or *Bgl*II adapter, *Bam*HI adapter, *Eco*RI adapter) (20 pmol) and
 96 the *Sal*I adapter (20 pmol) were added for a total volume of 25 µl. The mixture was incubated
 97 at 16 °C for 2 h and then at 60 °C for 20 min to inactivate T4 DNA ligase. The sample was
 98 redigested with 5 U of *Pst*I (or *Bgl*II or *Bam*HI or *Eco*RI) and 5 U of *Sal*I at 37 °C for 30 min
 99 to cleave any restriction sites reformed by ligation, and then was submitted to amplification.
 100 Each PCR mixture included 2.5 µl of a 1/10 dilution of template DNA, 0.5 U of *GoTaq* DNA
 101 polymerase (Promega), deoxynucleoside triphosphates (200 µM each) (Promega), and the
 102 oligonucleotide primers in 1x PCR buffer (Promega). Typically, the oligonucleotides *Ps*1 (or

BG1) and either PsaIA, PsaIC, PsaIG, or PsaIT were used together as primers. Primer sequences are indicated in Table 2. Amplification was performed in an iCycler thermocycler (BioRad) with an amplification profile that consisted of an initial denaturation step at 94 °C for 5 min, five cycles of denaturation at 94 °C for 30 s, primer annealing at 55 °C for 1 min, and extension at 72 °C for 90 s, and then 30 cycles of denaturation at 94 °C for 30 s, primer annealing at 60 °C for 30 s, and extension at 72 °C for 90 s. All experiments included negative controls which were processed with the samples. The IRS-PCR reaction products were run on 2% (w/v) agarose gels containing 0.5 µg of ethidium bromide per ml.

2.3. Specific PCRs

Specific IRS-PCR fragments (V, VI) of the marine mammal *Brucella* isolates were gel-purified and their nucleotide sequence was determined to design specific primers. The primers used for specific PCRs of DNA fragments V and VI are listed in Table 2. Moreover, specific PCRs for DNA fragments I, II, III, and IV were performed on all the marine mammal strains of this study. PCR was performed on extracted DNAs as described previously. Briefly, amplification reaction mixtures were prepared in volumes of 100 µl containing 10 mM Tris-HCl (pH 9.0), 50 mM KCl, 1.5 mM MgCl₂, 0.1% Triton X-100 (1 x PCR buffer; Promega), a 200 µM concentration of each deoxynucleoside triphosphate, a 1 µM concentration of each primer, 100 ng of genomic DNA, and 5 U of *Taq* DNA polymerase (Promega). The temperature cycling for the amplification was performed in an iCycler thermocycler (BioRad) as follows: cycle 1 was 94 °C for 5 min (denaturation); the next 30 cycles were 62 °C for 30 s (annealing), 70 °C for 30 s (extension), and 94 °C for 30 s (denaturation); and the last cycle was 62 °C for 30 s (annealing) and 70 °C for 10 min (extension). The PCR products were run on 0.8% (w/v) agarose gels containing 0.5 µg of ethidium bromide per ml.

2.4. Cloning and partial nucleotide sequence of the putative *B. pinnipedialis* genomic island

PCR was performed with primers used to amplify fragment I as described previously (Cloekaert et al., 2003). The PCR product of 319 bp was purified using the QIAquick PCR purification Kit (QiagenTM). This probe was then labelled with the NEBlot-Phototope kit (BiolabsTM), following the manufacturer's protocol.

A *B. pinnipedialis* strain B2/94 (reference strain) genomic library was constructed in lambdaGEM-11 *Bam*HI half-site arms (Promega, Madison, Wis.) by following the instructions of the manufacturer. Briefly, *B. pinnipedialis* B2/94 DNA, extracted and purified as described previously by Cloekaert et al. (2003), was partially digested for 30 min at 37 °C with *Sau*3AI (Promega) at 0.025 U/μg of DNA, the enzyme concentration giving the highest percentage of fragments ranging from 15 to 23 kb. DNA fragments were ligated with T4 DNA ligase (Promega) to lambdaGEM-11 *Bam*HI half-site arms. Recombinant phage DNA was packaged in vitro with the Packagene System (Promega), and the library was titrated by determination of the number of PFU that appeared after infection of *E. coli* KW251 cells (Promega). Recombinant phages were transferred to nitrocellulose filters, and phages were screened by hybridization of the lysis plaques with the probe described above. DNA of a positive phage was extracted from culture supernatants of *E. coli* KW251 cells infected with the phage and cultured until lysis was observed. Phage DNA was then cut with *Sac*I, and restriction fragments were ligated into pGEM-7Zf+ (Promega) cut with *Sac*I. Competent *E. coli* JM109 cells (Promega) were transformed with recombinant plasmid DNA as described previously (Vizcaíno *et al.*, 1996), and bacteria were spread on Luria-Bertani (LB) broth-ampicillin (50 μg/ml) plates containing isopropyl-1-thio-β-D-galactopyranoside (IPTG) and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal). *E. coli* JM109 colonies bearing recombinant plasmids were transferred to nitrocellulose, lysed with 10% sodium dodecyl sulfate (SDS), and screened with the specific probe mentioned above. One positive colony was found bearing a plasmid with a *Sac*I insert of about 6 kb. This plasmid was named

pMZ4501. The *SacI* insert was sequenced by the GENOME expressTM society (Meylan, France).

2.5. Nucleotide accession numbers

The nucleotide sequence of the novel *B. pinnipedialis* DNA fragment V and the partial sequence of the *B. pinnipedialis* genomic island have been deposited in GenBank under accession numbers EU373600 and AY174073 respectively.

3. Results and discussion

3.1. Identification of specific DNA fragments of *B. pinnipedialis*

IRS-PCR was performed on 18 terrestrial mammal reference (species and biovar) strains and 74 marine mammal strains and one human isolate from New Zealand likely from marine mammal origin (McDonald et al., 2006). Three new couples of restriction enzymes were tested relative to the previous IRS-PCR study: *BglII/SalI*, *BamHI/SalI*, *EcoRI/SalI*. The screening of 50 new marine mammal isolates and the use of these different couples of restriction enzymes revealed five DNA fragments specific to *Brucella* isolated from marine mammals (Fig 1). Nucleotide sequencing of the fragment identified by the couple of restriction enzymes *SalI/EcoRI* with the primers PSalG and BG1 corresponded to fragment II with portion of IS711 identified in the previous IRS-PCR study (Fig. 1) (Cloeckaert et al., 2003). Furthermore, the fragment identified by the couple of restriction enzymes *SalI/BglII* with primers PSalA and BG1 corresponded to fragment III of this previous IRS-PCR study. The other fragment identified by the couple of restriction enzymes *SalI/BglII* with the primers PSalG and BG1, revealed that it corresponded to the *B. ceti* B1/94 *bp26* gene with portion of IS711 found downstream of *bp26* (AF242533) (Cloeckaert et al., 2000). Fragments II and III corresponded to part of genes encoding the *B. ceti* D-β-hydroxybutyrate dehydrogenase

(AY174070) and the *B. ceti* cytochrome B561 like protein (AY174071) respectively. Thus in the present IRS-PCR study, we confirmed two fragments previously identified, containing part of the IS711 element.

The two other DNA fragments, identified respectively by the restriction enzymes *SalI/BamHI* with primers PSaIG and BG1 and by the restriction enzymes *SalI/PstI* with primers PSaIC and PS1, of 305 bp and 780 bp were specific of *B. pinnipedialis* strains identified in this study and were named V and VI respectively. Nucleotide sequencing of fragment V revealed nucleotide and amino acid sequence identity of 96 % with an hypothetical protein from *Ochrobactrum anthropi* (GenBank accession number YP_001371022). Nucleotide sequencing of fragment VI revealed a deduced amino acid sequence with an amino acid sequence identity of 76 % to an aldehyde dehydrogenase from other *Brucella* species (GenBank accession number YP_001257642). Specific PCRs were performed with primers designed on the basis of nucleotide sequences of fragments V and VI. Like for the IRS-PCR results these DNA fragments were confirmed to be absent in all terrestrial mammal and cetacean strains studied (Table 1). Interestingly both fragments were shown to be present in all pinniped (seal and otter) isolates except in hooded seal isolates (Table 1). Nevertheless, hooded seal isolates belong likely also to *B. pinnipedialis* according to other characteristics like their oxidative metabolism profiles and their *omp2a* and *omp2b* gene characteristics (Jacques et al., 2007; Tryland et al., 2005). Fragments V and VI were also confirmed to be present in the human isolate from New Zealand.

3.2. Identification of a putative genomic island specific of *B. pinnipedialis*

A 5853 bp *SacI* fragment comprising the *B. pinnipedialis*-specific fragment I was cloned and sequenced. A search in databases using blastn revealed that this sequence was novel. Nucleotide sequencing revealed that its overall G+C content was lower i.e. 53 % than

the *Brucella* overall G+C content which is 57.3 % (Lavigne et al., 2005). The lower G+C content and the fact that this DNA fragment is absent in all other *Brucella* species including four *Brucella* species for which the genome sequences are available (*B. abortus*, *B. melitensis*, *B. suis*, and *B. ovis*) indicates that it is likely part of a genomic island. Four ORFs were identified in this fragment (Fig 1C). ORF1 (G+C content of 51.7 %) codes for a putative membrane protein with 35 % amino acid sequence identity to that of *Oceanicola batsensis* HTCC2597 (GenBank accession number EAQ04147.1), ORF2 (G+C content of 53 %) codes for a putative enoyl-CoA hydratase with 46 % amino acid sequence identity to that of *Burkholderia xenovorans* LB400 (GenBank accession number ABE36201.1), ORF3 (G+C content of 54.5 %) codes for a putative aldehyde dehydrogenase highly similar to that of other *Brucella* species and *Ochrobactrum anthropi* (73 % amino acid sequence identity; GenBank accession number AAX75972.1), and ORF4 (G+C content of 54 %) codes for a deoxycytidylate deaminase-related protein from *Stigmatella aurantiaca* (33 % amino acid sequence identity; GenBank accession number EAU69662). The genes were named *bpiM*, *bpiE*, *bpiA*, and *bpiD* respectively (Fig. 1C).

Among the two novel IRS-PCR DNA fragments identified for *B. pinnipedialis*, fragment VI was identified in the almost 6 kb nucleotide sequence as part of *bpiA*. Fragment V was not found within this sequence. However, the fact that strains positive for fragments I and VI are also positive for fragment V, and an identical size of fragments obtained by Southern blot of *Xba*I digested DNA with probes of fragment I and V (data not shown), suggest that fragment V could also be part of this genomic island.

Brucella genome sequencing have revealed the existence of several genomic islands ranging from 3.8 to 44.1 kb in size. Thus, nine genomic islands, called GI-1 to GI-9, have been identified to date in the *Brucella* sequenced genomes (Paulsen et al., 2002; Rajashekara et al., 2004; Vizcaino et al., 2004). All genomic islands display a different G+C content to the

overall G+C content of *Brucella* species. Most of them were likely acquired by horizontal transfer (Rajashekara et al., 2004). According to the nucleotide sequence the *B. pinnipedialis*-genomic island could be a metabolic island but further sequencing is required to identify the putative functions of this island and also to identify more specific features of genomic islands like presence of genes involved in integration, excision, and transfer. This will be reached with the genome sequencing of the *B. pinnipedialis* reference strain which is underway.

Although hooded seal isolates are considered as *B. pinnipedialis* strains, they do not carry this genomic island. Possibly acquisition of this genomic island in *B. pinnipedialis* may be posterior to separation between hooded seal isolates and other seal isolates and therefore hooded seal isolates could form an ancestor inside the *B. pinnipedialis* species. This hypothesis is currently being investigated in our laboratory by analyzing IS711 distribution and *omp2a* and *omp2b* gene diversity. Preliminary results indicate that hooded seal isolates carry less IS711 elements in their genomes than the *B. pinnipedialis* reference strain (unpublished results). There are also less *omp2a*-specific nucleotides at the 3' end of their *omp2b* genes compared to that of the *B. pinnipedialis* reference strain indicative of a lesser extent of *omp2b* gene conversion to *omp2a* (unpublished results). These preliminary data conforsts the hypothesis of hooded seal isolates being ancestors inside the *B. pinnipedialis* species.

Acknowledgements

We thank G. Foster, M. Tryland and W. McDonald for the gift of the 50 new *Brucella* strains of marine mammal origin.

References

- Alton, G.G., Jones, L.M., Angus, R.D., Verger, J.M., 1998. Techniques for the brucellosis laboratory, INRA, Paris, France.
- Bricker, B.J., Ewalt, D.R., MacMillan, A.P., Foster, G., Brew, S. 2000. Molecular characterization of *Brucella* strains isolated from marine mammals. J. Clin. Microbiol. 38, 1258-1262.
- Clavareau, C., Wellemans, V., Walravens, K., Tryland, M., Verger, J.M., Grayon, M., Cloeckaert, A., Letesson, J.J., Godfroid, J. 1998. Phenotypic and molecular characterization of a *Brucella* strain isolated from a minke whale (*Balaenoptera acutorostrata*). Microbiology 144, 3267-3273.
- Cloeckaert, A., Grayon, M., Grepinet, O. 2000. An IS711 element downstream of the *bp26* gene is a specific marker of *Brucella* spp. isolated from marine mammals. Clin. Diagn. Lab. Immunol. 7, 835-839.
- Cloeckaert, A., Verger, J.M., Grayon, M., Paquet, J.Y., Garin-Bastuji, B., Foster, G., Godfroid, J. 2001. Classification of *Brucella* spp. isolated from marine mammals by DNA polymorphism at the *omp2* locus. Microbes Infect. 3, 729-738.
- Cloeckaert, A., Grayon, M., Grépinet, O., Boumedine, K.S. 2003. Classification of *Brucella* strains isolated from marine mammals by infrequent restriction site-PCR and development of specific PCR identification tests. Microbes Infect. 5, 593-602.
- Corbel, M.J., Brinley-Morgan, W.J. 1984. Genus *Brucella* Meyer and Shaw 1920. Bergey's Manual of Systematic Bacteriology, Vol 1 (Krieg NR & Holt JG, eds.), pp. 377-388. Williams and Wilkins, Baltimore, MD.
- Foster, G., Osterman, B.S., Godfroid, J., Jacques, I., Cloeckaert, A. 2007. *Brucella ceti* sp.

- 275 nov. and *Brucella pinnipedialis* sp. nov. for *Brucella* strains with cetaceans and seals as their
276 preferred hosts. Int. J. Syst. Evol. Microbiol. 57, 2688-2693.
- 277 Jacques, I., Grayon, M., Verger, J.M. 2007. Oxidative metabolic profiles of *Brucella* strains
278 isolated from marine mammals: contribution to their species classification. FEMS Microbiol.
279 Lett. 270, 245-249.
- 280 Lavigne, J.P., Vergunst, A.C., Bourg, G., O'Callaghan, D. 2005. The IncP island in the
281 genome of *Brucella suis* 1330 was acquired by site-specific integration. Infect. Immun. 73,
282 7779-7783.
- 283 McDonald, W.L., Jamaludin, R., Mackereth, G., Hansen, M., Humphrey, S., Short, P., Taylor,
284 T., Swinger, J., Dawson, C.E., Whatmore, A.M., Stubberfield, E., Perrett, L.L., Simmons, G.
285 2006. Characterization of a *Brucella* sp. strain as a marine-mammal type despite isolation
286 from a patient with spinal osteomyelitis in New Zealand. J. Clin. Microbiol. 44, 4363-4370.
- 287 Moreno, E., Cloeckaert, A., Moriyón, I. 2002. *Brucella* evolution and taxonomy. Vet.
288 Microbiol. 90, 209-227.
- 289 Paulsen, I.T., Seshadri, R., Nelson, K.E., Eisen, J.A., Heidelberg, J.F., Read, T.D., Dodson,
290 R.J., Umayam, L., Brinkac, L.M., Beanan, M.J., Daugherty, S.C., Deboy, R.T., Durkin, A.S.,
291 Kolonay, J.F., Madupu, R., Nelson, W.C., Ayodeji, B., Kraul, M., Shetty, J., Malek, J., Van
292 Aken, S.E., Riedmuller, S., Tettelin, H., Gill, S.R., White, O., Salzberg, S.L., Hoover, D.L.,
293 Lindler, L.E., Halling, S.M., Boyle, S.M., Fraser, C.M. 2002. The *Brucella suis* genome
294 reveals fundamental similarities between animal and plant pathogens and symbionts. Proc.
295 Natl. Acad. Sci. USA 99, 13148-13153.

- 296 Rajashekara, G., Glasner, J.D., Glover, D.A., Splitter, G.A. 2004. Comparative whole-
 297 genome hybridization reveals genomic islands in *Brucella* species. J. Bacteriol. 186, 5040-
 298 5051.
- 299 Scholz, H.C., Hubalek, Z., Sedlacek, I., Vergnaud, G., Tomaso, H., Al Dahouk, S., Melzer, F.,
 300 Kampfer, P., Neubauer, H., Cloeckaert, A., Maquart, M., Zygmunt, M.S., Whatmore, A.M.,
 301 Falsen, E., Bahn, P., Gollner, C., Pfeffer, M., Huber, B., Busse, H.J., Nockler, K. 2008.
 302 *Brucella microti* sp. nov., isolated from the common vole *Microtus arvalis*. Int. J. Syst. Evol.
 303 Microbiol. 58, 375-382.
- 304 Sohn, A.H., Probert, W.S., Glaser C.A., Gupta N., Bollen A.W., Wong J.D., Grace, E.M.,
 305 McDonald, W.C. 2003. Human neurobrucellosis with intracerebral granuloma caused by a
 306 marine mammal *Brucella* spp. Emerg. Infect. Dis. 9, 485-488.
- 307 Tryland, M., Sørensen, K.K., Godfroid, J. 2005. Prevalence of *Brucella pinnipediae* in
 308 healthy hooded seals (*Cystophora cristata*) from the North Atlantic Ocean and ringed seals
 309 (*Phoca hispida*) from Svalbard. Vet. Microbiol. 105, 103-111.
- 310 Vizcaíno, N., Cloeckaert, A., Zygmunt, M.S., Dubray, G. 1996. Cloning, nucleotide sequence,
 311 and expression of the *Brucella melitensis omp31* gene coding for an immunogenic major
 312 outer membrane protein. Infect. Immun. 64, 3744-3751.
- 313 Vizcaíno, N., Caro-Hernández, P., Cloeckaert, A., Fernández-Lago, L. 2004. DNA
 314 polymorphism in the *omp25/omp31* family of *Brucella* spp.: identification of a 1.7-kb
 315 inversion in *Brucella cetaceae* and of a 15.1-kb genomic island, absent from *Brucella ovis*,
 316 related to the synthesis of smooth lipopolysaccharide. Microbes Infect 6, 821-834.

317

318

319 FIGURE LEGENDS

320 Fig 1. A. IRS-PCR patterns of *Brucella* strains identifying the five DNA fragments specific of
 321 marine mammal *Brucella* strains. Lane A, IRS-PCR pattern of *B. ceti* strain B1/94 obtained
 322 with restriction enzymes *SalI/EcoRI* and primers PsaG and BG1; lane B, IRS-PCR pattern of
 323 *B. ceti* strain B1/94 obtained with restriction enzymes *SalI/BglII* and primers PsaA and BG1;
 324 lane C, IRS-PCR pattern of *B. ceti* strain B1/94 obtained with restriction enzymes *SalI/BglII*
 325 and primers PsaG and BG1; lane D, IRS-PCR pattern of *B. pinnipedialis* strain B2/94
 326 obtained with restriction enzymes *SalI/PstI* and primers PsaC and PS1; lane E, IRS-PCR
 327 pattern of *B. pinnipedialis* strain B2/94 obtained with restriction enzymes *SalI/BamHI* and
 328 primers PsaG and BG1. Lane M: molecular mass marker. The specific DNA fragments 1, 2,
 329 3, 4 and 5 for which the nucleotide sequence has been determined are indicated by arrows.

330 B. Schematic representation of the nucleotide sequences of specific DNA fragments 1, 2, 3, 4
 331 and 5 from marine mammal *Brucella* isolates identified in this study with the new couples of
 332 restriction enzymes. Arrows indicate location of the primers used for specific PCRs II, III, V
 333 and VI (see also Table 2). BMEI0268 and BMEII1073 are annotations according to the
 334 sequenced genome of *B. melitensis* 16M and correspond respectively to D- β -hydroxybutyrate
 335 dehydrogenase and cytochrome B561 genes (GenBank accession numbers NC 003317 and
 336 NC 003318).

337 C. Schematic view and annotation of the 5853 bp partial sequence of the *B. pinnipedialis*-
 338 specific genomic island according to homology in databases. Plain arrow represents a gene
 339 coding for a putative membrane protein; Striped arrows represent genes coding for putative
 340 enzymatic proteins. BpiM: putative membrane protein; BpiE: putative enoyl-CoA hydratase;
 341 BpiA: putative aldehyde dehydrogenase; BpiD: deoxycytidylate deaminase related-protein.

Table 1. Marine mammal *Brucella* strains studied

Species	Strain	Host or source	Latin name	Geographic origin	Specific PCRs ^a					
					I	II	III	IV	V	VI
<i>B. pinnipedialis</i>	B2/94 ^b	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M2466/93/4	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M445/99/2	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M13/01/1	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M292/94/1	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M336/94/1	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M339/94/2	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M972/94/1	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M514/96/4	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M449/02/2	Common seal	<i>Phoca vitulina</i>	Scotland	+	-	-	-	+	+
	M621/99/2	Grey seal	<i>Halichoerus grypus</i>	Scotland	+	-	-	-	+	+
	M194/00/1	Grey seal	<i>Halichoerus grypus</i>	Scotland	+	-	-	-	+	+
	M2375/94/3	Grey seal	<i>Halichoerus grypus</i>	Scotland	+	-	-	-	+	+
	M89/03/05	Porpoise	<i>Phocoena phocoena</i>	Scotland	+	-	-	-	+	+
	M192/00/1	Minke whale	<i>Balaenoptera acutorostrata</i>	Scotland	+	-	-	-	+	+
	M1771/94/1	Otter	<i>Lutra lutra</i>	Scotland	+	-	-	-	+	+
	22a-2	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	23a-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	24a-2	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	25a-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	30a-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	17a-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	37a-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	38g-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	39a-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	53c-1	Hooded seal	<i>Cystophora cristata</i>	Norway	-	-	-	-	-	-
	M163/99/10	Hooded seal	<i>Cystophora cristata</i>	Scotland	-	-	-	-	-	-
	M603/99/7	Hooded seal	<i>Cystophora cristata</i>	Scotland	-	-	-	-	-	-
	02/611	Human	<i>Homo sapiens</i>	New Zealand	+	-	-	-	+	+
<i>B. ceti</i>	B1/94 ^b	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M1570/94/1	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M1661/94/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M39/94/1	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M854/98/8	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M1747/98/3	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M499/99/10	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M12/00/3	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M23/03/4	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M165/03/6	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M78/05/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M40/05/1	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M291/03/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M93/04/3	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M234/05/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M199/04/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M58/05/1	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M38/04/3	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M51/04/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M493/99/1	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M103/99/1	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
	M615/99/2	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-

M195/03/10	Porpoise	<i>Phocoena phocoena</i>	Scotland	-	+	+	-	-	-
7763/2	Bottlenose dolphin	<i>Tursiops truncatus</i>	France	-	+	+	-	-	-
M452/97/2	Common dolphin	<i>Delphinus delphis</i>	Scotland	-	+	+	-	-	-
M870/97/1	White-beaked dolphin	<i>Lagenorhynchus albirostris</i>	Scotland	-	+	+	-	-	-
M997/94/2	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
M2438/95/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
M2788/97/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
M181/97/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
M2/00/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
M187/00/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
M52/01/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	+	+	-	-	-
B202R	Minke whale	<i>Balaenoptera acutorostrata</i>	Norway	-	+	+	-	-	-
M490/95/1	Common seal	<i>Phoca vitulina</i>	Scotland	-	+	+	-	-	-
B14/94	Common dolphin	<i>Delphinus delphis</i>	Scotland	-	-	-	+	-	-
M644/93/1	Common dolphin	<i>Delphinus delphis</i>	Scotland	-	-	-	+	-	-
M9/02/01	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M13/05/1	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M40/95/1	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M2194/94/1	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M642/99/2	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M656/99/1	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M22/02/01	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M654/99/1	Striped dolphin	<i>Stenella coeruleoalba</i>	Scotland	-	-	-	+	-	-
M61/05/1	White-sided dolphin	<i>Lagenorhynchus acutus</i>	Scotland	-	-	-	+	-	-

^a PCRs using specific primers designed from the nucleotide sequences of IRS-PCR fragments I, II, III, IV (Cloeckaert et al., 2003), V and VI (this study) specific to marine mammal *Brucella* isolates (see Fig. 1).

^b Reference strain.

Table 2. Primers and adapters used in this study

Primers	Restriction enzyme	Adapter sequence (5' to 3')	Oligonucleotide sequence (5' to 3')	Annealing temp (°C)
Primers and adapters for IRS-PCR				
<i>SalI</i>		Sal1: PO ₄ -TCGATACTGGCAGACTCT AX2: GCCAGTA	Pas1A: AGAGTCTGCCAGTATCGACA	55
			PsalT: AGAGTCTGCCAGTATCGACT	55
			PsalC: AGAGTCTGCCAGTATCGACC	55
			PsalG: AGAGTCTGCCAGTATCGACG	55
<i>PstI</i>		PS1: GACTCGACTCGCATGCA AH2: TGCAGT	PS1: GACTCGACTCGCATGCA	55
<i>EcoRI</i>		BG1: GACTCGACTCGCA ER2: AATTTGCGAGT	BG1: GACTCGACTCGCA	55
<i>BglII/BamHI*</i>		BG1: GACTCGACTCGCA BG2: GATCTGCGAGT	BG1: GACTCGACTCGCA	55
Primers for specific PCRs				
Fragment I			I1: AGGTACTTCATTTCGTTCCGG	58
			I2: GCGACTCAGATGATCCCACC	58
Fragment II			II1: GCGGCTGGACGGCAGAATAA	58
			II2: AATGGACAGCGGTTTCATGCC	58
Fragment III			III1: GCAAATCCCAGTTGGAAATG	58
			III2: TGCTCATCTGTAAGGCTTCG	58
Fragment IV			IV1: GCTGGAGGATTTCTTCTTG	58
			IV2: CGGAAGATATGCTTCGATCC	58
Fragment V			V1: AGGGGCTGGATGGCTTTGTT	58
			V2: CGGATAGAGAAGATGCGGTC	58
Fragment VI			VI1: TCCAGCTCTTGAAGGAGATC	58
			VI2: CAGATCCGGTTTGGACATAA	58

* Adapter and primer sequences are the same for both enzymes *BglII* and *BamHI*.

Fig. 1

A.

B.

1(II)

2 (III)

3 (*bp26*-IS711)

4 (V)

5 (VI)

C.