Role of coagulase-negative staphylococci in human disease

A. Piette, G. Verschraegen

► To cite this version:

A. Piette, G. Verschraegen. Role of coagulase-negative staphylococci in human disease. *Veterinary Microbiology*, 2009, 134 (1-2), pp.45. 10.1016/j.vetmic.2008.09.009 . hal-00532477

HAL Id: hal-00532477

<https://hal.science/hal-00532477>

Submitted on 4 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Role of coagulase-negative staphylococci in human disease

Authors: A. Piette, G. Verschraegen

PII: S0378-1135(08)00363-5

DOI: doi:10.1016/j.vetmic.2008.09.009

Reference: VETMIC 4148

To appear in: *VETMIC*

Please cite this article as: Piette, A., Verschraegen, G., Role of coagulase-negative staphylococci in human disease, *Veterinary Microbiology* (2008), doi:10.1016/j.vetmic.2008.09.009

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Role of coagulase-negative staphylococci in human disease.

A. Piette*, G. Verschraegen

Department of Microbiology, University Hospital Ghent, Belgium

De Pintelaan 185, 9000 Gent

* Corresponding author. Tel.: +32 9 3326639; Fax: +32 9 3324985

E-mail address: anne.piette@ugent.be

9 Abstract

10
11 Coagulase-negative staphylococci (CNS) are normal inhabitants of human skin and mucous
12 membranes. They have long been dismissed as culture contaminants, but now the potentially
13 important role of CNS as pathogens and their increasing incidence has been recognized.
14 Approximately 55%-75% of nosocomial isolates is methicillin resistant. CNS were the first
15 organisms in which glycopeptide resistance was recognized. In the immunocompetent host, CNS
16 endocarditis and urinary tract infections with *Staphylococcus saprophyticus* are the most common
17 CNS infections. Other patients are usually immunocompromised, with indwelling or implanted
18 foreign bodies. CNS account for approximately 30% of all nosocomial blood stream infections. The
19 majority of these concern catheter-related sepsis. Other important infections due to CNS include
20 central nervous system shunt infections, endophthalmitis, surgical site infections, peritonitis in
21 patients with continuous ambulatory peritoneal dialysis and foreign body infections. CNS are rarely
22 associated with mastitis in humans. *Staphylococcus lugdunensis* is more pathogenic than other CNS
23 as it expresses several potential virulence factors. The distinction between clinically significant,
24 pathogenic and contaminating isolates is a major problem. Several studies show clonal intra and inter
25 hospital spread of *Staphylococcus epidermidis* strains which suggests that infection control measures
26 may be necessary for multiresistant CNS isolates as for methicillin resistant *Staphylococcus aureus*.
27 As a result of medical progress, mainly due to the use of invasive and indwelling medical devices,
28 CNS are now a major cause of nosocomial and health-care related infections.

29 **Keywords:** coagulase-negative staphylococci, human medicine, species distribution, antibiotic
30 resistance, endocarditis, urinary tract infections, blood stream infections, nosocomial infection
31

1. Introduction

Coagulase-negative staphylococci (CNS) are normal inhabitants of human skin and mucous membranes. They were first described in 1884 by Rodenbach as *Staphylococcus albus*, an avirulent staphylococcus. They have long been dismissed as culture contaminants, even in type 1 samples (samples obtained from a normally sterile site by needle aspiration or surgery). Only in 1958 the first report on the potential pathogenicity of CNS in patients with septicaemia was published (Smith et al., 1958). Later on, casuistic reports of CNS endocarditis, wound and urinary tract infections (UTI) appeared (Kloos et al., 1994). Since the 1970s, CNS are recognised as etiologic agents of a wide variety of infections. Patients with CNS infections are usually immunocompromised, with indwelling or implanted foreign bodies. CNS play a role in bacteraemia, central nervous system shunt infection, endocarditis, urinary tract infection, surgical site infections, endophthalmitis, foreign body infection and many other infections. The distinction between clinically significant, pathogenic and contaminating isolates is difficult and remains a major problem.

An overview of the medical literature on CNS is given, with special focus on identification, species distribution, virulence factors, antibiotic resistance, epidemiology and specific infections caused by CNS in humans.

2. Identification

Before 1975, *S. albus* or *S. epidermidis* was distinguished from *S. aureus* by the inability to clot blood plasma. Today, tests based on coagulase production and the thermonuclease reaction are used for rapid differentiation of *S. aureus* from other staphylococcal species (Huebner et al., 1999). Currently, 39 species of CNS are recognized (Euzéby, 2007). Sixteen of these species have been found in specimens of human origin. They are grouped in novobiocin-resistant species (*S. cohnii*, *S. saprophyticus*, *S. sciuri*, *S. xylois*) and novobiocin-susceptible species (*S. auricularis*, *S. capitis*, *S.*

57 *caprae*, *S. epidermidis*, *S. haemolyticus*, *S. hominis*, *S. lugdunensis*, *S. pasteurii*, *S. saccharolyticus*, *S.*
 58 *schleiferi*, *S. simulans*, *S. warneri*) (von Eiff et al., 2001).

59 Accurate identification of CNS isolates to species level is difficult to perform and expensive.
 60 According to Huebner et al., 1999, identification to species level is, in general, not necessary for
 61 good patient management. However, recent literature indicates accurate species identification is
 62 necessary to provide a better understanding of pathogenic potential of various CNS species (Heikens
 63 et al., 2005; Sivadon et al., 2005).

64 Before 1975, CNS were classified in different biotypes, according to Baird-Parker. In 1975,
 65 Kloos and Schleifer proposed a scheme for identification of staphylococci in the routine laboratory.
 66 This scheme is based on 13 key characteristics, such as coagulase activity, hemolysis, nitrate
 67 reduction and aerobic acid production from carbohydrates (Kloos et al., 1975). Nowadays, besides
 68 phenotypic methods, a variety of genotypic methods (16S rRNA, *sodA*, *gap*, *rpoB*, *tuf* gene
 69 sequencing) have been developed (Heikens et al., 2005; Layer et al., 2006; Mellmann et al., 2006).
 70 Heikens et al. compared phenotypic with genotypic (16S rRNA gene and *tuf* gene sequencing)
 71 identification (Heikens et al., 2005). The results showed that *tuf* gene sequencing is the best
 72 identification method (5 of 57 isolates misidentified). The API staph ID test (BioMérieux, Marcy
 73 l'Etoile, France) appeared to be a reliable alternative (7 of 57 isolates misidentified). The
 74 performance of the BD Phoenix system (Becton Dickinson Diagnostic Systems, Sparks, MD, USA)
 75 was poor. A recent study (Layer et al., 2006) compared 3 phenotypic, commercial identification
 76 methods with a molecular identification method (*gap* based terminal restriction fragment length
 77 polymorphism analysis) as reference method. The BD Phoenix ID-13 system (Becton Dickinson
 78 Diagnostic Systems) identified 18 of 27 reference strains and 70 of 86 clinical isolates correctly, the
 79 VITEK 2 ID-GP system (BioMérieux) 20 of 27 reference strains and 80 of 86 clinical isolates, the ID
 80 32 STAPH system (BioMérieux) 23 of 27 reference strains and 19 of 20 clinical isolates. Thus,
 81 genotypic methods are superior to phenotypic methods, but some phenotypic methods have a highly

acceptable level of identification accuracy when used in routine practice (Heikens et al., 2005; Layer et al., 2006).

3. Species distribution

Interpretation and comparison of the species distribution found in different studies is very difficult. The type of identification method used, may influence the species distribution found and some *Staphylococcus* species demonstrate habitat or niche preference. *S. capitis* for example is mainly found on the adult head, *S. cohnii* on the feet. *S. saprophyticus* is found in urine of young women (Kloos et al., 1994; Bannerman, 2003).

S. epidermidis, *S. haemolyticus* and *S. hominis* are the most frequently encountered CNS species in clinical samples. The isolation frequency of the different CNS species in clinical, human specimens is shown in Table 1. Usually, more than 50% of CNS isolates belong to the species *S. epidermidis*. In India, low percentages of *S. epidermidis* were found (13.2%) and high percentages of *S. haemolyticus* (71.9%) (Chaudhury et al., 2007). A Japanese study (Kawamura et al., 1998) found a very high percentage of *S. caprae* (14%), while the frequency of isolation in other studies is very low. In the study by Kawamura et al., 1998, all isolates were identified by a DNA-DNA hybridisation and by conventional, phenotypic identification based on the Kloos and Schleifer method. None of the *S. caprae* isolates was identified correctly by the phenotypic method.

In Spanish point prevalence studies in a large group of hospitals, performed from 1986 to 2002, showed no major shifts in the distribution of CNS species (Cuevas et al., 2004). When the distribution in 2002 is compared with 1986, there is a decrease in isolation of *S. epidermidis*, *S. simulans*, *S. cohnii* and *S. xylois* and an increase in isolation of *S. hominis* and *S. saprophyticus*. *S. epidermidis* is the predominant pathogen in intravascular catheter-related infections, nosocomial bacteremia, endocarditis, urinary tract and surgical wounds infections, central nervous system shunt infections, ophthalmologic infections, peritoneal dialysis-related infections and infections of

108 prosthetic joints (Bannerman, 2003). *S. haemolyticus* has been implicated in native valve
 109 endocarditis, septicemia, urinary tract infections, peritonitis and wound, bone and joint infections
 110 (Bannerman, 2003). *S. saprophyticus* is associated with urinary tract infections in young females. *S.*
 111 *lugdunensis* has been implicated in arthritis, catheter infections, bacteremia, urinary tract infections,
 112 prosthetic joint infections and endocarditis (Bannerman, 2003). Other CNS species have been
 113 implicated in a variety of infections.

114

115 4. Virulence factors

116

117 Already in 1972, the “slime” production of CNS was noted as an important factor in the
 118 pathogenesis of infections (Huebner et al., 1999). This “slime” or biofilm is the most important
 119 virulence factor of *S. epidermidis*. The biofilm formation enables attachment and persistence of the
 120 bacteria on foreign materials. Moreover, bacteria organized in biofilms are protected from the action
 121 of antibiotics and the immune system (Costerton et al., 1999; Mack et al., 2007). The first phase in
 122 biofilm formation is attachment, in which staphylococcal binding proteins such as the
 123 autolysin/adhesins AtlE and Aae, the fibrinogen-binding protein Fbe/Sdrg, the fibronectin-binding
 124 protein Embp and the lipase GehD, play an important role (Mack et al., 2007). The most important
 125 adhesion molecule is AtlE. The second phase in biofilm formation is accumulation, in which the
 126 polysaccharide intercellular protein PIA, encoded by the *icaADBC* locus, is a major functional
 127 component (von Eiff et al., 2002; Mack et al., 2007). In strains lacking the *icaADBC* locus, biofilm
 128 formation is mediated by the accumulation-associated protein (Aap) or by the biofilm-associated
 129 protein (Bap/Bhp) (Ziebuhr et al., 2006). Recently, 2 biofilm producing strains, negative for PIA,
 130 Aap and Bhp, have been described (Qin et al., 2007). The regulation of biofilm formation is very
 131 complicated and best studied in *S. epidermidis*. Different regulatory systems, such as *sae*, alternative
 132 sigma factor σ^B , *sar* and quorum sensing (this is inter and intra species communication, possible
 133 when a certain bacterial density has been established) systems play a role (Xu et al., 2006; Mack et

134 al., 2007). The 2 quorum sensing systems in *S. epidermidis*, *agr* and *luxS/AI-2*, repress the biofilm
 135 formation: *luxS* system by downregulating PIA production, *agr* system by downregulating AtlE
 136 production. *Agr* controls the expression of toxins and secreted virulence factors and the interaction
 137 with the immune system (Kong et al., 2006). Quorum sensing systems have potential as therapeutic
 138 targets for control of staphylococcal infections (Harraghy et al., 2007). Another important virulence
 139 factor of CNS is the antibiotic resistance. Regulation of biofilm formation and methicillin resistance
 140 seem to use similar pathways, as insertion of a certain transposon influences both biofilm formation
 141 and the expression of methicillin resistance (Mack et al., 2007). Methicillin resistance was found to
 142 be significantly higher in slime positive isolates (81%) than in slime negative isolates (57%) (Koksal
 143 et al., 2007).

144 The phenotypic variability and heterogeneous gene expression observed in CNS, especially in
 145 *S. epidermidis*, is an advantage for adaptation to changing environmental conditions (Ziebuhr et al.,
 146 2006).

147 Other potential virulence factors of *S. epidermidis* include the following: extracellular
 148 enzymes and toxins: metalloprotease with elastase activity, cysteine protease, serine protease, lipase,
 149 fatty acid modifying enzymes (FAME) and the δ -toxin. CNS also produce lantibiotics (von Eiff et
 150 al., 2002; Vuong et al., 2002). Lantibiotics are bacteriocins such as epidermin, which are active
 151 against other gram-positive bacteria.

152 Several studies tried to discriminate between invasive and commensal *S. epidermidis* strains
 153 by the detection of virulence associated genes (Frebourg et al., 2000; Vandecasteele et al., 2003;
 154 Rohde et al., 2004). In the study by Rohde et al (Rohde et al., 2004) the virulence genes *icaADBC*,
 155 *mecA* and IS256 were all highly prevalent in invasive strains. Commensal strains from healthy
 156 individuals almost completely lacked *icaADBC*, *mecA* and IS256. However in the commensal
 157 strains isolated from hospitalized bone marrow transplant patients the studied genes had the same
 158 distribution as the invasive strains, suggesting replacement of their own commensal flora by hospital
 159 strains. As a consequence these genes cannot discriminate between invasive and commensal strains.

160 *S. lugdunensis* and *S. schleiferi* are more pathogenic than other CNS species. Like *S. aureus*,
 161 they may express a clumping factor and/or produce a thermostable DNase (von Eiff et al., 2002). A
 162 variable percentage of these strains produces extracellular slime or glycocalyx, esterase, FAME,
 163 protease, lipase and a haemolysin. *S. lugdunensis* is more likely than other CNS, to be considered a
 164 pathogen until proven otherwise (Poutanen et al., 2001; Koziol-Montewka et al., 2006). *S.*
 165 *lugdunensis* is often misidentified as *S. aureus*, because slide agglutination tests are often positive.

166 The main virulence factor of *S. saprophyticus* in UTI is the capacity to adhere to uroepithelial
 167 cells by means of surface-associated proteins: the autolysin/adhesin Aas, Ssp and Sdrl. Once
 168 colonization has been established several invasion factors are produced: urease, elastase, lipase and
 169 FAME (von Eiff et al., 2002).

170

171 **5. Antibiotic resistance**

172

173 Resistance to antibiotics in CNS is of major concern. Penicillin resistance in CNS is very
 174 high: even in 1968 the resistance rate was 60% (Corse et al., 1968). Nowadays, resistance is around
 175 91% in clinical strains (Cuevas et al., 2004; Koksai et al., 2007). Methicillin resistance in
 176 staphylococci is caused by expression of PBP2a encoded by the *mecA* gene. *MecA* is located on a
 177 genetic element called the staphylococcal cassette chromosome (SCC). There is evidence of
 178 horizontal transfer of SCC cassettes between staphylococcal species (Hanssen et al., 2004), which
 179 implies that CNS could serve as a reservoir for the spread of resistance genes to *S. aureus*. In clinical
 180 samples, rates of methicillin resistance of 55% - 77% and even 86% in ICU settings, have been
 181 reported (Del' Alamo et al., 1999; Agvald-Öhman et al., 2004; Cuevas et al., 2004; Chaudhury et al.,
 182 2007; Gatermann et al., 2007; Jones et al., 2007a; Jones et al., 2007b; Koksai et al., 2007; Sader et
 183 al., 2007;). In continuous ambulatory peritoneal dialysis (CAPD) patients and medical students 38%
 184 and 27%, respectively, of colonizing CNS strains were methicillin resistant (de Mattos et al., 2003;
 185 Koziol-Montewka et al., 2006; Higuchi et al., 2007). Several studies show an increase of MR CNS

over time. In Spain, methicillin or oxacillin resistance remained stable (26%-34% of CNS) between 1986 and 1994, increased significantly in 1996 (51%) and reached 61% in 2002 (Cuevas et al., 2004). An increase of methicillin resistant CNS strains was also found in other studies: from 38% in 1996 to 68% in 2007 in blood of patients with true bacteremia (Koksal et al., 2007), from 19% in 1991 to 74% in 1998 in peritoneal dialysis patients (Zelenitsky et al., 2000). It is generally assumed that approximately 80% of nosocomial isolates and 30-40% of isolates obtained from healthy carriers or patients from the community, demonstrate resistance to methicillin (Huebner et al., 1999; de Mattos et al., 2003; Koziol-Montewka et al., 2006).

In Spain, resistance to ciprofloxacin and erythromycin increased progressively in CNS from 1% and 41% in 1986 to 45% and 63% in 2002 (Cuevas et al., 2004). Susceptibility percentages to different antibiotics are given in Table 2. Two surveillance studies, performed in 2006, one in the US and one in Europe, Asia, Australia and Latin America show comparable susceptibility results. Susceptibility rates to fluoroquinolones, erythromycin, clindamycin, trimethoprim/sulfamethoxazole, gentamicin, teicoplanin and linezolid are 42%-47%, 30%-34%, 60%-61%, 62%-59%, 71%-58%, 99% and 98-99%, respectively (Jones et al., 2007a, Jones et al., 2007b). Methicillin resistant strains have high rates of resistance to other classes of antibiotics. Resistance rates in methicillin susceptible strains are much lower (Reynolds et al., 2004; Biedenbach et al., 2007; Koksal et al., 2007; Sader et al., 2007).

CNS were the first organisms in which acquired glycopeptide resistance (vancomycin MIC ≥ 8 $\mu\text{g/mL}$, teicoplanin MIC ≥ 16 $\mu\text{g/mL}$) was recognized (Biavasco et al., 2000). Most of the clinical glycopeptides resistant isolates are resistant to teicoplanin, but susceptible to vancomycin, indicating a heterogeneous expression of glycopeptides resistance (Nunes et al., 2006). The resistance mechanisms to glycopeptides in CNS are not yet fully understood. Most reports show that resistance mechanism in CNS is similar to that described in VISA and hetero-VISA strains (Nunes et al., 2006). This mechanism is multifactorial, expression related and can be selected, in vitro, by exposure to teicoplanin and vancomycin (Biavasco et al., 2000; Nunes et al., 2006). Laboratory detection of

glycopeptide-resistance may be problematic, as there is an influence of various technical factors. Susceptibility testing by disk diffusion results in a very high percentage of false susceptible results (very major errors in 80% of the nonsusceptible to teicoplanin strains) (Del' Alamo et al., 1999). Susceptibility rates to vancomycin and teicoplanin of respectively 99%-100% and 68%-100% are found in clinical studies (Del' Alamo et al., 1999; Cuevas et al., 2004; Reynolds et al., 2004; Chaudhury et al., 2007; Koksai et al., 2007; Sader et al., 2007).

Novel antibiotics have been developed to overcome the resistance problem in CNS. These antibiotics include for example linezolid, daptomycin, tigecycline and quinupristin/dalfopristin. Susceptibility of CNS isolates towards these agents is usually 100% (Garrison et al., 2005; Sader et al., 2007). In recent surveillance studies the lowest reported susceptibility percentages are 98,4% for linezolid (Jones et al., 2007a), 99% for quinupristin/dalfopristin (Jones et al., 2007b) and 99,9% for daptomycin (Sader et al., 2007). In one institution, 4% of CNS, originating from 25 patients, were found to be linezolid resistant. The majority of these resistant strains showed some genetic relatedness, previous linezolid use was an independent predictor of linezolid resistance and the majority of patients had been accommodated in a single ward (Potoski et al., 2006).

The prevalence of resistance varies widely among staphylococcal species. The highest rates of resistance are found in *S. haemolyticus*: 76-96% oxacillin resistant, 80-90% erythromycin resistant, 26-29% teicoplanin nonsusceptible (Gill et al., 1983; Del' Alamo et al., 1999; Chaudhury et al., 2007; Gatermann et al., 2007). Oxacillin resistance rates are also high in *S. hominis* (80%) (Gatermann et al., 2007) and *S. epidermidis* (38-81%) (Gill et al., 1983; Del' Alamo et al., 1999; Gatermann et al., 2007). Teicoplanin nonsusceptible isolates occur in 3% of *S. epidermidis* and in none of the *S. hominis* isolates (Del' Alamo et al., 1999). *S. lugdunensis* has nearly uniform in vitro susceptibility to most antimicrobials, including penicillins, cephalosporins and macrolides (Poutanen et al., 2001; Gatermann et al., 2007).

6. Epidemiology

238

239 Several studies show that most CNS infections are hospital-acquired or health-care related
 240 (Rupp et al., 1994; Huebner et al., 1999). Colonization of patients and hospital staff with antibiotic
 241 resistant *S. epidermidis* precedes infection with these organisms (Kloos et al., 1994; Nouwen et al.,
 242 1998; Widerström et al., 2006). Some clones are probably endemic in the hospital environment. CNS
 243 have the ability to survive in the intensive care unit (ICU) surroundings on medical devices and
 244 medical equipment for weeks to months (Neely et al., 2000). The clonal spread of identical or closely
 245 related methicillin-resistant CNS strains within hospitals and even between hospitals has been
 246 demonstrated (Monsen et al., 2000; Widerström et al., 2006). Frequent transmission of CNS between
 247 intubated patients was demonstrated in a multidisciplinary ICU (Agvald-Öhman et al., 2004). The
 248 spread of CNS strains is well studied in neonatal ICUs, where CNS are the major causative organism
 249 of sepsis. A relatively small number of molecular types of CNS can persist in the neonatal ICU for
 250 many years. The *mecA* gene carriage in these clusters is usually very high, which suggests that
 251 antibiotic resistance is one of the major selective forces (Krediet et al., 2004). Multi locus sequence
 252 typing identified a sequence type (ST 27), which contained exclusively *ica*-positive isolates and
 253 represented the majority of clinical strains within different hospitals in Germany. The ST 27 clone
 254 was also detected in a Norwegian hospital and in medical facilities in the United States (Kozitskaya
 255 et al., 2005). The combination of biofilm formation, antibiotic resistance and genetic flexibility (due
 256 to multiple copies of IS256) in the ST27 clone may explain why this clone is so predominant in
 257 hospital settings. It is tempting to assume that patients who are admitted to hospital are rapidly
 258 colonized with this clone and that this newly acquired microflora might represent the origin for later
 259 infection (Kozitskaya et al., 2005; Ziebuhr et al., 2006). Hospital personnel can also be responsible
 260 for spreading multi resistant CNS isolates (Hira et al., 2007). Clonal types can only be distinguished
 261 by genotypic methods and not by phenotypic appearance of the colonies or by antibiotype (Nouwen
 262 et al., 1998; Miragaia et al., 2002). Risk factors for the emergence and spread of CNS clones in
 263 hospitals include the duration of hospital stay (especially ICU stay), duration of antibiotic treatment,

antibiotic pressure in the environment, and hygienic standards (Widerström et al., 2006). The use of contact and hand hygiene precautions is extremely important for preventing nosocomial colonization and infections. Maybe periodic surveillance of patients and staff in specific wards could be useful to avoid the spread of multi resistant CNS. A good antibiotic policy, in order to reduce the antibiotic pressure, will help to reduce the incidence of multi resistant CNS.

7. Specific infections

7.1. Urinary tract infections (UTI)

S. saprophyticus is the second (after *Escherichia coli*) most frequent causative organism of uncomplicated urinary tract infections (UTI) in women. Complications, such as recurrent infection, acute pyelonephritis, nephrolithiasis, septicaemia and endocarditis have been documented but are rare (Raz et al., 2005; Widerström et al., 2007). The vast majority of infections occur in young, sexually active women. *S. saprophyticus* can also cause UTI in males of all ages (Raz et al., 2005). *S. saprophyticus* is found in 3-9% of cases of uncomplicated acute cystitis (Christiaens et al., 1998; Grude et al., 2005; Nys et al., 2006). Frequency of isolation is age dependent. In a recent study, *S. saprophyticus* was isolated from 7%, 4% and 0.5% of subjects in age categories 11-20 years, 21-50 years and 51-70 years, respectively (Nys et al., 2006). *S. saprophyticus* can be pathogenic in low numbers ($<10^5$ cfu/mL) (Rupp et al., 1994). *S. saprophyticus* is probably often missed as causative organism of UTI, as the bacteriuria is considered to be nonsignificant, especially when bacterial counts are low.

S. saprophyticus has also been isolated from 7% of rectal swabs taken from carcasses of cattle and pigs. The micro-organism is a common contaminant of various food samples, especially of raw beef and pork (Raz et al., 2005). In humans, the major reservoir of *S. saprophyticus* is the gastrointestinal tract. The following risk factors have been identified: recent sexual intercourse,

290 outdoor swimming and meat processing (Huebner et al., 1999). Colonization and infection is more
 291 frequent during summer and fall.

292 Other CNS species, mainly *S. epidermidis*, are occasionally found as causative organisms in
 293 UTI (<5% of UTI in hospital environment). *S. epidermidis* is usually isolated from hospitalized,
 294 elderly patients with urinary catheters or other manipulations to the urinary tract (Rupp et al., 1994;
 295 Raz et al., 2005).

296

297 7.2. Endocarditis

298

299 The incidence and characteristics of CNS endocarditis were studied in a large cohort of
 300 patients from the International Collaboration on Endocarditis Merged Database (Chu et al., 2004).
 301 CNS accounted for 7% of cases of definite native valve endocarditis (NVE), excluding endocarditis
 302 associated with injection drug use (Chu et al., 2004). Most CNS NVE is caused by *S. epidermidis*
 303 (85%). The remainder is caused by *S. hominis* (6%), *S. lugdunensis* (5%), *S. capitis*, *S. caprae* and *S.*
 304 *simulans*. Rates of heart failure and mortality were found to be similar between patients with CNS
 305 and *S. aureus* NVE (Chu et al., 2004). Patients with NVE caused by staphylococci were significantly
 306 more likely to have health-care associated endocarditis than patients with viridans streptococcal
 307 NVE.

308 In a study from 1999, streptococci are responsible for 48% of cases of infective endocarditis,
 309 *S. aureus* for 23% and CNS for 6% (Hoen et al., 2002). These frequencies were also found in a study
 310 conducted between 1951 and 1965: *S. epidermidis* was the third most frequent isolated organism (in
 311 10% of cases) in cases of bacterial endocarditis, after viridans streptococci (47%) and *S. aureus*
 312 (24%) (Dalton et al., 1967). Injecting heroin users are susceptible to right-sided endocarditis due to *S.*
 313 *epidermidis* (von Eiff et al., 2001).

314 Characteristics of patients with prosthetic valve endocarditis (PVE) were studied and
 315 compared with NVE patients in the International Collaboration on Endocarditis – Prospective Cohort

Study (Wang et al., 2007). Patients with PVE had a higher rate of CNS infection (16.9%) than patients with NVE (8.3%). In literature, higher rates (than in the study by Wang et al., 2007) of CNS in infections of prosthetic valves are found (40-50% of cases) (Huebner et al., 1999; von Eiff et al., 2001).

S. lugdunensis endocarditis has more features of *S. aureus* endocarditis than of CNS endocarditis. The infection is often aggressive, most cases are community-acquired and only a minority of patients (approximately 25%) has prosthetic valves (Poutanen et al., 2001).

7.3. Blood stream infections (BSI)

Almost all cases of bacteremia due to CNS are nosocomial or at least health-care related in origin. Intravascular and cardiovascular devices, such as peripheral and central catheters, vascular grafts, implanted defibrillators and coronary stents, are widely used. Catheter related infections are by far the most common cause of bacteremia due to CNS (Huebner et al., 1999).

Throughout the 1960s and 1970s, Gram-negative organisms were the most common pathogens isolated from patients with nosocomial blood stream infections (BSI) (Wisplinghoff et al., 2004), but in one study (Dalton et al., 1967), between 1951 and 1965, *S. epidermidis* accounted for 23% of positive blood cultures in patients with suspected bacteremia. Second and third most frequent isolated organism was *S. aureus* (12%) and *Corynebacterium* species (12%). The high rates of common skin bacteria (such as *Corynebacterium* species and *S. epidermidis*) in the study by Dalton et al., 1967, raise concern about the clinical significance. The incidence of bacteremia caused by CNS rose between 161% and 754%, respectively in large teaching and small non-teaching hospitals, between 1980 and 1989 (Rupp et al., 1994). During 1986 - 1989 CNS were the most frequently reported cause of BSI, accounting for 27% of BSI (Schaberg et al., 1991).

In a nation wide surveillance study in the USA, performed between 1995 and 2002, CNS accounted for 31% of all nosocomial BSI (Wisplinghoff et al., 2004). The crude mortality of BSI due

342 to CNS was 25.7% in an intensive care unit (ICU) and 13.8% in a non-ICU ward. In Flanders,
 343 Belgium, CNS account for 25.9% of all nosocomial BSI. Forty-eight percent of BSI cases in
 344 Flanders were due to catheter related sepsis, 32% to sepsis of unknown origin, and 20% were
 345 associated with another infection or an invasive manipulation (Suetens et al., 2002).

346 CNS are the most common pathogens in catheter-related infections (50-70% of cases). In
 347 addition to bacteremia, CNS, mainly *S. epidermidis*, can cause exit-site infections, tunnel infections,
 348 infected thrombophlebitis, endocarditis and abscesses (Rupp et al., 1994; Huebner et al., 1999). At
 349 the time of removal, 8-40% of all catheters are colonized by CNS (Rupp et al., 1994).
 350 Positive blood cultures with CNS can be explained by contamination, true bacteremia or transient
 351 benign bacteremia (Rupp et al., 1994). It may be very difficult to distinguish between infection and
 352 contamination. As the positive predictive value of a blood culture growing CNS varies from 6 to
 353 12% (Correa et al., 1999), the criteria used for defining true bacteremia influence reported CNS
 354 prevalence. The CDC criteria for defining bloodstream infection (Garner et al., 1988) are used in
 355 most of the studies. The intensity of surveillance also influences the reported BSI incidence and
 356 prevalence of CNS.

357 There is increasing evidence that the mucosa, rather than skin, is the likely source of CNS
 358 bacteremia in cancer patients (Costa et al., 2004; Costa et al., 2006). A recent study showed that
 359 among cancer patients with CNS positive blood cultures and an indwelling central venous catheter
 360 (CVC), the mucosal sites were the most frequently colonised by CNS (nasal mucosa in 86%, rectal
 361 mucosa in 40%). The skin at the CVC site was colonised by CNS in 32% of patients. Among patients
 362 with true bacteremia, 6 mucosal isolates and only 1 skin isolate were genetically related to the isolate
 363 recovered from blood based on pulsed-field gel electrophoresis (Costa et al., 2006). A review of the
 364 literature confirms that CNS colonise mucosal sites including gut, nares and throat and may
 365 translocate from mucosal sites to the bloodstream (Costa et al., 2004). This is supported by
 366 epidemiological, experimental, clinical and molecular relatedness studies. The assumption that the
 367 skin is the primary source of CNS infection has been the reason to try to achieve a reduction of CNS

catheter related bacteremia with skin decontamination and the use of maximum barrier precautions at the time of CVC insertion. However, not all studies show a positive effect of these two interventions (Costa et al., 2004).

7.4. Foreign body-related infections (FBRIs)

Insertion or implantation of medical devices is associated with a risk of bacterial and fungal infections. Medical devices are increasingly used in almost all fields of medicine for diagnostic and/or therapeutic procedures. The contamination of the device occurs most likely by inoculation with only a few micro-organisms from the patient's skin or mucous membranes during implantation or subsequent manipulations (Rupp et al., 1994; von Eiff et al., 2001; von Eiff et al., 2005). The pathogens may also be acquired from the hands of the surgical or clinical staff (von Eiff et al., 2005). Staphylococci, particularly *S. epidermidis*, account for the majority of FBRIs (von Eiff et al., 2005). Most CNS isolates causing foreign body-related infections (FBRI) are methicillin resistant.

Neurosurgical devices (ventricular shunts, implantable stimulators, intracranial pressure devices) are frequently used. Of the pathogens isolated from ventriculoperitoneal shunt infections in pediatric patients less than 8 years old, 53% were CNS (Filka et al., 1999). Clinical symptoms can be subtle and nonspecific. (Huebner et al., 1999). In a review of CSF shunt infections, caused by a single pathogen, CNS and *S. aureus* were isolated with equal frequency, i. e. in 23% of cases each (Wang et al., 2004). Others reported that CNS are the most common cause of shunt infections (40-60% of infections) (Rupp et al., 1994). In a retrospective study, pure culture of CNS accounted for 11% of cases of adult bacterial meningitis (Huang et al., 2005). All cases had post-neurosurgical states as underlying condition.

CNS cause 17-56% of episodes of continuous ambulatory peritoneal dialysis-associated peritonitis (Rupp et al., 1994, Monsen et al., 2000). *S. epidermidis* is the predominant species (78%) followed by *S. haemolyticus* (11%) (Monsen et al., 2000). From 1991 to 1998, the incidence of

394 dialysis-related peritonitis caused by *S. epidermidis* decreased, probably due to the introduction of a
 395 new dialysis system. Even so, *S. epidermidis* was still the most common cause of peritonitis in 1998,
 396 causing 28% of culture positive cases (Zelenitsky et al., 2000).

397 Up to 60% of vascular graft infections and 20-40% of prosthetic joint infections are caused
 398 by CNS (Rupp et al., 1994).

399 CNS are also associated with low grade foreign body infections such as aseptic loosening of hip or
 400 other joint prostheses, or fibrous capsular contracture syndrome after mammary augmentation with
 401 silicone prostheses (von Eiff et al., 2001).

402

403 7.5. Surgical site infection

404

405 *S. aureus* is the most common pathogen isolated from surgical site infections (SSI) (20%),
 406 followed by CNS (14%) and *Enterococcus* species (12%) (CDC, 1996). In a secondary care hospital,
 407 CNS were the third (10%) most frequently isolated micro-organism, after *S. aureus* (21%) and
 408 *Streptococcus* species (11.2%) (Cantlon et al., 2006). Outbreaks of SSI originating from operating
 409 room personnel have been reported, but probably most infections are indogenous (Huebner et al.,
 410 1999). Sternal osteomyelitis following median sternotomy is caused predominantly by CNS (Rupp et
 411 al., 1994).

412

413 7.6. Endophthalmitis

414

415 CNS are by far the most common pathogens isolated from vitreous biopsies in cases of
 416 endophthalmitis. In a review of 67 cases of endophthalmitis, 31% of the culture positive biopsies
 417 (only 44%) yielded CNS as pathogen (Karacal et al., 2007). Most cases of endophthalmitis were
 418 postoperative (73.5%), mainly after cataract surgery. Other cases were associated with endogenous
 419 origin (13%), post-intravitreal injection (6%), keratitis (4%) and post-traumatic (3%). Post-traumatic

endophthalmitis, a complication of a penetrating eye injury, is caused by CNS in 28% of cases (Abu El-Asrar et al., 1999).

7.7. Mastitis

CNS are rarely associated with mastitis in humans. In lactation mastitis, the agents most frequently cultured from milk are *S. aureus* and CNS (Barbosa-Cesnik et al., 2003). In non lactational breast abscesses, CNS are rarely isolated as sole pathogen. Some cases of non lactational breast abscesses, especially with *S. lugdunensis* have been published in literature (Asnis et al., 2003). *S. aureus* and anaerobes are the most common organisms to cause breast abscesses (Surani et al., 1993).

8. Conclusions and Outlook

CNS are a group of micro-organisms that are increasingly implicated as a cause of significant infection. The infection rate has been correlated with the increase in the use of invasive and indwelling medical devices and the growing number of immunocompromised patients. Community acquired, primary CNS infections are rare, except for *S. saprophyticus* UTI, NVE and *S. lugdunensis* infections. The increase of CNS infections is particularly high in bacteraemia. CNS are now a major cause of nosocomial or health care related infections. The consequences of these infections are substantial, both in morbidity and mortality and in financial costs. Clonal spread of CNS clusters within and even between hospitals has been demonstrated. This is of major concern as CNS are often multi-drug resistant and can serve as a reservoir of resistance genes. Implementation of infection control measures similar to those used for MRSA may be necessary to prevent further spread of methicillin and multi-drug resistant CNS.

Conflict of Interest

None of the authors (A. Piette, G. Verschraegen) has a financial or personal relationship with other people or organizations that could inappropriately influence or bias the paper entitled “Role of coagulase-negative staphylococci in human disease”.

References

- Abu el-Asrar, A.M., al-Amro, S.A., al-Mosallam, A.A., al-Obeidan, S., 1999. Post-traumatic endophthalmitis: causative organisms and visual outcome. *Eur. J. Ophthalmol.* 9, 21-31.
- Agvald-Ohman, C., Lund, B., Edlund, C., 2004. Multiresistant coagulase-negative staphylococci disseminate frequently between intubated patients in a multidisciplinary intensive care unit. *Crit. Care.* 8, R42-R47.
- Asnis, D.S., St John, S., Tickoo, R., Arora, A., 2003. *Staphylococcus lugdunensis* breast abscess: is it real? *Clin. Infect. Dis.* 36, 1348.
- Bannerman, T., 2003. *Staphylococcus*, *Micrococcus* and other catalase-positive cocci that grow aerobically. In Murray, P., Baron, E., Jorgenson, J., Tenover, M., Tenover, R., (Eds.) *Manual of clinical microbiology*, ASM Press, Washington.
- Barbosa-Cesnik, C., Schwartz, K., Foxman, B., 2003. Lactation mastitis. *JAMA.* 289, 1609-1612.
- Biavasco, F., Vignaroli, C., Varaldo, P.E., 2000. Glycopeptide resistance in coagulase-negative staphylococci. *Eur. J. Clin. Microbiol. Infect. Dis.* 19, 403–417.
- Biedenbach, D.J., Bell, J.M., Sader, H.S., Fritsche, T.R., Jones, R.N., Turnidge, J.D., 2007. Antimicrobial susceptibility of Gram-positive bacterial isolates from the Asia-Pacific region and an in vitro evaluation of the bactericidal activity of daptomycin, vancomycin, and teicoplanin: a SENTRY Program Report (2003–2004). *Int. J. Antimicrob. Agents.* 30, 143-149.

- 472 Cantlon, C.A., Stemper, M.E., Schwan, W.R., Hoffman, M.A., Qutaishat, S.S., 2006.
 473 Significant pathogens isolated from surgical site infections at a community hospital in the Midwest.
 474 Am. J. Infect. Control. 34, 526-529.
- 475 CDC, 1996. National nosocomial infections surveillance (NNIS) report, data summary from
 476 october 1986 - april 1996, issued may 1996. Am. J. Infect. Control. 24, 380-388.
- 477 Chaudhury, A., Kumar, A.G., 2007. In vitro activity of antimicrobial agents against oxacillin
 478 resistant staphylococci with special reference to *Staphylococcus haemolyticus*. Indian J. Med.
 479 Microbiol. 25, 50-52.
- 480 Christiaens, T.H., Heytens, S., Verschraegen, G., De Meyere, M., De Maeseneer, J., 1998.
 481 Which bacteria are found in Belgian women with uncomplicated urinary tract infections in primary
 482 health care, and what is their susceptibility pattern anno 95-96. Acta Clin. Belg. 53, 184-188.
- 483 Chu, V.H., Cabell, C.H., Abrutyn, E., Corey, G.R., Hoen, B., Miro, J.M., Olaison, L.,
 484 Stryjewski, M.E., Pappas, P., Anstrom, K.J., Eykyn, S., Habib, G., Benito, N., Fowler, V.G Jr, 2004.
 485 Native valve endocarditis due to coagulase-negative staphylococci: report of 99 episodes from the
 486 International Collaboration on Endocarditis Merged Database. Clin. Infect. Dis. 39, 1527-30.
- 487 Correa, L., Pittet, D., 2000. Problems and solutions in hospital-acquired bacteraemia. J. Hosp.
 488 Infect. 46, 89-95.
- 489 Corse, J., Williams, R.E., 1968. Antibiotic resistance of coagulase-negative staphylococci and
 490 micrococci. J. Clin. Pathol. 21, 722-728.
- 491 Costa, S.F., Miceli, M.H., Anaissie, E.J., 2004. Mucosa or skin as source of coagulase-
 492 negative staphylococcal bacteraemia? Lancet Infect. Dis. 4, 278-286.
- 493 Costa, S.F., Barone, A.A., Miceli, M.H., van der Heijden, I.M., Soares, R.E., Levin, A.S.,
 494 Anaissie, E.J., 2006. Colonization and molecular epidemiology of coagulase-negative
 495 Staphylococcal bacteremia in cancer patients: a pilot study. Am. J. Infect. Control. 6, 34, 36-40.
- 496 Costerton, J.W., Stewart, P.S., Greenberg, E.P., 1999. Bacterial biofilms: a common cause of
 497 persistent infections. Science. 284, 1318-1322.

- 498 Cuevas, O., Cercenado, E., Vindel, A., Guinea, J., Sanchez-Conde, M., Sanchez-Somolinos,
 499 M., Bouza, E., 2004. Evolution of the antimicrobial resistance of *Staphylococcus* spp. in Spain: Five
 500 nationwide prevalence studies, 1986 to 2002. *Antimicrob. Agents Chemother.* 48, 4240–4245.
- 501 Dalton, H.P., Allison, M.J., 1967. Etiology of bacteremia. *Appl. Microbiol.* 15, 808-814.
- 502 Del' Alamo, L., Cereda, R.F., Tosin, I., Miranda, E.A., Sader, H.S., 1999. Antimicrobial
 503 susceptibility of coagulase-negative staphylococci and characterization of isolates with reduced
 504 susceptibility to glycopeptides. *Diagn. Microbiol. Infect. Dis.* 34, 185-191.
- 505 de Mattos, E.M., Teixeira, L.A., Alves, V.M., Rezenda e Resende, C.A., da Silva Coimbra,
 506 M.V., da Silva-Carvalho, M.C., Ferreira-Carvalho, B.T., Figueiredo, A.M., 2003. Isolation of
 507 methicillin-resistant coagulase-negative staphylococci from patients undergoing continuous
 508 ambulatory peritoneal dialysis (CAPD) and comparison of different molecular techniques for
 509 discriminating isolates of *Staphylococcus epidermidis*. *Diagn. Microbiol. Infect. Dis.* 45, 13-22.
- 510 Euzéby, J.P., 2007, LSPN List of prokaryotic names with standing in nomenclature,
 511 <http://www.bacterio.cict.fr/>
- 512 Filka, J., Huttova, M., Tuharsky, J., Sagat, T., Kralinsky, K., Kremery V., 1999. Nosocomial
 513 meningitis in children after ventriculoperitoneal shunt insertion. *Acta Paediatr.* 88, 576-578.
- 514 Frebourg, N.B., Lefebvre, S., Baert, S., Lemeland, J.F., 2000. PCR-based assay for
 515 discrimination between invasive and contaminating *Staphylococcus epidermidis* strains. *J. Clin.*
 516 *Microbiol.* 38, 877–880.
- 517 Garner, J.S., Jarvis, W.R., Emory, T.G., Horan, T.C., Hughes, J.M., 1988. CDC definitions
 518 for nosocomial infections. *Am. J. Infect. Control.* 16, 128-140.
- 519 Garrison, M.W., Neumiller, J.J., Setter, S.M., 2005. Tigecycline: an investigational
 520 glycylcycline antimicrobial with activity against resistant gram-positive organisms. *Clin. Ther.* 27,
 521 12-22.
- 522 Gatermann, S.G., Koschinski, T., Friedrich, S., 2007. Distribution and expression of
 523 macrolide resistance genes in coagulase-negative staphylococci. *Clin. Microbiol. Infect.* 13, 777-781.

- Gill, V.J., Selepak, S.T., Williams, E.C., 1983. Species identification and antibiotic susceptibilities of coagulase-negative staphylococci isolated from clinical specimens. *J. Clin. Microbiol.* 18, 1314-1319.
- Grude, N., Tveten, Y., Jenkins, A., Kristansen, B.E., 2005. Uncomplicated urinary tract infections. *Scand. J. Prim. Health care.* 23, 115-119.
- Hanssen, A.M., Kjeldsen, G., Sollid, J.U., 2004. Local variants of Staphylococcal cassette chromosome *mec* in sporadic methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative Staphylococci: evidence of horizontal gene transfer? *Antimicrob. Agents Chemother.* 48, 285-296.
- Harraghy, N., Kerdudou, S., Herrmann, M., 2007. Quorum-sensing systems in staphylococci as therapeutic targets. *Anal. Bioanal. Chem.* 387, 437-444.
- Heikens, E., Flier, A., Paauw, A., Florijn, A.C., Fluit, A., 2005. Comparison of genotypic and phenotypic methods for species-level identification of clinical isolates of coagulase-negative staphylococci. *J. Clin. Microbiol.* 43, 2286-2290.
- Higuchi, W., Isobe, H., Iwao, Y., Dohmae, S., Saito, K., Takano, T., Otsuka, T., Baranovich, T., Endo, C., Suzuki, N., Tomiyama, Y., Yamamoto, T., 2007. Extensive multidrug resistance of coagulase-negative staphylococci in medical students. *J. Infect. Chemother.* 13, 63-66.
- Hira, V., Sluijter, M., Estevão, S., Horst-Kreft, D., Ott, A., de Groot, R., Hermans, P.W., Kornelisse, R.F., 2007. Clinical and molecular epidemiologic characteristics of coagulase-negative staphylococcal bloodstream infections in intensive care neonates. *Pediatr. Infect. Dis. J.* 26, 607-612.
- Hoen B., Alla, F., Selton-Suty, C., Béguinot, I., Bouvet, A., Briançon, S., Casalta, J.P., Danchin, N., Delahaye, F., Etienne, J., Le Moing, V., Leport, C., Mainardi, J.L., Ruimy, R., Vandenesch, F., 2002. Changing profile of infective endocarditis, results of a 1-year survey in France. *JAMA.* 288, 75-81.

- Huang, C.R., Lu, C.H., Wu, J.J., Chang, H.W., Chien, C.C., Lei, C.B., Chang, W.N., 2005. Coagulase-negative staphylococcal meningitis in adults: clinical characteristics and therapeutic outcomes. *Infection*. 33, 56-60.
- Huebner, J., Goldmann, D.A., 1999. Coagulase-negative staphylococci: Role as Pathogens. *Annu. Rev. Med.* 50, 223-236.
- Jarløv, J.O., Højbjerg, T., Busch-Sørensen, C., Scheibel, J., Møller, J.K., Kolmos, H.J., Wandall, D.A., 1996. Coagulase-negative Staphylococci in Danish blood cultures: species distribution and antibiotic susceptibility. *J. Hosp. Infect.* 32, 217-227.
- Jones, R.N., Fritsche, T.R., Sader, H.S., Ross, J.E., 2007. LEADER surveillance program results for 2006: an activity and spectrum analysis of linezolid using clinical isolates from the United States (50 medical centers). *Diagn. Microbiol. Infect. Dis.* 59, 309-317.
- Jones, R.N., Fritsche, T.R., Sader, H.S., Ross, J.E., 2007. Zyvox® annual appraisal of potency and spectrum program results for 2006: an activity and spectrum analysis of linezolid using clinical isolates from 16 countries. *Diagn. Microbiol. Infect. Dis.* 59, 199-209.
- Karacal, H., Kymes, S.M., Apte, R.S., 2007. Retrospective analysis of etiopathogenesis of all cases of endophthalmitis at a large tertiary referral center. *Int. Ophthalmol.* 27, 251-259.
- Kawamura, Y., Hou, X.G., Sultana, F., Hirose, K., Miyake, M., Shu, S.E., Ezaki, T., 1998. Distribution of Staphylococcus species among human clinical specimens and emended description of *Staphylococcus caprae*. *J. Clin. Microbiol.* 36, 2038-2042.
- Kleeman, K.T., Bannerman, T.L., Kloos, W.E., 1993. Species distribution of coagulase-negative staphylococcal isolates at a community hospital and implications for selection of staphylococcal identification procedures. *J. Clin. Microbiol.* 31, 1318-1321.
- Kloos, W.E., Schleifer, K.J., 1975. Simplified scheme for routine identification of human *Staphylococcus* species. *J. Clin. Microbiol.* 1, 82-88.
- Kloos, W.E., Bannerman, T.L., 1994. Update on clinical significance of coagulase-negative staphylococci. *Clin. Microbiol. Rev.* 7, 117-140.

- 574 Koksai, F., Yasar, H., Samasti, M., 2007. Antibiotic resistance patterns of coagulase-negative
 575 staphylococcus strains isolated from blood cultures of septicemic patients in Turkey. Microbiol. Res.
 576 DOI:10.1016/j.micres.2007.03.004.
- 577 Kong, K.F., Vuong, C., Otto, M., 2006. *Staphylococcus quorum* sensing in biofilm formation
 578 and infection. Int. J. Med. Microbiol. 296, 133-139.
- 579 Koziol-Montewka, M., Szczepanik, A., Baranowicz, I., Jozwiak, L., Ksiazek, A., Kaczor, D.,
 580 2006. The investigation of *Staphylococcus aureus* and coagulase-negative staphylococci nasal
 581 carriage among patients undergoing haemodialysis. Microbiol. Res. 161, 281-287.
- 582 Kozitskaya, S., Olson, M.E., Fey, P.D., Witte, W., Ohlsen, K., Ziebuhr, W., 2005. Clonal
 583 analysis of *Staphylococcus epidermidis* isolates carrying or lacking biofilm-mediating genes by
 584 multilocus sequence typing. J. Clin. Microbiol. 43, 4751-4757.
- 585 Krediet, T.G., Mascini, E.M., van Rooij, E., Vlooswijk, J., Paauw, A., Gerards, L.J., Fleer,
 586 A., 2004. Molecular epidemiology of coagulase-negative staphylococci causing sepsis in a neonatal
 587 intensive care unit over an 11-year period. J. Clin. Microbiol. 42, 992-995.
- 588 Layer, F., Ghebremedhin, B., Moder, K., König, W., König, B., 2006. Comparative study
 589 using various methods for identification of *Staphylococcus* species in clinical specimens. J. Clin.
 590 Microbiol. 44, 2824-2830.
- 591 Mack, D. Davies, A.P., Harris, L.G., Rohde, H., Horstkotte, M.A., Knobloch, J.K., 2007.
 592 Microbial interactions in *Staphylococcus epidermidis* biofilms. Anal. Bioanal. Chem. 387, 399-408.
- 593 Mellmann, A., Becker, K., von Eiff, C., Keckevoet, U., Schumann, P., Harmsen, D., 2006.
 594 Sequencing and staphylococci identification. Emerg. Infect. Dis. 12, 333-336.
- 595 Miragaia, M., Couto, I., Pereira, S.F., Kristinsson, K.G., Westh, H., Jarløv, J.O., Carriço, J.,
 596 Almeida, J., Santos-Sanches, I., de Lencastre, H., 2002. Molecular characterization of methicillin-
 597 resistant *Staphylococcus epidermidis* clones: evidence of geographic dissemination. J. Clin.
 598 Microbiol. 40, 430-438.

- 599 Monsen, T., Olofsson, C., Rönmark, M., Wiström, J., 2000. Clonal spread of staphylococci
600 among patients with peritonitis associated with continuous ambulatory peritoneal dialysis. *Kidney*
601 *Int.* 57, 613-618.
- 602 Neely, A.N., Maley, M.P., 2000. Survival of Enterococci and Staphylococci on hospital
603 fabrics and plastic. *J. Clin. Microbiol.* 38, 724–726.
- 604 Nouwen, J.L., van Belkum, A., de Marie, S., Sluijs, J., Wielenga, J.J., Kluytmans, J.A.,
605 Verbrugh, H.A., 1998. Clonal expansion of *Staphylococcus epidermidis* strains causing Hickman
606 catheter-related infections in a hemato-oncologic department. *J. Clin. Microbiol.* 36, 2696-2702.
- 607 Nunes, A.P., Teixeira, L.M., Iorio, N.L., Bastos, C.C., de Sousa Fonseca, L., Souto-Pradón,
608 T., dos Santos, K.R., 2006. Heterogeneous resistance to vancomycin in *Staphylococcus epidermidis*,
609 *Staphylococcus haemolyticus* and *Staphylococcus warneri* clinical strains: characterisation of
610 glycopeptide susceptibility profiles and cell wall thickening. *Int. J. Antimicrob. Agents.* 27, 307-315.
- 611 Nys, S., van Merode, T., Bartelds, A.I., Stobberingh, E.E., 2006. Urinary tract infections in
612 general practice patients: diagnostic tests versus bacteriological culture. *J. Antimicrob. Chemother.*
613 57, 955-958.
- 614 Potoski, B.A., Adams, J., Clarke, L., Shutt, K., Linden, P.K., Baxter, C., Pasculle, A.W.,
615 Capitano, B., Peleg, A.Y., Szabo, D., Paterson, D.L., 2006. Epidemiological profile of linezolid-
616 resistant coagulase-negative staphylococci. *Clin. Infect. Dis.* 43, 165-171.
- 617 Poutanen, S.M., Baron, E.J., 2001. *Staphylococcus lugdunensis*: a notably distinct coagulase-
618 negative staphylococcus. *Clin. Microbiol. Newsletter.* 23, 147-150.
- 619 Qin, Z., Yang, X., Yang, L., Jiang, J., Ou, Y., Molin, S., Qu, D., 2007. Formation and
620 properties of in vitro biofilms of *ica*-negative *Staphylococcus epidermidis* clinical isolates. *J. Med.*
621 *Microbiol.* 56, 83-93.
- 622 Raz, R., Colodner, R., Kunin, C.M., 2005. Who Are You—*Staphylococcus saprophyticus*?
623 *Clin. Infect. Dis.* 40, 896-898.

- Reynolds, R., Potz, N., Colman, M., Williams, A., Livermore, D., MacGowan, A., 2004. Antimicrobial susceptibility of the pathogens of bacteraemia in the UK and Ireland 2001-2002: the BSAC bacteraemia resistance surveillance programme. *J. Antimicrob. Chemother.* 53, 1018-1032.
- Rohde, H., Kalitzky, M., Kröger, N., Scherpe, S., Horstkotte, M.A., Knobloch, J.K., Zander, A.R., Mack, D., 2004. Detection of virulence-associated genes not useful for discriminating between invasive and commensal *Staphylococcus epidermidis* strains from a bone marrow transplant unit. *J. Clin. Microbiol.* 42, 5614-5619.
- Rupp, M.E., Archer, G.L., 1994. Coagulase-negative staphylococci: pathogens associated with medical progress. *Clin. Infect. Dis.* 19, 231-245.
- Sader, H.S., Watters, A.A., Fritsche, T.R., Jones, R.N., 2007. Daptomycin antimicrobial activity tested against methicillin-resistant staphylococci and vancomycin-resistant enterococci isolated in European medical centers (2005). *BMC Infect. Dis.* 7, 29 DOI:10.1186/1471-2334-7-29.
- Schaberg, D.R., Culver, D.H., Gaynes, R.P., 1991. Major trends in the microbial etiology of nosocomial infection. *Am. J. Med.* 91, S72-S75.
- Singhal, R., Dhawan, S., Mohanty, S., Sood, S., Dhawan, B., Das, B., Kapil, A., 2006. Species distribution and antimicrobial susceptibility of coagulase negative staphylococci in a tertiary care hospital. *Indian J. Med. Res.* 123, 569-470.
- Sivadon, V., Rottman, M., Chaverot, S., Quincampoix, J.C., Avettand, V., de Mazancourt, P., Bernard, L., Trieu-Cuot, P., Féron, J.M., Lortat-Jacob, A., Piriou, P., Judet, T., Gaillard, J.L., 2005. Use of genotypic identification by *sodA* sequencing in a prospective study to examine the distribution of coagulase-negative *Staphylococcus* species among strains recovered during septic orthopedic surgery and evaluate their significance. *J. Clin. Microbiol.* 43, 2952-2954.
- Suetens, C., Leens, E., Versporten, A., Jans, B., 2002. Hogere incidentie septicemieën met coagulase-negatieve stafylokokken na introductie kwaliteitsdecreet in Vlaanderen. *Noso-info.* 1, 11-13.

- 649 Surani, S., Chandna, H., Weinstein, R.A., 1994. Breast abscess: coagulase-negative
650 staphylococcus as a sole pathogen. Clin. Infect. Dis. 19, 814-815.
- 651 Vandecasteele, S.J., Peetermans, W.E., Merckx, R., Rijnders B.J., Van Eldere, J., 2003.
652 Reliability of the *ica*, *aap* and *atlE* genes in the discrimination between invasive, colonizing and
653 contaminant *Staphylococcus epidermidis* isolates in the diagnosis of catheter-related infections. Clin.
654 Microbiol. Infect. 9, 114–119.
- 655 von Eiff C., Proctor, R.A., Peters, G., 2001. Coagulase-negative staphylococci: pathogens
656 have major role in nosocomial infections. Postgrad. Med. 110, 63-76.
- 657 von Eiff, C., Peters, G., Heilmann, C., 2002. Pathogenesis of infections due to coagulase
658 negative staphylococci. Lancet Infect. Dis. 2, 677–85.
- 659 von Eiff, C., Jansen, B., Kohnene, W., Becker, K., 2005. Infections associated with medical
660 devices, pathogenesis, management and prophylaxis. Drugs. 65, 179-214.
- 661 Vuong, C., Otto, M., 2002. *Staphylococcus epidermidis* infections. Microbes Infect. 4, 481-
662 489.
- 663 Wang, K.W., Chang, W.N., Shih, T.Y., Huang, C.R., Tsai, N.W., Chang, C.S., Chuang, Y.C.,
664 Liliang, P.C., Su, T.M., Rau, C.S., Tsai, Y.D., Cheng, B.C., Hung, P.L., Chang, C.J., Lu, C.H., 2004.
665 Infection of cerebrospinal fluid shunts: causative pathogens, clinical features and outcomes. Jpn. J.
666 Infect. Dis. 57, 44-48.
- 667 Wang, A., Athan, E., Pappas, P.A., Fowler, V.G., Olaison, L., Paré, C., Almirante, B.,
668 Muñoz, P., Rizzi, M., Naber, C., Logar, M., Tattevin, P., Iarussi, D.L., Selton-Suty, C., Jones, S.B.,
669 Casabé, J., Morris, A., Corey, G.R., Cabell, C.H., 2007. Contemporary clinical profile and outcome
670 of prosthetic valve endocarditis. JAMA. 12, 1354-1361.
- 671 Widerström, M., Monsen, T., Karlsson C., Wiström, J., 2006. Molecular epidemiology of
672 methicillin-resistant coagulase-negative staphylococci in a Swedish county hospital: evidence of
673 intra and interhospital clonal spread. J. Hosp. Infect. 64, 177-183.

- 674 Widerström, M., Wiström, J., Ferry, S., Karlsson, C., Monsen, T., 2007. Molecular
675 epidemiology of *Staphylococcus saprophyticus* isolated from women with uncomplicated
676 community-acquired urinary tract infection. *J. Clin. Microbiol.* 45, 1561-1564.
- 677 Wisplinghoff, H., Bischoff, T., Tallent, S.M., Seifert, H., Wenzel, R.P., Edmond, M.B., 2004.
678 Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective
679 nationwide surveillance study. *Clin. Infect. Dis.* 39, 309-17.
- 680 Xu, L., Li, H., Vuong, C., Vadyvaloo, V., Wang, J., Yao, Y., Otto, M., Gao, Q., 2006. Role
681 of the *luxS* quorum-sensing system in biofilm formation and virulence of *Staphylococcus*
682 *epidermidis*. *Infect. Immun.* 74, 488-496.
- 683 Zelenitsky, S., Barns, L., Findlay, I., Alfa, M., Ariano, R., Fine, A., Harding, G., 2000.
684 Analysis of microbiological trends in peritoneal dialysis-related peritonitis from 1991 to 1998. *Am.*
685 *J. Kidney Dis.* 36, 1009-1013.
- 686 Ziebuhr, W., Hennig, S., Eckart, M., Kränzler, H., Batzilla, C., Kozitskaya, S., 2006.
687 Nosocomial infections by *Staphylococcus epidermidis*: how a commensal bacterium turns into a
688 pathogen. *Int. J. Antimicrob. Agents.* 28S, S14-S20.

689 Table 1

690 Species distribution (%), sample origin, used identification method of CNS in clinical samples in different studies (country and publication date in
 691 heading).

	USA 1983 (1)	USA 1993 (2)	Denmark 1996 (3)	Japan 1998 (4)	Brazil 1999 (5)	Spain 2004 (6)	France 2005 (7)	India 2006 (8)	India 2007 (9)	Germany 2007 (10)	Turkey 2007 (11)
<i>S. epidermidis</i>	75	65	58	50	50	56	71	34	13	67	44
<i>S. haemolyticus</i>	7	13	9	16	10	5	2	13	72	12	12
<i>S. hominis</i>	5	7	12	5	12	18	4	2	6	9	9
<i>S. simulans</i>	1	2		6		1	1		2	1	2
<i>S. capitis</i>	2	4	<1	5		3	6	10	1	1	8
<i>S. saprophyticus</i>	1	1	1	5		5				<1	3
<i>S. warneri</i>	3	4	2	3		4	7	1	1	2	4
<i>S. lugdunensis</i>		3	<1	2			3	13	1	3	9
<i>S. caprae</i>				14			2			1	
<i>S. cohnii</i>	<1	1	1	<1	6	0	<1			1	<1
<i>S. auricularis</i>		<1		<1							<1
<i>S. schleiferi</i>								12		1	<1
<i>S. xylois</i>						0		6		1	5
<i>S. pasteurii</i>							2				
sample origin	CS	CS	BSI	CS	BSI		BJI	CS	CS	BSI	BSI
identification method	SP	Vitek1	SP	DNA	Microsc	SP	sodA	SP	Vitek2	SP + sodA	API

692 (1) Gill et al. (2) Kleeman et al. (3) Jarlöv et al. (4) Kawamura et al. (5) Del' Alamo et al. (6) Cuevas et al. (7) Sivadon et al. (8) Singhal et al. (9)
 693 Chaudhury et al. (10) Gattermann et al. (11) Koksai et al.; CS: clinical specimens, BJI: bone and joint infections; SP: standard procedures (based on
 694 Kloos and Schleifer), Vitek1: Vitek GP card (Biomérieux), *sodA*: *sodA* sequencing, DNA: DNA-DNA hybridization, Microsc: Microscan positive
 695 combo panel type 6 (DADE MicroScan Inc., Sacramento, CA), Vitek2: Vitek ID 32 Staph (Biomérieux), API: API ID 32 Staph (Biomérieux)

696

696 Table 2

697 Percentage susceptibility of CNS in different studies (in heading country and publication date); MR: % susceptibility in methicillin resistant strains, MS:

698 % susceptibility in methicillin susceptible strains

% susceptibility	Spain 2002 (1)	UK, Ireland 2004 (2)		Asia_Pacific region 2007 (3)		USA 2007 (4)	Europe, Asia, Australia, Latin America 2007 (5)	Turkey 2007 (6)		Europe, Turkey, Israel 2007 (7)	
	39	24		15,2		23,1	25,7	32,5		28	
		MR	MS	MR	MS			MR	MS	MR	MS
methicillin/oxacillin											
penicillin	21	0	24.5			12.1	15.3	0	9		
ceftriaxone						51.1	47.6				
ciprofloxacin	55	40.8	81.6			42.2	46.5	33	77	27.8	86.9
levofloxacin				50.2	89.1	42.3	47.5			28.1	87.3
cotrimoxazole	76			60.8	89.1	61.9	59.3	32	62	52.6	91.8
gentamicin	72	25.7	78.6			71.3	57.7	10	83		
tetracycline		39.2	67.3	83.5	89.1	86.9	83.7	40	55		
erythromycin	37	27.3	62.2	31.8	78.3	28.8	33.8	20	63	27.2	65.7
clindamycin	66	69.1	93.9	68.6	95.7	59.9	60.7	28	82	67	94.8
chloramphenicol	91			87.1	91.3			44	87	86.6	97.4
fusidic acid								75	85		
rifampicin	93									85.3	96.1
teicoplanin	100	65	77.6	96.1	89.1	99.3	99.3	100	100	97.3	99.6
vancomycin	100	99.7	100	100	100	100	100	100	100	100	100
linezolid	100	100	100	100	100	98.4	99.5			100	100
quinupristin/dalfopristin	100			99.6	100	99.3	99			99.3	100
daptomycin						100				100	99.6

699 (1) Cuevas et al. (2) Reynolds et al. (3) Biedenbach et al. (4) Jones et al., 2007a (5) Jones et al., 2007b (6) Koksall et al. (7) Sader et al.