Cryptogein, a fungal elicitor, remodels the phenylpropanoid metabolism of tobacco cell suspension cultures in a calcium-dependent manner.

Nicolas Amelot, Audrey Carrouche, Saïda Danoun, Stéphane Bourque, Jacques Haiech, Alain Pugin, Raoul Ranjeva, Jacqueline Grima Pettenati, Christian Mazars, Christian Brière

► To cite this version:

Nicolas Amelot, Audrey Carrouche, Saïda Danoun, Stéphane Bourque, Jacques Haiech, et al.. Cryptogein, a fungal elicitor, remodels the phenylpropanoid metabolism of tobacco cell suspension cultures in a calcium-dependent manner.. Plant, Cell and Environment, 2011, 34 (1), pp.149-161. 10.1111/j.1365-3040.2010.02233.x . hal-00532805

HAL Id: hal-00532805

<https://hal.science/hal-00532805>

Submitted on 4 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cryptogein, a fungal elicitor, remodels the phenylpropanoid metabolism of tobacco cell suspension cultures in a calcium-dependent manner.

Running title: Ca²⁺-dependent remodelling of phenylpropanoid metabolism

Nicolas Amelot¹, Audrey Carrouché¹, Saïda Danoun¹, Stéphane Bourque³, Jacques Haiech⁴, Alain Pugin³, Raoul Ranjeva², Jacqueline Grima-Pettenati², Christian Mazars², Christian Brière².

¹Université de Toulouse; UPS ; UMR 5546, Surfaces Cellulaires et Signalisation chez les Végétaux ; BP 42617, F-31326, Castanet-Tolosan, France

²CNRS ; UMR 5546 ; BP 42617, F-31326, Castanet-Tolosan, France

³UMR INRA 1088 / CNRS 5184 / Université de Bourgogne Plante-Microbe-Environnement, 17 Rue Sully, BP 86510, 21065 Dijon cedex, France;

⁴UMR CNRS 7200, Université de Strasbourg, Faculté de Pharmacie 74, route du Rhin, F-67401 Illkirch, France

Author for correspondence:

C. Brière

Pôle de Biotechnologie végétale, UMR 5546 ; BP 42617, F-31326, Castanet-Tolosan, France

Tel: +33(0)5 34 32 38 90

email: briere@scsv.ups-tlse.fr

ABSTRACT

Plant cells use calcium-based signalling pathways to transduce biotic and /or abiotic stimuli into adaptive responses. However, little is known about the coupling between calcium signalling, transcriptional regulation and the downstream biochemical processes. To understand these relationships better, we challenged tobacco BY-2 cells with cryptogein and evaluated how calcium transients (monitored through the calcium sensor aequorin) impact (i) transcript levels of phenylpropanoid genes (assessed by RT-qPCR) and (ii) derived-phenolic compounds (analysed by mass spectrometry). Most genes of the phenylpropanoid pathway were up-regulated by cryptogein and cell wall bound phenolic compounds accumulated (mainly 5-hydroxyferulic acid). Accumulation of both transcripts and phenolics was calcium-dependent. The transcriptional regulation of phenylpropanoid genes was correlated in a non-linear manner with stimulus intensity and with components of the cryptogein-induced calcium signature. In addition, calmodulin inhibitors increased the sensitivity of cells to low concentrations of cryptogein. These results led us to propose a model of coupling between the cryptogein signal, calcium signalling and the transcriptional response, exerting control of transcription through the coordinated action of two decoding modules exerting opposite effects.

1 **Keywords:** calcium signalling, transcription regulation, phenylpropanoid, modelling,
2 *Nicotiana tabacum* (tobacco BY-2), cryptogein

3 **Abbreviations**

4 PAL, phenylalanine ammonia-lyase; C4H, cinnamate 4-hydroxylase; 4CL, 4-coumarate-CoA
5 ligase; C3H, *p*-coumarate 3-hydroxylase; HCT, hydroxycinnamoyl CoA:quinate/shikimate
6 hydroxycinnamoyltransferase; CCoAOMT, caffeoyl CoA 3-*O*-methyltransferase; COMT
7 caffeic acid/5-hydroxyferulic acid *O*-methyltransferase; F5H, ferulate 5-hydroxylase; CCR,
8 cinnamoyl CoA reductase; CAD, cinnamylalcohol dehydrogenase.

9

1 INTRODUCTION

2 Plant cells use calcium-based signalling pathways to transduce a variety of external and/or
3 internal stimuli into adaptive responses (for review, Hetherington & Brownlee 2004, Kudla,
4 Batistic & Hashimoto 2010, Lecourieux, Ranjeva & Pugin 2006, McAinsh & Pittman 2009,
5 Sudha & Ravishankar 2002, White & Broadley 2003). A paradigm establishing the central
6 role of calcium in a complex regulatory network with multiple interacting components has
7 been inferred, at the single cell level, from the multifaceted approach used to study guard-cell
8 physiology (Allen *et al.* 2001a, Allen & Schroeder 2001b, Israelsson *et al.* 2006, Schroeder *et*
9 *al.* 2001). As a general feature, in response to variations of abiotic (light, temperature,
10 drought, etc.) and/or biotic (symbionts, pathogens, etc.) environment including stressful
11 conditions that plants have to face, free calcium concentration varies in cells (Lecourieux *et*
12 *al.* 2002, Pauly *et al.* 2000, Ward, Pei & Schroeder 1995, Xiong *et al.* 2007). These changes
13 control basic cellular functions including ion and molecule transport and most probably gene
14 expression (Kaplan *et al.* 2006, Kim & Kim 2006, Lecourieux *et al.* 2002, Moscatiello *et al.*
15 2006). Indeed, the identification of transcriptional regulators modulated by calcium and
16 calmodulin (Kim *et al.* 2009) supports the importance of calcium in the transcriptional
17 process during plant responses to endogenous and exogenous stimuli.

18 Also interesting, is the evidence that the calcium signature itself and some of its components
19 (e.g. amplitude, duration) are correlated to gene expression. For instance, during the
20 establishment of symbiosis in root hairs Miwa *et al.* (2006) showed a correlation between the
21 number of Ca^{2+} spikes and the transcript level of the Early Nodulation 11 (ENOD11) gene.
22 They suggested that a minimum of 36 spikes is required for *ENOD11* induction. Similarly, the
23 shape of the calcium signal in response to salt stress was shown to be very important for the
24 control of salt-stress genes. Using the Ca^{2+} -ATPase loss of function mutant in the moss
25 *Physcomitrella patens*, a sustained elevation of free calcium was observed instead of a

transient elevation. This sustained level of calcium down-regulates salt-stress-induced genes rendering the mutant plants more susceptible to salt stress (Qudeimat *et al.* 2008). These examples illustrate how the calcium-dependent processes underlying the calcium signature can control gene expression. It may thus be hypothesized that calcium, as a signal, interferes with the various steps involved in the conversion of a stimulus into a biological output. Surprisingly little is known about the relationships linking calcium to transcriptional regulation, protein synthesis and enzyme activities, and *in fine* the remodelling of the biochemical processes downstream (Kudla *et al.*, 2010).

For a better understanding of the connections between these different events and to identify critical regulatory steps (Weckwerth 2003) or signalling hubs, we took advantage of the biological model consisting of tobacco cell suspension cultures challenged with cryptogein. Cryptogein, a polypeptide elicitor secreted by the oomycete *Phytophthora cryptogea*, induces free calcium changes in the cytosol ($\Delta[\text{Ca}^{2+}]_{\text{cyt}}$) in tobacco cells and enhances the transcription of the gene encoding phenylalanine ammonia lyase (PAL), the enzyme catalysing the first committed step of the phenylpropanoid pathway (Lecourieux *et al.* 2002). Although this biological model has been extensively studied at different levels of the signalling pathway (Garcia-Brügger *et al.* 2006), it is still not known if other genes of the multi-branched pathway are differentially regulated in response to cryptogein, and there are no data showing changes in enzyme activities or increases in the amounts/concentrations of phenolic compounds.

In the present paper, we made use of this simplified biological system to evaluate qualitative and quantitative changes in successive steps of elicitor-induced signalling pathways, including calcium-transient generation, transcriptional regulation and metabolism reorientation. Our work aimed at bridging the different steps involved in the signalling pathway that converts the initial perception of cryptogein (input) into the final response (output) that

consists here of the changes induced in the metabolism of phenolic compounds (Fig. 1). We reasoned that quantitative parameters and time-dependence are essential to get a deeper insight into the regulatory processes and may shed light on overlooked properties. In this way, we have addressed the following questions:

- How do free calcium concentrations vary with time when cells are challenged with increasing concentrations of cryptogein?
- Are all genes coding for the enzymes of the phenylpropanoid pathway induced by cryptogein in a calcium-dependent manner?
- How does cryptogein remodel phenylpropanoid metabolism?

On these grounds we propose an integrated model that could explain key features of the input/output network linking cryptogein signal perception to gene expression through calcium signalling.

MATERIAL AND METHODS

Products

Cryptogein was purified according to Bonnet *et al.* (1996) and dissolved in water as a 100 μ M stock solution. [3 H]-S-adenosyl-L-methionine (specific activity 0.37 TBq/mmol) was from MP Biochemicals (Illkirch, France) and [14 C(U)]-L-phenylalanine (specific activity 6.8 GBq/mmol) was from Perkin-Elmer (France). A 5 mM stock solution of native coelentarazine (Interchim, Montluçon, France) in methanol was stored at -20°C in 10 μ L aliquots. All other compounds were from Sigma-Aldrich.

Radiolabelled trans-cinnamic acid was obtained by the enzymatic conversion of 0.185 MBq of [14 C(U)]-L-phenylalanine by 0.007 units of phenylalanine ammonia lyase (Sigma-Aldrich) in Tris-HCl buffer pH 8.5 at 30°C for 2 hours. After stopping the reaction with 6 μ L 6M HCl,

cinnamic acid was extracted in 2x1 mL diethyl ether, evaporated to dryness and taken up in 50µL MeOH.

Cell culture and luminescence measurements

Transgenic tobacco cells (*Nicotiana tabacum* L. cv Bright Yellow BY-2 lines) expressing cytosolic apo-aequorin (Takahashi *et al.* 1997) were grown and processed according to Pauly *et al.* (2001). Cells harvested at late exponential growth phase were washed twice and resuspended at a density of 20% (Packed Cell Volume, PCV) in the suspension buffer (175 mM mannitol, 0.5 mM K₂SO₄, 0.5 mM CaCl₂, 2 mM MES KOH, pH 5.8), supplemented with 2 µM coelenterazine to reconstitute active aequorin for luminescence measurements. Subsequently, cells were incubated at room temperature overnight in the dark on a rotary shaker (130 rpm). Measurements of luminescence and calcium calibration were carried out as previously described (Pauly *et al.* 2001).

Elicitor treatments

Cells in suspension medium were transferred into a glass vial and challenged with the indicated concentrations of effectors. After given times of incubation, cell aliquots were collected and filtered under vacuum on a 37 µm mesh nylon membrane and immediately frozen in liquid nitrogen.

Quantitative real-time RT-PCR

Tobacco BY-2 cell samples were ground in liquid nitrogen and total RNAs isolated using the Nucleospin II RNA kit (Macherey-Nagel, Düren, Germany) according to the manufacturer's instructions. First-strand cDNA was synthesized from 5µg total RNA in a 20µl reaction volume containing 50ng random primers, 10 mM dNTPs, 1µl RNasOUT (40 units) and 200 units of SuperScript III reverse transcriptase, according to the manufacturer's instructions (Invitrogen, Carlsbad, USA). After 50 minutes at 50°C followed by 5 minutes at 85°C, cDNAs were purified using the PCR Clean-Up system (Promega, Madison, USA). Two µL

cDNAs were used as templates in a quantitative real-time PCR assay (10 μ L) performed on the sequence detection system ABI PRISM 7900HT (Applied Biosystems, Foster City, USA) using the SYBR Green PCR Master Mix (Applied Biosystems). After an initial denaturation of 10 minutes at 95°C, 40 cycles of 15 sec at 95°C followed by 1 minute at 60°C were performed. Amplification specificity was checked by melting-curve analysis, and PCR efficiency was determined using standard curves constructed with serial dilutions of PCR products as templates. 18S rRNA was used as an internal control. Quantification of expression ratios was performed according to the mathematical model developed by Pfaffl (2001). Primers and amplicon sizes are given in Supporting Information (table S1). In the case of multigene families, primers have been designed to hybridize either to all known genes of the class postulated to have a role in lignin biosynthesis and/or specifically to one gene member (table S1).

Measurement of enzymatic activities of PAL and CCoAOMT

Cell samples were ground in liquid nitrogen and weighed (approx. 2g/sample) before being extracted for 30 minutes in 4 mL of protein extraction buffer (2 mM MgCl_2 , 10 mM DDT, antiprotease (P9599 Sigma), 10% glycerol, 0.1M Tris-HCl pH 7.5) at 4°C under gentle shaking. After a 30 minute centrifugation (10 000 x g, 4°C) the supernatant was collected and desalted on a PD10 column equilibrated with the same buffer at pH 7.5 for CCoAOMT activity or pH 8.8 for PAL activity measurements. PAL activity was assayed by spectrophotometry. The reaction mixture (1 mL) contained 50 μ g protein extract and 1 mM L-phenylalanine (Sigma) in 0.1M Tris-HCl, pH 8.8 buffer. The formation of cinnamic acid was monitored for 10 minutes by measuring absorption at 290 nm and PAL activity was determined from the slope of the absorption curve using the extinction coefficient of cinnamic acid at 290 nm = 10000 $\text{M}^{-1}\text{cm}^{-1}$

CCoAOMT activity was determined by radiometric assay. The reaction mixture contained 10 μ L of protein extract, 3.7kBq [3 H]-S-adenosyl-L-methionine, 40 μ M S-adenosyl-L-methionine, 40 μ M caffeoyl-CoA in 0.1M Tris-HCl, pH 7.5 (total volume 50 μ L). After 30 minutes of incubation at 30°C the reaction was stopped with 10 μ L 3M NaOH and incubated for a further 10 minutes at 37°C to allow feruloyl-CoA hydrolysis before adding 40 μ L 1M HCl to stop the reaction. Ferulic acid was extracted 10 minutes in 200 μ L of hexane/ethyl acetate (1:1 v/v) and 150 μ L of the organic phase was collected and evaporated overnight before addition of 8 mL of scintillation cocktail for radioactivity counting.

Radioactivity measurements

[14 C]-cinnamic acid was added to the cells (0.007MBq. g $^{-1}$ fresh cells) immediately before treatment with cryptogein. After 4 hours of incubation at 20°C in the dark and under shaking, cell aliquots were filtered under vacuum and frozen immediately. A 1 mL aliquot of filtrate was collected and transferred into a scintillation vial containing 8 mL of Ready Safe cocktail (Beckman Instr., Fullerton CA USA) to measure residual radioactivity in the medium. Frozen cell samples were weighed before grinding in 80% EtOH with glass beads (Fastprep system, MP Biomedicals, Illkirch, France). Ground cells were then extracted for 30 minutes at 4°C in 5 mL 80% EtOH under gentle shaking, and centrifuged 10 minutes at 3000 x g. A 1 mL aliquot of supernatant was collected for measuring the radioactivity of ethanol-soluble compounds (total radioactivity). Another 1 mL aliquot of the supernatant was added to a PVPP column, which selectively complexes phenolic compounds (radioactive soluble phenolics). Radioactive non-phenolic products were recovered in the pass through (radioactive soluble non-phenolics) and the distribution of the radiolabel was calculated as: radioactive soluble phenolics = total radioactivity – radioactive non-phenolics. The remaining supernatant was discarded and the cell pellet resuspended in water, and a 200 μ L aliquot collected for counting the radioactivity incorporated into the non-soluble fraction.

1 Extraction and analysis of soluble and wall-bound phenolics

2 Frozen cell samples were ground in liquid nitrogen, and the powder weighed before being
3 extracted in 5 mL 80% EtOH for 30 minutes at 4°C under gentle shaking in the dark. After 10
4 minute centrifugation at 3000 x g the supernatant was collected for analysis of soluble
5 phenolics. Phenolic material contained in the ethanol-insoluble fraction was extracted
6 according to Bolwell *et al.* (1985). The pellets were resuspended in 5 mL of 4.25M NaOH and
7 hydrolyzed for 6 days at 4°C followed by 1 day at 37°C in the dark under gentle shaking.
8 After 15 minute centrifugation at 4500 x g the supernatant was collected, acidified to pH 1
9 with 12M HCl, then extracted twice in ethyl acetate. The organic phase was collected, and
10 evaporated to dryness under N₂. The residue was re-dissolved in 100 µL of water/methanol
11 (1/1 v/v) for analysis by UPLC-QTOF using a Waters ACQUITY UPLC System (Waters,
12 Milford MA, USA). Separation was achieved using an ACQUITY UPLC BEH C18 column
13 (100 mm, 2.1 mm i.d., 1.7 µm particle size; Waters) maintained at 40°C, with a mobile phase
14 flow rate of 0.30 mL.min⁻¹ in isocratic mode. The mobile phase contained 32 % methanol and
15 0.1 % formic acid in water. The injection volume was 2 µL with partial loop. A time-of-flight
16 mass spectrometer with an electrospray ionization (ESI-MS) interface was used for mass
17 spectrometry analysis (Micromass, Manchester, UK). Data acquisition was performed using
18 negative ion mode over a m/z range 50–1000. The general conditions were: source
19 temperature 130 °C, capillary voltage -2.8 kV and cone voltage 40 V, extraction cone -4 V,
20 ion guide -1 V, the collision energy 5 eV, desolvation temperature 350 °C, desolvation
21 flow 750 L.h⁻¹, the nebulizer 30 L.h⁻¹ and data were acquired using a scan time of 0.5 s.
22 Leucine-enkephalin (M-H)⁻ (m/z 554.2615) was used for the lock mass of the lock spray.
23 MassLynx 4.1 software (Waters, Manchester, UK) was used for data analysis.

Phenolic compound identification was based on comparison of retention times of commercial standards, accurate mass measurements, and MS/MS fragmentation analysis (collision energy: 15 eV).

Statistical analysis

Statistical tests and multivariate analysis (Principal Component Analysis) were carried out using the R statistical software (R Development Core Team 2009).

RESULTS

Cryptogein induces a biphasic calcium signature in tobacco BY-2 cells

Cryptogein induces biphasic calcium variations in the cytosol of BY2 cells (Fig. 2a). This response is characterized by an increase of free cytosolic calcium $\Delta[\text{Ca}^{2+}]_{\text{cyt}}$ starting as early as 2 minutes after elicitation, reaching a maximum after about 5 min, followed by a sustained elevation (plateau) of cytosolic free calcium lasting for more than an hour. This cytosolic calcium variation was inhibited in cells co-incubated with 1 mM La^{3+} , illustrating the absolute requirement of an external calcium influx. The maximum calcium elevation (peak value) and the plateau value increased in a dose-dependent manner but with different EC_{50} values (Fig. 2b). Thus, the maximum plateau value is reached for very low cryptogein concentrations (less than 100 nM), while the peak value continues to increase for cryptogein concentrations above 1 μM . Fitting a simple one-site binding model to experimental data enabled us to estimate the EC_{50} values at 128 nM cryptogein for the calcium peak ($\text{B}_{\text{max}} = 0.9 \mu\text{M}$) and to 17 nM cryptogein for the calcium plateau ($\text{B}_{\text{max}} = 0.1 \mu\text{M}$), a value close to the measured Kd of cryptogein binding to cell membranes (Bourque *et al.* 1999). This suggests either the existence of two cryptogein receptors, with low and high affinities, and involved in the generation of the calcium-peak or the sustained plateau respectively, or the existence of only

one type of cryptogein receptor, but interacting with two types of protein complexes with different affinities.

Cryptogein strongly increased the number of transcripts of genes of the common phenylpropanoid biosynthetic pathway

We measured the transcript levels of genes encoding enzymes involved in the synthesis of phenylpropanoids and lignin (*PAL*, *C4H*, *4CL*, *HCT*, *C3H*, *COMT*, *CCoAOMT*, *F5H*, *CCR* and *CAD*), as well as of three genes known to be involved in defence responses: *HSR203J*, a cell death response marker (Pontier *et al.* 1994), and two PR genes, *PR1* and *PR3* (Ward *et al.* 1991). Transcripts levels in tobacco BY-2 cells were measured using RT-qPCR after treatments of cells with various concentrations of cryptogein (from 10 nM to 1 μ M) and different durations (from 30 minutes to 8 hours). The ratios of transcripts levels in treated to that in control cells are presented in Fig. 3.

After a cryptogein treatment of 3 hr, the transcript levels of the majority of the genes tested were strongly enhanced, but to different extent, ranging from 4-fold for *C4H* to about 200-fold for *CCoAOMT* and *F5H*. In the core pathway, *PAL*, *C4H* and *4CL* transcript accumulation was increased on average by 26-, 4- and 15-fold respectively. In contrast transcript accumulation of *CAD*, the last gene of the monolignol biosynthesis pathway, was decreased by about 2-fold. Interestingly, cryptogein had no effect on transcription of the two defence marker genes (*PR1* and *PR3*) while the expression of the cell death response marker *HSR203J* was strongly up-regulated (55-fold on average).

Transcript accumulation of tested genes was strongly inhibited when cells were challenged with cryptogein in the presence of 1 mM La^{3+} , a calcium channel blocker (Fig. 3a), or in the presence of the calcium chelator EGTA (Supplemental information S2), showing that calcium influx was required for the transcriptional response.

1 *De novo* synthesis of proteins was also required to allow the transcriptional response to
2 cryptogein, as shown by its complete suppression by a 30 minute pre-treatment of the cells
3 with cycloheximide prior to the addition of cryptogein (Supporting Information Fig. S3).

4 Because many genes of the phenylpropanoid pathway belong to small gene families, we
5 examined transcript accumulation in response to cryptogein for each known isoform of three
6 strongly induced genes. For PAL and 4CL, only the expression of one gene member was up-
7 regulated, PAL A and 4CL2, respectively. Transcript levels of *CCoAOMT* genes were highly
8 and equally up-regulated upon cryptogein treatment except *CCoAOMT5* which was only
9 weakly activated in comparison to the other family members (Supporting Information Fig.
10 S4). These data show that the intensity of the transcriptional response within a gene family
11 may be specific either for a particular gene or for a set of genes.

12 The magnitude of the transcriptional response to cryptogein increased with time and depended
13 upon the intensity of the stimulus. As exemplified with *PAL*, *CCoAOMT* and *HCT*, the
14 transcript levels of the most activated genes increased after the cryptogein treatment as early
15 as 30 minute and continued increasing for several hours (Fig. 3b). Different induction kinetics
16 patterns were observed depending on the gene considered. Transcripts levels of some genes
17 like *CCoAOMT* reached a maximum after 4 hours of treatment then decreased, while
18 transcripts of other genes like *COMT* and *HSR203J* continued to accumulate for 8 hours.

19 Furthermore, the intensity of the transcriptional response was strongly dependent upon the
20 intensity of the stimulus (Fig. 3c). Similar dose-response curves were observed for the
21 different genes of the phenylpropanoid pathway, showing a strong correlation between these
22 genes in terms of dose dependence. Dose-response curves were non-linear and biphasic: the
23 effect of cryptogein began to be significant at 20 nM, reached a maximum between 50 nM
24 and 100 nM (depending on the gene considered), and then decreased fairly sharply while

1 increasing the cryptogein concentration (Fig. 3c). As observed for the kinetics patterns,
2 differences in the dose-response curves were observed between genes.

3 Thus, in order to analyse the correlations between gene transcript ratios in dose-response
4 experiments, we performed a Principal Component Analysis (PCA) of the correlations
5 between kinetic data obtained for all studied genes. Briefly, this multivariate analysis allows
6 representing graphically the correlations between variables and/or the proximities between
7 observations on planes with a minimal loss of information. “Calcium-peak” and “Calcium-
8 plateau” values measured at different concentrations of cryptogein were included in this
9 analysis in order to analyze the correlations between transcript ratios and these two
10 characteristics of the calcium signature. This analysis led to the definition of two groups of
11 up-regulated genes in addition to the group formed by *CAD*, *PR1* and *PR3* that are not up- or
12 even down-regulated (Fig. 4). The first group (*4CL*, *C4H*, *C3H*, *COMT*, *F5H*, *CCR*) is
13 correlated with the “Ca-peak” variable and is colocalized with observations made at 100nM,
14 200nM and 1µM cryptogein., It corresponds to genes whose transcript accumulation reached
15 a maximum for cryptogein concentrations above 100 nM. The second group (*PAL*, *HCT*,
16 *CCoAOMT*, and *HSR203J*), correlated to the “Ca-plateau” variable, corresponds to genes
17 whose activation was maximal at 60 nM cryptogein and then decreased for higher
18 concentrations.

20 **Cryptogein-induced enhancement of transcription correlates with increased** 21 **activities of enzymes of the phenylpropanoid pathway**

22 In order to evaluate whether transcriptional activation was correlated with the activation of the
23 corresponding enzymes, we assayed the enzymatic activities of PAL and CCoAOMT whose
24 corresponding genes were among the most activated. These activities were strongly increased
25 (2.5x for PAL, 6x for CCoAOMT) in BY2 cells challenged with 0.1µM cryptogein for 4

hours as compared to control cells (Fig. 5). These increases were totally inhibited when the cells were treated by 1mM La^{3+} (Fig. 5 inset). A kinetic study showed that the cryptogein-induced increase of these enzyme activities began as early as 30 minute after elicitation for CCoAOMT (2 hours for PAL) and increased continuously during the whole experiment (8 hours). This continuous increase is consistent with the increases in transcript levels of the corresponding genes (Fig. 5), indicating that cryptogein-induced enhancement of transcript accumulation resulted in an increase of enzymatic activities.

Cryptogein induces accumulation of cell-wall associated phenolic compounds

From the results presented above, we reasoned that the distribution of phenolics (metabolic remodelling) might be significantly altered in cells challenged with cryptogein. To address this question, cells were fed with [$^{14}\text{C}(\text{U})$]-cinnamic acid and incubated with cryptogein. After 4 hours of incubation, the cell suspension was filtered and radioactivity was measured in different fractions: external medium, phenolic soluble fraction, non-phenolic soluble fraction, and insoluble fraction (containing insoluble phenolics). Compared to control cells, cryptogein-treated cells showed a slight decrease of residual radioactivity in the external medium and in the soluble fraction containing non-phenolic compounds (including radiolabelled free cinnamic acid), but an increase in the insoluble fraction (Fig. 6). The variations of radioactivity in the different fractions were inhibited by 1mM La^{3+} suggesting that, in line with what we showed for gene expression and enzymatic activities, the putative remodelling of phenolic metabolism is a calcium-dependent process. The decrease of radioactivity observed in the soluble non-phenolic fraction and the concomitant increase measured in the insoluble fraction in response to cryptogein strongly suggest an accumulation of phenolics in the cell walls.

In order to identify these insoluble phenolic compounds, cell wall pellets were hydrolyzed with NaOH in order to release associated esterified phenolics that were further analyzed by

MS/MS. Intermediate compounds of the “core” phenylpropanoid pathway were found to be significantly increased in cryptogein-treated samples as compared to controls *i.e.*: *p*-coumaric acid, ferulic acid, and 5-hydroxy-ferulic acid which was found only as traces in control samples but at a high level in cryptogein-treated cells. In addition, a small increase in 4-hydroxybenzaldehyde was also observed in treated cells (table 1).

Calcium controls the cryptogein-induced response

The data reported above are consistent with the existence of non-linear relationships between different steps of cryptogein signalling in tobacco cells and highlight the crucial role of the steps associated with the generation of the calcium signature. They further suggest the possible co-existence of regulatory modules differing in their sensitivity to calcium and/or effector concentration, and able to modulate the response at the transcriptional level. These results raise questions on the respective roles of the two phases of the cryptogein-induced calcium signature and on the potential involvement of calcium sensor proteins.

To address these questions, we disturbed the cryptogein-induced response either by modulating the calcium response at a key step of the time course, or by using inhibitors of calcium-sensor proteins and we further investigated the effects of these perturbations on the transcript levels of representative genes.

Sustained calcium elevation plays a key role in the cryptogein-induced response

In order to determine the relative importance of the transient peak and of the sustained phase of calcium elevation (plateau) in cryptogein-induced transcriptional regulation, we applied 1mM La³⁺ either just before the cryptogein treatment, thus suppressing the whole calcium response (light grey curve, Fig. 7a), or 10 minutes after cryptogein treatment (arrow) in order to maintain the first peak while suppressing the sustained plateau (dark grey curve, Fig. 7a). After a 3-hour treatment, we measured the accumulated level of transcripts of phenylpropanoid genes in each condition (Fig. 7b).

Adding lanthanum just before or 10 minutes after the cryptogein treatment resulted in both cases in a strong reduction of transcript levels of *PAL*, *HCT* or *CCoAOMT* (Fig. 7b) as compared to cryptogein treatment alone (see also Fig 3a). Furthermore no significant difference ($p=0.05$) was observed between the inhibitory effects of the two treatments with lanthanum. In contrast, cryptogein-induced accumulation of transcripts of *C4H*, *4CL*, *C3H*, *COMT* and *CCR* was only slightly modified by addition of lanthanum after 10 min, suggesting that transcriptional regulation of these genes is mainly dependent of the early calcium influx. It is noteworthy that these two subsets of genes are identical to those identified by the PCA analysis, suggesting a possible link between the intensity of the stimulus (PCA analysis; figure 4), the two different phases of the calcium response (figure 7) and gene expression.

Regulators with low and high affinity for calcium control the transcriptional response

To assess the possibility of regulatory effects downstream of the calcium signal, we evaluated the effects of compounds able to interact with calmodulins or CaM-like proteins on cryptogein-induced responses. We used W-13 known as a calmodulin inhibitor ($K_d = 50 \mu\text{M}$) and W-12 (a molecule of the same family as W-13 with a H/Cl substitution and a lower affinity for CaM, $K_d = 250 \mu\text{M}$) as negative control.

By themselves, these drugs provoked changes in cellular calcium (Dagher *et al.* 2008) but in a differential manner (Supporting Information Fig. S4). Within two minutes, the calmodulin inhibitor W-13 induced a transient dose-dependent increase in cytosolic calcium, followed by a rapid decrease to the resting level. In contrast, W-12 here considered as the negative counterpart of W-13, induced significantly lower calcium variations in the cytosol. In subsequent experiments these two compounds were used at a final concentration of $50 \mu\text{M}$ for which the calcium level returned to the basal level within 10 minutes. In order to investigate the effect of these two drugs either on the calcium response or on the transcriptional response

induced by cryptogein, W-13 or W-12 were added to the cell suspension 10 minutes before cryptogein treatment. Then, cytosolic calcium variations were recorded for 15 min, and the cells were collected after 4 hours for transcriptional analysis. The calcium response induced by high concentrations of cryptogein (0.25 μ M and above) was not modified by either inhibitor, but when cryptogein was added at low concentrations (< 50 nM) the sustained calcium elevation (plateau value), and to a lesser extent the peak value, were significantly increased by W13 (Fig. 8a), but not by W12 (not shown). When added alone, these compounds had no effect on the transcript level of phenylpropanoid genes (not shown). When added before cryptogein-treatment, they had no effect on the transcript increase induced by concentrations of cryptogein above 100 nM, but a strong enhancement of the cryptogein effect was mediated by the CaM inhibitors at low concentrations of cryptogein i.e. 25 nM (Fig. 8b). As expected, this effect was higher with the calmodulin inhibitor W-13 than with its negative counterpart W-12.

These results confirm the importance of the plateau phase of the calcium signature in the increase of transcript levels induced by cryptogein. Furthermore, the differential effect of anti-CaM compounds, between low and high cryptogein treatments, is consistent with the hypothesis that two calcium decoding modules, one with low affinity and the other with high affinity for calcium, are involved in the regulation of this response.

Modelling the coupling between cryptogein-induced calcium and transcriptional responses

The results presented above show the existence of correlations between the different phases of the cryptogein-induced calcium signature and the transcriptional response: the calcium-plateau value correlates positively with transcript accumulation, while an increase of the calcium-peak value above a threshold correlates negatively with transcript accumulation. Furthermore they support the hypothesis of two decoding modules (involving different

1 calcium sensors) able to modulate the response to cryptogein and which could explain the
2 observed correlations.

3 To test this hypothesis and verify its coherence with the experimental data we used computer
4 modelling and built a model of the non-linear system coupling the cryptogein signal to
5 calcium and transcriptional regulation. The model can be described as follows (see scheme
6 Fig. 9).

7 Perception of cryptogein triggers opening of calcium channels, which results in an influx of
8 external calcium. This calcium influx activates calcium-dependent mechanisms that depends
9 upon the strength of the stimulus (= cryptogein concentration). At low concentrations of
10 cryptogein (10 nM – 50 nM), the concerted activation of calcium channels and transporters
11 results in a small but sustained elevation of free cytosolic calcium concentration (=plateau
12 value). This increase (or influx) is nevertheless sufficient to activate calcium-sensors
13 belonging to a high affinity decoding module (module 1), which in turn will enhance
14 transcription of response genes. For higher cryptogein concentrations (> 50 nM), the larger
15 calcium influx triggers the release of calcium from internal stores and/or the opening of
16 additional plasma-membrane calcium channels, resulting in a large transient increase of
17 cytosolic calcium (=peak value) followed by a slow return to the plateau value through the
18 action of homeostatic mechanisms. This has two consequences: 1) the activation of high
19 affinity module 1 due to sustained calcium elevation, and thus differential expression of
20 responsive genes, and 2) the activation of other calcium responsive effectors of a low-affinity
21 module (module 2), which have an inhibitory effect on the transcription of the same genes.
22 This results in equilibrium between activation-inhibition mechanisms through the activation
23 of these two modules, linking the transcriptional response to the cryptogein concentration in a
24 non-linear manner.

We used a simple phenomenological approach to formalize this model using a two-component system derived from chemical binding kinetics (Supporting Information S7). In this model each component (modules 1 and 2) is characterized by an “apparent” affinity constant for cryptogein (K1 and K2 resp.), a parameter defining its maximum activity (S1 and S2 resp.), and a Hill coefficient (n1 and n2 resp.) to account for cooperativity. This model integrates, in a simple way, all the steps between cryptogein binding and transcriptional activation, and thus its parameters depend implicitly on the characteristics of the calcium decoding modules. This model fits well to experimental dose-response curves obtained for transcription of *PAL*, *HCT*, and *CCoAOMT* (Fig.10a). Interestingly, the K values for cryptogein estimated for each component were both very close to the K_m value of the dose-response curve observed for the calcium plateau (30 nM), which is in the range of IC_{50} values measured for cryptogein binding to its receptor. In order to test the model further, we used it to fit the effect of calmodulin inhibitors on cryptogein-induced transcription. A good fit to experimental curves was obtained (Fig.10b) by increasing the affinity parameter K1 that corresponds to the “apparent” affinity of the “activation” module (the other parameters being held constant). Interestingly, the K1 value estimated for the W13 experimental curve was greater than its value estimated for W12.

The hypothetical mechanisms at the basis of this simple model seem to be sufficient not only to fit dose-response curves but also the effect of anti-calmodulin compounds on the transcriptional response to cryptogein. It is worth noting that this model was built on assumptions derived from observed correlations between calcium and expression data only. However, by modifying only two parameter values, the model was able to simulate (at least qualitatively) the effect of anti-calmodulin compounds. Considering that data from anti-CaM experiments were not used to derive the model, this last result is in some extent a first step in the validation of this model.

DISCUSSION

In this work our aim was to build an integrated view of the relationships linking the calcium response to the short-term responses induced at the transcriptional and metabolic levels by the necrotizing elicitor cryptogein in BY-2 cells. For this purpose, we first carried out a quantitative analysis of the biological responses at different levels, and then exploited these data to propose a simple coupling model between calcium variations and transcriptional regulation of genes of the phenylpropanoid pathway.

Cryptogein increases transcription of most genes of the phenylpropanoid pathway

In BY-2 cells cryptogein induced an increase of transcript levels for most of the genes of the phenylpropanoid pathway, with expression ratios ranging from 4 (e.g. *C4H*) to more than 100 (e.g. *CCoAOMT*), but with the noticeable exception of *CAD*, which catalyses the last step leading to lignin monomers, that was not induced or was even repressed for high concentrations of elicitor. These transcriptional responses occurred early after elicitation (less than 30 minutes for the most activated gene *CCoAOMT*) and transcript accumulation continued increasing for several hours, reaching a maximum after 4 hours for *PAL*, *HCT* and *CCoAOMT*. For other activated phenylpropanoid genes like *COMT*, their transcripts continued to accumulate for at least 8 hours after treatment. Depending on the gene, these responses were totally or partially, suppressed when calcium influx was blocked by lanthanum. This is consistent with the calcium-dependence of the biological responses to cryptogein shown in previous works (Lecourieux *et al.* 2002).

An increased accumulation of transcripts of genes coding for key enzymes of the phenylpropanoid pathway (*PAL*, *4CL*, *CCoAOMT*, *C4H*) in response to elicitors has been previously reported in various plant species (Dixon & Paiva 1995, Hano *et al.* 2006, Kauss *et*

1 *al.* 1993, Shinya *et al.* 2007, Suzuki *et al.* 2003). In particular, cryptogein activates
2 transcription of the phenylalanine-ammonia lyase (PAL) gene in tobacco cells (Lecourieux *et*
3 *al.* 2002, Suty *et al.* 1995). In this work, we extended the study to a larger set of genes
4 belonging to the phenylpropanoid pathway and quantified corresponding transcript levels.
5 Transcription of *HSR203J*, a marker of hypersensitive response in *tobacco* (Pontier *et al.*
6 1994), was strongly increased by the cryptogein treatment as previously shown (Lecourieux *et*
7 *al.* 2002). In contrast *PR1* and *PR3*, two genes involved in defence responses through the
8 salicylic acid pathway (Keller *et al.* 1996a, Ward *et al.* 1991), were not differentially
9 expressed in BY-2 cells. This is in agreement with the known biological effect of cryptogein
10 in tobacco, which induces cell death through a hypersensitive response, while the salicylic
11 pathway has been shown not to be involved in the HR response (Keller *et al.*, 1996).

12 **Phenolic metabolism changes in response to cryptogein**

13 In tobacco BY-2 cells, cryptogein induced an early accumulation of cell-wall bound phenolic
14 compounds (*p*-coumaric acid, ferulic acid, 5-hydroxyferulic acid, 4-hydroxybenzaldehyde),
15 whereas no consistent variation of free soluble phenolics was detected after a 4-hour
16 treatment. The accumulation of cell-wall-bound phenylpropanoid derivatives has often been
17 described as a form of biochemical defence in response to elicitors. For instance, Bolwell *et*
18 *al.* (1985) reported an accumulation of phenolic material bound to cellulosic and hemi-
19 cellulosic fractions and of free hydroxycinnamic acids in bean cells. Some of the phenolic-
20 bound esters identified in this study were also found in cells treated with elicitor fractions
21 from other oomycetes. It is the case of extracts from *Phytophthora megasperma* that induced
22 accumulation of cell-wall-bound 4-coumaric acid, ferulic acid, 4-hydroxybenzoic acid, 4-
23 hydroxybenzaldehyde and vanillin (Kauss *et al.* 1993). Cell-wall-bound 4-
24 hydroxybenzaldehyde and tyramine amides were also detected in suspension cultures of
25 potato treated with an elicitor extract of *Phytophthora infestans* (Keller *et al.* 1996b).

1 Since in response to cryptogein, CAD transcript levels are slightly down-regulated whereas
2 most upstream genes are up-regulated, it is conceivable that the CAD substrate
3 (coniferaldehyde) could be (at least in part) not converted into coniferyl alcohol, but rather
4 redirected to ferulic acid by oxidation. Such a hydroxycinnamaldehyde dehydrogenase
5 (*REF1*, *AtALDH2C4*) activity has been demonstrated in *Arabidopsis thaliana* (Nair *et al.*
6 2004). The identification of this enzyme inverted the conventional model of flux, ferulate and
7 sinapate being products of coniferaldehyde and sinapaldehyde oxidation, rather than being
8 precursors in their synthesis. In support of the hypothesis that cell-wall-bound ferulate could
9 originate from coniferaldehyde, we recently identified within the publicly available tobacco
10 ESTs, a sequence highly homologous to *AtALDH2C4*. Furthermore, we have shown that this
11 gene was significantly over-expressed by 5 times in BY-2 cells challenged with cryptogein
12 (Amelot *et al.*, in preparation).

13 An increased amount of 4-hydroxybenzaldehyde was also observed in cryptogein-treated
14 cells. Although the origin of this compound is unknown, one could hypothesize the existence
15 of a 4-hydroxybenzaldehyde synthase activity in tobacco. Such a synthase activity, which
16 converts 4-coumaric acid to hydroxybenzaldehyde, was found in tissue cultures of *Vanilla*
17 *planifolia* (Podstolski *et al.* 2002) and in hairy root cultures of *Daucus carota* (Sircar & Mitra
18 2008).

19 Taken together, our results clearly show a reorientation of the phenylpropanoid metabolism in
20 response to cryptogein, resulting in the incorporation of new cell wall phenolic compounds
21 which likely reinforce the cell wall barrier. Moreover they suggest a new possible route
22 accounting for the accumulation of ferulic acid in tobacco BY-2 cells.

23 **Calcium controls the cryptogein-induced responses**

24 Our results showed that the phenylpropanoid genes belong to two distinct subsets that respond
25 differentially to cryptogein treatment. The regulation of these two subsets differs by their

sensitivity to stimulus intensity (i.e. cryptogein concentration) as well as by their correlation with one particular phase of the calcium signature (peak and/or plateau). Indeed, the long lasting influx of calcium responsible for the sustained increased level of cytosolic calcium (plateau) is necessary for the appropriate transcriptional response of genes induced by low concentrations of cryptogein (i.e. *PAL*, *HCT* and *CCoAOMT*). Thus, a positive correlation is observed at low cryptogein concentrations between the sustained calcium elevation and transcript accumulation of these genes, while at higher concentrations the calcium peak value was negatively correlated with transcription. On the other hand, *C4H*, *4CL*, *C3H*, *COMT* and *CCR* genes activation seem to rely on higher cryptogein concentrations. Furthermore, it appears that the early phase of the calcium response is mainly responsible for their transcriptional response. Such differential responsiveness to calcium response would allow the plant to selectively activate genes of the phenylpropanoid pathway.

To explain these results one might consider that different calcium stores involved in the generation of the cryptogein-induced calcium signature (Garcia-Brügger *et al.* 2006, Kadota *et al.* 2004, Lecourieux *et al.* 2002) contribute differently to the subsequent biological responses. However, computer simulations of our two-component model rather suggest the existence of two different calcium decoding modules, with low and high affinity for calcium, associated with the main calcium peak and the sustained phase (plateau) respectively, and displaying either an inhibiting or an activating effect.

This model is also supported by the specific effect of calmodulin inhibitor W13 on the transcriptional response to low cryptogein concentrations only, which suggests the involvement of calmodulins or calmodulin-like proteins in the transduction process. This effect could be simulated with our model by increasing the value of the “apparent” affinity coefficient of the “activation” component (Fig.10). One may thus hypothesize that W13 inhibits calcium-binding proteins (such as CaMs, CMLs or potentially other EF-hand

proteins) involved either in the control of calcium homeostasis (channels or pumps) and/or in signal transduction. It is known that some Ca-channels are regulated by calmodulin (Tadross, Dick & Yue 2008), Ca^{2+} -CaM is also involved in the control of calcium pumps (Boursiac & Harper 2007). Thus, according to our model, increasing the channel activity or inhibiting pump activity would result in an increase of the calcium resting level (plateau value), which could in turn activate calcium-dependent transcription complexes. When the cryptogein concentration is higher ($> 0.1 \mu\text{M}$), activation is already at its maximum, and thus cannot be enhanced further, but for genes like *PAL* or *CCoAOMT* inhibition can still be increased, leading to a transcriptional response lower than the one obtained at low cryptogein. Globally, anti-CaMs appear to shift the cryptogein concentration at which the transcriptional response is maximal towards lower values.

The non-linear relationship between the calcium signal and downstream responses highlights the complex regulatory properties of calcium and reinforces the concept of calcium signature, considered here as an “observable” giving an overview at the whole population level of the underlying calcium-dependent processes thought to encode/decode information required for the specificity of the final responses. As the calcium signature observed at the level of a population represents the integral of calcium signals occurring in each individual cell, the possibility to measure calcium signals at the single cell level will be invaluable to further interpret the observed data at the mechanistic level. Finally, the identification of calcium- or CaM-dependent transcription factors involved in the regulation of genes of the phenylpropanoid pathway in tobacco should bring new insights in understanding cryptogein signalling.

ACKNOWLEDGEMENTS

The authors thank Sabine Grat for her technical assistance in cell culture maintenance and Nathalie Ladouce for her help in RT-qPCR analysis. Raoul Ranjeva thanks the Miller Institute for Basic Research (UC Berkeley) for supporting him with a Miller Visiting Professorship. Nicolas Amelot thanks the *Association pour la Recherche sur les Nicotianae* for financial support. This work was supported by a grant from the *Agence Nationale pour la Recherche* (ANR-06-BLAN-370) and by the *Centre National de la Recherche Scientifique*.

REFERENCES

- Allen G.J., Chu S.P., Harrington C.L., Schumacher K., Hoffmann T., Tang Y.Y., Grill E. & Schroeder J.I. (2001a) A defined range of guard cell calcium oscillation parameters encodes stomatal movements. *Nature*, **411**, 1053-1057.
- Allen G.J. & Schroeder J.I. (2001b) Combining genetics and cell biology to crack the code of plant cell calcium signaling. *Science STKE*, **2001**, RE13.
- Binet M.N., Humbert C., Lecourieux D., Vantard M. & Pugin A. (2001) Disruption of microtubular cytoskeleton induced by cryptogein, an elicitor of hypersensitive response in tobacco cells. *Plant Physiology*, **125**, 564-572.
- Bolwell G.P., Robbins M.P. & Dixon R.A. (1985) Metabolic changes in elicitor-treated bean cells. Enzymic responses associated with rapid changes in cell wall components. *European Journal of Biochemistry*, **148**, 571-578.
- Bonnet P., Bourdon E., Ponchet M., Blein J.-P. & Ricci P. (1996) Acquired resistance triggered by elicitors in tobacco and other plants. *European Journal of Plant Pathology*, **102**, 181-192.
- Bourque S., Binet M.N., Ponchet M., Pugin A. & Lebrun-Garcia A. (1999) Characterization of the cryptogein binding sites on plant plasma membranes. *Journal of Biological Chemistry*, **274**, 34699-34705.

- 1 Boursiac Y. & Harper J.F. (2007) The origin and function of calmodulin regulated Ca^{2+}
2 pumps in plants. *Journal of Bioenergetics and Biomembranes*, **39**, 409-414.
- 3 Dagher R., Briere C., Feve M., Zeniou M., Pigault C., Mazars C., Chneiweiss H., Ranjeva R.,
4 Kilhoffer M.C. & Haiech J. (2008) Calcium fingerprints induced by Calmodulin
5 interactors in eukaryotic cells. *Biochimica Biophysica Acta*, **1793**, 1068-1077.
- 6 Dixon R.A. & Paiva N.L. (1995) Stress-induced phenylpropanoid metabolism. *The Plant Cell*,
7 **7**, 1085-1097.
- 8 Garcia-Brügger A., Lamotte O., Vandelle E., Bourque S., Lecourieux D., Poinssot B.,
9 Wendehenne D. & Pugin A. (2006) Early signaling events induced by elicitors of plant
10 defenses. *Molecular Plant Microbe Interactions*, **19**, 711-724.
- 11 Hano C., Addi M., Bensaddek L., Crônier D., Baltora-Rosset S., Doussot J., Maury S.,
12 Mesnard F., Chabbert B., Hawkins S., Lainé E. & Lamblin F. (2006) Differential
13 accumulation of monolignol-derived compounds in elicited flax (*Linum*
14 *usitatissimum*) cell suspension cultures. *Planta*, **223**, 975-989.
- 15 Hetherington A.M. & Brownlee C. (2004) The generation of Ca^{2+} signals in plants. *Annual*
16 *Review of Plant Biology*, **55**, 401-427.
- 17 Israelsson M., Siegel R.S., Young J., Hashimoto M., Iba K. & Schroeder J.I. (2006) Guard
18 cell ABA and CO_2 signaling network updates and Ca^{2+} sensor priming hypothesis.
19 *Current Opinion in Plant Biology*, **9**, 654-663.
- 20 Kadota Y., Goh T., Tomatsu H., Tamauchi R., Higashi K., Muto S. & Kuchitsu K. (2004)
21 Cryptogein-induced initial events in tobacco BY-2 cells: pharmacological
22 characterization of molecular relationship among cytosolic Ca^{2+} transients, anion
23 efflux and production of reactive oxygen species. *Plant and Cell Physiology*, **45**, 160-
24 170.

- 1 Kaplan B., Davydov O., Knight H., Galon Y., Marc R. Knight, Fluhr R. & Fromm H. (2006)
- 2 Rapid transcriptome changes induced by cytosolic Ca^{2+} transients reveal ABRE-
- 3 related sequences as Ca^{2+} -responsive cis elements in Arabidopsis. *The Plant Cell*, **18**,
- 4 2733–2748.
- 5 Kauss H., Franke R., Krause K., Conrath U., Jeblick W., Grimmig B. & Matern U. (1993)
- 6 Conditioning of parsley (*Petroselinum crispum* L.) suspension cells increases elicitor-
- 7 induced incorporation of cell wall phenolics. *Plant Physiology*, **102**, 459-466.
- 8 Keller H., Bonnet P., Galiana E., Pruvot L., Friedrich L., Ryals J. & Ricci P. (1996a) Salicylic
- 9 acid mediates elicitor-induced systemic acquired resistance, but not necrosis in
- 10 tobacco. *Molecular Plant-Microbe Interactions*, **9**, 696–703.
- 11 Keller H., Hohlfeld H., Wray V., Hahlbrock K., Scheel D. & Strack D. (1996b) Changes in
- 12 the accumulation of soluble and cell wall-bound phenolics in elicitor-treated cell
- 13 suspension cultures and fungus-infected leaves of *Solanum tuberosum*.
- 14 *Phytochemistry*, **42**, 389-396.
- 15 Kim J. & Kim H.Y. (2006) Functional analysis of a calcium-binding transcription factor
- 16 involved in plant salt stress signaling. *FEBS Letters*, **580**, 5251-5256.
- 17 Kim M.C., Chung W.S., Yun D.J. & Cho M.J. (2009) Calcium and Calmodulin-Mediated
- 18 Regulation of Gene Expression in Plants. *Molecular Plant*, **2**, 13-21.
- 19 Kudla J., Batistic O. & Hashimoto K. (2010) Calcium Signals: The Lead Currency of Plant
- 20 Information Processing. *The Plant Cell*, **in press**.
- 21 Lecourieux D., Mazars C., Pauly N., Ranjeva R. & Pugin A. (2002) Analysis and effects of
- 22 cytosolic free calcium increases in response to elicitors in *Nicotiana plumbaginifolia*
- 23 cells. *The Plant Cell*, **14**, 2627-2641.
- 24 Lecourieux D., Ranjeva R. & Pugin A. (2006) Calcium in plant defence-signalling pathways.
- 25 *New Phytologist*, **171**, 249-269.

- 1 McAinsh M.R. & Pittman J.K. (2009) Shaping the calcium signature. *New Phytologist*, **181**,
2 275-294.
- 3 Miwa H., Sun J., Oldroyd G.E. & Downie J.A. (2006) Analysis of calcium spiking using a
4 cameleon calcium sensor reveals that nodulation gene expression is regulated by
5 calcium spike number and the developmental status of the cell. *Plant Journal*, **48**, 883-
6 894.
- 7 Moscatiello R., Mariani P., Sanders D. & Maathuis F.J.M. (2006) Transcriptional analysis of
8 calcium-dependent and calcium-independent signalling pathways induced by
9 oligogalacturonides. *Journal of Experimental Botany*, **57**, 2847-2865.
- 10 Nair R.B., Bastress K.L., Ruegger M.O., Denault J.W. & Chapple C. (2004) The Arabidopsis
11 thaliana REDUCED EPIDERMAL FLUORESCENCE1 gene encodes an aldehyde
12 dehydrogenase involved in ferulic acid and sinapic acid biosynthesis. *The Plant Cell*,
13 **16**, 544-554.
- 14 Pauly N., Knight M.R., Thuleau P., Graziana A., Muto S., Ranjeva R. & Mazars C. (2001)
15 The nucleus together with the cytosol generates patterns of specific cellular calcium
16 signatures in tobacco suspension culture cells. *Cell Calcium*, **30**, 413-421.
- 17 Pauly N., Knight M.R., Thuleau P., Van der Luit A.H., Moreau M., Trewavas A.J., Ranjeva
18 R. & Mazars C. (2000) Control of free calcium in plant cell nuclei. *Nature*, **405**, 754-
19 755.
- 20 Pfaffl M.W. (2001) A new mathematical model for relative quantification in real-time RT-
21 PCR. *Nucleic Acids Research*, **29**, e45.
- 22 Podstolski A., Havkin-Frenkel D., Malinowski J., Blount J.W., Kourteva G. & Dixon R.A.
23 (2002) Unusual 4-hydroxybenzaldehyde synthase activity from tissue cultures of the
24 vanilla orchid *Vanilla planifolia*. *Phytochemistry*, **61**, 611-620.

- 1 Pontier D., Godiard L., Marco Y. & Roby D. (1994) hsr203J, a tobacco gene whose activation
2 is rapid, highly localized and specific for incompatible plant/pathogen interactions.
3 *Plant Journal*, **5**, 507-521.
- 4 Qudeimat E., Faltusz A.M., Wheeler G., Lang D., Brownlee C., Reski R. & Frank W. (2008)
5 A PIIB-type Ca²⁺-ATPase is essential for stress adaptation in *Physcomitrella patens*.
6 *Proceedings of the National Academy of Sciences of the United States of America*,
7 **105**, 19555-19560.
- 8 R Development Core Team (2009) R: A language and environment for statistical computing.
9 R Foundation for Statistical Computing, Vienna, Austria.
- 10 Schroeder J.I., Allen G.J., Hugouvieux V., Kwak J.M. & Waner D. (2001) Guard Cell Signal
11 Transduction. *Annual Review of Plant Physiology and Plant Molecular Biology*, **52**,
12 627-658.
- 13 Shinya T., Galis I., Narisawa T., Sasaki M., Fukuda H., Matsuoka H., Saito M. & Matsuoka
14 K. (2007) Comprehensive analysis of glucan elicitor-regulated gene expression in
15 tobacco BY-2 cells reveals a novel MYB transcription factor involved in the
16 regulation of phenylpropanoid metabolism. *Plant and Cell Physiology*, **48**.
- 17 Sircar D. & Mitra A. (2008) Evidence for p-hydroxybenzoate formation involving enzymatic
18 phenylpropanoid side-chain cleavage in hairy roots of *Daucus carota*. *Journal of Plant*
19 *Physiology*, **165**, 407-414.
- 20 Sudha G. & Ravishankar G. (2002) Involvement and interaction of various signaling
21 compounds on the plant metabolic events during defense response, resistance to stress
22 factors, formation of secondary metabolites and their molecular aspects. *Plant Cell*,
23 *Tissue and Organ Culture*, **71**, 181-212.
- 24 Sutty L., Blein J., Ricci P. & Pugin A. (1995) Early changes in gene expression in tobacco
25 cells elicited with cryptogein. *Molecular Plant Microbe Interactions*, **8**, 644-651.

- 1 Suzuki A., Yano A., Nishiuchi T., Nakano T., Kodama H., Yamaguchi K. & Shinshi H.
2 (2003) Comprehensive analysis of early response genes to two different microbial
3 elicitors in tobacco cells. *Plant Science*, **173**, 291-301.
- 4 Tadross M.R., Dick I.E. & Yue D.T. (2008) Mechanism of local and global Ca^{2+} sensing by
5 calmodulin in complex with a Ca^{2+} channel. *Cell*, **133**, 1228-1240.
- 6 Takahashi K., Isobe M., Knight M.R., Trewavas A.J. & Muto S. (1997) Hypoosmotic shock
7 induces increases in cytosolic Ca^{2+} in tobacco suspension-culture cells. *Plant*
8 *Physiology*, **113**, 587-594.
- 9 Ward E.R., Uknes S.J., Williams S.C., Dincher S.S., Wiederhold D.L., Alexander D.C., Ahl-
10 Goy P., Metraux J.P. & Ryals J.A. (1991) Coordinate gene activity in response to
11 agents that induce systemic acquired resistance. *The Plant Cell*, **3**, 1085-1094.
- 12 Ward J.M., Pei Z.-M. & Schroeder J.I. (1995) Roles of ion channels in initiation of signal
13 transduction in higher plants. *The Plant Cell*, **7**, 833-844.
- 14 Weckwerth W. (2003) Metabolomics in systems biology. *Annual Review of Plant Biology*, **54**,
15 669-689.
- 16 White P.J. & Broadley M.R. (2003) Calcium in plants. *Annals of Botany*, **92**, 487-511.
- 17 Xiong T.C., Bourque S., Lecourieux D., Amelot N., Grat S., Briere C., Mazars C., Pugin A. &
18 Ranjeva R. (2007) Calcium signaling in plant cell organelles delimited by a double
19 membrane. *Biochimica and Biophysica Acta*, **1763**, 1209-1215.

Table 1

Cryptogein effect on the accumulation of insoluble phenolic compounds in tobacco BY-2 cells. For each compound, ratio between treated *versus* control cells was calculated based on their relative amounts measured by MS/MS. Data are means and standard-errors from 4 independent experiments. Paired-t-test was used for statistical comparisons between amounts (Log transformed).

Phenolic compound	Ratio (mean $\pm$ SE)	P value
<i>p</i> -coumaric acid	3.3 $\pm$ 1.0	0.04
ferulic acid	1.9 $\pm$ 0.3	0.05
5-hydroxyferulic acid	184 $\pm$ 112	0.005
4-hydrobenzaldehyde	1.8 $\pm$ 0.3	0.06

Figures legends

Figure 1. Biosynthetic routes for phenolic compounds adapted from the model proposed by Nair *et al.* (2004) for *A. thaliana*. Encircled compounds accumulate in tobacco BY-2 cell walls in response to cryptogein. Light grey arrows indicate routes that are unlikely to occur *in planta*. Dotted arrow indicates a hypothetic route for the biosynthesis of 5-hydroxyferulate (see results section). PAL, phenylalanine ammonia-lyase; C4H, cinnamate 4-hydroxylase; 4CL, 4-coumarate-CoA ligase; C3H, *p*-coumarate 3-hydroxylase; HCT, hydroxycinnamoyl CoA:quinate/shikimate hydroxycinnamoyltransferase; CCoAOMT, caffeoyl-CoA 3-*O*-methyltransferase; COMT caffeic acid/5-hydroxyferulic acid *O*-methyltransferase; F5H, ferulate 5-hydroxylase; CCR, cinnamoyl CoA reductase; CAD, cinnamyl alcohol dehydrogenase. REF1, Reduced Epidermal Fluorescence1.

Figure 2. Cryptogein-induced calcium signatures in tobacco BY-2 cells. (a) Time-course of calcium variation in BY-2 cells induced either by H₂O (trace) or cryptogein (1 μ M , bullets) (data from one representative experiment). (b) Dose-dependence of (●) peak value and (■) plateau value. (means $\pm$ s.e. of 3 independent experiments) .

Figure 3. Effects of cryptogein on transcript accumulation of phenylpropanoid genes in tobacco BY-2 cells. (a) Relative transcript levels of genes of phenylpropanoid in cells challenged for 4h with 0.1 μ M cryptogein in the absence (grey) or presence (black) of 1mM La³⁺. (b) Time-course of *PAL* (white), *HCT* (grey), and *CCoAOMT* (black) transcript accumulation induced by 0.1 μ M cryptogein. (c) Dose-dependence of transcript levels of *PAL* (white), *HCT* (grey) and *CCoAOMT* (black) on a 4-hour cryptogein treatment. Transcript levels assessed by RT qPCR are expressed as mean $\pm$ s.e. (n=3) of transcript ratios relative to control cells (H₂O treatment).

Figure 4. Principal Component Analysis (PCA). Analysis of transcription ratio data plus calcium-peak and calcium-plateau values from dose-response experiments. Symbols 10nM to 1µM represent the cryptogein concentration used in experiments. Cry+la = Co-treatment with La^{3+} . Planes defined by the first two principal components are displayed.

Figure 5. Time course of relative enzymatic activities of PAL and CCoAOMT. Activities were measured in total protein extracts from cells challenged with 0.1µM cryptogein: PAL (white bars) and CCoAOMT (black bars). Traces: relative transcript levels of CCoAOMT (closed symbols) and PAL (open symbols). Inset: Effect of 1mM La^{3+} on PAL and CCoAOMT activities measured after a 4-hour treatment. Enzyme activities are expressed relative to control cells (H_2O treatment). Means±s.e. (n= 3).

Figure 6. Distribution of radioactivity in tobacco BY-2 cell suspensions incubated for 4 hours with [$^{14}\text{C}(\text{U})$]-cinnamic acid. At t=0 hour, cells were treated with water (control, white bar), 0.1µM cryptogein (black bar), or 0.1µM cryptogein + 1mM La^{3+} (grey bar). Relative radioactivity is expressed as the ratio of the activity measured in one compartment over the total radioactivity added to the cell suspension. Means±s.e. (n=3).

Figure 7. Effect of inhibiting calcium influx on cryptogein-induced calcium variations (a) and transcript accumulation (b) in tobacco BY-2 cells. 1mM La^{3+} was added at t=0 (black line and black bars) or t=10 minutes (grey line and grey bars) after treating cells with 0.5µM cryptogein. In (b) data are expressed as the percentage of inhibition of the transcript accumulation induced by cryptogein alone (means±s.e. of 3 experiments).

Figure 8. Effects of calmodulin inhibitors W12 and W13 on the cryptogein-response. (a) Effects of W13 preincubation on the cryoptogein-induced calcium signature. Bullets: 50 nM cryptogein, empty bullets: 50 µM W13+50 nM cryptogein, square: 250 nM cryptogein, empty square: 50 µM W13 + 250 nM cryptogein (means±s.e. of 2 replicates). (b) Effects of W12

(negative control) and W13 preincubation on cryptogein-induced *CCoAOMT* transcript accumulation. Control (white bar), W12 (grey bar) and W13 (black bar). Inhibitors (50 μ M) were added to cells 10 minutes before cryptogein. Data are representative of two independent experiments.

Figure 9. Proposed model for regulation of cryptogein-induced transcriptional response by calcium.

Figure 10. Fits of the two-component model described in Fig. 9 to experimental data. (a) Dose-response curves for transcript accumulation of *PAL* (circle), *HCT* (triangle) and *CCoAOMT* (square) induced by a 4-hour treatment with cryptogein. (b) Effect of anti-CaM compounds on *CCoAOMT* transcript accumulation induced by cryptogein: control (+), W-12 pre-treatment (x), W-13 pre-treatment (*). Experimental data (symbols) and theoretical curves (lines). Parameter values are given in Supporting Information S7.

1 **Supporting Information:**

2 Table S1: primers and amplicons sizes.

3 Figure S2: Comparison between lanthanum chloride and EGTA pre-treatments on cryptogein-
4 induced phenylpropanoid genes.

5 Figure S3: Effects of a cycloheximide pre-treatment on cryptogein-induced transcript
6 accumulation of phenylpropanoid genes.

7 Figure S4: Effect of cryptogein on transcript accumulation of genes coding for isoforms of
8 *PAL*, *4CL* and *CCoAOMT*.

9 Figure S5: Cytosolic calcium responses to calmodulin inhibitors in tobacco BY-2 cells.

10 Figure S6: Assessment of cryptogein-induced cell death in tobacco BY-2 cells.

11 Figure S7: Model formulation.

12

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Table S1: Primers and amplicon sizes:

Gene	Amplicon size	upper 5'	lower 5'
<i>18S rRNA</i>	171bp	cgcgctacactgatgtattc	gtacaaagggcaggacgta
<i>PAL</i>	94bp	gacaaagtgttcacagcaatg	aaacagatwggagaggagca
<i>PAL A</i>	228bp	cctcgtcgaccacgccttgc	taattcagctccaagttctc
<i>PAL B</i>	242bp	acacattgccacattcagca	caccattgggaccaacagcc
<i>C4H</i>	124bp	tcaacacaatggtggaatgc	actttgggacgtttggttca
<i>4CL</i>	89bp	cttctcaaccatcccaacatt	ctaacaacaaaagccactgga
<i>4CL1</i>	178bp	ttcctggagttaatacagaag	gggaatttggttctcacagcg
<i>4CL2</i>	179bp	tttcttggaattgataaacg	ggaaatttggtcgaacagta
<i>COMT</i>	132bp	cctgcaaatgggaaggtgat	cagtcctttcttgcctcct
<i>C3H</i>	142bp	tggctgaggtgatcaagaac	tatgggaggaaggggaagtc
<i>HCT</i>	127bp	ggctgccaatccatgatgct	gcaacagattgactgccatca
<i>CCoAOMT</i>	96bp	acaccctatggaatggatca	ccttggtgagttccaatacga
<i>CCoAOMT 1</i>	89bp	aatttgccagctacctgttgg	ttggtgccacaaatagacgat
<i>CCoAOMT 2</i>	92bp	gaactattgtccatttgggca	acgaatgatacagaacagggga
<i>CCoAOMT 3</i>	130bp	tttacagggctagttcatggca	ggtaacatcgacaatgcaacct
<i>CCoAOMT 4</i>	102bp	cattggctgctgattctaga	aaatcctccattgcctgtaa
<i>CCoAOMT 5</i>	113bp	acgtatgatttcatttgggat	caatccttgatcaacggctagg
<i>CCoAOMT6</i>	121bp	cacatatgatttcatttcgtgg	caattctgggatcagctgcaaaa
<i>F5H</i>	100bp	agcccggaaggaagat	cccatccaagccaaggta
<i>CCR</i>	99bp	gacttctgcaaaaacaccaa	caccaaategactcctttt

S2. Comparison between lanthanum chloride and EGTA pre-treatments on cryptogein-induced phenylpropanoid genes expression in tobacco BY-2 cells.

Figure S2. Comparison between lanthanum chloride and EGTA pre-treatments on cryptogein-induced phenylpropanoid genes expression in tobacco BY-2 cells.

(a) Time course of cytosolic calcium variation in response to 0.1 μM cryptogein in the absence (dark grey trace) or presence of either 1 mM La^{3+} (black trace) or 2 mM EGTA (light grey trace). Data show one representative experiment out of three. (b) Relative transcript levels assessed by RT-qPCR in cells treated for 3 hours with 0.1 μM cryptogein in the absence (dark grey bars) or presence of either 1 mM La^{3+} (black bars) or 2 mM EGTA (light grey bars). Transcript levels are expressed as means $\pm SE$ (n=3) of transcript ratios relative to control cells (H_2O treatment).

S3. Effects of a cycloheximide pre-treatment on cryptogein-induced phenylpropanoid genes in BY-2 tobacco cells

Figure S3. Effect of a cycloheximide pre-treatment on the cryptogein-induced accumulation of phenylpropanoid gene transcripts. Transcript levels were assessed by RT-qPCR in tobacco BY-2 cells treated for 3 hours with 0.1 μ M cryptogein. Relative transcript levels are ratios of transcript numbers assessed in treated cells to those of cycloheximide only-treated cells. Black bars: cells pre-incubated during 30-minutes with 50 μ M cycloheximide before cryptogein treatment. Grey bars: control (no pre-treatment) cells (data from one experiment).

S4. Effect of cryptogein on expression of genes coding for isoforms of PAL, 4CL and CCoAOMT

Figure S4. Relative transcript levels of genes coding for PAL, 4CL and CCoAOMT isoforms. Transcript levels were assessed by RT-qPCR in tobacco BY-2 cells treated for 3 hours with 0.1 μ M cryptogein. Relative transcript levels are ratios of transcript levels assessed in treated cells versus transcript levels obtained in control (H_2O treated) cells (data from one experiment).

S5. Cytosolic calcium responses to calmodulin inhibitors in tobacco BY-2 cells

Figure S5. Cytosolic calcium variations induced in tobacco BY-2 cells by calmodulin interactors: W12 (low affinity, $K_d = 250\mu\text{M}$) and W13 (high affinity, $K_d = 50\mu\text{M}$).

S6. Assessment of cryptogein-induced cell death in tobacco BY-2 cells.

Figure S6. Effect of a 24-hour treatment with cryptogein and/or 1mM La^{3+} on the viability of tobacco BY-2 suspension cultured cells (relative viability of treated cells to that of control cells (data from one experiment with three replicates).

S7. Model formulation

The model used to describe the relationship between cryptogein concentration “C” and transcript ratio “R” (measured 4 hours after treatment) was inspired from enzymatic kinetic laws for one ligand and two binding sites (Adair-Klotz model) and Hill phenomenological equation. Here the ligand is cryptogein, the observable variable is the transcript ratio, and the binding sites correspond by analogy to the two putative decoding modules. Thus the equilibrium reads as follows:

$$R = \frac{1 + S1 (K1.C)^{n1} + S2 (K1.K2.C^2)^{n2}}{1 + (K1.C)^{n1} + (K1.K2.C^2)^{n2}} \quad (1)$$

Where K1, S1, and n1 are the association constant, the maximum response and the Hill coefficient for component 1 (respectively K2, S2, and n2 for component 2). For a given condition Ki values and Hill coefficients were supposed to be the same for all genes, while coefficients S1 and S2 were gene dependent.

Estimated values of the coefficients for three genes (data from Fig. 10a):

$$K1 = 23.3 \mu\text{M}^{-1}; K2 = 24.9 \mu\text{M}^{-1}; n1 = 3.9; n2 = 2.7$$

	PAL	HCT	CCoAOMT
S1	58.2	101.9	678
S2	13.5	77.6	11.6

Estimated values of the coefficients for CCoAOMT (data from Fig. 10b):

$$S1=411.2; S2=123.3; n1=3.9; n2=2.7; K2=30.5$$

	Cryptogein	W12 + Cryptogein	W13 + Cryptogein
K1 (μM^{-1})	33.1	41.4	52.8