Prevalence of renal insufficiency in breast cancer patients and related pharmacological issues

Vincent Launay-Vacher, Joseph Gligorov, Christophe Tourneau, Nicolas Janus, Jean-Philippe Spano, Isabelle Ray-Coquard, Stéphane Oudard, Xavier Pourrat, Jean-François Morere, Gilbert Deray, et al.

► To cite this version:

Vincent Launay-Vacher, Joseph Gligorov, Christophe Tourneau, Nicolas Janus, Jean-Philippe Spano, et al.. Prevalence of renal insufficiency in breast cancer patients and related pharmacological issues. Breast Cancer Research and Treatment, 2008, pp.745-753. 10.1007/s10549-008-0131-1 . hal-00534546

HAL Id: hal-00534546

<https://hal.science/hal-00534546>

Submitted on 10 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PREVALENCE OF RENAL INSUFFICIENCY IN BREAST CANCER PATIENTS AND RELATED PHARMACOLOGICAL ISSUES.

VINCENT LAUNAY-VACHER, PHARM.D.¹, JOSEPH GLIGOROV, M.D.², CHRISTOPHE LE TOURNEAU, M.D.³, NICOLAS JANUS, PHARM.D.¹, JEAN-PHILIPPE SPANO, M.D.⁴, ISABELLE RAY-COQUARD, M.D.⁵, STÉPHANE OUDARD, M.D.⁶, XAVIER POURRAT, PHARM.D.⁷, JEAN-FRANÇOIS MORERE, M.D.⁸, GILBERT DERAY, M.D.¹, PHILIPPE BEUZEBOC, M.D.³, AND THE RENAL INSUFFICIENCY AND ANTICANCER MEDICATIONS (IRMA) STUDY GROUP

¹NEPHROLOGY, HÔPITAL PITIÉ-SALPÊTRIÈRE, PARIS, FRANCE;

²MEDICAL ONCOLOGY, HÔPITAL TENON, PARIS, FRANCE;

³MEDICAL ONCOLOGY, INSTITUT CURIE, PARIS, FRANCE;

⁴MEDICAL ONCOLOGY, HÔPITAL PITIÉ-SALPÊTRIÈRE, PARIS, FRANCE;

⁵MEDICAL ONCOLOGY, CENTRE LÉON BÉRARD, LYON, FRANCE;

⁶MEDICAL ONCOLOGY, HÔPITAL EUROPÉEN GEORGES POMPIDOU, PARIS, FRANCE;

⁷CLINICAL PHARMACY, HÔPITAL TROUSSAU, TOURS, FRANCE;

⁸MEDICAL ONCOLOGY, HÔPITAL AVICENNE, BOBIGNY, FRANCE;

Corresponding Author:

Vincent Launay-Vacher, Pharm.D., Department of Nephrology Pr. Deray, Hôpital Pitié-Salpêtrière, 47-83 Boulevard de l'Hôpital, 75013 Paris, France.

Tel: +33 (0)1 42 17 72 30; Fax: +33 (0)1 42 17 72 12

E-mail: vincent.launay-vacher@psl.aphp.fr

ABSTRACT: 1ST PROPOSITION

The Renal Insufficiency and Anticancer Medications (IRMA) study is a French national, observational study which demonstrated the high prevalence of abnormal renal function in a population of 4684 solid tumour patients. Among them, 50 to 60% had decreased renal function defined as CrCl below 90, and 80% were treated with anticancer drugs that either necessitated dosage adjustment in case of RI or were potentially nephrotoxic drugs. Since patients and drugs used differ depending on the type of tumour, the IRMA Study Group started analyses in different subgroups of patients.

In the 1898 IRMA patients with breast cancer, the prevalence of RI was still very high in spite of a normal serum creatinine in almost all cases. Some anticancer drugs, as in particular some bisphosphonates, capecitabine and platinum salts, may be nephrotoxic and/or need dosage adjustment. However other important drugs in breast cancer do not require dose reduction, and do not present with potential nephrotoxicity (anthracyclines, taxanes, trastuzumab). Both issues seem to be slightly but significantly more important in patients with bone metastases as compared to patients with a non-metastatic disease.

ABSTRACT: 2ND PROPOSITION

BACKGROUND. The Renal Insufficiency and Anticancer Medications (IRMA) study was a French national observational study. The results from this study of nearly 5000 patients demonstrated the high prevalence of renal impairment in a population of patients with solid tumors with 50 to 60% presenting with a decreased renal function defined as CrCl below 90, and 80% being treated with anticancer drugs that either necessitated dosage adjustment in case of RI or were potentially nephrotoxic drugs. Here are presented the results on patients with breast cancer.

METHODS. Every cancer patient who presented at oncology departments that participated in the study over at least 1 of 2 predefined periods during 2004 was included in the IRMA study. Data on patients with breast cancer were analyzed. Renal function was calculated using Cockcroft-Gault and abbreviated Modification of Diet in Renal Disease (aMDRD) formulae to estimate the prevalence of renal insufficiency (RI) according to the KDOQI-KDIGO definition of kidney disease. Anticancer drugs were studied with regard to their potential renal toxicity and need for dosage adjustment.

RESULTS. Of the 1898 IRMA breast cancer patients, 1.6% had a SCr level >110 $\mu\text{mol/L}$. However, when they were assessed using Cockcroft-Gault and aMDRD formulae, 51.8% and 50.8% had abnormal renal function, respectively. Of the 3465 anticancer drug prescriptions, 53.2% required dose adjustments for RI or were drugs with no available data on their administration in patients with RI. Of the patients treated, 90.4% received at least 1 such drug, and 76.9% received potentially nephrotoxic drugs.

CONCLUSIONS. In the 1898 IRMA patients with breast cancer, the prevalence of RI was still very high in spite of a normal serum creatinine in almost all cases. Some anticancer drugs, as in particular some bisphosphonates, capecitabine and platinum salts, may be nephrotoxic and/or need dosage adjustment. However other important drugs in breast cancer do not require dose reduction, and do not present with potential nephrotoxicity (anthracyclines, taxanes, trastuzumab). Both issues seem to be slightly but significantly more important in patients with bone metastases as compared to patients with a non-metastatic disease.

INTRODUCTION

The increased incidence of malignancies in patients with chronic renal failure has been discussed since the mid-1970s.^{1,2} In their study, Cengiz *et al.* reported that 188 (6.7%) organ tumors of varying pathologies were found in 2817 patients with chronic renal failure³. Although it is known to be a common pathology in the general population⁴, the prevalence of renal insufficiency (RI) among patients with cancer has only been recently assessed in the “Renal Insufficiency and Anticancer Medications” (IRMA) study⁵, where we found a high prevalence of renal insufficiency among 4684 solid tumor patients from 15 French cancer centers. Only 7.2% had elevated serum creatinine (SCR) but 50 to 60% had creatinine clearance (CrCl) below 90 mL/min, with the majority of them between 60 to 90, which defines a stage 2 kidney disease according to the K/DOQI-KDIGO official international definition^{6,7}. Such a reduction in renal function may be too low to be of significant clinical importance in the oncology setting. This should however be considered, especially for patients near 60, as a risk factor for developing more severe renal insufficiency. Furthermore, markers of renal damage, which are part of the criteria required to define a stage 2 renal disease, were not recorded in the IRMA study. When present, *ie.* when a stage 2 kidney disease is actually diagnosed, nephrological investigations may be needed. This high prevalence of RI is an important issue in clinical practice for the handling of anticancer drugs in those patients. Since some anticancer drugs used in breast cancer are excreted predominantly in the urine as unchanged drug or active metabolite(s), any reduction in renal clearance results in accumulation of potentially toxic species and overdosage. The dosage of chemotherapeutic agents used in these patients with RI will, therefore, require dosage reduction to avoid severe toxicities. Furthermore, using potentially nephrotoxic anticancer drugs will also require specific monitoring and, when available, specific prevention methods in order to help reduce the risk for renal toxicity, especially in patients with baseline abnormal renal function before anticancer therapy is started.

The development of platinum-based chemotherapy for triple-negative⁸ and HER2 positive^{9,10,11,12} breast cancers, in particular the emerging and promising strategies associating a taxane, a platinum salt and trastuzumab, and the growing use of IV bisphosphonates (BP) emphasise the need for regular monitoring of renal function, with particular attention to patients with preexisting impaired renal function.

In breast cancer patients with bone metastases (BM), there is a paucity of data for optimal use of BP mainly regarding initiation and treatment duration. Benefit with zoledronate and ibandronate has been shown for a treatment duration of up to 2 years^{13,14}. Renal safety must be considered because over the past few years, case reports and randomized trials¹⁵ have raised cases of renal toxicity with IV BP, for instance with zoledronic acid-associated renal impairment leading to renal failure¹⁶.

Monitoring must include an appropriate evaluation of CrCl in patients receiving I.V BP because SCR values can be misleading.

Here are presented the results from the subgroup analysis of the 1898 Breast Cancer (BC) patients in the IRMA population.

METHODS

Study Design for the IRMA study

The IRMA study included a total of 4684 patients being treated for solid tumors (either in hospital or as outpatients) in an oncology department. The study consisted of a retrospective patient data collection on one of two, specific, 15-day time periods between either the 1st and the 15th of February 2004 or the 1st and the 15th of October 2004. These time periods were selected to avoid summer vacations and to be representative of the whole year, with one period in winter/spring and the other in fall/winter. Patients were included regardless of disease pathology, treatment (antineoplastic drugs used/to be used, pretreated or not). Patients were excluded if they were <18 years of age, had a diagnosis of myeloma, or presented with end-stage renal disease requiring renal replacement therapy (either hemodialysis or peritoneal dialysis) since the aim of the IRMA study was to determine the prevalence of potentially undiagnosed RI, meaning non-dialysis patients, those latter obviously presenting with this disease.

The following data were collected for each patient: sex, age, weight, serum creatinine, blood urea nitrogen, hemoglobinemia, type of tumor, metastasis, and anticancer drugs prescribed. Patients who were known to present with acute renal failure were excluded in order to determine the prevalence of potentially chronic abnormal renal function. Estimations of renal function were made by calculation from SCR using the Cockcroft-Gault (CG) formula¹⁷ and the abbreviated Modification of Diet in Renal Disease (aMDRD) formula¹⁸.

Cockcroft-Gault formula:

$$\text{CrCl (mL/min)} = k \times [(140 - \text{age}) \times \text{weight (kg)}] / \text{SCR } (\mu\text{mol/L})$$

k = 1.23 (male) or 1.04 (female); CrCl = creatinine clearance;

SCR = serum creatinine.

aMDRD formula:

$$\text{GFR (mL/min/1.73 m}^2\text{)} = k \times 186 \times [\text{SCR}]^{-1.154} \times [\text{age}]^{-0.203}$$

k = 1 (male) or 0.742 (female); GFR = glomerular filtration rate;

SCR = serum creatinine (mg/dL).

Renal function, calculated using either formula, was staged in accordance with the international clinical practice guidelines from the K/DOQI and the KDIGO^{6,7}:

- Stage 1: GFR $\geq$ 90 mL/min
- Stage 2: GFR 60 to 89 mL/min
- Stage 3: GFR 30 to 59 mL/min
- Stage 4: GFR 15 to 29 mL/min
- Stage 5: GFR <15 mL/min

With regard to anticancer therapies prescribed to study patients, those requiring dosage adjustment were identified in accordance with their pharmacokinetics and available recommendations from both their individual Summary of Product Characteristics (SmPC), and from two specific reference books on drug dosage adjustment in patients with RI: *Drug prescribing in renal failure. Dosing guidelines for adults. 4th edition*¹⁹ and *GPR Anticancéreux. Guide de prescription des médicaments chez le patient insuffisant rénal. 3^{ème} édition*²⁰. Anticancer therapies were then classified as “Yes” when adjustment was required, “No” when adjustment was not necessary, and “ND” when no data were available in the literature. To obtain profiles of anticancer therapies with regard to renal tolerance, an exhaustive literature search was performed using *PubMed* to identify any potential renal side effects of the therapies. If, at least, some cases of nephrotoxicity were retrieved, the therapy was classified as “Yes”, meaning “potentially nephrotoxic.” Therapies were labeled “No” when no cases had been found or there were no suggestions of potential renal toxicity. As a result, some drugs with different risks for renal toxicity may be similarly classified as “Yes”. For instance, Tamoxifen and Paclitaxel are both reported as potentially nephrotoxic because those drugs have been reported with one case of nephrotic syndrome for Tamoxifen²¹, and evidence of increased renal toxicity of Cisplatin + Paclitaxel versus Cisplatin alone²², respectively.

RESULTS OF THE BREAST CANCER SUBGROUP ANALYSIS

Patient Demographics

A total of 1898 patients with breast cancer were included in the study. Of these, 20 patients were men, and the mean age of all patients was 55.1 years (range, 23 to 95). Nearly one third of the patients (30.4%) had BM.

Renal Insufficiency

Table 1 shows the percentages of patients among the 5 stages of RI. In Figure 1 are presented the number of patients in each stage according to the formula used to estimate renal function, either CG or aMDRD.

For 10.5% of patients, no SCr could be retrieved in the medical file, and among the whole population of breast cancer patients, only 31 (1.63%) had a SCR level $\geq 110 \mu\text{mol/L}$. In spite of this low proportion of patients with elevated SCR, a majority of patients had in fact a decreased CrCl or eGFR: 51.8% and 50.8% of patients had abnormal renal function ($< 90 \text{ mL/min}$) when calculated using the CG or aMDRD formulae, respectively (Table 1). This high prevalence of RI was also observed in the 1667 patients (87.8%) whose SCR levels were normal, *ie.* $< 110 \mu\text{mol/L}$: 57.1% and 56.0% of those patients had abnormal renal function ($< 90 \text{ mL/min}$) when calculated using the CG or aMDRD formulae, respectively.

The prevalence of RI in IRMA breast cancer patients was also studied in patients with BM as compared to patients without BM. Table 2 describes the two populations of patients: patients with breast cancer and BM and patients with breast cancer without BM. Serum creatinine was elevated ($\geq 110 \mu\text{mol/L}$) in only 2.8% of patients with BM and in 1.1% of patients without BM ($p=0.0036$). Patients with BM were significantly older (57.7 vs. 53.9 years, $p<0.0001$) and had a higher (but not significantly, $p=0.06$) mean SCR (70.4 vs. 68.0 $\mu\text{mol/L}$). Mean CG estimates of renal function differed between groups (84.1 vs 89.7 mL/min , $p=0.0004$). However, they were not statistically different when estimating renal function with aMDRD (88.1 vs 89.2, $p>0.05$).

However, when calculating the prevalence of RI in those two groups of patients (Table 4), some significant differences appeared. The prevalence of renal dysfunction was as follows: 62.2% of breast cancer patients with BM had a CG creatinine clearance below 90 mL/min as compared to 56.8% of patients without BM ($p=0.0381$) and 60.0% of breast cancer patients with BM had an aMDRD estimated glomerular filtration rate below 90 $\text{mL/min}/1.73\text{m}^2$ versus 55.4% in patients without BM ($p>0.05$). There was thus a statistically significant difference in the prevalence of renal insufficiency with CG but the difference was not significant with aMDRD.

Anemia, according to the WHO definition ($<12 \text{ g/dL}$ in women and $<13 \text{ g/dL}$ in men) was observed in 44.1% of IRMA breast cancer patients. Further analysis showed that 2.1% and 18.4% of the patients had a hemoglobin level lower than 9 g/dL and within 9 to 11 g/dL , respectively. For 5.1%, no hemoglobin value was available in the medical file of the patient. Comparing patients with or without BM and for whom a hemoglobin value was available (Table 5), it appeared that patients with BM had statistically significantly more often hemoglobin levels within 9 and 11 and lower than 9 as compared to patients without BM: 25.1% vs. 15.5% and 3.6% vs. 1.4%, respectively.

Anticancer Drugs

Among the whole study population of 1898 breast cancer patients, 89.7% were receiving anticancer drugs during the tie period studied. This resulted in 1702 treated patients for a total number of anticancer drugs prescriptions of 3465, *ie.* 2.0 drugs/patient.

The prescriptions comprised 40 different drugs (Table 6) of which 35% are drugs that necessitate dosage adjustment in RI and 30% are drugs for which no data were available in the literature on whether it would be necessary or not to adjust their dosage in case of renal impairment. Of the 3465 prescriptions, 53.2% were for drugs for which a dosage adjustment was necessary, or for which there were no available data concerning administration in patients with RI, 44.8% and 8.4%, respectively. Finally, 80.2% of the patients were receiving at least one drug that necessitates dosage adjustment in RI and 10.2% at least one drug for which no data were available, resulting in a total 90.4% of breast cancer patients in IRMA with at least one drug for which caution is mandatory in RI (either dosage adjustment required or no data available). Regarding nephrotoxicity, 65% of the drugs were potentially toxic to the kidneys (57.5%) or for which there were no data in the literature on their renal tolerance (7.5%). Among the prescriptions, 41.7% were prescriptions of drugs that may be nephrotoxic, and 76.9% of breast cancer patients in IRMA were receiving at least one drug that may exhibit renal toxicity.

DISCUSSION

In this study we found that RI is highly prevalent in breast cancer patients. However, the frequency of RI clearly is routinely underestimated in clinical practice because physicians most often base their diagnosis on SCR measurements. It is crucial to outline that SCR is not appropriate for evaluating renal function. We observed that 10.5% of our patients had no report of SCR determination in their file. Even if those patients with no available SCR may be those who were only seen for a routine control and/or patients who were not treated or treated with non-nephrotoxic drugs (such as trastuzumab), it remains important to 1) measure SCR on a regular basis, the frequency being determined according to the patient's renal function, and 2) to report it in the patient's file.

It is furthermore of major importance that renal function is appropriately evaluated in all breast cancer patients by estimating CrCl or GFR calculated using either the CG or aMDRD formulae. This includes patients who have normal SCR levels since even in those with a so-called "normal" serum creatinine level, the frequency of stage 2 or more RI was 57.1% and 56.0% with CG or aMDRD formulae, respectively.

To date there are no data allowing recommending the use of one formula over the other, in oncology patients as well as in the general population. Most often, differences between the two formulae in terms of renal function are weak. Some differences may be observed in the ranging of the patients in the different stages of RI. As shown on Figure 1, such differences between the two formulae were not relevant in our patients. However, when the CG and the aMDRD estimates differ in a particular patient, it may be useful to consider a measure of the actual GFR, after discussions with a nephrologist. Such methods include a 24-hour urine collection to measure creatinine clearance or a measure of the actual GFR using a specific marker of renal filtration such as $^{51}\text{Cr-EDTA}$, for instance. However, such methods necessitate a trained staff, time, and are not cost-effective for a systematic evaluation of patients' renal function. They must be used only in some specific cases, and patients who may benefit from such a determination of renal function should be closely identified together with nephrologists, according to the patient's profile and its estimated renal function with CG and aMDRD.

In patients with stage 2 RI, potential drug-nephrotoxicity is the main issue. Many studies have demonstrated that pre-existing abnormal renal function is a risk factor for drug-induced nephrotoxicity²³. As a result, in those patients with mildly decreased renal function, anticancer drugs, antineoplastic or supportive care, should be cautiously selected in order to administer drugs that are not or are less nephrotoxic.

When renal function is lower than 60 mL/min, the risk for nephrotoxicity is even higher, and moreover, the clinical consequences are more severe since any further deterioration of renal function may precipitate end-stage renal disease. In those patients, in addition to nephrotoxicity, the question of drug dosage adjustment is crucial to avoid overdose due to accumulation of the drug from reduced excretion and related toxicities, such as neurological, hematological, skeletal, cardiological, and hepatological toxicities. Indeed, when renal function declines and is lower than 60 mL/min, pharmacokinetic changes often necessitate modification of the drug dosage to ensure efficacy and safety.

Over 90% of breast cancer patients in the IRMA population received at least one drug that requires dosage adjustment in patients with RI, or for which there were no data available regarding their use in patients with RI (in either the literature or the SmPC), in spite of the fact that the most important drugs in breast cancer do not require dose reduction, and do not present with potential nephrotoxicity (anthracyclines, taxanes, trastuzumab). In patients who are receiving drugs that require dosage adjustment, renal function should be calculated (with CG or aMDRD formulae) to prescribe drugs at their adjusted dosage in accordance with their renal function, preferably prior to each course of anticancer therapy, chemotherapy or other type of treatment. Where information is not available on dosage adjustments in patients with RI, the prescription of these drugs should be approved by a pharmacologist or a nephrologist, together with the oncologist in charge of the patient. When an alternative treatment exists for which recommendations are available, this should be preferably used.

The use of potentially nephrotoxic therapies in patients at high risk for drug renal toxicity due to pre-existing renal impairment should be avoided if possible and alternative treatments should be considered. However, in IRMA breast cancer patients the frequency of nephrotoxic drug prescriptions was high with 76.85% of the treated patients receiving at least one such drug.

There were significantly ($p < 0.05$) more prescriptions of anticancer drugs necessitating dosage adjustment in RI or drugs being potentially nephrotoxic in breast cancer patients with BM as compared to breast cancer patients without BM (Tables 7, 8 and 9). However, the percentage of patients receiving at least one such drug was higher in patients without BM, 82.9% vs. 74.5% of treated patients in each group

($p < 0.05$). Furthermore, 78.7% of treated patients without BM received at least one potentially nephrotoxic drug versus 72.2% of patients with BM ($p < 0.05$). This means that patients with BM receive more frequently several drugs that may necessitate dosage adjustment and/or which can be nephrotoxic.

The IRMA study did not allow to evaluate whether the prescription of such drugs was or may have been responsible for the higher frequency of decreased renal function in those breast cancer patients with BM when renal function is calculated with aMDRD. For instance, the frequent administration of contrast media for CT scan, which is known to be a potential cause of renal dysfunction, may be one of the etiologies, in addition to or rather than chemotherapy. However, the message that came out was that, once patients are diagnosed with decreased renal function, drug therapy should be re-evaluated, dosages adjusted where necessary, and some potentially nephrotoxic drugs changed for less or non-nephrotoxic ones, when efficacy has been proven to be of a similar range.

In breast cancer therapy, the choice of the anticancer drug to use obeys multiple rules and criteria, indications and protocols. It may not be always possible not to use potentially nephrotoxic drugs. However, it remains very important to be aware of the renal function of patients who receive such drugs and to monitor renal function on a regular basis, prior to each course of therapy during treatment. Depending on the level of renal function of the patient once in remission, a twice-a-year monitoring may be sufficient during follow-up. Furthermore, in those patients who require a nephrotoxic antineoplastic chemotherapy, and especially those with baseline decreased renal function, cautious selection and analysis of associated drugs should be performed. For example, non-steroidal anti-inflammatory drugs, where possible, should be avoided since they may potentate the renal toxicity of chemotherapy.

Regarding the treatment of BM, when a bisphosphonate is to be prescribed, the same selection rules should be applied according to renal tolerance and dosage adjustment. There are no specific limitations for the use of BP in the elderly. Task forces from the International Society of Geriatric Oncology recommended a monitoring of CrCl in every patient, preferably before each course, the use of an agent with the best possible renal tolerability, and the assessment of hydration status^{24,25,26}.

In patients with evidence of renal deterioration during treatment, IV BP should be withheld and only resumed when serum creatinine returns to within 10% baseline²⁷. In case of mild to moderate renal impairment (CrCl:30-60 ml/min) regulations recommend lower doses of zoledronate and longer infusion for pamidronate. Treatment with zoledronate is contraindicated in patients with severe dysfunction (CrCl < 30 ml/min).

A recent label of ibandronate approved by European regulatory authorities allows a dosing regimen of 6 mg over 60 minutes instead of 15 minutes when CrCl is 30-50 ml/min. Furthermore, a dose reduction for patients whose CrCl is below 30 has been recommended for ibandronate (2 mg instead of 6 mg).

As a result, where available, ibandronate should be preferred to any other IV bisphosphonate since it appears to be non nephrotoxic and only necessitates dosage adjustment in patients with a CrCl or a eGFR lower than 30^{28,15}. Oral bisphosphonates may be an option of choice since they are renally well-tolerated.

Under the direction of the IRMA Scientific Committee, the IRMA Study has been coordinated by ICAR, a National Medical Advisory Service on the interactions between drugs and the kidney (*i.e.* drug dosage adjustment, drug nephrotoxicity, drug-drug interactions with immunosuppressive therapies) located in the Department of Nephrology at Pitié-Salpêtrière Hospital in Paris, France.

The authors are indebted to the physicians who took time from their busy work schedules to participate in the IRMA Study Group and to Roche France, thanks to whom ICAR services are available to French oncologists through an unrestricted educational grant.

IRMA Scientific Committee (Alphabetical):

P. Beuzeboc, Paris; G. Deray, Paris; J. Gligorov, Paris; V. Launay-Vacher, Paris; J.-F. Morere, Paris; S. Oudard, Paris; X. Pourrat, Tours; I. Ray-Coquard, Lyon, J.-P. Spano, Paris.

IRMA Study Group (Alphabetical):

Pr A. Adenis, Lille; Dr E. Banu, Paris; Dr P. Beuzeboc, Paris; Dr H. Boostandoost, Paris; Dr S. Clisant, Lille; Dr N. Colbert, Paris; Dr L-M. Dourthe, Metz; Dr D. Egret, Nantes; Dr J. Egreteau, Lorient; Dr J. Gligorov, Paris; Dr J-P. Guastalla, Lyon; Dr M-F. Jaeger, Metz; Dr F. Joly, Caen; Dr J. Ayllon, Paris; Dr C. Le Tourneau, Paris; Dr C.-B. Levache, Périgueux; Dr A. Lortholary, Nantes; Mr C. Maguire, Périgueux; Dr A. Monnier, Montbéliard; Pr J-F. Morere, Paris; Dr E. Ouahrani, Paris; Pr S. Oudard, Paris; Dr I. Ray-Coquard, Lyon; Pr O. Rixe, Paris; Dr E. Sevin, Caen; Dr L. Stefani, Grenoble, France.

TABLE 1. RENAL FUNCTION OF THE 1898 IRMA PATIENTS WITH BREAST CANCER USING THE COCKCROFT-GAULT AND THE AMDRD FORMULAE.

	Cockcroft & Gault CrCl (mL/min)		aMDRD eGFR (mL/min/1.73m2)	
	N	%	N	%
≥ 90	697	36.7%	733	38.6%
89-60	713	37.6%	817	43.0%
59-30	261	13.7%	141	7.4%
29-15	7	0.4%	5	0.3%
< 15	2	0.1%	2	0.1%
ND	218	11.5%	200	10.6%
		100%		100%

CrCl = Creatinine Clearance; eGFR = estimated glomerular filtration rate; N = Number of patients.

FIGURE 1. .COMPARISON OF SCREENING BREAST CANCER PATIENTS FOR RENAL INSUFFICIENCY WITH COCKCROFT-GAULT OR AMDRD.

aMDRD = abbreviated Modification of Diet in Renal Disease; RI = Renal Insufficiency
To be published online only

TABLE 2. CHARACTERISTICS OF BREAST CANCER PATIENTS IN THE IRMA STUDY WITH OR WITHOUT BONE METASTASES.

	Patients with BM (N=577)		Patients without BM (N=1321)		p-value
	Mean	SD	Mean	SD	
Age (years)	57.7*	11.2	53.9*	11.4	p<0.0001
Weight (kg)	63.6	13.5	64.5	13.3	p>0.05
SCR (μmol/L)	70.4	27.2	68.0	15.4	p>0.05
CG (mL/min)	84.1*	30.8	89.7*	29.5	p=0.0004
aMDRD (mL/min/1.73m²)	88.1	27.7	89.2	23.3	p>0.05
Hb (g/dL)	11.7*	1.5	12.1*	1.3	p<0.0001

BM = Bone Metastases; SCR = Serum Creatinine; CG = Cockcroft-Gault formula; aMDRD = abbreviated Modification of Diet in Renal Disease formula; Hb = Hemoglobinemia.

TABLE 3. RENAL FUNCTION OF IRMA BREAST CANCER PATIENTS WITH OR WITHOUT BONE METASTASES.

	Breast cancer patients with BM (N=577)		Breast cancer patients without BM (N=1321)	
	N	%	N	%
Cockcroft-Gault CrCl (mL/min)				
≥ 90	198	34.3%	499	37.8%
89-60	209	36.2%	504	38.2%
59-30	112	19.4%	149	11.3%
29-15	3	0.5%	4	0.3%
< 15	2	0.4%	0	0%
ND	53	9.2%	165	12.5%
aMDRD GFR (mL/min/1.73m2)				
≥ 90	212	36.7	521	39.4
89-60	260	45.1	557	42.2
59-30	53	9.2	88	6.7
29-15	3	0.5	2	0.2
< 15	2	0.4	0	0.0
ND	47	8.2	153	11.6

BM = Bone Metastases; N = Number of patients; CrCl = Creatinine clearance; GFR = Glomerular filtration rate

TABLE 4. COMPARISON OF THE PREVALENCE OF RI IN BREAST CANCER PATIENTS WITH OR WITHOUT BONE METASTASES, FOR PATIENTS WITH AVAILABLE DATA.

Renal Function	Patients with BM (N=524)	Patients without BM (N=1156)	p-value
CG < 90	62.2%	56.8%	p=0,0381
CG ≥ 90	37.8%	43.2%	
	(N=530)	(N=1168)	
aMDRD < 90	60.0%	55.4%	p>0,05
aMDRD ≥ 90	40.0%	44.6. %	

BM = Bone Metastases; CG = Cockcroft-Gault formula (mL/min); aMDRD = abbreviated Modification of Diet in Renal Disease formula (mL/min/1.73m²)

To be published online only

TABLE 5. HEMOGLOBIN VALUES IN IRMA BREAST CANCER PATIENTS WITH OR WITHOUT BONE METASTASES, FOR PATIENTS WITH AVAILAIBLE.

Hb Value	Patients with BM (N=539)	Patients without BM (N=1262)
>=11	69.2%	82.3%
9-11	26.9%	16.2%
<9	3.9%	1.5%

Hb = Hemoglobin (g/dL); BM = Bone Metastases

TABLE 6. ANTICANCER DRUGS PRESCRIBED TO BREAST CANCER PATIENTS IN THE IRMA STUDY (1702 TREATED PATIENTS – 3465 PRESCRIPTIONS).

INN	NUMBER OF PRESCRIPTIONS	PERCENTAGE OF THE TOTAL NUMBER OF PRESCRIPTIONS	NEED FOR DOSAGE ADJUSTMENT	POTENTIAL NEPHROTOXICITY
Cyclophosphamide	631	18.21%	Yes	No
5-FU	618	17.84%	No	No
Epirubicin	544	15.70%	No	Yes (isolated cases)
Docetaxel	379	10.94%	Yes	No
Vinorelbine	271	7.82%	Yes	No
Trastuzumab	251	7.24%	ND	Yes (isolated cases)
Doxorubicin	175	5.05%	No	Yes (isolated cases)
Paclitaxel	125	3.61%	No	Yes (isolated cases)
Zoledronate	112	3.23%	Yes	Yes
Gemcitabine	79	2.28%	No	Yes
Capecitabine	60	1.73%	Yes	No
Methotrexate	40	1.15%	Yes	Yes
Carboplatin	28	0.81%	Yes	Yes
Mitomycin C	19	0.55%	Yes	Yes
Oxaliplatin	13	0.38%	No	Yes
Tamoxifen	12	0.35%	No	Yes (isolated cases)

Vindesine	12	0.35%	No	No
Pirarubicin	11	0.32%	ND	Yes (isolated cases)
Letrozole	9	0.26%	No	No
Vinblastine	9	0.26%	No	No
Anastrozole	8	0.23%	No	No
Cisplatin	8	0.23%	Yes	Yes
Mitoxantrone	8	0.23%	No	No
Pamidronate	7	0.20%	ND	Yes
Exemestane	6	0.17%	No	No
Thiotepa	6	0.17%	ND	Yes
Irinotecan	4	0.12%	ND	Yes
Goserelin	4	0.12%	ND	No

Anticancer drugs prescribed less than 0.1% in IRMA breast cancer patients were (decreasing number of prescriptions): DHER vaccine, Fotemustin, Temozolomide, Etoposide, Ixabepilone, Clodronate, Edotecarine, Estramustine, Interferon, Interleukin-2, Melphalan, and Megestrol.
INN = International Non-proprietary Name; 5-FU = 5-fluorouracile.

TABLE 7. ANTICANCER DRUGS PRESCRIBED TO BREAST CANCER PATIENTS WITH BONE METASTASES IN THE IRMA STUDY (529 TREATED PATIENTS – 912 PRESCRIPTIONS).

INN	NUMBER OF PRESCRIPTIONS	PERCENTAGE OF THE TOTAL NUMBER OF PRESCRIPTIONS	NEED FOR DOSAGE ADJUSTMENT	POTENTIAL NEPHROTOXICITY
Docetaxel	125	13.7%	Yes	No
Vinorelbine	121	13.3%	Yes	No
Zoledronate	107	11.7%	Yes	Yes
Trastuzumab	87	9.5%	ND	Yes (isolated cases)
5-FU	78	8.6%	No	No
Paclitaxel	62	6.8%	No	Yes (isolated cases)
Cyclophosphamide	57	6.3%	Yes	No
Epirubicin	47	5.2%	No	Yes (isolated cases)
Gemcitabine	40	4.4%	No	Yes
Doxorubicin	35	3.8%	No	Yes (isolated cases)
Capecitabine	34	3.7%	Yes	No
Methotrexate	19	2.1%	Yes	Yes
Carboplatin	16	1.8%	Yes	Yes
Mitomycin C	12	1.3%	Yes	Yes
Oxaliplatin	7	0.8%	No	Yes
Letrozole	6	0.7%	No	No
Pamidronate	6	0.7%	ND	Yes

Vindesine	6	0.7%	No	No
Cisplatin	5	0.5%	Yes	Yes
Thiotepa	5	0.5%	ND	Yes
Vinblastine	5	0.5%	No	No
Irinotecan	4	0.4%	ND	Yes
Mitoxantrone	4	0.4%	No	No
Exemestane	4	0.4%	No	No
Pirarubicin	4	0.4%	ND	Yes (isolated cases)
Anastrozole	3	0.3%	No	No
Tamoxifen	3	0.3%	No	Yes (isolated cases)
Goserelin	3	0.3%	ND	No
Etoposide	2	0.2%	Yes	No

Anticancer drugs prescribed less than 0.1% in IRMA breast cancer patients with bone metastases were (decreasing number of prescriptions): Clodronate, Edotecarine, Melphalan, Megestrol, Temozolomide.

INN = International Non-proprietary Name; 5-FU = 5-fluorouracile.

To be published online only

TABLE 8. ANTICANCER DRUGS PRESCRIBED TO BREAST CANCER PATIENTS WITHOUT BONE METASTASES IN THE IRMA STUDY (1173 TREATED PATIENTS – 2553 PRESCRIPTIONS).

INN	NUMBER OF PRESCRIPTIONS	PERCENTAGE OF THE TOTAL NUMBER OF PRESCRIPTIONS	NEED FOR DOSAGE ADJUSTMENT	POTENTIAL NEPHROTOXICITY
Cyclophosphamide	574	22.48%	Yes	No
5-FU	540	21.15%	No	No
Epirubicin	497	19.47%	No	Yes (isolated cases)
Docetaxel	254	9.95%	Yes	No
Trastuzumab	164	6.42%	ND	Yes (isolated cases)
Vinorelbine	150	5.88%	Yes	No
Doxorubicin	140	5.48%	No	Yes (isolated cases)
Paclitaxel	63	2.47%	No	Yes (isolated cases)
Gemcitabine	39	1.53%	No	Yes
Capecitabine	26	1.02%	Yes	No
Methotrexate	21	0.82%	Yes	Yes
Carboplatin	12	0.47%	Yes	Yes
Tamoxifen	9	0.35%	No	Yes (isolated cases)
Mitomycin C	7	0.27%	Yes	Yes
Pirarubicin	7	0.27%	ND	Yes (isolated cases)
Oxaliplatin	6	0.24%	No	Yes

Vindesine	6	0.24%	No	No
Anastrozole	5	0.20%	No	No
Zoledronate	5	0.20%	Yes	Yes
Mitoxantrone	4	0.16%	No	No
Vinblastine	4	0.16%	No	No
Cisplatin	3	0.12%	Yes	Yes
Letrozole	3	0.12%	No	No

Anticancer drugs prescribed less than 0.1% in IRMA breast cancer patients without Bone Metastases were (decreasing number of prescriptions): DHER vaccine, Fotemustin, Ixabepilone, Estramustin, Interferon, Interleukin-2, Pamidronate, Temozolomide, Thiotepa, Goserelin.

INN = International Non-proprietary Name; 5-FU = 5-fluorouracile.

To be published online only

TABLE 9. COMPARISON OF THE PRESCRIPTION PATTERN OF DRUGS THAT NECESSITATE DOSAGE ADJUSTMENT IN RI OR MAY BE NEPHROTOXIC IN BREAST CANCER PATIENTS WITH OR WITHOUT BONE METASTASES.

	Patients with BM	Patients without BM	
Prescriptions	N=912	N=2553	
<i>Dosage adjustment in RI</i>			
Yes	54.82%	41.28%	
ND	12.28%	7.01%	
<i>Yes or ND</i>	<i>67.10%</i>	<i>48.29%</i>	<i>p<0.0001</i>
No	32.89%	51.70%	
<i>Nephrotoxicity</i>			
Yes	50.66%	38.39%	
ND	0.11%	0.07%	
<i>Yes or ND</i>	<i>50.77%</i>	<i>38.46%</i>	<i>p<0.0001</i>
No	49.23%	61.54%	
Patients (treated)	N=529	N=1173	
<i>Dosage adjustment in RI</i>			
Yes	74.48%	82.86%	
ND	10.96%	9.80%	

Yes or ND	85.44%	92.66%	p<0.0001
No	14.56%	7.33%	
<i>Nephrotoxicity</i>			
Yes	72.20%	78.69%	
ND	0.20%	0.17%	
Yes or ND	72.4%	78.86%	p=0.0035
No	27.6%	21.14%	

BM = Bone Metastases; RI = Renal Insufficiency; ND = No Data

The authors have no conflict of interest to declare which would be relevant to this article.

References

1. Matas AJ, Simmons RL, Kjellstrand CM, Buselmeier TJ, Najarian JS (1975) Increased incidence of malignancy during chronic renal failure. *Lancet* 1:883–886 [Medline](#). doi:10.1016/S0140-6736(75)91684-0
2. Sutherland GA, Glass J, Gabriel R (1977) Increased incidence of malignancy in chronic renal failure. *Nephron* 18:182–184 [Medline](#). doi:10.1159/000130762
3. Cengiz K (2002) Increased incidence of neoplasia in chronic renal failure (20-year experience). *Int Urol Nephrol* 33:121–126 [Medline](#). doi:10.1023/A:1014489911153
4. Jones CA, McQuillan GM, Kusek JW, et al (1998) Serum creatinine levels in the US population: third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 32:992–999 [Medline](#). doi:10.1016/S0272-6386(98)70074-5
5. Launay-Vacher V, Oudard S, Janus N, Gligorov J, Pourrat X, Rixe O, et al, Renal Insufficiency and Cancer Medications (IRMA) Study Group. (2007) Prevalence of Renal Insufficiency in cancer patients and implications for anticancer drug management: the renal insufficiency and anticancer medications (IRMA) study. *Cancer* 110(6):1376–1384 [Medline](#). doi:10.1002/cncr.22904
6. National Kidney Foundation (2002) K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 39(2)(Suppl 1):S1–S266 [Medline](#).
7. Levey AS, Eckardt KU, Tsukamoto Y, et al (2005) Definition and classification of chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 67:2089–2100 [Medline](#). doi:10.1111/j.1523-1755.2005.00365.x
- ⁸ Torrisi R, Balduzzi A, Ghisini R, Rocca A, Bottiglieri L, Giovanardi F, Veronesi P, Luini A, Orlando L, Viale G, Goldhirsch A, Colleoni M. Tailored preoperative treatment of locally advanced triple negative (hormone receptor negative and HER2 negative) breast cancer with epirubicin, cisplatin, and infusional fluorouracil followed by weekly paclitaxel. *Cancer Chemother Pharmacol* 2007 Dec 7; [Epub ahead of print]
9. Coudert BP, Largillier R, Arnould L, Chollet P, Campone M, Coeffic D, et al (2007) Multicenter phase II trial of neoadjuvant therapy with trastuzumab, docetaxel, and carboplatin for human epidermal growth factor receptor-2-overexpressing stage II or III breast cancer: results of the GETN(A)-1 trial. *J Clin Oncol* 25(19):2678–2684 [Medline](#). doi:10.1200/JCO.2006.09.9994

10. Robert N, Leyland-Jones B, Asmar L, Belt R, Ilegbodun D, Loesch D, et al (2006) Randomized phase III study of trastuzumab, paclitaxel, and carboplatin compared with trastuzumab and paclitaxel in women with HER-2-overexpressing metastatic breast cancer. *J Clin Oncol* 24(18):2786–2792 [Medline](#). doi:10.1200/JCO.2005.04.1764
11. Hurley J, Doliny P, Reis I, Silva O, Gomez-Fernandez C, Velez P, et al (2006) Docetaxel, cisplatin, and trastuzumab as primary systemic therapy for human epidermal growth factor receptor 2-positive locally advanced breast cancer. *J Clin Oncol* 24(21):3515.
12. Slamon D, Eiermann W, Robert N, et al Phase III randomized trial comparing doxorubicin and cyclophosphamide followed by docetaxel and trastuzumab (AC TH) with docetaxel, carboplatin and trastuzumab (TCH) in HER2-positive early breast cancer patients: BCIRG 006 study. 28th Annual San Antonio Breast Cancer Symposium. San Antonio 2005, abstract 1
13. Body JJ (2006) Breast cancer: bisphosphonate therapy for metastatic bone disease. *Clin Cancer Res* 12:6258s–6263s [Medline](#). doi:10.1158/1078-0432.CCR-06-0840
14. Tripathy AD, Diel I, Body JJ (2004) Ibandronate reduces skeletal morbidity in patients with breast cancer. *Semin Oncol* 31(5)(Suppl 10):64–66. doi:10.1053/j.seminoncol.2004.07.025
15. Rosen LS, Gordon D, Kaminski M, Howell A, Belch A, Mackey J, et al (2003) Long-term efficacy and safety of zoledronic acid compared with pamidronate disodium in the treatment of skeletal complications in patients with advanced multiple myeloma or breast carcinoma: a randomized, double-blind, multicenter, comparative trial. *Cancer* 98(8):1735–1744 [Medline](#). doi:10.1002/cncr.11701
16. Chang JT, Green L, Beitz J (2003) Renal failure with the use of zoledronic acid. *N Engl J Med* 349(17):1676–1679 [Medline](#). doi:10.1056/NEJM200310233491721
17. Cockcroft DW, Gault MH (1976) Prediction of CrCl from serum creatinine. *Nephron* 16:31–41 [Medline](#). doi:10.1159/000130554
18. Levey AS, Greene T, Kusek JW, et al. A simplified equation to predict glomerular filtration rate from serum creatinine. *J Am Soc Nephrol* 2000;11:0828. abstract.
19. Aronoff GR, Berns JS, Brier ME, et al Drug prescribing in renal failure. Dosing guidelines for adults. 4th ed. USA: American College of Physicians – American Society of Internal Medicine; 1999.
20. Launay-Vacher V, Karie S, Deray G GPR Anticancéreux. Guide de prescription des médicaments chez le patient insuffisant rénal. 3^{ème} édition. France: Méditations International; 2005.

21. Lauro S, Lalle M, D'Andrea MR, Vecchione A, Frati L (1994) Nephrotic syndrome and adjuvant treatment with tamoxifen for early breast cancer. Case report and review of the literature. *Anticancer Res* 4(5B):171–172.

22. Merouani A, Davidson SA, Schrier RW (1997) Increased nephrotoxicity of combination taxol and cisplatin chemotherapy in gynecologic cancers as compared to cisplatin alone. *Am J Nephrol* 17(1):53–58 [Medline](#).

23. Nikolsky E, Mehran R, Lasic Z, et al (2005) Low hematocrit predicts contrast-induced nephropathy after percutaneous coronary interventions. *Kidney Int* 67:706–713 [Medline](#). doi:10.1111/j.1523-1755.2005.67131.x

24. Body JJ, Coleman R, Clezardin P, Ripamonti C, Rizzoli R, Aapro M, International Society of Geriatric Oncology. (2007) International Society of Geriatric Oncology (SIOG) clinical practice recommendations for the use of bisphosphonates in elderly patients. *Eur J Cancer* 43:852–858 [Medline](#). doi:10.1016/j.ejca.2006.12.006

25. Lichtman S, Wildiers H, Launay-Vacher V, Steer C, Chatelut E, Aapro M (2007) International Society of Geriatric Oncology (SIOG) recommendations for the adjustment of dosing in elderly cancer patients with renal insufficiency. *Eur J Cancer* 43:14–34 [Medline](#). doi:10.1016/j.ejca.2006.11.004

26. Launay-Vacher V, Chatelut E, Lichtman S, Wildiers H, Steer C, Aapro M (2007) Renal insufficiency in elderly cancer patients: International Society of Geriatric Oncology (SIOG) clinical practice recommendations. *Ann Oncol* 18:1314–1321 [Medline](#). doi:10.1093/annonc/mdm011

27. Aapro M, Abrahamsson PA, Body JJ, Coleman RE, Colomer R, Costa L, et al (2008) Guidance on the use of bisphosphonates in solid tumours: recommendations of an international expert panel. *Ann Oncol* 19(3):420–432 [Medline](#). doi:10.1093/annonc/mdm442

28. Body JJ, Diel IJ, Tripathy D, Bergstrom B (2006) Intravenous ibandronate does not affect time to renal function deterioration in patients with skeletal metastases from breast cancer: phase III trial results. *Eur J Cancer Care (Engl)* 15(3):299–302 [Medline](#). doi:10.1111/j.1365-2354.2005.00641.x