Delirium, thrombocytopenia, insomnia, and mild liver damage associated with MAOI withdrawal

Alfredo Bellon, John H. Coverdale

► To cite this version:

Alfredo Bellon, John H. Coverdale. Delirium, thrombocytopenia, insomnia, and mild liver damage associated with MAOI withdrawal. *European Journal of Clinical Pharmacology*, 2009, 65 (12), pp.1269-1270. 10.1007/s00228-009-0711-y . hal-00534988

HAL Id: hal-00534988

<https://hal.science/hal-00534988>

Submitted on 11 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Delirium, thrombocytopenia, insomnia, and mild liver damage associated with MAOI withdrawal

Alfredo Bellon • John H. Coverdale

Received: 24 June 2009 / Accepted: 21 July 2009 / Published online: 5 August 2009
© Springer-Verlag 2009

Dear Sir,

Monoamine oxidase inhibitors (MAOIs) are effective second-line treatment options for depression [1, 2]. A recent study showed that tranylcypromine (TCP), a nonselective and irreversible MAOI, remains commonly used in standard psychiatric clinics [3]. The risk of hypertensive crisis due to drug–food interactions when MAOIs are prescribed is widely known. Less well appreciated is the fact that MAOIs can cause addiction and that a particular patient population may be at high risk. Here we present a case of TCP dependence that brings together all signs and symptoms so far described after TCP withdrawal, including delirium, thrombocytopenia, insomnia, and mild liver damage.

Case

A 30-year-old woman diagnosed with major depressive disorder and borderline personality disorder and with past history of alcohol, marihuana, amphetamine, and cocaine abuse, presented to the emergency room. For the preceding 1–2 years, she had been treated with TCP and titrated to 60 mg a day, along with chlorpromazine 350 mg at night for “sleep.” Two months after the initiation of TCP, she steadily increased her dose to 170 mg a day. In order to maintain her TCP intake, the patient visited multiple

physicians in search for prescriptions. Three days prior to admission, she ran out of TCP and gradually became agitated, irritable, and paranoid. When she presented to the emergency room, she was confused, unable to recognize her mother, and oriented to person only. She was described as responding to internal stimuli and laughing inappropriately. On physical examination, she had a mild tachycardia, a severe tremor, and extensive bruising involving upper and lower extremities. Laboratory evaluation revealed a thrombocytopenia of 72,000/ μ l.

On further inpatient evaluation, her cognitive status improved. By the second day, her platelets had decreased to 55,000/ μ l, and on the third day, these cells dropped further to 25,000/ μ l. During her third day of hospitalization, her partial thromboplastin time (PTT) extended to 40.9 s and her liver function tests showed an aspartate transaminase (AST) of 106 IU/l. A liver ultrasound revealed fatty-acid infiltrates. On day 6, her platelets returned to normal limits. Nine days after admission, the patient was able to sleep 6 h without any medication, and her tremor was completely resolved.

Discussion

This abrupt change in behavior and cognition coincides with previously reported cases of delirium after acute withdrawal from TCP [4–6]. It has been proposed that delirium and paranoia could result from acute removal of the sympathomimetic component of MAOIs on presynaptic receptors [7]. Under normal conditions, this presynaptic effect aims to diminish the release of catecholamines. It is likely that when MAOIs are suddenly stopped, norepinephrine and dopamine acutely increase in the synapse, rendering patients susceptible to psychosis and delirium [6, 7].

A. Bellon (✉)
Centre de Psychiatrie et Neurosciences, Psychiatry,
Paris, France
e-mail: alfredobellon@yahoo.com

J. H. Coverdale
Menninger Department of Psychiatry & Behavioral Sciences,
Baylor College of Medicine,
Houston, TX, USA

Thrombocytopenia has also been previously described after a sudden removal of TCP [4, 5, 8]. Even though the reason for the thrombocytopenia is unclear, this has been explained as a sympathomimetic effect secondary to TCP withdrawal that results in spleen venous pooling and also as a consequence of the long-term inhibition of MAO in platelets which, in turn, may lead to a reduced life span of these cells [9].

The AST increase, PPT extension, as well as the fatty-acid infiltrate could have originated from MAOI abuse [4, 10]. This possibility is further supported by the AST and PTT normalization after being off TCP for several consecutive days. On the other hand, insomnia could be associated to a rebound from rapid eye movement sleep suppression, as it has been previously observed when MAOIs are abruptly stopped [11].

The combination of a past history of substance abuse and a diagnosis of either borderline or histrionic personality disorder seems to render patients at higher risk of developing MAOI dependence [11–17], perhaps because these medications are either structurally similar to amphetamines [6, 17] or are, like selegiline, metabolized into excitatory compounds [18].

Clinicians are exhorted to ponder the addictive potential of MAOIs when prescribing these drugs.

Acknowledgements AB receives support from FRM France, CONACYT #74641 and The Young Minds in Psychiatry Award.

References

1. Parker GRK, Wilhelm K, Mitchell P (2001) Assessing the comparative effectiveness of antidepressant therapies: a prospective clinical practice study. *J Clin Psychiatry* 62(2):17–125
2. Stewart JW, Thase ME (2007) Treating DSM-IV depression with atypical features. *J Clin Psychiatry* 68(4):e10
3. Adli M, Pilhatsch M, Bauer M, Koberle U, Ricken R, Janssen G et al (2008) Safety of high-intensity treatment with the irreversible monoamine oxidase inhibitor tranylcypromine in patients with treatment-resistant depression. *Pharmacopsychiatry* 41(6):252–257
4. Briggs NC, Jefferson JW, Koenecke FH (1990) Tranylcypromine addiction: a case report and review. *J Clin Psychiatry* 51(10):426–429
5. Davids E, Roschke J, Klawe C, Grunder G, Schmoldt A (2000) Tranylcypromine abuse associated with delirium and thrombocytopenia. *J Clin Psychopharmacol* 20(2):270–271
6. Dilsaver SC (1988) Monoamine oxidase inhibitor withdrawal phenomena: symptoms and pathophysiology. *Acta Psychiatr Scand* 78(1):1–7
7. Dilsaver SC (1990) Heterocyclic antidepressant, monoamine oxidase inhibitor and neuroleptic withdrawal phenomena. *Prog Neuropsychopharmacol Biol Psychiatry* 14(2):137–161
8. Chatterjee A, Tosyali MC (1995) Thrombocytopenia and delirium associated with tranylcypromine overdose. *J Clin Psychopharmacol* 15(2):143–144
9. Szelenyi A, Albrecht J (1998) Tranylcypromine abuse associated with an isolated thrombocytopenia. *Pharmacopsychiatry* 31(6):238–240
10. Bandt C, Hoffbauer FW (1964) Liver injury associated with tranylcypromine therapy. *JAMA* 188:752–753
11. Le Gassicke JL, Ashcroft GW ED, Evans JJ, Oswald I, Riston EB (1965) The clinical state, sleep and amine metabolism of tranylcypromine ('Parnate') addict. *Br J Psychiatry* 111:357–364
12. Ben-Arie O, George GC (1979) A case of tranylcypromine ('Parnate') addiction. *Br J Psychiatry* 135:273–274
13. Brady KT, Lydiard RB, Kellner C (1991) Tranylcypromine abuse. *Am J Psychiatry* 148(9):1268–1269
14. Davidson J, Pelton S (1986) Forms of atypical depression and their response to antidepressant drugs. *Psychiatry Res* 17(2):87–95
15. Kayser A, Robinson DS, Nies A, Howard D (1985) Response to phenelzine among depressed patients with features of hysteroid dysphoria. *Am J Psychiatry* 142(4):486–488
16. Shopsin B, Kline NS (1976) Monoamine oxidase inhibitors: potential for drug abuse. *Biol Psychiatry* 11(4):451–456
17. Vartzopoulos D, Krull F (1991) Dependence on monoamine oxidase inhibitors in high dose. *Br J Psychiatry* 158:856–857
18. Fujita Y, Takahashi K, Takei M et al (2008) Detection of levorotatory methamphetamine and levorotatory amphetamine in urine after ingestion of an overdose of selegiline. *Yakugaku Zasshi* 128(10):1507–1512