Use of polymerase chain reactions for the detection of *M. gallisepticum*, *M. imitans*, *M. iowae*, *M. meleagridis* and *M. synoviae* in birds of prey

Michael Lierz, Nils Hagen, Dörte Lüschoff, Hafez Mohamed Hafez

► To cite this version:

Michael Lierz, Nils Hagen, Dörte Lüschoff, Hafez Mohamed Hafez. Use of polymerase chain reactions for the detection of *M. gallisepticum*, *M. imitans*, *M. iowae*, *M. meleagridis* and *M. synoviae* in birds of prey. *Avian Pathology*, 2008, 37 (05), pp.471-476. 10.1080/03079450802272952 . hal-00540126

HAL Id: hal-00540126

<https://hal.science/hal-00540126>

Submitted on 26 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Use of polymerase chain reactions for the detection of *M. gallisepticum*, *M. imitans*, *M. iowae*, *M. meleagridis* and *M. synoviae* in birds of prey

Journal:	<i>Avian Pathology</i>
Manuscript ID:	CAVP-2008-0031.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	23-Apr-2008
Complete List of Authors:	Lierz, Michael; Free University of Berlin, Institute for Poultry Diseases Hagen, Nils Lüschow, Dörte; Institute of Poultry Diseases, Free University Berlin Hafez, Hafez; Institute of Poultry Diseases, Free University of Berlin, Koserstr. 21, 14195 Berlin; Germany., Poultry Diseases
Keywords:	Poultry, falconiformes, mycoplasma, molecular biology

SCHOLARONE™
Manuscripts

Cavp-2008-0031.R1

Use of polymerase chain reactions to detect *Mycoplasma gallisepticum*, *Mycoplasma imitans*, *Mycoplasma iowae*, *Mycoplasma meleagridis* and *Mycoplasma synoviae* in birds of prey

Lierz, M. *, N. Hagen, D. Lueschow and H.M. Hafez

Institute for Poultry Diseases, Free University of Berlin, Koenigsweg 63, 14163 Berlin, Germany

Short title: *Mycoplasma* PCRs for raptor samples

Received: 25 March 2008

* To whom correspondence should be addressed. Te.: +49 30 83862699. Fax: +49 30 83862690. E-mail: lierz.michael@vetmed.fu-berlin.de;

Cavp-2008-0031.R1

Use of polymerase chain reactions to detect *Mycoplasma gallisepticum*, *Mycoplasma imitans*, *Mycoplasma iowae*, *Mycoplasma meleagridis* and *Mycoplasma synoviae* in birds of prey

Lierz, M. *, N. Hagen, D. Lueschow and H.M. Hafez

Abstract

Certain *Mycoplasma* spp. are pathogens of poultry, but little is known of the role of mycoplasmas in disease of birds of prey. Therefore species-specific PCRs for the detection of the poultry pathogens *Mycoplasma gallisepticum*, *Mycoplasma imitans*, *Mycoplasma iowae*, *Mycoplasma meleagridis* and *Mycoplasma synoviae* were evaluated for use in birds of prey. The specificities of the PCR methods were established using avian and other mycoplasmas and also selected walled bacteria. The sensitivities of the different PCR assays varied between 100 fg and 10 pg DNA. Fifty three tracheal swabs from healthy captive and free ranging birds of prey were then investigated using these PCRs and in no case was an amplicon obtained for *M. gallisepticum*/*M. imitans*, *M. iowae* or *M. synoviae*. Species-specific primers for *M. meleagridis* amplified a product from eight birds of prey but restriction enzyme analysis as well as sequencing of PCR products demonstrated these results to be false positives. Alignment studies of the sequenced products with the 16S rRNA gene sequence of various *Mycoplasma* species in GenBank demonstrated an identity of 91% to *M. meleagridis* but of 98% to *M. buteonis* or *M. gallopavonis*. Isolation and attempted identification of these mycoplasmas suggested that it may be a previously unrecognised species.

Introduction

Four *Mycoplasma* species are recognised pathogens of domestic poultry, causing economic loss. These are *M. gallisepticum*, *M. iowae*, *M. meleagridis* and *M. synoviae*. Traditionally mycoplasmas were thought to have a restricted host range but the more recent literature has cast doubt this and even a mammalian *Mycoplasma* (*Mycoplasma capricolum*-like) was isolated from a chicken (Benčina *et al.*, 2006).

M. gallisepticum has been isolated from a variety of different avian species including galliformes (Benčina *et al.*, 2003), anseriformes (Ibrahim *et al.*, 2000), psittaciformes (Bozeman *et al.*, 1984), columbiformes (Benčina *et al.*, 1987), passeriformes (Dhondt *et al.*, 1998; Hartup *et al.*, 1998; Mikaelian *et al.*, 2001) and birds of prey (Poveda *et al.*, 1990 a,b; Morishita *et al.*, 1997). *M. iowae*, usually found in turkeys, has been isolated from a range of free-flying birds (Amin, 1977), geese (Lin *et al.*, 1995), a yellow napped amazon (*Amazona ochrocephala auropalliata*) (Bozeman *et al.*, 1984) and even an apple seed (Grau *et al.*, 1991).

In addition to its usual host the turkey, there have been repeated isolations of *M. meleagridis* from birds of prey (Lierz *et al.*, 2000a, 2002) and there have also been unconfirmed reports of isolations from quail and peacocks (Simecka *et al.*, 1992). Apart from chicken and turkey, *M. synoviae* was isolated from other galliformes (Benčina *et al.*, 1987; Poveda *et al.*, 1990b) as well as from anseriformes (Benčina *et al.*, 1988a,b; Tjong, 1990), pigeons (Benčina *et al.*, 1987), a house sparrow (*Passer domesticus*) (Poveda *et al.*, 1990b).

The wide host range of the poultry pathogenic *Mycoplasma* species may present a risk for transmission of these pathogens to poultry. One host group of interest in this context is birds of prey, because some of these mycoplasmas have already been found in them. These birds live in close proximity to humans and can be found regularly near poultry farms because they feed on rodents (common buzzards (*Buteo buteo*), kestrels (*Falco tinnunculus*)) or on the poultry (Northern goshawks (*Accipiter gentilis*), peregrines (*Falco peregrinus*)). In captivity, birds of prey are regularly fed with poultry by-products, especially day old cockerels, which may represent a source of infection.

In general mycoplasmas are common in birds of prey (Morishita *et al.*, 1997, Lierz *et al.*, 2000b, 2002, 2008a,b) but the *Mycoplasma* species that are pathogenic for poultry are only occasionally reported. Therefore the presence of *M. gallisepticum*, *M. iowae*, *M. meleagridis* and *M. synoviae* in captive and free-ranging raptors from Germany was investigated using polymerase chain reactions (PCRs). Such PCRs are already described within the literature but have focussed on their use in poultry and specificity testing is therefore usually limited to certain

reference strains. The aim of the present investigation was to evaluate the PCRs extensively before using them in an avian population of unknown *Mycoplasma* status.

The PCR described by Marois *et al.* (2000) was evaluated for the detection of *M. synoviae* and that described by Boyle *et al.* (1995) for *M. meleagridis*. The PCRs to detect *M. gallisepticum* and *M. iowae* were designed as part of this study. *M. gallisepticum* and *M. imitans* share a 99% identity in their 16S rRNA gene and the PCR was designed to allow their differentiation using restriction enzyme analysis of the PCR product as suggested earlier (Kempf, 1997). This was considered important, as the presence of *M. imitans* cannot be excluded in birds of prey.

Material and Methods

PCR primer design. Species-specific primers for the detection of *M. gallisepticum*/*M. imitans* and *M. iowae* were designed based on comparison and alignment of GenBank sequences (16S rRNA gene) of the 24 avian *Mycoplasma* species using the Clustal W software program (European Molecular Biology Laboratory, Heidelberg, Germany) (Higgins *et al.*, 1996). Species-specific primers for the detection of *M. synoviae* and *M. meleagridis* were taken from the literature (Marois *et al.*, 2000; Boyle *et al.*, 1995). The specificity of each PCR was confirmed using avian *Mycoplasma* strains, certain other *Mycoplasma* species and also walled bacteria occurring commonly in bird tracheas (Table 1). The optimal annealing temperature of each PCR was determined using a gradient-PCR (iCycler iQ™, Biorad, Hilden, Germany). The sensitivity of each PCR was evaluated using 10 fold serial dilutions of specific *Mycoplasma*-DNA as well as establishing the number of colony-forming units (CFU) of the corresponding *Mycoplasma* strain (Albers & Fletcher 1982).

Field sample DNA. The DNA extracts from tracheal swabs from 53 healthy captive and free-ranging birds of prey (Table 2) of two previous studies (Lierz *et al.*, 2008 a,b) were investigated. The samples selected had been positive for mycoplasmal DNA using a *Mycoplasma* genus-specific PCR. DNA was originally extracted from the swabs using the QIAmp®DNA Mini Kit (Qiagen, Hilden, Germany) using the manufacturer's instructions, and as modified by Lüscho *et al.* (2004). Briefly, 200 µl of sample (swab diluted in PBS buffer) was mixed with 20 µl proteinase K and 200 µl lysis buffer (AL). After incubation for 1 h at 56 °C and 10 min at 70 °C, 200µl ethanol was added. The mixture was put on a spin column and centrifuged at 6000 g for 1

min. The spin column was washed with 500 µl wash buffer (AW1) and centrifuged as before. After a second wash step with 500 µl with buffer (AW2) and centrifugation at 20,000 g for 3 min, the DNA was eluted from the column by the addition of 100 µl elution buffer (AE) before further incubation for 1 min, and a final centrifugation at 6000 g for 1 min.

DNA extraction from *Mycoplasma* cultures for specificity and sensitivity investigations was carried out using the QIAmp®DNA Mini Kit (Qiagen, Hilden, Germany). The concentration of DNA was measured photometrically (wavelength: 260 nm).

PCRs. These were carried out using Taq PCR Master Mix (Qiagen, Hilden, Germany) according to the manufacturer's instructions, but with only half of the recommended amount of reaction sample. Briefly, a 50 µl reaction sample contained 25 µl Taq PCR Master Mix, plus 12.5 pmol of primers (*M. gallisepticum*/ *M. imitans*, *M. iowae*) or 25 pmol of primers (*M. synoviae*, *M. meleagridis*), and 50 ng DNA and nuclease-free water up to 50 µl. Amplification was conducted after initial denaturation for 4 min at 94 °C followed by 35 cycles consisting of denaturation (30 sec, 94°C), different annealing temperatures according to the primers used (Table 3), synthesis (30 sec, 72°C) and a final polymerization for 10 min at 72°C. The amplified PCR product (10 µl) was then separated by agarose gel electrophoresis (0.8% for *M. gallisepticum*/*M. imitans*; *M. meleagridis* and 2% for *M. synoviae* and *M. iowae*), stained with ethidium bromide and viewed by ultraviolet transillumination.

Specificity of PCR results from field samples. To ensure that amplicons from field samples were specific, they were digested by different restriction enzymes (Table 4) and the results compared to the restriction enzyme pattern of amplicons obtained from the corresponding positive control strain which had been established earlier. As the *M. gallisepticum* PCR is also expected to amplify a product with *M. imitans*, amplicons of this PCR restriction enzymes were selected to provide differentiation (Table 4).

In addition selected amplicons of field samples were sequenced. The PCR product was gel purified with the Invisorb Spin DNA Extraction Kit (Invitek, Berlin, Germany) and sequenced using the same forward and reverse primers (Agowa GmbH, Berlin, Germany). The sequence of the product was compared to the 16S rRNA gene sequences of *Mycoplasma* spp. in GenBank using the software program Blast (NCBI, USA) (Altschul *et al.*, 1990).

Isolation and identification. *Mycoplasma* isolates from the tracheal swabs of the birds whose amplicon was sequenced were used for single colony subcultures. These isolates were obtained

in previous studies (Lierz *et al.*, 2008a, b). The fifth single colony subculture was produced using liquid and agar media produced according to Bradbury (1998) without adding thallium acetate and tested by the above PCRs. If positive, an immunobinding assay (IBA) was conducted according to Kotani & McGarrity (1985), using antiserum against the following avian *Mycoplasma* reference strains: *M. buteonis*, *M. corogypsi*, *M. falconis*, *M. gallisepticum*, *M. gypis*, *M. iowae*, *M. lipofaciens*, *M. meleagridis*, *M. synoviae*.

One isolate was subjected to a sequence analysis of its 16S rRNA gene. DNA extraction of the pure culture as well as PCR of the 16S rRNA gene was performed according to Tanner *et al.* (2000) and Blackwood *et al.* (2003), using the 27F (forward) and 1492R (reverse) primers. After purification with the Invisorb Spin DNA Extraction Kit (Invitek, Berlin, Germany) the PCR products were sequenced by using the same forward and reverse primers (Agowa GmbH, Berlin, Germany). The sequences were compared to the 16S rRNA gene sequences of *Mycoplasma* spp. provided by GenBank as well as to the sequences obtained directly from the amplicons in the above-used PCRs, using the software programs Blast (NCBI, USA) (Altschul *et al.*, 1990) and Clustal W (Higgins *et al.*, 1996).

Results

Establishment of PCRs. The primers for the species-specific PCRs, their binding locations within the 16S rRNA gene, the optimal annealing temperatures, the amplicon sizes and the PCR sensitivities are shown in Table 3. The *Mycoplasma* species-specific PCRs amplified fragments of the expected size from the DNA of the corresponding *Mycoplasma* strain only, and not from other mycoplasmas, acholeplasmas or the walled bacteria tested. The *M. gallisepticum* PCR amplified a product of same size from *M. imitans* as expected but it was possible to differentiate between the two species using the *MseI* and *AseI* restriction enzymes (Figure 1).

Investigation of field samples. No sample gave a positive result using the species-specific PCRs for *M. gallisepticum*/*M. imitans*, *M. iowae* or *M. synoviae*. The PCR to detect *M. meleagridis* amplified a product from eight samples (three peregrines; one kestrel, one Barbary falcon (*Falco pelegrinoides*), one saker (*Falco cherrug*), one gyr-saker hybrid and one Marsh harrier (*Circus aeruginosus*)).

Restriction enzyme and sequence analysis. None of the eight amplicons obtained by the *M. meleagridis* PCR was digested using the restriction enzyme (*Hind*III) whereas the amplicon of the *M. meleagridis* reference strain cleaved into the expected size (Figure 2). Sequence analysis of three amplicons (334 bp) obtained from the *M. meleagridis*-specific PCR (one Barbary falcon, GD 9, GenBank Acc. No: EU339238); one kestrel, M 14, GenBank Acc.No: EU339239; one peregrine, M 26, GenBank Acc.No: EU339240) revealed a 100% identity between the three products. Alignment analysis with the 16s rRNA gene sequence of *M. meleagridis* (GenBank Acc. No: L24106) demonstrated only a 91% identity, but a 98.5% match with the 16S rRNA gene sequence of *M. buteonis* (GenBank Acc.No: AF412971) or *M. gallopavonis* (GenBank Acc.No: AF412980) (98.1%).

Isolation and identification. *Mycoplasma* isolates obtained from the same three birds where the amplicon was sequenced, gave a positive result in the *M. meleagridis* PCR, but could not be identified in the IBA. As all above sequenced amplicons showed a similar sequence, the 16S rRNA gene of only one of these isolates (M 26 from the peregrine) was sequenced, yielding a product of 1335bp (GenBank Acc.No: EU660528). The sequence included the sequences obtained from the *M. meleagridis* PCR amplicons and showed a 98% match with the 16S rRNA gene sequence of *M. buteonis* (GenBank Acc.No: AF412971) or *M. gallopavonis* (GenBank Acc.No: AF412980).

Discussion

The poultry pathogens *M. gallisepticum*, *M. iowae*, *M. meleagridis* and *M. synoviae* have been isolated from a variety of different hosts (Bradbury, 2006) including *M. gallisepticum* and *M. meleagridis* from birds of prey (Poveda *et al.*, 1990 a, b; Morishita *et al.*, 1997; Lierz *et al.*, 2000, 2002). Mycoplasmas are common in birds of prey and certain species (*M. buteonis*, *M. gypis*, *M. falconis*) are regularly detected in addition to a number of unidentified species (Lierz *et al.*, 2008b). As free-ranging birds of prey are often sited close to poultry farms and captive raptors are regularly fed with poultry by-products, the occurrence of the poultry pathogen *Mycoplasma* species within healthy raptors could implicate them as a potential source of infection. Therefore, this avian group was investigated using specific PCRs for the poultry pathogenic *Mycoplasma* spp.

All PCRs used were first evaluated for their specificity using most of the known avian *Mycoplasma* species, plus some other mycoplasmas and walled bacteria commonly isolated from tracheal swabs of birds. The PCRs amplified only a specific product although, as expected, the PCR for *M. gallisepticum* amplified an additional product with the *M. imitans* reference strain. Differentiation was achieved by restriction enzyme analysis of the PCR product.

The sensitivities of the PCRs ranged between 0.1 and 10 CFU and appear to be adequate. However, it should be considered that the sensitivity was tested using DNA from pure cultures and might be lower when used on field samples.

Investigation of field samples that had previously tested positive using a generic *Mycoplasma* PCR (Lierz *et al.*, 2008a,b), yielded no PCR products for *M. gallisepticum*/*M. imitans*, *M. iowae* or *M. synoviae*. Inhibition of the PCRs seems an unlikely explanation in view of the earlier positive results with the same samples. For *M. iowae* tracheal swab material may not have provided the best sample since this organism is often detected in gastrointestinal samples. The *M. meleagridis* PCR amplified a product from eight field samples, but restriction enzyme analysis as well as sequencing three of those products demonstrated the results to be false positives.

This study has indicated that the pathogenic poultry *Mycoplasma* species are uncommon in birds of prey, compared to other *Mycoplasma* species such as *M. buteonis*, *M. gypis* and *M. falconis*. The demonstration of mycoplasmas in diseased raptors had regularly resulted in a diagnosis of 'mycoplasmosis' which is questionable considering the high prevalence of these organisms in healthy birds of prey (Lierz *et al.*, 2008b). However, as the pathogenic poultry mycoplasmas do not seem to occur regularly in healthy raptors, then their demonstration in diseases bird of prey might be of clinical value.

Although the PCRs were extensively evaluated using almost all known recognised avian *Mycoplasma* species and appeared to be specific, the *M. meleagridis* PCR gave false positive results. This underlines that positive results from PCRs must be carefully interpreted, especially if the tests are used in avian populations such as birds of prey, where there is regular detection of unidentified *Mycoplasma* species (Lierz *et al.*, 2000a, 2002, 2008a,b). The putative *Mycoplasma* species which caused the false positive results seems to be closely related to *M. buteonis* as it has a 98% identity with the 16S rRNA gene of *M. buteonis* reference strain. However, as the isolate did not give a positive result with *M. buteonis* antiserum in immunobinding assays, it may represent a new species. Interestingly, it seems to be common within the bird of prey population in Germany as it was detected in eight out of 53 (15%) of the birds investigated. These birds were not connected epidemiologically as they originated from captivity and the wild. Further

studies are necessary to describe these isolates in more detail. Furthermore the ability of this organism to transmit to poultry should be investigated since it could potentially cause diagnostic problems if the above described PCR were to be used in turkey breeders. Any positive results obtained using this PCR, including those from poultry, should be confirmed using restriction enzyme analysis or sequencing.

Acknowledgement

The authors would like to thank Prof. Dr. J. Bradbury, University of Liverpool, Dr. Mark Grant, British United Turkeys, UK, Prof. Dr. Wieler, Institute for Microbiology, Free University of Berlin and the International Organization of Mycoplasma for the provision of strains and antisera.

References

- Albers, A.C. & Fletcher, R.D. (1982). Simple method for quantitation of viable mycoplasmas. *Applied and Environmental Microbiology*, 43, 958-960.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215, 403-410.
- Amin, M.M. (1977). Avian Mycoplasmas: studies on isolation, infection and control. PhD Thesis. University of Liverpool.
- Benčina, D., Dorrer, D. & Tadina, T. (1987). Mycoplasma species isolated from six avian species. *Avian Pathology*, 16, 653-664.
- Benčina, D., Tadina, T., & Dorrer, D. (1988a). Natural infection of ducks with *Mycoplasma synoviae* and *Mycoplasma gallisepticum* and mycoplasma egg transmission. *Avian Pathology*, 17, 441-449.

- Benčina, D., Tadina, T. & Dorrer, D. (1988b). Natural infection of geese with *Mycoplasma gallisepticum* and *Mycoplasma synoviae* and egg transmission of the mycoplasmas. *Avian Pathology*, 17, 925-928.
- Benčina, D., Mrzel, I., Rojs, O. Z., Bidovec, A. & Dovč, A. (2003). Characterisation of *Mycoplasma gallisepticum* strains involved in respiratory disease in pheasants and peafowl. *Veterinary Record*, 152, 230-234.
- Benčina, D., Bradbury, J.M., Stipkovits, L., Varga, Z., Razpet, A., Bidovec, A. & Dovč, P. (2006). Isolation of *Mycoplasma capricolum*-like strains from chickens. *Veterinary Microbiology*, 112, 23-31.
- Blackwood, C.B., Marsh, T., Kim, S.-A., & Paul, E.A. (2003). Terminal restriction fragment length polymorphism data analysis for quantitative comparison of microbial communities. *Applied and Environmental Microbiology*, 69, 926- 932.
- Bozeman, L.H., Kleven, S.H. & Davis, R.B. (1984). Mycoplasma challenge studies in budgerigars (*Melopsittacus undulatus*) and chickens. *Avian Diseases*, 28, 426-434.
- Boyle, J.S., Good, R.T. & Morrow, C.J. (1995) Detection of the turkey pathogens *Mycoplasma meleagridis* and *M. iowae* by amplification of genes coding for rRNA. *Journal of Clinical Microbiology*, 33, 1335-1338.
- Bradbury, J.M.. (1998). Recovery of Mycoplasmas from birds. In R. Miles & R. Nicholas (Eds). *Mycoplasma Protocols* (pp. 45-51). New Jersey: Humana Press Inc.
- Bradbury, J.M. (2006). Mycoplasmas: a century and not out. *Turkeys*, 54, 9-11.
- Dhondt, A.A., Tessaglia, D.L. & Slothower R.L. (1998). Epidemic mycoplasmal conjunctivitis in house finches from eastern North America. *Journal of Wildlife Diseases*, 34, 265-280.
- Grau, O., Laigret, F., Carle, P., Tully, J.G., Rose, D.L. & Bové, J.M. (1991). Identification of a plant-derived mollicute as a strain of an avian pathogen, *Mycoplasma iowae*, and its

- implications for mollicute taxonomy. *International Journal of Systematic Bacteriology*, 41, 473- 478.
- Hartup, B.K., Mohammed, H.O., Kollias, G.V. & Dhondt, A.A. (1998). Risk factors associated with mycoplasmal conjunctivitis in house finches. *Journal of Wildlife Diseases*, 34, 281-288.
- Higgins, D.G., Thompson, J.D. & Gibson T.J. (1996). Using CLUSTAL for multiple sequence alignments. *Methods in Enzymology*, 266, 383-402.
- Ibrahim, R.S., Sohair, A. & Hussein, Z. (2000). Bacterial agents associated with sinusitis in waterfowls and turkeys in Assiut and El-Minia Governorates. *Assiut Veterinary Medicine Journal*, 44, 185-195.
- Kempf, I. (1997). DNA amplification methods for diagnosis and epidemiological investigations of avian mycoplasmosis. *Acta Veterinaria Hungarica*, 45, 373-386.
- Kotani, H. & Mc Garrity, G.J. (1985). Identification of mycoplasma colonies by immunobinding. *Journal of Clinical Microbiology*, 23, 783-785.
- Lierz, M., Schmidt, R., Brunnberg, L. & Runge, M. (2000a). Isolation of *Mycoplasma meleagridis* from free-ranging birds of prey in Germany. *Journal of Veterinary Medicine B*, 47, 63-67.
- Lierz, M., Schmidt, R., Goebel, T., Ehrlein, J. & Runge, M. (2000b). Detection of *Mycoplasma* spp. in raptorial birds in Germany. In J.T. Lumeij, J.D. Remple, P.T. Redig, M. Lierz & J.E. Cooper (Eds). *Raptor Biomedicine III* (pp. 25-33). Lake Worth: Zoological Education Network Inc.
- Lierz, M., Schmidt, R. & Runge, M. (2002). *Mycoplasma* species isolated from falcons in the Middle East. *Veterinary Record*, 151, 92-93.
- Lierz, M., Hagen, N., Lueschow, D. & Hafez, H.M. (2008a). Species-specific polymerase chain reactions for the detection of *Mycoplasma buteonis*, *Mycoplasma falconis*,

- Mycoplasma gypis* and *Mycoplasma corogypsi* in captive birds of prey. *Avian Diseases*, 52, 94-99.
- Lierz, M., Hagen, N., Hernandez, S.J. & Hafez, H.M. (2008b). Occurrence of mycoplasmas in free-ranging birds of prey in Germany. *Journal of Wildlife Diseases*, in press.
- Lin, M., Liu, S., Su, W., Lan, Y., Chung, I., Lin, M.Y., Liu, S.S., Su, W.S., Lan, Y.C. & Chung, I.C. (1995). Isolation of avian mycoplasmas from geese in Taiwan. *Journal of the Chinese Society of Veterinary Medicine*, 21, 347-353.
- Lüschow, D., Hoffmann, T. & Hafez, H.M. (2004). Differentiation of avian poxvirus strains on the basis of nucleotide sequences of 4b gene fragment. *Avian Diseases*, 48, 453-462.
- Marois, C., Dufour-Gesbert, F. & Kempf, I. (2000) Detection of *Mycoplasma synoviae* in poultry environment samples by culture and polymerase chain reaction. *Veterinary Microbiology*, 73, 311-318.
- Mikaelian, I., Ley, D.H., Claveau, R., Lemieux, M. & Berube J.P. (2001). Mycoplasmosis in evening and pine grosbeaks with conjunctivitis in Quebec. *Journal of Wildlife Diseases*, 37, 826-830.
- Morishita, T.Y., Aye, P.P. & Brooks, D.C. (1997). A survey of diseases of raptorial birds. *Journal of Avian Medicine and Surgery*, 11, 77-92.
- Poveda, J.B., Giebel, J., Kirchhoff, H. & Fernandez, A. (1990a). Isolation of mycoplasmas from buzzard, falcons and vultures. *Avian Pathology*, 19, 779-783.
- Poveda, J.B., Carranza, J., Miranda, A., Garrido, A., Hermoso, M., Fernandez, A. & Domenech, J. (1990b). An epizootiological study of avian mycoplasmas in southern Spain. *Avian Pathology*, 19, 627-633.
- Simecka, J.W., Davis, J.K., Davidson, M.K., Ross, S.E., Stadtländer, C.T.K.-H. & Cassell, G. H. (1992). Mycoplasma diseases of animals. In: J. Maniloff, R. N. McElhaney, L. R. Finch & J. B. Baseman (1992). *Mycoplasmas: Molecular Biology and Pathogenesis* (pp. 391-415). Washington D.C: American Society for Microbiology.

- Tanner, M.A., Everett, C.L., Coleman, W.J., Yang, M.M. & Youvan, D.C. (2000). Complex microbial communities inhabiting sulfide-rich black mud from marine coastal environments. *Biotechnology et alia*, 8, 1-16; www.et-al.com.
- Tiong, S.K. (1990). Mycoplasmas and acholeplasmas isolated from ducks and their possible association with pasteurellas. *Veterinary Record*, 127, 64-66.

For Peer Review Only

Table 1. Strain designations and sources of mycoplasmas, acholeplasmas and walled bacteria used to establish specificity and sensitivity of PCRs

<i>M. anatis</i> 1340 ^T (ATCC 25524) ^a	<i>M. sturni</i> UCMF ^T (ATCC 51945) ^a
<i>M. anseris</i> 1219 ^T (ATCC 49234) ^a	<i>M. synoviae</i> WVU 1853 ^T (ATCC 25204) ^a
<i>M. buteonis</i> Bb/T2g ^T (ATCC 51371) ^a	<i>M. gallisepticum</i> 3322 (FI) ^b
<i>M. corogypsi</i> BV1 ^T (ATCC 51148) ^a	<i>M. iowae</i> 2786 (FI) ^c
<i>M. cloacale</i> 383 ^T (ATCC 35276) ^a	<i>M. arginini</i> NCTC 10129 ^T (ATCC 23838) ^d
<i>M. columborale</i> MMP-4 ^T (ATCC 29258) ^a	<i>M. arthritidis</i> NCTC 10162 ^d
<i>M. columbinasale</i> 694 ^T (ATCC 33549) ^a	<i>M. canis</i> NCTC 10146 ^d
<i>M. columbinum</i> MMP-1 ^T (ATCC 29257) ^a	<i>M. equigenitalium</i> T 37 ^T (ATCC 29869) ^d
<i>M. falconis</i> H/T1 ^T (ATCC 51372) ^a	<i>M. felis</i> CO ^{Td}
<i>M. gallopavonis</i> 1197 ^a	<i>M. fermentans</i> NCTC 10117 ^T (ATCC 19989) ^d
<i>M. gallinarum</i> 1504 ^a	<i>M. hyorhinis</i> GDL (ATCC 23839) ^d
<i>M. gallisepticum</i> A5969 ^a	<i>M. salivarium</i> NCTC 10113 ^T (ATCC 23064) ^d
<i>M. gallinaceum</i> 887 ^a	<i>M. spumans</i> PG 13 ^T (ATCC 19526) ^d
<i>M. glycyphilum</i> 486 ^T (ATCC 35277) ^a	<i>A. laidlawii</i> NCTC 10116 ^T (ATCC 23206) ^d
<i>M. gypis</i> B1/T1 ^T (ATCC 51370) ^a	<i>Av. avium</i> 1158 ^d
<i>M. iowae</i> DK – CPA ^a	<i>Av. gallinarum</i> 1586 ^d
<i>M. imitans</i> 4229 ^T (ATCC 51306) ^a	<i>P. multocida</i> 338 ^d
<i>M. iners</i> PG30 ^T (ATCC 19705) ^a	<i>Ps. aeruginosa</i> 9380 ^d
<i>M. lipofaciens</i> R171 ^T (ATCC 35015) ^a	ORT 1280/91 (FI) ^b
<i>M. meleagridis</i> 17529 ^T (ATCC 25294) ^a	<i>S. enterica</i> serovar Typhimurium 778/05 (FI) ^b
<i>M. pullorum</i> 111R ^a	<i>E. coli</i> B6-05 (FI) ^b

A.= *Acholeplasma*; Av. = *Avibacterium*; E.= *Escherichia*; FI= Field isolate; M. = *Mycoplasma*; ORT= *Ornithobacterium rhinotracheale*; P. *Pasteurella*; Ps.= *Pseudomonas*; S.= *Salmonella*; ^T = Type strain; ^aprovided by International Organization for Mycoplasma, Department of Veterinary Pathobiology, Purdue University, IN, USA; ^bprovided by Institute of Poultry Diseases, Free University of Berlin, Germany; ^cprovided by Mark Grant, British United Turkey, UK; ^dprovided by Institute of Microbiology, Free University Berlin, Germany.

Table 2. Species, age and source of birds of prey sampled for PCR studies in Germany

Species	Total	Age		Source	
		Adult	Juvenile	Captivity	Wild
Peregrine (<i>Falco peregrinus</i>)	8	8	0	8	0
Kestrel (<i>Falco tinnunculus</i>)	5	3	2	0	5
Barbary Falcon (<i>Falco pelegrinoides</i>)	2	2	0	2	0
Saker (<i>Falco cherrug</i>)	2	2	0	2	0
Hobby (<i>Falco subbuteo</i>)	1	1	0	0	1
Gyr Falcon (<i>Falco rusticolus</i>)	1	1	0	1	0
Gyr Falcon-saker hybrid	2	2	0	2	0
Northern goshawk (<i>Accipiter gentilis</i>)	18	10	8	6	12
Common buzzard (<i>Buteo buteo</i>)	7	3	4	0	7
Red kite (<i>Milvus milvus</i>)	5	0	5	0	5
Marsh harrier (<i>Circus aeruginosus</i>)	1	1	0	0	1
Barn owl (<i>Tyto alba</i>)	1	1	0	0	1
Total	53	34	19	21	32

Table 3. Details of species-specific PCRs amplifying products within the mycoplasmal 16S rRNA gene

<i>Mycoplasma</i> species	Primer sequence (5'-3')	Polarity	Position ^a	Annealing	Size of amplicon (bp)	Sensitivity
<i>M. gallisepticum/M. imitans</i>	MGF CCA AGG CGA TGA CGT GTA GTT	Forward	884-904 / 239-259	30 sec	591	100 fg 0.1 CFU
	MGR CTG CAG CAC CGA AGT ATT CGC TC	Reverse	1451-1475 / 806-830	60°C		
	MIF AGT CGA ACG GGG TGC TTG CAC	Forward	62-82	30 sec		
	MIR TGC CTC TCA GCA CAG CTA TGC	Reverse	277-299	60°C		
<i>M. iowae</i>	MMF ^b CGA GCG AAG TTT TTC GGA AC	Forward	36-55	30 sec	237	100 fg 0.1 CFU
	MMR ^b GGT ACC GTC AGG ATA AAT GC	Reverse	438-458	56°C		
	MSF ^c GAG AAG CAA AAT AGT GAT ATC A	Forward	1192-1213	30 sec		
	MSR ^c CAG TCG TCT CCG AAG TTA ACA A	Reverse	1381-1403	56°C		
<i>M. meleagridis</i>	MMF ^b CGA GCG AAG TTT TTC GGA AC	Forward	36-55	30 sec	422	10 pg 10 CFU
	MMR ^b GGT ACC GTC AGG ATA AAT GC	Reverse	438-458	56°C		
	MSF ^c GAG AAG CAA AAT AGT GAT ATC A	Forward	1192-1213	30 sec		
	MSR ^c CAG TCG TCT CCG AAG TTA ACA A	Reverse	1381-1403	56°C		
<i>M. synoviae</i>	MMF ^b CGA GCG AAG TTT TTC GGA AC	Forward	36-55	30 sec	211	1 pg 1 CFU
	MMR ^b GGT ACC GTC AGG ATA AAT GC	Reverse	438-458	56°C		
	MSF ^c GAG AAG CAA AAT AGT GAT ATC A	Forward	1192-1213	30 sec		
	MSR ^c CAG TCG TCT CCG AAG TTA ACA A	Reverse	1381-1403	56°C		

^aPosition according to 16S rRNA gene provided by GenBank for the corresponding reference strains. Accession Nos: *M. gallisepticum* (L08897), *M. imitans* (L24103), *M. iowae* (M24293), *M. meleagridis* (L24106), *M. synoviae* (X52083).

^bAccording to Boyle *et al.* (1995); ^cAccording to Marois *et al.* (2000)

<i>Mycoplasma</i> species	Amplicon size (bp)	Enzyme	Fragment size (bp)	GenBank accession no.
<i>M. gallisepticum</i>	591	<i>Mse</i> I	14, 19, 33, 131, 188, 206	L08897
<i>M. imitans</i>	591	<i>Mse</i> I	14, 33, 206, 338	L24103
<i>M. gallisepticum</i>	591	<i>Ase</i> I	206, 385	L08897
<i>M. imitans</i>	591	<i>Ase</i> I	591	L24103
<i>M. iowae</i>	237	<i>Mse</i> I	104, 133	M24293
<i>M. meleagridis</i>	422	<i>Hind</i> III	145, 277	L24106
<i>M. synoviae</i>	211	<i>Hinf</i> I	86, 125	X52083

Figure 1. *Restriction enzyme analysis of M. gallisepticum and M. imitans amplicons. Lanes 1, & 5: 100 bp ladder, lane 2: M. gallisepticum product undigested, lanes 3 & 7: M. gallisepticum product after digestion with MseI (lane 3) and AseI (lane 7), lanes 4 & 6: M. imitans product after digestion with MseI (lane 4) and AseI (lane 6).*

Figure 2. *Restriction enzyme analysis (using HindIII) of amplicons obtained from the M. meleagridis PCR. Lane M: 100 bp ladder, lane 1: M. meleagridis type strain 17529 (ATTC 25294), lanes 2-9: amplicons from field samples.*

Fig. 1: Electrophoresis of the restriction enzyme analysis of *M. gallisepticum* and *M. imitans* amplicons. Lane 1,5: 100bp-ladder, line 2: *M. gallisepticum*- product undigested, lane 3,7: *M. gallisepticum* product after digestion with *MseI* (lane 3) and *AseI* (lane 7), line 4,6: *M. imitans* product after digestion with *MseI* (lane 4) and *AseI* (lane 6).

Fig. 2: Electrophoresis of the restriction enzyme analysis (using HindIII) of amplicons obtained from the *M. meleagridis* – PCR. Lane M: 100bp-ladder, lane 1: *M. meleagridis* type strain 17529 (ATCC 25294), lane 2-9: amplicons from field samples
61x23mm (600 x 600 DPI)