Progress in the production of medicinally important secondary metabolites in recombinant microorganisms or plants-Progress in alkaloid biosynthesis

Michael Wink, Holger Schäfer

► To cite this version:

Michael Wink, Holger Schäfer. Progress in the production of medicinally important secondary metabolites in recombinant microorganisms or plants-Progress in alkaloid biosynthesis. *Biotechnology Journal*, 2009, 4 (12), pp.1684. 10.1002/biot.200900229 . hal-00540529

HAL Id: hal-00540529

<https://hal.science/hal-00540529>

Submitted on 27 Nov 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Progress in the production of medicinally important
secondary metabolites in recombinant microorganisms or
plants-Progress in alkaloid biosynthesis**

Journal:	<i>Biotechnology Journal</i>
Manuscript ID:	biot.200900229.R1
Wiley - Manuscript type:	Review
Date Submitted by the Author:	28-Oct-2009
Complete List of Authors:	Wink, Michael; Heidelberg University, Institute for Pharmacy and Molecular Biotechnology Schäfer, Holger; Heidelberg University, Institute of Pharmacy and Molecular Biotechnology
Keywords:	Secondary metabolites, In-vitro production, Metabolic engineering, Alkaloid biosynthesis, Alkaloid production

Review

**Progress in the production of medicinally important secondary metabolites
in recombinant microorganisms or plants: Progress in alkaloid biosynthesis**

Holger Schäfer and Michael Wink

Institute of Pharmacy and Molecular Biotechnology, Heidelberg University, Heidelberg,
Germany

Correspondence: Professor Dr. Michael Wink, Institute of Pharmacy and Molecular
Biotechnology, Heidelberg University, Im Neuenheimer Feld 364, 69120 Heidelberg,
Germany

E-mail: wink@uni-hd.de

Fax: +49-6221-544-884

Keywords: Alkaloids/ Biosynthesis / Genes / *In-vitro* production / Terpenoids

Abbreviations: **BIA**, benzyloisoquinoline alkaloids; **MIA**, monoterpene indole alkaloid; **PA**,
pyrrolizidine alkaloid; **SM**, secondary metabolite

Plants produce a high diversity of natural products or secondary metabolites which are
important for the communication of plants with other organisms. A prominent function is the
protection against herbivores and/or microbial pathogens. Some natural products are also
involved in defence against abiotic stress, *e.g.* UV-B exposure. Many of the secondary
metabolites have interesting biological properties and quite a number are of medicinal

importance. Because the production of the valuable natural products, such as the anticancer drugs paclitaxel, vinblastine or camptothecin in plants is a costly process, biotechnological alternatives to produce these alkaloids more economically become more and more important. This review provides an overview of the state of art to produce alkaloids in recombinant microorganisms, such as bacteria or yeast. Some progress has been made in the metabolic engineering usually employing a single recombinant alkaloid gene. More importantly, for benzyloquinoline, monoterpene indole and diterpene alkaloids (taxanes) as well as some terpenoids and phenolics the proof of concept for the production of complex alkaloids in recombinant *Escherichia coli* and yeast has already been achieved. In a long-term perspective, it will probably be possible to generate gene cassettes for complete pathways, which could then be used for the production of valuable natural products in bioreactors or for metabolic engineering of crop plants to improve their resistance against herbivores and/or microbial pathogens.

Received: 10-Sep-2009

Revised: 28-Oct-2009

Accepted: 02-Nov-2009

1 Introduction

Plants produce a wide variety and high diversity of secondary metabolites (SM), which are not needed for primary or energy metabolism. They are not useless waste compounds, however, as previously assumed, but important for the ecological fitness of a plant producing them. Secondary metabolites have apparently evolved as a means for plants to protect themselves against insects, mammals and other herbivores, against bacteria, fungi, viruses and

even other competing plants. Some plants use SM in addition to attract pollinating and seed dispersing animals or for UV protection [1–6].

Plants usually synthesize, transport and store SM in a specific and particular way [3, 4]. Even a single plant produces a complex mixture of SM, which often derives from different types of SM; *e.g.* most plants produce phenolics, such as flavonoids but concomitantly terpenoids, such as saponins. The types of SM produced are sometimes but not always typical for a certain systematic group of plants [3, 5, 6]. Among more than 100 000 structures of SM that have been identified so far, we can distinguish between nitrogen-containing and nitrogen-free SM. Among nitrogen-containing SM, alkaloids are the largest group with more than 20 000 structures, many of them with pronounced pharmacological and toxic properties [7–9]. Also important are non-protein amino acids (700 structures), amines (100 structures), cyanogenic glucosides (60 structures), glucosinolates (100 structures), alkamides (150 structures), as well as lectins and other peptides (2000 structures). In the class of nitrogen-free SM, even more structures have been determined. The largest class is terpenoids with more than 20 000 known compounds, among them mono-, sesqui-, di-, and triterpenes with interesting bioactivities. Another bioactive group of SM, the polyphenols, is characterized by the presence of several phenolic hydroxyl groups, which can dissociate into O[−] ions under physiological conditions. Members of polyphenols are flavonoids, anthocyanins, and tannins. In addition, phenylpropanoids, coumarins, lignans and anthraquinones often possess phenolic OH-groups [4–8].

Structures of most SM are not random but the consequence of millions of years of selection during evolution. Therefore, it is not surprising that pharmacologists have discovered that a number of SM exhibit significant biological properties and can interact with molecular targets in human cells or microorganisms [4–9]. Consequently, many of the drugs used in medicine today derive directly from plants or indirectly in that structures of bioactive SM were used as a lead for the chemical synthesis with improved activities. SM from plants either are used as

isolated chemical entities or as complex extracts (typical for phytomedicine/phytotherapy) [7–9]. The latter approach has interesting aspects as the extracts apparently contain not only SM with additive but also synergistic properties [9]. Examples of isolated SM, which are being used in medicine include vinblastine, vincristine, paclitaxel (taxol), camptothecin, demecolcine, and podophyllotoxin (used in cancer therapy as probably the most important drugs), but also emetine, serpentine, ajmaline, reserpine, yohimbine, strychnine, ergobasine, ergotamine, quinine, quinidine, sparteine, ephedrine, lobeline, caffeine, berberine, sanguinarine, tubocurarine, papaverine, morphine, codeine, thebaine, noscapine, atropine, scopolamine, cardiac glycosides, artemisinin, anthraquinones and several others [7, 8]. As can be seen from this list, most of the interesting drugs are alkaloids; especially the anticancer drugs have a large market. So far, these drugs derive from plants, usually grown in plantations. Since their production in plants is usually low, the production costs are high and in consequence these drugs are costly.

Plant biotechnologists have explored possibilities to produce these valuable drugs using various *in vitro* systems, including bioreactors (Fig. 1). Callus cultures, suspension cell cultures, organ cultures (root and hairy root cultures) and even large-scale fermentation of suspended cells were successfully established over the last 40 years [10–21]. Thus, technically the *in-vitro* production should be feasible. However, the employment of callus and suspension cultures of medicinal plants often encountered the problem of very low or insufficient product yields. Apparently, the genes encoding the proteins necessary for biosynthesis, transport and storage of SM are not adequately expressed in most undifferentiated cell cultures [22]. There are a few notable exceptions with ginsenosides in *Panax ginseng*, shikonin in *Lithospermum erythrorhizon*, berberine in *Coptis japonica*, rosmarinic acid in *Coleus blumei*, anthraquinones in *Morinda citrifolia* or paclitaxel in *Taxus* sp. [21, 23]. More encouragingly, root and hairy root cultures, which are differentiated tissues, show excellent product yields for those SM that are produced in roots (which is unfortunately

not the case for all SM) [21, 23, 24]. However, the large-scale fermentation of roots and hairy roots still is a challenge, although bioreactors have been developed for a sustainable *in vitro* cultivation [22–24].

((Figure 1))

An alternative to the production of valuable plant drugs in plant cells was discussed already more than 30 years ago but even more so these days [25–39]. Once the genes are known that encode the enzymes of a biosynthetic pathway, it should be possible to functionally express these genes in a microbial system, such as bacteria or yeasts, and let them do the production (Fig. 1). As a first step, it could be shown in 1989 that *Escherichia coli* transformed with the plant gene encoding phenylalanine ammonium lyase (PAL) would convert phenylalanine into cinnamic acid, an important intermediate in the biosynthesis of flavonoids and some phenylpropanoids [39]. As a prerequisite for the production of complex natural products we need to isolate and characterize all the genes involved in the synthesis and storage of SM and then find a way to co-express them concomitantly in a microbial system. The search for the genes of plant secondary metabolism turned out to be very difficult and slow, because the genes of a biosynthetic pathway are usually not clustered as in bacteria, but apparently well dispersed over the plant genome. Secondly, mutants to select the genes were also not available. Therefore, each individual enzyme of a pathway had to be isolated beforehand, sequenced and then the genes could be isolated by employing corresponding primers for PCR or cDNA synthesis. Once such a gene became known it was usually much easier to find homologous genes in other plants. In the last 20 years, an impressive number of genes of secondary metabolism has been isolated (see next Sections). Several of the genes could be expressed in recombinant microorganisms and plants (reviewed in [25–39]). Most excitingly, researchers were successful to functionally co-express two or even more genes of a pathway. Thus, through metabolic engineering it was possible to produce a few selected benzyloquinoline alkaloids and key intermediates of artemisinin or taxane biosynthesis in

1
2
3 recombinant *E. coli* or *Saccharomyces cerevisiae* [28, 33–35]. Although these processes are
4
5 still incomplete and not yet commercial, they are important steps forward to the *in vitro*
6
7 production of SM.
8
9

10 This review will mainly focus on the progress made in finding the genes involved in alkaloid
11
12 biosynthesis and in expressing these genes in recombinant systems. Alkaloids were selected
13
14 as this group contains many candidates of medical importance and are therefore especially
15
16 interesting for biotechnology. The few success stories that have been reported so far on the *in-*
17
18 *vitro* production of alkaloids will be discussed in more detail as well as an outlook for the
19
20 future developments in this challenging field of biotechnology.
21
22

23
24 For the purpose of this review the weekly updated database Biosis Previews® (Thomson
25
26 Scientific, Inc.) was searched systematically for relevant publications using the names of
27
28 enzymes involved in alkaloid biosynthesis, of substrates or products as keywords. With the
29
30 goal of giving a comprehensive summary of any progress made in metabolic engineering,
31
32 literature was cited only, if a gene had been both cloned and functionally characterized, *e.g.* in
33
34 a heterologous system.
35
36
37
38
39
40

41 **2 Biosynthesis of alkaloids**

42
43
44
45

46 The enzymes that catalyze the biosynthesis of SM are usually substrate-specific, whereas
47
48 enzymes of breakdown or turnover (such as esterases, glycosidases) have a much broader
49
50 substrate spectrum. The nitrogen atom in alkaloids derives from amino acids in most cases
51
52 (except steroidal alkaloids) with phenylalanine, tyrosine, tryptophan, lysine, and ornithine
53
54 being a precursor most often involved. In a first reaction, the amino acids are decarboxylated
55
56 by decarboxylases with phenylethylamine, dopamine, tryptamine, cadaverine, and putrescine
57
58 as the corresponding amines. Further reactions often involve the reaction of the amino group
59
60 of the amine with an aldehyde function in the same or a second molecule. Aldehydes and

amines are known to spontaneously form a Schiff's base under physiological conditions. Other reactions include ring closure, oxidation or reductions of double bonds, the addition of functional groups (hydroxyl-, methyl, methylene dioxy groups) and the further modifications of OH-groups (esterification, glycosylation, methoxylation) (reviewed in [40–46]). A scheme of the pathways leading to major groups of alkaloids is illustrated in Fig. 2. ((Figure 2))

The search for the enzymes involved in the biosynthesis of alkaloids is an ongoing process and so far has been successful (at least partially) for a number of alkaloidal groups, such as morphinane-, protoberberine-, monoterpene indole-, Taxus-, ergot-, purine-, tropane-, Nicotiana-, pyrrolizidine and furanoquinoline alkaloids (see reviews [40–55]). However, considering the high structural diversity of alkaloids and other SM, much work still needs to be done until the picture is complete. Pathways have usually been explored in a few plants that produce a certain type of SM and it is silently assumed that pathways are identical in all plants that produce such compounds. This assumption still needs to be tested, but it is likely that different organisms evolved different solutions for the same task.

3 Genes of alkaloid biosynthesis

Once the enzymes had been isolated and purified, they could be sequenced and using the genetic code, primers for PCR and cDNA synthesis could be deduced. With some luck full-length cDNA clones could be generated encoding specific enzymes in a particular biosynthetic pathway. The corresponding genes could then be characterized and expressed in recombinant systems. This topic will be explored in more detail for major alkaloid groups and the relevant newer literature is summarized in the following. Earlier and recent work has been summarized in [40–55].

3.1 Benzyloquinoline alkaloids

Benzyloquinoline alkaloids form one of the major groups with several drugs of medical importance, such as morphine, thebaine, codeine, papaverine, berberine or sanguinarine. The biosynthesis of benzyloquinoline alkaloids, which include structural types such as tetrahydroisoquinoline, morphinan, protoberberine, benzophenanthridine and aporphine alkaloids (Fig. 3) has been reviewed recently [47, 48, 51, 55–57]. The pathway leading to tetrahydroisoquinoline, morphinane, and protoberberine alkaloids has been elucidated almost completely by now and most of the responsible genes have been cloned and characterized (Table 1). For a few enzymes X-ray data are available, such as berberine bridge enzyme (BBE) [58, 59] and norcoclaurine synthase [60].

((Figure 3)) ((Table 1))

3.2 Monoterpene indole alkaloids

Monoterpene indole alkaloids are especially abundant in Apocynaceae and contain several drugs of medicinal importance, such as the dimeric vinblastine and vincristine from *Catharanthus roseus* (important anticancer drugs), reserpine, ajmaline, ajmalicine, strychnine and yohimbine. Recent reviews of the biosynthesis of monoterpene indole alkaloids, which combine tryptamine and a monoterpene (secologanin) in their skeleton, have recently been published [47, 52, 55, 87, 88]. Strictosidine synthase (STR) is the key enzyme of this pathway and has been studied by many groups. It was the first alkaloid gene to be cloned in 1988 by T. Kutchan and M. Zenk [89]. The pathway leading to a few complex monoterpene indole-alkaloids (ajmalicine, serpentine, ajmaline, tabersonine, vindoline, catharanthine, and dimeric vinblastine) has been elucidated partially by now (Fig. 4) and several of the responsible genes

have been cloned and characterized (Table 2). For a few enzymes X-ray data are available, such as STR [90], strictosidine beta-D-glucosidase (SGD) [91, 92] and vinorine synthase [110, 122] from *Rauvolfia serpentina*.

((Figure 4)) ((Table 2))

3.3 Ergot alkaloids

Claviceps purpureus and other fungi produce indole alkaloids of the ergot alkaloid type with pronounced activities in neuronal signal transduction (such as found in LSD, ergotamine, ergometrine). Ergot alkaloids have also been detected in a few plants of the family Convolvulaceae. It could be shown recently, that the ergot alkaloid production in *Ipomoea* is due to an endophytic fungus that lives in symbiosis with its host plant [113]. The biosynthesis of the lysergic acid skeleton starts from tryptamine to which a unit of active isoprene is added. The corresponding 4-dimethylallyltryptophan synthase or 7-dimethylallyltryptophan synthase have been cloned, characterized (Table 3) and heterogeneously expressed in *E. coli* [114, 115].

((Table 3))

3.4 Purine alkaloids

Purine alkaloids, such as caffeine, theobromine and theophylline mediate the known stimulating bioactivities of coffee and tea (inhibition of phosphodiesterase, adenosine receptor antagonist) [4, 7]. The biosynthesis of purine alkaloids, such as caffeine, starts with xanthosine as a precursor. The pathway to xanthosine is assumed to follow the general pathway to purines, which is required for the purine bases adenine and guanine of DNA. The modifications of xanthosine are relatively simple and involve consecutive *N*-methylations

with SAM as a methyl donor [40, 41, 123–125]. The responsible genes have been cloned and characterized (Table 4).

((Table 4))

3.5 Paclitaxel (Taxol®)

Paclitaxel is a complex diterpene alkaloid, which has been introduced as a powerful anticancer drug (Taxol®) during the last 20 years [130]. Originally isolated from the bark of the Pacific yew tree (*Taxus brevifolia*), the production of this important drug was not sustainable at first. Later it was found that leaves from other *Taxus* species could be used to isolate taxanes in a more sustainable way: The taxanes can be converted into paclitaxel in a semisynthetic process [40, 41, 130]. Nevertheless, researchers have started programs for the chemical and biotechnological production of paclitaxel or its precursors. Several steps in the biosynthesis of taxanes have been elucidated [40, 41, 46] (Fig. 5) and some of the corresponding genes have been isolated and characterized (Table 5).

((Figure 5)) ((Table 5))

3.6 Tropane alkaloids and nicotine

Hyoscyamine and scopolamine are tropane alkaloids of medical importance since they are antagonist of the muscarinic acetylcholine receptor [4, 7, 40, 41]. Nicotine is a major alkaloid of tobacco; it is an agonist at the nicotinic acetylcholine receptor [4, 7, 40, 41] and had been used as a natural insecticide for many years [151]. The biosynthetic pathway leading to tropane alkaloids (including the polyhydroxyalkaloids of the calystegine type) and nicotine has been intensely studied and the initial steps have been elucidated (Fig. 6), but the corresponding synthase genes are still enigmatic. In nicotine biosynthesis two pathways can

lead to the intermediary putrescine: (i) via ornithine decarboxylation and (ii) from ornithine to arginine and after decarboxylation (ADC) via agmatine. Several genes have been isolated, expressed and characterized (Table 6).

((Figure 6)) ((Table 6))

3.7 Pyrrolizidine alkaloids

Pyrrolizidine alkaloids (PA), which occur in most taxa of the Boraginaceae and several Asteraceae, are of general toxicological importance, since most PA have mutagenic and even carcinogenic properties [40, 41]. Only one enzyme (homospermidine synthase) of PA biosynthesis and the corresponding gene have been isolated and characterized so far (Fig. 2 and Table 7) [177, 178]. When overexpressed in tobacco cells, an overproduction of homospermidine was observed [179].

((Table 7))

3.8 Acridone alkaloids

Acridone alkaloids are common in Rutaceae and use anthranilate as a key intermediate (Fig. 2). So far, two enzymes plus corresponding genes have been characterized (Table 8). The key enzyme acridone synthase has high sequence similarity to chalcone synthase (a key enzyme in the formation of flavonoids); only three amino acids differ between both enzymes [180].

((Table 8))

3.9 Betalains

Betalains are nitrogen-containing red-orange or yellow colored SM (and therefore alkaloids), which replace anthocyanins in some taxonomic groups (*e.g.* Caryophyllales) as flower pigments. They also occur in some fungi. A few steps of their biosynthesis have been elucidated in detail (Table 9).

((Table 9))

4 Metabolic engineering and alkaloid production in recombinant systems

The enzymes mentioned in Section 3 were characterized either as proteins purified from plant sources, or after the corresponding genes had been expressed in recombinant bacteria or yeasts in order to have enough material. Usually the focus of such work was not the functional expression of these genes in terms of metabolic engineering. A few experiments have been reported in which a single key enzyme was cloned in a heterologous system, usually another plant species, such as tobacco, and a change in the profile of primary and secondary metabolites was recorded. Sometimes the recombinant plants were fed with the appropriate precursor and it was analyzed if a biotransformation took place.

Ultimately, it will be necessary to transform plants or microbes with a series of genes or complete pathways and co-express them in order to produce a given compound in a recombinant system (Fig. 1). For the synthesis of antibiotics, such an approach has been shown to be feasible [190, 191]. However, here the situation is somehow easier, as the genes come in a cassette already and contain all the appropriate signal elements. In this Section, the progress in the line of metabolic engineering of natural products from plants will be discussed.

4.1 Benzyloquinoline alkaloids

In terms of metabolic engineering two alkaloid genes from *Coptis japonica* (6OMT and 4'OMT) were overexpressed in another alkaloid plant, *Eschscholzia californica*. Consequently, the alkaloid content was elevated 7.5 times [192]. When berberine bridge enzyme was knocked down by RNAi in *Eschscholzia californica*, an enhanced reticuline accumulation was observed [193]. In another approach salutaridinol 7-O-acetyltransferase in opium poppy was overexpressed or suppressed by RNAi [194].

When multiple genes in the BIA pathway were functionally co-expressed in yeast and *E. coli* the production of simple benzyliosquinoline alkaloids such as reticuline and related alkaloids [195], or reticuline, magnoflorine, and scoulerine was demonstrated [33]. These findings are especially encouraging because they are a proof of concept that the production of complex alkaloids is possible by using recombinant microorganisms. However, this approach is not optimal yet, since the production was not carried out in a single recombinant system with some steps done in *E. coli*, some *in vitro* and others through transformations in yeast.

4.2. Monoterpene indole alkaloids (MIA)

Metabolic engineering with genes of the MIA pathway in homologous and heterologous cell culture systems has been carried out with tryptophan decarboxylase and strictosidine synthase. The different experiments and results have been summarized in Table 10. The production of MIA in recombinant microorganisms is more complicated than that of BIA since the biosynthesis of the terpenoid precursor, secologanin is complex and the corresponding biosynthesis genes are not available.

((Table 10))

4.3 Purine alkaloids

When tobacco plants were transformed with the genes of caffeine biosynthesis, such as XMT, MXMT and DXMT a caffeine production was recorded; it has been argued that such a transformation could be interesting for crop plants, because caffeine apparently functions as a natural pest repellent [205]. The down-regulation of the genes of caffeine biosynthesis produced coffee plants without caffeine; this approach is interesting for the production of decaffeinated coffee [206].

4.4 Paclitaxel

Because of the importance of paclitaxel as an anticancer drug, research with regard of metabolic engineering of the taxane pathway was quite active during the last 10 years [29, 30]. The key enzyme taxadiene synthase, which converts geranyl-geranyl diphosphate into the diterpene skeleton, could be functionally expressed and taxadiene was detected in *Arabidopsis thaliana* [207], in transgenic tomato with carotenoid deficiency [208] and in the moss *Physcomitrella patens* [209].

A few groups were already successful in co-expressing several genes of paclitaxel biosynthesis in recombinant microbes: The expression of isopentenyl diphosphate isomerase, geranylgeranyl diphosphate synthase and taxadiene synthase in *E. coli* made it possible to synthesize taxadiene from isopentenyl diphosphate both in cell-free extracts and in recombinant bacteria [28, 31]. The co-expression of reductase and oxygenase enhances the formation of hydroxylated taxanes in recombinant yeast cells [210]. A significant step in the same direction was reported from the Croteau lab [28], which functionally expressed eight genes of taxane biosynthesis in yeast and could demonstrate a formation of taxadiene-5 α -acetoxy-10 β -ol from precursors of primary metabolism. However, a cytochrome p450 hydroxylation step seems to be a bottleneck in the present process [28].

5 Conclusions and outlook

Research towards the production of valuable alkaloids has seen much progress during the last two decades [38, 211, 212], but we are still not at a stage where these results can be used directly in the biotech industry. The proofs of concept obtained with isoquinoline alkaloids and taxanes are very encouraging but also show the problems that have to be overcome. It should be mentioned that similar approaches have been done with pathways leading to mono- and sesquiterpenes (several terpene synthases have been isolated and functionally expressed) [25, 26, 34, 35, 38, 211–213]. Research in the biosynthesis of flavonoids and related polyphenols has the longest tradition so far and many papers show the impact of metabolic engineering and practical applications [26, 27, 32, 37, 38, 211, 212].

In the long run, once the expression cassettes for different pathways have been established, these systems are likely to be used for the production of valuable drugs in fermented microorganisms, for the biotransformation of critical steps in a chemical synthesis or when transferred into crop plants for the enhancement of resistance against microbes and/or herbivores.

The authors have declared no conflict of interest.

6 References

- [1] Harborne, J. B., *Introduction to Ecological Biochemistry*, Academic Press, London 1993, 4th Edn.
- [2] Wink, M., Plant breeding: importance of plant secondary metabolites for protection against pathogens and herbivores. *Theor. Appl. Gen.* 1988, 75, 225–233.
- [3] Wink, M. (Ed.), *Biochemistry of Plant Secondary Metabolism*, Annual Plant Reviews Vol. 2, Sheffield Academic Press, Sheffield 1999.

- [4] Wink, M. (Ed.), *Function of Plant Secondary Metabolites and their Exploitation in Biotechnology*, Annual Plant Reviews Vol. 3, Sheffield Academic Press, Sheffield 1999.
- [5] Wink, M., Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry* 2003, 64, 3–19.
- [6] Wink, M., Plant secondary metabolism: Diversity, function and its evolution. *Natural Products Comm.* 2008, 3, 1205–1216.
- [7] van Wyk, B.-E., Wink, M., *Medicinal Plants of the World*. Briza, Pretoria 2004.
- [8] Wink, M., van Wyk, B.-E., *Mind-altering and Poisonous Plants of the World*. Briza, Pretoria 2008.
- [9] Wink, M., Evolutionary advantage and molecular modes of action of multi-component mixtures used in phytomedicine. *Current Drug Metab.* 2008, 9, 996–1009.
- [10] Bourgaud, F., Gravot, A., Milesi, S., Gontier, E., Production of plant secondary metabolites: a historical perspective. *Plant Sci.* 2001, 161, 839–851.
- [11] Canter, P. H., Thomas, H., Ernst, E., Bringing medicinal plants into cultivation: Opportunities and challenges for biotechnology. *Trends Biotechnol.* 2005, 23, 180–183.
- [12] Chattopadhyay, S., Farkya, S., Srivastava, A. K., Bisaria, V. S., Bioprocess considerations for production of secondary metabolite by plant cell suspension cultures. *Biotechnol. Bioprocess Eng.* 2002, 7, 138–149.
- [13] Kolewe, M. E., Gaurav, V., Roberts, S. C., Pharmaceutically active natural product synthesis and supply via plant cell culture technology. *Mol. Pharmaceutics* 2008, 5, 243–256.
- [14] Matkowski, A., Plant *in vitro* culture for the production of antioxidants – a review. *Biotechnol. Adv.* 2008, 26, 548–560.
- [15] McCoy, E., O'Connor, S. E. Natural products from plant cell cultures. *Progress Drug Res.* 2008, 65, 330–370.

- [16] Rao, R., Ravishankar, G. A., Plant cell cultures: Chemical factories of secondary metabolites. *Biotechnology Adv.* 2002, 20, 101–151.
- [17] Smetanska, I., Production of secondary metabolites using plant tissue cultures. *Food Biotechnol.* 2008, 11, 187–228.
- [18] Vanisree, M., Lee, C.-Y., Lo, S.-F., Nalawade, S. M. *et al.*, Studies on the production of some important secondary metabolites from medicinal plants by plant tissue culture. *Bot. Bull. Acad. Sin.* 2004, 45, 1–22.
- [19] Verpoorte, R., Contin, A., Memelink, J. Biotechnology for production of plant secondary metabolites. *Phytochem. Rev.* 2002, 1, 13–25.
- [20] Walton, N. J., Alfermann, A. W., Rhodes, M. J. C., Production of secondary metabolites in cell and differentiated organ cultures, in: Wink, M. (Ed.). *Functions of Plant Secondary Metabolites and their Exploitation in Biotechnology*, Sheffield Academic Press, Sheffield 1999, pp. 311–345.
- [21] Wink, M., Alfermann, A. W., Franke, R., Wetterauer, B. *et al.*, Sustainable bioproduction of phytochemicals by plant *in vitro* cultures: anticancer agents. *Plant Genetic Resources.* 2005, 3, 90–100.
- [22] Wink, M., Why do lupin cell cultures fail to produce alkaloids in large quantities? *Plant Cell, Tiss. Org. Cult.* 1987, 8, 103–111.
- [23] Alfermann, A. W., Production of natural products by cell and organ cultures, in: Wink, M. (Ed.), *Function and Biotechnology of Plant Secondary Metabolites*, Wiley-Blackwell, Oxford 2010 (in press).
- [24] Wildi, E., Wink, M., Biotechnology potential of hairy root culture, in: Govil, J. N., Ananda Kumar, P., Singh, V. K. (Eds.), *Biotechnology and Genetic Engineering*, Sci Tech Pub., USA 2002, Vol. 4, pp. 441–454.

- [25] Ajikumar, P. K., Tyo, K., Carlsen, P., Mucha, O. *et al.*, Terpenoids: Opportunities for biosynthesis of natural product drugs using engineered microorganisms. *Mol. Pharmaceut.* 2008, 5, 167–190.
- [26] Chemler, J. A., Koffas, M. A. G., Metabolic engineering for plant natural product biosynthesis in microbes. *Curr. Opinion Biotechnol.* 2008, 19, 597–605.
- [27] Leonard, E., Yan, Y., Fower, Z. L., Li, Z. *et al.*, Strain improvement of recombinant *Escherichia coli* for efficient production of plant flavonoids. *Mol. Pharm.* 2009, 5, 257–265.
- [28] DeJong, J. M., Liu, Y., Bollon, A. P., Long, R. M. *et al.*, Genetic engineering of taxol biosynthetic genes in *Saccharomyces cerevisiae*. *Biotechnol. Bioeng.* 2006, 93, 212–224.
- [29] Exposito, O., Bonfill, M., Moyano, E., Onrubis, M. *et al.*, Biotechnological production of Taxol and related taxoids: Current state and prospects. *Anti-Cancer Agents in Med. Chem.* 2009, 9, 109–121.
- [30] Frense, D., Taxanes: perspectives for biotechnological production. *Appl. Microbiol. Biotechnol.* 2007, 73, 1233–1240.
- [31] Huang, T.-K., Plesha, M. A., Falk, B. W., Dandekar, A. M., McDonald, K. A., Bioreactor strategies for improving production yield and functionality of a recombinant human protein in transgenic tobacco cell cultures. *Biotechnol. Bioengin.* 2009, 102, 508–520.
- [32] Leonard, E., Runguphan, W., O'Connor, S., Prather, K. J., Opportunities in metabolic engineering to facilitate scalable alkaloid production. *Nature Chem. Biol.* 2009, 5, 292–300.
- [33] Minami, H., Kim, J.-S., Ikezawa, N., Takemura, T. *et al.*, Microbial production of plant benzyloquinoline alkaloids. *Proc. Natl. Acad. Sci. USA* 2008, 105, 7393–7398.

- [34] Lindahl, A., Olsson, M. E., Mercke, P., Tollbom, O. *et al.*, Production of artemisinin precursor amorpha-4,11-diene by engineered *Saccharomyces cerevisiae*. Effect of *ERG9* repression on sesquiterpene biosynthesis. *Biotechnol. Lett.* 2006, 28, 571–580.
- [35] Ro, D-K., Paradise, E. M., Ouellet, M., Fisher, K. J. *et al.*, Production of antimalarial drug precursor artemisinic acid in engineered yeast. *Nature* 2006, 440, 940–943.
- [36] Marasco, E. K., Schmidt-Dannert, C., Biosynthesis of plant natural products and characterization of plant biosynthetic pathways in recombinant microorganisms, in: Verpoorte, R., Alfermann, A. W., Johnson, T. S. (Eds.), *Applications of Plant Metabolic Engineering*. Springer, Heidelberg 2007, pp. 1–43.
- [37] Oksman-Caldentey, K.-M., Häkkinen, S. T., Rischer, H., Metabolic engineering of the alkaloid biosynthesis in plants: Functional genomic approaches, in: Verpoorte, R., Alfermann, A. W., Johnson, T. S. (Eds.), *Applications of Plant Metabolic Engineering* Springer, Heidelberg 2007, pp. 109–143.
- [38] Wu, S., Chappell, J., Metabolic engineering of natural products in plants; tools of the trade and challenges for the future. *Curr. Opin. Biotechnol.* 2008, 19, 145–152.
- [39] Wink, M., Genes of secondary metabolism: differential expression in plants and *in vitro* cultures and functional expression in genetically-transformed microorganisms, in: Kurz, W. (Ed.), *Primary and Secondary Metabolism in Cell Cultures*, Springer, Heidelberg 1989, pp. 239–251.
- [40] Roberts, M. F., Wink, M., *Alkaloids:- Biochemistry, Ecological Functions and Medical Applications*. Plenum, New York 1998.
- [41] Dewick, P. M., *Medicinal Natural Products. A biosynthetic approach*. Wiley, New York 2002, 2nd edn.
- [42] Kutchan, T. M., Alkaloid biosynthesis: the basis for metabolic engineering of medicinal plants. *The Plant Cell* 1995, 7, 1959–1970.

- [43] Kutchan, T. M., Dittrich, H., Bracher, D., Zenk, M. H., Enzymology and molecular biology of alkaloid biosynthesis. *Tetrahedron* 1991, 47, 5945–54.
- [44] Saito, K., Murakoshi, I., Genes in alkaloid metabolism, in: Roberts, M. F., Wink, M., (Eds.), *Alkaloids: Biochemistry, Ecological Functions and Medical Applications*. Plenum, New York, 1998, pp. 147–57.
- [45] Facchini, P., Alkaloid biosynthesis in plants: biochemistry, cell biology, molecular regulation, and metabolic engineering applications. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 2001, 52, 29–66.
- [46] Roberts, M. F., Strack, D., Wink, M., Biosynthesis of alkaloids and betalains, in: Wink, M. (Ed.), *Biochemistry of plant secondary metabolism*, Annual Plant Reviews Wiley-Blackwell, Oxford 2010, Vol. 2, 2nd edn., (in press).
- [47] Facchini, P. J., De Luca, V., Opium poppy and Madagascar periwinkle: model non-model systems to investigate alkaloid biosynthesis in plants. *The Plant J.* 2008, 54, 763–784.
- [48] Zenk, M. H., Juenger, M., Evolution and current status of the phytochemistry of nitrogenous compounds, *Phytochemistry* 2007, 68, 2757–2772.
- [49] Facchini, P. J., Bird, D. A., St-Pierre, B., Can *Arabidopsis* make complex alkaloids? *Trends Plant Sci.* 2004, 9, 116–122.
- [50] Kutchan, T. M., Heterologous expression of alkaloid biosynthetic genes: a review. *Gene* 1996, 179, 73–81.
- [51] Sato, F., Inai, K., Hashimoto, T., Metabolic engineering in alkaloid biosynthesis: case studies in tyrosine and putrescine derived alkaloids, in: Verpoorte, R., Alfermann, A. W., Johnson, T. S. (Eds.), *Applications of Plant Metabolic Engineering*, Springer, Heidelberg 2007, pp. 145–173.

- [52] Stöckigt, J., Panjikar, S., Ruppert, M., Barleben, L. *et al.*, The molecular architecture of the major enzymes from ajmaline biosynthetic pathway. *Phytochem. Rev.* 2007, 6, 15–34.
- [53] Tydzynski, P., Correia, T., Keller, U., Biotechnology and genetics of ergot alkaloids. *Appl. Microbiol. Biotechnol.* 2001, 57, 593–605.
- [54] Zenk, M. H., Chasing the enzymes of biosynthesis, in: *Organic Reactivity: Physical and Biological Aspects*. The Royal Society of Chemistry, London 1995, pp. 89–109.
- [55] Ziegler, J., Facchini, P. J., Alkaloid biosynthesis: Metabolism and trafficking. *Annu. Rev. Plant Biol.* 2008, 59, 735–769.
- [56] Liscombe, D. K., Facchini, P. J., Molecular cloning and characterization of tetrahydroprotoberberine cis-*N*-methyltransferase, an enzyme involved in alkaloid biosynthesis in opium poppy. *J. Biol. Chem.* 2007, 282, 14741–14751.
- [57] Sato, F., Inui, T., Takemura, T., Metabolic engineering in isoquinoline alkaloid biosynthesis. *Current Pharm. Biotech.* 2007, 8, 211–218.
- [58] Winkler, A., Lyskowski, A., Riedl, S., Puhl, M., Kutchan, T. M., Macheroux, P., Gruber, K., A concerted mechanism for berberine bridge enzyme. *Nature Chem. Biol.* 2008, 4, 739–741.
- [59] Winkler, A., Motz, K., Riedl, S., Puhl, M. *et al.*, Structural and mechanistic studies reveal the functional role of bicovalent flavinylation in berberine bridge enzyme. *J. Biol. Chem.* 2009, 284, 19993–20001.
- [60] Ilari, A., Franceschini, S., Bonamore, A., Arengi, F. *et al.*, Structural basis of enzymatic (S)-norcoclaurine biosynthesis. *J. Biol. Chem.* 2009, 284, 897–904.
- [61] Facchini, P. J., De Luca, V., Differential and tissue-specific expression of a gene family for tyrosine/dopa decarboxylase in opium poppy. *J. Biol. Chem.* 1994, 269, 26684–26690.

- [62] Kang, S., Kang, K., Lee, K., Back, K., Characterization of rice tryptophan decarboxylases and their direct involvement in serotonin biosynthesis in transgenic rice. *Planta* 2007, 227, 263–272.
- [63] Samanani, N., Park, S.-U., Facchini, P. J., Cell type-specific localization of transcripts encoding nine consecutive enzymes involved in protoberberine alkaloid biosynthesis. *Plant Cell* 2005, 17, 915–926.
- [64] Lehmann, T., Pollmann, S., Gene expression and characterization of a stress-induced tyrosine decarboxylase from *Arabidopsis thaliana*. *FEBS Lett.* 2009, 583, 1895–1900.
- [65] Samanani, N., Liscombe, D. K., Facchini, P. J., Molecular cloning and characterization of norcoclaurine synthase, an enzyme catalyzing the first committed step in benzyloquinoline alkaloid biosynthesis. *Plant J.* 2004, 40, 302–313.
- [66] Minami, H., Dubouzet, E., Iwasa, K., Sato, F., Functional analysis of norcoclaurine synthase in *Coptis japonica*. *J. Biol. Chem.* 2007, 282, 6274–6282.
- [67] Frick, S., Kutchan, T. M., Molecular cloning and functional expression of O-methyltransferases common to isoquinoline alkaloid and phenylpropanoid biosynthesis. *Plant J.* 1999, 17, 329–339.
- [68] Morishige, T., Tsujita, T., Yamada, Y., Sato, F., Molecular characterization of the S-adenosyl-L-methionine:3'-hydroxy-N-methylcoclaurine 4'-O-methyltransferase involved in isoquinoline alkaloid biosynthesis in *Coptis japonica*. *J. Biol. Chem.* 2000, 275, 23398–23405.
- [69] Facchini, P. J., Park, S.-U., Developmental and inducible accumulation of gene transcripts involved in alkaloid biosynthesis in opium poppy. *Phytochemistry* 2003, 64, 177–186.
- [70] Ounaroon, A., Decker, G., Schmidt, J., Lottspeich, F., Kutchan, T. M., (R,S)-Reticuline 7-O-methyltransferase and (R,S)-norcoclaurine 6-O-methyltransferase of

- Papaver somniferum: cDNA cloning and characterization of methyl transfer enzymes of alkaloid biosynthesis in opium poppy. *Plant J.* 2003, 36, 808–819.
- [71] Choi, K.-B., Morishige, T., Shitan, N., Yazaki, K., Sato, F., Molecular cloning and characterization of coclaurine N-methyltransferase from cultured cells of *Coptis japonica*. *J. Biol. Chem.* 2002, 277, 830–835.
- [72] Kraus, P. F. X., Kutchan, T. M., Molecular cloning and heterologous expression of a cDNA encoding berbaminine synthase, a C-O phenol-coupling cytochrome P450 from the higher plant *Berberis stolonifera*. *Proc. Natl. Acad. Sci. USA* 1995, 92, 2071–2075.
- [73] Pauli, H. H., Kutchan, T. M., Molecular cloning and functional heterologous expression of two alleles encoding (S)-N-methylcoclaurine 3'-hydroxylase (CYP80B1), a new methyl jasmonate-inducible cytochrome P-450-dependent mono-oxygenase of benzyloquinoline alkaloid biosynthesis. *Plant J.* 1998, 13, 793–801.
- [74] Huang, F.-C., Kutchan, T. M., Distribution of morphinan and benzo[c]phenanthridine alkaloid gene transcript accumulation in *Papaver somniferum*. *Phytochemistry* 2000, 53, 555–564.
- [75] Ikezawa, N., Tanaka, M., Nagayoshi, M., Shinkyo, R. *et al.*, Molecular cloning and characterization of CYP719, a methylenedioxy bridge-forming enzyme that belongs to a novel P450 family, from cultured *Coptis japonica* cells. *J. Biol. Chem.* 2003, 278, 38557–38565.
- [76] Dittrich, H., Kutchan, T. M., Molecular cloning expression and induction of berberine bridge enzyme and enzymes essential to the formation of benzophenanthridine alkaloids in the response of plants to pathogenic attack. *Proc. Natl. Acad. Sci. USA* 1991, 88, 9969–9973.
- [77] Facchini, P. J., Penzes, C., Johnson, A. G., Bull, D., Molecular characterization of berberine bridge enzyme genes from opium poppy. *Plant Physiol.* 1996, 112, 1669–1677.

- [78] Ikezawa, N., Iwasa, K., Sato, F., CYP719A subfamily of cytochrome P450 oxygenases and isoquinoline alkaloid biosynthesis in *Eschscholzia californica*. *Plant Cell Rep.*, 2009, 28, 123–133.
- [79] Ikezawa, N., Iwasa, K., Sato, F., Molecular cloning and characterization of methylenedioxy bridge-forming enzymes involved in stylophine biosynthesis in *Eschscholzia californica*. *FEBS J.* 2007, 274, 1019–1035.
- [80] Ikezawa, N., Tanaka, M., Nagayoshi, M., Shinkyo, R. *et al.*, Molecular cloning and characterization of CYP719, a methylenedioxy bridge-forming enzyme that belongs to a novel P450 family, from cultured *Coptis japonica* cells. *J. Biol. Chem.* 2003, 278, 38557–38565.
- [81] Takeshita, N., Fujiwara, H., Mimura, H., Fitchen, J. H. *et al.*, Molecular cloning and characterization of *S*-adenosyl-*L*-methionine:Scoulerine-9-*O*-methyltransferase from cultured cells of *Coptis japonica*. *Plant & Cell Physiol.* 1995, 36, 29–36.
- [82] Morishige, T., Dubouzet, E., Choi, Yazaki, K., Sato, F., Molecular cloning of columbamine *O*-methyltransferase from cultured *Coptis japonica* cells. *Eur. J. Biochem.* 2002, 269, 5659–5667.
- [83] Ziegler, J., Voigtlaender, S., Schmidt, J., Kramell, R. *et al.*, Comparative transcript and alkaloid profiling in *Papaver* species identifies a short chain dehydrogenase/reductase involved in morphine biosynthesis. *Plant J.* 2006, 48, 177–192.
- [84] Grothe, T., Lenz, R., Kutchan, T. M., Molecular characterization of the salutaridinol 7-*O*-acetyltransferase involved in morphine biosynthesis in opium poppy *Papaver somniferum*. *J. Biol. Chem.* 2001, 276, 30717–30723.
- [85] Unterlinner, B., Lenz, R., Kutchan, T. M., Molecular cloning and functional expression of codeinone reductase: The penultimate enzyme in morphine biosynthesis in the opium poppy *Papaver somniferum*. *Plant J.* 1999, 18, 465–475.

- [86] Ikezawa, N., Iwasa, K., Sato, F., Molecular cloning and characterization of CYP80G2, a cytochrome p450 that catalyzes an intramolecular C - C phenol coupling of (S)-reticuline in magnoflorine biosynthesis, from cultured *Coptis japonica* cells. *J. Biol. Chem.* 2008, 283, 8810–8821.
- [87] Ruppert, M., , Ma, X., Stöckigt, J., Alkaloid biosynthesis in *Rauvolfia*- cDNA Cloning of major enzymes of the ajmaline pathway. *Curr. Org. Chem.* 2005, 9, 1431–1444.
- [88] Ganjewala, D., Kumar, S., Luthra, R., An account of cloned genes of methyl-erythritol-4-phosphate pathway of isoprenoid biosynthesis in plants. *Curr. Iss. Mol. Biol.* 2009, 1, 135–145.
- [89] Kutchan, T. M., Hampp, N., Lottspeich, F., Beyreuther, F., Zenk, M. H., The complementary DNA clone for strictosidine synthase from *Rauvolfia serpentina* DNA sequence determination and expression in *Escherichia coli*. *Febs Lett.* 1988, 237, 40–44.
- [90] Koepke, J., Ma, X., Frittsch, G., Michel, H., Stoeckigt, J., Crystallization and preliminary X-ray analysis of strictosidine synthase and its complex with the substrate tryptamine. *Acta Crystallographica Sect. D Biol. Crystall.* 2005, 61, 690–693.
- [91] Barleben, L., Ma, X., Koepke, J., Peng, G. *et al.*, Expression, purification, crystallization and preliminary X-ray analysis of strictosidine glucosidase, an enzyme initiating biosynthetic pathways to a unique diversity of indole alkaloid skeletons. *Biochim. Biophys. Acta* 2005, 1, 89–92.
- [92] Barleben, L., Panjikar, S., Ruppert, M., Koepke, J., Stoeckigt, J., Molecular architecture of strictosidine glucosidase: The gateway to the biosynthesis of the monoterpenoid indole alkaloid family. *Plant Cell* 2007, 19, 2886–2897.
- [93] De Luca, V., Marineau, C., Brisson, N., Molecular cloning and analysis of complementary DNA encoding a plant tryptophan decarboxylase. Comparison with animal Dopa decarboxylases. *Proc. Natl. Acad. Sci. USA* 1989, 86, 2582–2586.

- [94] Lopez-Meyer, M., Nessler, C. L., Tryptophan decarboxylase is encoded by two autonomously regulated genes in *Camptotheca acuminata* which are differentially expressed during development and stress. *Plant J.* 1997, 11, 1167–1175.
- [95] Yamazaki, Y., Sudo, H., Yamazaki, M., Aimi, N., Saito, K., Camptothecin biosynthetic genes in hairy roots of *Ophiorrhiza pumila*: Cloning, characterization and differential expression in tissues and by stress compounds. *Plant & Cell Physiol.* 2003, 44, 395–403.
- [96] Collu, G., Unver, N., Peltenburg-Looman, A. M. G., van der Heijden, R. *et al.*, Geraniol 10-hydroxylase, a cytochrome P450 enzyme involved in terpenoid indole alkaloid biosynthesis. *FEBS Lett.* 2001, 508, 215–220.
- [97] Irmeler, S., Schroeder, G., St-Pierre, B., Crouch, N. P. *et al.*, Indole alkaloid biosynthesis in *Catharanthus roseus*: New enzyme activities and identification of cytochrome P450 CYP72A1 as secologanin synthase. *Plant J.* 2000, 24, 797–804.
- [98] Murata, J., Roepke, J., Gordon, H., De Luca, V., The leaf epidermome of *Catharanthus roseus* reveals its biochemical specialization. *Plant Cell* 2008, 20, 524–542.
- [99] McKnight, T. D., Roessner, C. A., Devagupta, R., Scott, A. I., Nessler, C. L., Nucleotide sequence of a complementary DNA encoding the vacuolar protein strictosidine synthase from *Catharanthus roseus*. *Nucl. Acids Res.* 1990, 18, 4939.
- [100] Bracher, D., Kutchan, T. M., Strictosidine synthase from *Rauwolfia serpentina* Analysis of a gene involved in indole alkaloid biosynthesis. *Arch. Biochem. Biophys.* 1992, 294, 717–723.
- [101] Chen, R., Liao, Z. H., Chen, M., Wang, Q. *et al.*, Molecular cloning and characterization of the strictosidine synthase gene from *Rauwolfia verticillata*. *Russ. J. Plant Physiol.* 2008, 55, 670–675.
- [102] Geerlings, A., Ibanez, M. M.-L., Memelink, J., van der Heijden, R., Verpoorte, R., Molecular cloning and analysis of strictosidine beta-D-glucosidase, an enzyme in

- terpenoid indole alkaloid biosynthesis in *Catharanthus roseus*. *J. Biol. Chem.* 2000, 275, 3051–3056.
- [103] Gerasimenko, I., Sheludko, Y., Ma, X., Stoeckigt, J., Heterologous expression of a *Rauvolfia* cDNA encoding strictosidine glucosidase, a biosynthetic key to over 2000 monoterpenoid indole alkaloids. *Eur. J. Biochem.* 2002, 269, 2204–2213.
- [104] Schroeder, G., Unterbusch, E., Kaltenbach, M., Schmidt, J. *et al.*, Light-induced cytochrome P450-dependent enzyme in indole alkaloid biosynthesis: Tabersonine 16-hydroxylase. *FEBS Lett.* 1999, 458, 97–102.
- [105] Dylan, L., Jun, M., Won, S. K., De, L. V., Application of carborundum abrasion for investigating the leaf epidermis: molecular cloning of *Catharanthus roseus* 16-hydroxytabersonine-16-*O*-methyltransferase. *Plant J.* 2008, 53, 225–236.
- [106] Vazquez-Flota, F., De Carolis, E., Alarco, A.-M., De Luca, V., Molecular cloning and characterization of desacetoxylvindoline-4-hydroxylase, a 2-oxoglutarate dependent-dioxygenase involved in the biosynthesis of vindoline in *Catharanthus roseus* (L.) G. Don. *Plant Mol. Biol.* 1997, 34, 935–948.
- [107] St-Pierre, B., Laflamme, P., Alarco, A.-M., De Luca, V., A terminal *O*-acetyltransferase involved in vindoline biosynthesis defines a new class of proteins responsible for coenzyme A-dependent acyl transfer. *Plant J.* 1998, 14, 703–713.
- [108] Laflamme, P., St.-Pierre, B., De Luca, V., Molecular and biochemical analysis of a Madagascar periwinkle root-specific minovincinine-19-hydroxy-*O*-acetyltransferase. *Plant Physiol.* 2001, 125, 189–198.
- [109] Dogru, E., Warzecha, H., Seibel, F., Haebel, S. *et al.*, The gene encoding polynuridine aldehyde esterase of monoterpenoid indole alkaloid biosynthesis in plants is an ortholog of the alpha/beta hydrolase super family. *Eur. J. Biochem.* 2000, 267, 1397–1406.

- [110] Bayer, A., Ma, X., Stoeckigt, J., Acetyltransfer in natural product biosynthesis - functional cloning and molecular analysis of vinorine synthase. *Bioorg. Medic. Chem.* 2004, 12, 2787–2795.
- [111] Warzecha, H., Gerasimenko, I., Kutchan, T. M., Stoeckigt, J., Molecular cloning and functional bacterial expression of a plant glucosidase specifically involved in alkaloid biosynthesis. *Phytochemistry* 2000, 54, 657–666.
- [112] Sun, L., Ruppert, M., Sheludko, Y., Warzecha, H. *et al.*, Purification, cloning, functional expression and characterization of perakine reductase: the first example from the AKR enzyme family, extending the alkaloidal network of the plant *Rauvolfia*. *Plant Mol. Biol.* 2008, 67, 455–467.
- [113] Markert, A., Steffan, N., Ploss, K., Hellwig, S. *et al.*, Biosynthesis and accumulation of ergoline alkaloids in a mutualistic association between *Ipomoea asarifolia* (Convolvulaceae) and a Clavicipitalean fungus. *Plant. Physiol.* 2008, 147, 296–305.
- [114] Steffan, N., Unsoeld, I. A., Li, S.-M., Chemoenzymatic synthesis of prenylated indole derivatives by using a 4-dimethylallyltryptophan synthase from *Aspergillus fumigatus*. *Chem. BioChem.* 2007, 8, 1298–1307.
- [115] Kremer, A., Li, S.-M., Potential of a 7-dimethylallyltryptophan synthase as a tool for production of prenylated indole derivatives. *Appl. Microbiol. Biotech.* 2008, 79, 951–961.
- [116] Tsai, H.-F., Wang, H., Gebler, J. C., Poulter, C. D., Schardl, C. L., The *Claviceps purpurea* gene encoding dimethylallyltryptophan synthase, the committed step for ergot alkaloid biosynthesis. *Biochem. Biophys. Res. Comm.* 1995, 216, 119–125.
- [117] Wang, J., Machado, C., Panaccione, D. G., Tsai, H.-F., Schardl, C. L., The determinant step in ergot alkaloid biosynthesis by an endophyte of perennial ryegrass. *Fungal Genetics Biol.* 2004, 41, 189–198.

- [118] Unsoeld, I. A., Li, S.-M., Overproduction, purification and characterization of FgaPT2, a dimethylallyltryptophan synthase from *Aspergillus fumigatus*. *Microbiology* 2005, 151, 1499–1505.
- [119] Ding, Y., Williams, R. M., Sherman, D. H., Molecular analysis of a 4-dimethylallyltryptophan synthase from *Malbranchea aurantiaca*. *J. Biol. Chem.* 2008, 283, 16068–16076.
- [120] Kremer, A., Westrich, L., Li, S.-M., A 7-dimethylallyltryptophan synthase from *Aspergillus fumigatus*: overproduction, purification and biochemical characterization. *Microbiology* 2007, 153, 3409–3416.
- [121] Rigbers, O., Li, S.-M., Ergot alkaloid biosynthesis in *Aspergillus fumigatus* - Overproduction and biochemical characterization of a 4-dimethylallyltryptophan N-methyltransferase. *J. Biol. Chem.* 2008, 283, 26859–26868.
- [122] Ma, X., Koepke, J., Bayer, A., Fritzsche, G. *et al.*, Crystallization and preliminary X-ray analysis of native and selenomethionyl vinorine synthase from *Rauvolfia serpentina*. *Acta Cryst.* 2005. D61, 694–696.
- [123] Ashihara, H., Sano, H., Crozier, A., Caffeine and related purine alkaloids: Biosynthesis, catabolism, function and genetic engineering. *Phytochemistry* 2008, 69, 841–856.
- [124] Kato, M., Mizuno, K., Crozier, A., Fujimura, T., Ashihara, H., Plant biotechnology: Caffeine synthase gene from tea leaves. *Nature* 2000, 406, 956–957.
- [125] McCarthy, A. A., McCarthy, J. G., The structure of two N-methyltransferases from the caffeine biosynthetic pathway. *Plant Physiol.* 2007, 144, 879–889.
- [126] Mizuno, K., Kato, M., Irino, F., Yoneyama, N. *et al.*, The first committed step reaction of caffeine biosynthesis: 7-methylxanthosine synthase is closely homologous to caffeine synthases in coffee (*Coffea arabica* L.). *FEBS Lett.* 2003, 547, 56–60.

- [127] Uefuji, H., Ogita, S., Yamaguchi, Y., Koizumi, N., Sano, H., Molecular cloning and functional characterization of three distinct *N*-methyltransferases involved in the caffeine biosynthetic pathway in coffee plants. *Plant Physiol.* 2003, 132, 372–380.
- [128] Mizuno, K., Okuda, A., Kato, M., Yoneyama, N. *et al.*, Isolation of a new dual-functional caffeine synthase gene encoding an enzyme for the conversion of 7-methylxanthine to caffeine from coffee (*Coffea arabica* L.). *FEBS Lett.* 2003, 534, 75–81.
- [129] Ogawa, M., Herai, Y., Koizumi, N., Kusano, T., Sano, H., 7-Methylxanthine methyltransferase of coffee plants: Gene isolation and enzymatic properties. *J. Biol. Chem.* 2001, 276, 8213–8218.
- [130] Fu, Y., Li, S., Zu, Y., Yang, G. *et al.*, Medicinal chemistry of paclitaxel and its analogues. *Curr. Medicinal Chem.* In press.
- [131] Wildung, M. R., Croteau, R., A cDNA clone for taxadiene synthase, the diterpene cyclase that catalyzes the committed step of taxol biosynthesis. *J. Biol. Chem.* 1996, 271, 9201–9204.
- [132] Wang, W., Shi, Q., Zhu, P., Ouyang, T. *et al.*, cDNA cloning, expression and characterization of taxadiene synthase, a diterpene cyclase from *Taxus chinensis*. *Acta Botanica Sinica* 2002, 44, 181–187.
- [133] Kai, G., Zhao, L., Zhang, L., Li, Z. *et al.*, Characterization and expression profile analysis of a new cDNA encoding taxadiene synthase from *Taxus media*. *J. Biochem. Mol. Biol.* 2005, 38, 668–675.
- [134] Hefner, J., Ketchum, R. E. B., Croteau, R., Cloning and functional expression of a cDNA encoding geranylgeranyl diphosphate synthase from *Taxus canadensis* and assessment of the role of this prenyltransferase in cells induced for taxol production. *Arch. Biochem. Biophys.* 1998, 360, 62–74.

- [135] Walker, K., Croteau, R., Taxol biosynthesis: Molecular cloning of a benzoyl-CoA:taxane 2 α -O-benzoyltransferase cDNA from *Taxus* and functional expression in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 2000, 97, 13591–13596.
- [136] Walker, K., Croteau, R., Molecular cloning of a 10-deacetylbaccatin III-10-O-acetyl transferase cDNA from *Taxus* and functional expression in *Escherichia coli*. *Proc. Nat. Acad. Sci. USA* 2000, 9, 13591–13596.
- [137] Guo, B., Kai, G., Gong, Y., Jin, H. *et al.*, Molecular cloning and heterologous expression of a 10-deacetylbaccatin III-10-O-acetyl transferase cDNA from *Taxus x media*. *Mol. Biol. Rep.* 2007, 34, 89–95.
- [138] Walker, K., Schoendorf, A., Croteau, R., Molecular cloning of a taxa-4(20),11(12)-dien-5 α -ol-O-acetyl transferase cDNA from *Taxus* and functional expression in *Escherichia coli*. *Arch. Biochem. Biophys.* 2000, 374, 371–380.
- [139] Jennewein, S., Rithner, C. D., Williams, R. M., Croteau, R. B., Taxol biosynthesis: Taxane 13 α -hydroxylase is a cytochrome P450-dependent monooxygenase. *Proc. Natl. Acad. Sci. USA* 2001, 98, 13595–13600.
- [140] Schoendorf, A., Rithner, C. D., Williams, R. M., Croteau, R. B., Molecular cloning of a cytochrome P450 taxane 10 β -hydroxylase cDNA from *Taxus* and functional expression in yeast. *Proc. Natl. Acad. Sci. USA* 2001, 98, 1501–1506.
- [141] Kai, G., Jiang, J., Zhao, D., Zhao, L. *et al.*, Isolation and expression profile analysis of a new cDNA encoding 5- α -taxadienol-10- β -hydroxylase from *Taxus media*. *J. Plant Biochem. Biotech.* 2006, 15, 1–5.
- [142] Jennewein, S., Rithner, C. D., Williams, R. M., Croteau, R., Taxoid metabolism: Taxoid 14 β -hydroxylase is a cytochrome P450-dependent monooxygenase. *Arch. Biochem. Biophys.* 2003, 413, 262–270.

- [143] Chau, M., Croteau, R., Molecular cloning and characterization of a cytochrome P450 taxoid 2 α -hydroxylase involved in *Taxol* biosynthesis. *Arch. Biochem. Biophys.* 2004, 427, 48–57.
- [144] Chau, M., Jennewein, S., Walker, K., Croteau, R., Taxol biosynthesis: Molecular cloning and of a cytochrome p450 characterization taxoid 7 β -hydroxylase. *Chemistry & Biology* 2004, 11, 663–672.
- [145] Jennewein, S., Long, R. M., Williams, R. M., Croteau, R., Cytochrome P450 taxadiene 5 α -hydroxylase, a mechanistically unusual monooxygenase catalyzing the first oxygenation step of taxol biosynthesis. *Chemistry & Biology* 2004, 11, 379–387.
- [146] Walker, K., Fujisaki, S. ., Long, R., Croteau, R., Molecular cloning and heterologous expression of the C-13 phenylpropanoid side chain-CoA acyltransferase that functions in Taxol biosynthesis. *Proc. Natl. Acad. Sci. USA* 2002, 99, 12715–12720.
- [147] Walker, K., Long, R., Croteau, R., The final acylation step in Taxol biosynthesis: Cloning of the taxoid C13-side-chain *N*-benzoyltransferase from *Taxus*. *Proc. Nat. Acad. Sci. USA*, 2002, 99, 9166–9171.
- [148] Walker, K. D., Klettke, K., Akiyama, T., Croteau, R., Cloning, heterologous expression, and characterization of a phenylalanine aminomutase involved in Taxol biosynthesis. *J. Biol. Chem.* 2004, 279, 53947–53954.
- [149] Steele, C. L., Chen, Y., Dougherty, B. A., Li, W. *et al.*, Purification, cloning, and functional expression of phenylalanine aminomutase: The first committed step in Taxol side-chain biosynthesis. *Arch. Biochem. Biophys.* 2005, 438, 1–10.
- [150] Chau, M., Walker, K., Long, R., Croteau, R., Regioselectivity of taxoid-*O*-acetyltransferases: heterologous expression and characterization of a new taxadien-5 α -ol-*O*-acetyltransferase. *Arch. Biochem. Biophys.* 2004, 430, 237–246.

- [151] Wink, M., Production and application of phytochemicals from an agricultural perspective, in: van Beek, T. A., Breteler, H. (Eds.) *Phytochemistry and agriculture, Proc. Phytochem. Soc. Europe*, Oxford University Press, 1993, 34, pp.171–213.
- [152] Michael, A. J., Furze, J. M., Rhodes, M. J. C., Burtin, D., Molecular cloning and functional identification of a plant ornithine decarboxylase cDNA. *Biochem. J.* 1996, 314, 241–248.
- [153] Lee, Y.-S., Cho, Y.-D., Mutation of cysteine-115 to alanine in *Nicotiana glutinosa* ornithine decarboxylase reduces enzyme activity. *J. Biochem. Mol. Biol.* 2001, 34, 472–477.
- [154] Xu, B., Sheehan, M. J., Timko, M. P., Differential induction of ornithine decarboxylase (ODC) gene family members in transgenic tobacco (*Nicotiana tabacum* L. cv. Bright Yellow 2) cell suspensions by methyl-jasmonate treatment. *Plant Growth Regulation* 2004, 44, 101–116.
- [155] Yoo, T. H., Park, C.-J., Ham, B.-K., Kim, K.-J., Paek, K.-H. Ornithine decarboxylase gene (CaODC1) is specifically induced during TMV-mediated but salicylate-independent resistant response in hot pepper. *Plant Cell Physiology* 2004, 45, 1537–1542.
- [156] Hibi, N., Higashiguchi, S., Hashimoto, T., Yamada, Y., Gene expression in tobacco low-nicotine mutants. *Plant Cell* 1994, 6, 723–735.
- [157] Suzuki, K., Yamada, Y., Hashimoto, T., Expression of *Atropa belladonna* putrescine *N*-methyltransferase gene in root pericycle. *Plant Cell Physiol.* 1999, 40, 289–297.
- [158] Liu, T., Zhu, P., Cheng, K.-d., Meng, C., Zhu, H., Molecular cloning and expression of putrescine *N*-methyltransferase from the hairy roots of *Anisodus tanguticus*. *Planta Medica* 2005, 71, 987–989.
- [159] Stenzel, O., Teuber, M., Draeger, B., Putrescine *N*-methyltransferase in *Solanum tuberosum* L., a calystegine-forming plant. *Planta* 2006, 223, 200–212.

- [160] Teuber, M., Azemi, M. E., Namjoyan, F., Meier, A.-C. *et al.*, Putrescine *N*-methyltransferases - a structure-function analysis. *Plant Mol. Biol.* 2007, 63, 787–801.
- [161] Kai, G., Zhang, Y., Chen, J., Li, L., Yan, X., Zhang, R., Liao, P., Lu, X., Wang, W., , Zhou, G., Molecular characterization and expression analysis of two distinct putrescine *N*-methyltransferases from roots of *Anisodus acutangulus*. *Physiologia Plant.*, 2009, 135, 121–129.
- [162] Katoh, A., Shoji, T., Hashimoto, T., Molecular cloning of *N*-methylputrescine oxidase from tobacco. *Plant Cell Physiol.* 2007, 48, 550–554.
- [163] Heim, W. G., Sykes, K. A., Hildreth, S. B., Sun, J. *et al.*, Cloning and characterization of a *Nicotiana tabacum* methylputrescine oxidase transcript. *Phytochemistry* 2007, 68, 454–463.
- [164] Nakajima, K., Hashimoto, T., Yamada, Y., Two tropinone reductases with different stereospecificities are short-chain dehydrogenases evolved from a common ancestor. *Proc. Natl. Acad. Sci. USA* 1993, 90, 9591–9595.
- [165] Portsteffen, A., Draeger, B., Nahrstedt, A., The reduction of tropinone in *Datura stramonium* root cultures by two specific reductases. *Phytochemistry* 1994, 37, 391–400.
- [166] Kaiser, H., Richter, U., Keiner, R., Brabant, A. *et al.*, Immunolocalisation of two tropinone reductases in potato (*Solanum tuberosum* L.) root, stolon, and tuber sprouts. *Planta* 2006, 225, 127–137.
- [167] Keiner, R., Kaiser, H., Nakajima, K., Hashimoto, T., Draeger, B., Molecular cloning, expression and characterization of tropinone reductase II, an enzyme of the SDR family in *Solanum tuberosum* (L.). *Plant Mol. Biol.* 2002, 48, 299–308.
- [168] Li, R., Reed, D. W., Liu, E., Nowak, J. *et al.*, Functional genomic analysis of alkaloid biosynthesis in *Hyoscyamus niger* reveals a cytochrome P450 involved in littorine rearrangement. *Chemistry & Biology* 2006, 13, 513–520.

- [169] Matsuda, J., Okabe, S., Hashimoto, T., Yamada, Y., Molecular cloning of hyoscyamine 6-beta hydroxylase a 2 oxoglutarate-dependent dioxygenase from cultured roots of *Hyoscyamus niger*. *J. Biol. Chem.* 1991, 266, 9460–9464.
- [170] Suzuki, K.-i., Yun, D.-J., Chen, X.-Y., Yamada, Y., Hashimoto, T., An *Atropa belladonna* hyoscyamine 6beta-hydroxylase gene is differentially expressed in the root pericycle and anthers. *Plant Mol. Biol.* 1999, 40, 141–152.
- [171] Liu, T., Zhu, P., Cheng, K., Meng, C., He, H., Molecular cloning, expression and characterization of hyoscyamine 6 beta-hydroxylase from hairy roots of *Anisodus tanguticus*. *Planta Medica* 2005, 71, 249–253.
- [172] Kai, G., Chen, J., Zhou, L. Zhou, L. *et al.*, Molecular cloning and characterization of a new cDNA encoding hyoscyamine 6 beta-hydroxylase from roots of *Anisodus acutangulus*. *J. Biochem. Mol. Biol.* 2007, 40, 715–722.
- [173] Cardillo, A. B., Talou, J. R., Giulietti, A. M., Expression of *Brugmansia candida* hyoscyamine 6beta-hydroxylase gene in *Saccharomyces cerevisiae* and its potential use as biocatalyst. *Microbial Cell Factories*. 2008, 7:17; doi:10.1186/1475-2859-7-17.
- [174] El Jaber-Vazdekis, N., Gonzalez, C., Ravelo, A. G., Zarate, R., Cloning, characterization and analysis of expression profiles of a cDNA encoding a hyoscyamine 6 beta-hydroxylase (H6H) from *Atropa baetica* Willk. *Plant Physiol. Biochem.* 2009, 47, 20–25.
- [175] Siminszky, B., Gavilano, L., Bowen, S. W., Dewey, R. E., Conversion of nicotine to normicotine in *Nicotiana tabacum* is mediated by CYP82E4, a cytochrome P450 monooxygenase. *Proc. Natl. Acad. Sci. USA* 2005, 102, 14919–14924.
- [176] Wang, J., Sheehan, M., Timko, M. P., Differential expression of two members of the nuclear gene family encoding arginine decarboxylase in tobacco (*Nicotiana tabacum* L.). *J. Plant Biochem. Biotech.* 2000, 9, 57–65.

- [177] Nurhayati, N., Gonde, D., Ober, D., Evolution of pyrrolizidine alkaloids in *Phalaenopsis* orchids and other monocotyledons: Identification of deoxyhypusine synthase, homospermidine synthase and related pseudogenes. *Phytochemistry* 2009, 70, 508–516.
- [178] Reimann, A., Nurhayati, N., Backenkoehler, A., Ober, D., Repeated evolution of the pyrrolizidine alkaloid-mediated defense system in separate angiosperm lineages. *Plant Cell* 2004, 16, 2772–2784.
- [179] Abdelhady, M. I. S., Beuerle, T., Ober, D., Homospermidine in transgenic tobacco results in considerably reduced spermidine levels but is not converted to pyrrolizidine alkaloid precursors. *Plant Mol. Biol.* 2009, 71, 145–155.
- [180] Lukacin, R., Schreiner, S., Matern, U., Transformation of acridone synthase to chalcone synthase. *FEBS Lett.* 2001, 508, 413–417.
- [181] Junghanns, K. T., Kneusel, R. E., Baumert, A., Maier, W. *et al.*, Molecular cloning and heterologous expression of acridone synthase from elicited *Ruta graveolens* L. cell suspension cultures. *Plant Mol. Biol.* 1995, 27, 681–692.
- [182] Wanibuchi, K., Zhang, P., Abe, T., Morita, H. *et al.*, An acridone-producing novel multifunctional type III polyketide synthase from *Huperzia serrata*. *FEBS J.* 2007, 274, 1073–1082.
- [183] Rohde, B., Hans, J., Martens, S., Baumert, A. *et al.*, Anthranilate *N*-methyltransferase, a branch-point enzyme of acridone biosynthesis. *Plant J.* 2008, 55, 541–553.
- [184] Girod, P. A., Zryd, J. P., Biogenesis of betalains. Purification and partial characterization of Dopa 4 5-dioxygenase from *Amanita muscaria*. *Phytochemistry* 1991, 30, 169–174.
- [185] Christinet, L., Burdet, F. X., Zaiko, M., Hinz, U., Zryd, J.-P., Characterization and functional identification of a novel plant 4,5-extradiol dioxygenase involved in betalain pigment biosynthesis in *Portulaca grandiflora*. *Plant Physiol.* 2004, 134, 265–274.

- [186] Takahashi, K., Sakuta, M., Cloning and characterization of DOPA dioxygenase from *Phytolacca americana*. *Plant Cell Physiol.* 2007, 48, 224–230.
- [187] Sasaki, N., Abe, Y., Goda, Y., Adachi, T. *et al.*, Detection of DOPA 4,5-dioxygenase (DOD) activity using recombinant protein prepared from *Escherichia coli* cells harboring cDNA encoding DOD from *Mirabilis jalapa*. *Plant Cell Physiol.* 2009, 50, 1012–1016.
- [188] Joy, R. W. I., Sugiyama, M., Fukuda, H., Komamine, A., Cloning and characterization of polyphenol oxidase cDNAs of *Phytolacca americana*. *Plant Physiol.* 1995, 107, 1083–1089.
- [189] Vogt, T., Grimm, R., Strack, D., Cloning and expression of a cDNA encoding betanidin 5-*O*-glucosyltransferase, a betanidin- and flavonoid-specific enzyme with high homology to inducible glucosyltransferases from the Solanaceae. *Plant J.* 1999, 19, 509–519.
- [190] Katz, L., Hutchinson, R., Genetic engineering of antibiotic producing organisms. *Annu. Rep. Medic. Chem.* 1992, 27, 129–38.
- [191] McDaniels, R., Ebert-Koshla, S., Hopwood, D.A., Koshla, C, Engineered biosynthesis of novel polyketides. *Science* 1993, 262, 1546–50.
- [192] Inui, T., Tamura, K.-I., Fujii, N., Morishige, T., Sato, F., Overexpression of *Coptis japonica* norcoclaurine 6-*O*-methyltransferase overcomes the rate-limiting step in benzyloquinoline alkaloid biosynthesis in cultured *Eschscholzia californica*. *Plant & Cell Physiol.*, 2007, 48, 252–262.
- [193] Fujii, N., Inui, T., Iwasa, K., Morishige, T., Sato, F., Knockdown of berberine bridge enzyme by RNAi accumulates (S)-reticuline and activates a silent pathway in cultured California poppy cells. *Transgenic Res.* 2007, 16, 363–375.
- [194] Allen, R. S, Miller, J. A. C, Chitty, J. A., Fist, A. J. *et al.*, Metabolic engineering of morphinan alkaloids by over-expression and RNAi suppression of salutaridinol 7-*O*-acetyltransferase in opium poppy. *Plant Biotechnol. J.* 2008, 6, 22–30.

- [195] Hawkins, K. M., Smolke, C. D., Production of benzyloquinoline alkaloids in *Saccharomyces cerevisiae*. *Nat. Chem. Biol.* 2008, 4, 564–573.
- [196] Chavadej, S., Brisson, N., McNeil, J. N., De Luca, V., Redirection of tryptophan leads to production of low indole glucosinolate canola. *Proc. Natl. Acad. Sci. USA* 1994, 91, 2166–2170.
- [197] Wang, M., Li, Q.-R., Di Fiore, S., Fischer, R., Expression of recombinant tryptophan decarboxylase in different subcellular compartments in tobacco plant. *Acta Botanica Sinica*. 2002, 44, 314–317.
- [198] Gill, R. I. S., Ellis, B. E., Over-expression of tryptophan decarboxylase gene in poplar and its possible role in resistance against *Malacosoma disstria*. *New Forests* 2006, 31, 195–209.
- [199] Goddijn, O. J. M., Pennings, E. J. M., Van Der Helm, P., Schilperoort, R. A. *et al.*, Overexpression of a tryptophan decarboxylase cDNA in *Catharanthus roseus* crown gall calluses results in increased tryptamine levels but not in increased terpenoid indole alkaloid production. *Transgenic Res.* 1995, 4, 315–323.
- [200] Leech, M. J., May, K., Hallard, D., Verpoorte, R. *et al.*, Expression of two consecutive genes of a secondary metabolic pathway in transgenic tobacco: Molecular diversity influences levels of expression and product accumulation. *Plant Mol. Biol.* 1998, 38, 765–774.
- [201] Geerlings, A., Hallard, D., Caballero, A. M., Cardoso, I. L. *et al.*, Alkaloid production by a *Cinchona officinalis* 'Ledgeriana' hairy root culture containing constitutive expression constructs of tryptophan decarboxylase and strictosidine synthase cDNAs from *Catharanthus roseus*. *Plant Cell Reports* 1999, 19, 191–196.
- [202] Canel, C., Lopes-Cardoso, M. I., Whitmer, S., Van Der Fits, L. *et al.*, Effects of over-expression of strictosidine synthase and tryptophan decarboxylase on alkaloid production by cell cultures of *Catharanthus roseus*. *Planta* 1998, 205, 414–419.

- [203] Whitmer, S., Canel, C., Van Der Heijden, R., Verpoorte, R., Long-term instability of alkaloid production by stably transformed cell lines of *Catharanthus roseus*. *Plant Cell Tissue & Organ Culture* 2003, 74, 73–80.
- [204] Hallard, D., Van Der Heijden, R., Verpoorte, R., Lopes Cardoso, M. I. *et al.*, Suspension cultured transgenic cells of *Nicotiana tabacum* expressing tryptophan decarboxylase and strictosidine synthase cDNAs from *Catharanthus roseus* produce strictosidine upon secologanin feeding. *Plant Cell Reports* 1997, 17, 50–54.
- [205] Uefuji, H., Tatsumi, Y., Morimoto, M., Kaothien-Nakayama, P. *et al.*, Caffeine production in tobacco plants by simultaneous expression of three coffee N-methyltransferases and its potential as a pest repellent. *Plant Mol. Biol.* 2005, 59, 221–227.
- [206] Ogita, S., Uefuji, H., Yamaguchi, Y., Koizumi N., Sano, H., RNA interference: Producing decaffeinated coffee plants. *Nature* 2003, 423, 823–823.
- [207] Besumbes, O., Sauret-Gueto, S., Phillips, M. A., Imperial, S., Rodriguez-Concepcion, M., Boronat, A., Metabolic engineering of isoprenoid biosynthesis in *Arabidopsis* for the production of taxadiene, the first committed precursor of Taxol. *Biotechnol. Bioeng.* 2004, 88, 168–175.
- [208] Kovacs, K., Zhang, L., Linforth, R. S. T., Whittaker, B. *et al.*, Redirection of carotenoid metabolism for the efficient production of taxadiene [taxa-4(5),11(12)-diene] in transgenic tomato fruit. *Transgenic Research* 2007, 16, 121–126.
- [209] Anterola, A., Shanle, E., Perroud, P.-F., Quatrano, R., Production of taxa-4(5),11(12)-diene by transgenic *Physcomitrella patens*. *Transgenic Research* 2009, 18, 655–660.
- [210] Jennewein, S., Park, H., DeJong, J. M., Long, R. M. *et al.*, Coexpression in yeast of *Taxus* cytochrome P450 reductase with cytochrome P450 oxygenases involved in taxol biosynthesis. *Biotechnology Bioengineering*, 2005, 89, 588–598.

[211] Petersen, M., Current status of metabolic phytochemistry. *Phytochemistry* 2007, 68, 2847–2860.

[212] Chappell, J., Production platforms for the molecular pharming of alkaloid diversity. *Proc. Natl. Acad. Sci. USA* 2008, 105, 7897–7898.

[213] Kirby, J., Keasling, J. D., Biosynthesis of plant isoprenoids: Perspectives for microbial engineering. *Annu. Rev. Plant Biol.* 2009, 60, 335–355.

Figure 1. Some strategies for the production of secondary metabolites.

Figure 2. Overview of biosynthetic pathways of major groups of alkaloids.

Figure 3. Biosynthesis of benzyloquinoline alkaloids. (A) Pathway from tyrosine to S-reticuline; (B) pathway from reticuline to protoberberine alkaloids; (C) pathway from reticuline to morphinan alkaloids, (D) pathway from scoulerine to benzophenanthridine alkaloids.

Figure 4. Biosynthesis of monoterpene indole alkaloids. (A) Pathway from tryptophan to ajmalicine; (B) pathway from strictosidine to ajmalin; (C) pathway from strictosidine to vinblastine.

Figure 5. Biosynthesis of paclitaxel.

Figure 6. Biosynthetic pathway from ornithine to tropane alkaloids.

Table 1. Cloned genes and characterized enzymes involved in the biosynthesis of benzylisoquinoline alkaloids (see also [42–46, 48, 50, 54] for earlier publications). *, These plants do not produce benzylisoquinoline alkaloids.

Enzyme	Plant	Reference
Tyrosine decarboxylase (TYDC)	<i>Papaver somniferum</i> , <i>Oryza sativa</i> *, <i>Thalictrum flavum</i> , <i>Arabidopsis thaliana</i> *	[61-64]
Norcoclaurine synthase (NCS)	<i>Thalictrum flavum</i> , <i>Coptis japonica</i>	[65, 66]
Norcoclaurine 6- <i>O</i> -methyltransferase (6OMT)	<i>Thalictrum tuberosum</i> , <i>Thalictrum flavum</i> , <i>Coptis japonica</i> , <i>Papaver somniferum</i> ,	[63, 67-70]
Coclaurine <i>N</i> -methyltransferase (CNMT)	<i>Thalictrum flavum</i> , <i>Papaver somniferum</i> , <i>Coptis japonica</i>	[63,69,71]
Berberamine synthase (Cyp80A1)	<i>Berberis stolonifera</i>	[72]
<i>N</i> -methylcocclaurine 3'-hydroxylase (Cyp80B)	<i>Thalictrum flavum</i> , <i>Eschscholzia californica</i>	[63, 73-75]
3-Hydroxy- <i>N</i> -methylcocclaurine 4- <i>O</i> -Methyltransferase (4'OMT)	<i>Thalictrum flavum</i> , <i>Coptis japonica</i> , <i>Papaver somniferum</i>	[63, 68, 69]
Reticuline 7- <i>O</i> -methyltransferase (7OMT)	<i>Papaver somniferum</i>	[70]
Berberine bridge enzyme (BBE)	<i>Thalictrum flavum</i> , <i>Eschscholzia californica</i> , <i>Papaver somniferum</i>	[63, 76,77]
Cheilanthalifoline synthase (Cyp719A5)	<i>Eschscholzia californica</i>	[78]
Stylophine synthase (Cyp719A2, 3)	<i>Eschscholzia californica</i>	[79]

Tetrahydroprotoberberine <i>N</i> -methyltransferase (TNMT)	<i>Papaver somniferum</i>	[56]
Cyp719A1	<i>Thalictrum flavum</i> , <i>Coptis japonica</i>	[63, 80]
Scoulerine 9- <i>O</i> -methyltransferase (SOMT)	<i>Thalictrum flavum</i> , <i>Coptis japonica</i>	[63, 81]
Columbamine <i>O</i> -methyltransferase (CoOMT)	<i>Coptis japonica</i>	[82]
Salutaridine reductase (SalR)	<i>Papaver somniferum</i>	[83]
Salutaridinol 7- <i>O</i> -acetyltransferase (SalAT)	<i>Papaver somniferum</i>	[84]
Codeinone reductase (COR)	<i>Papaver somniferum</i>	[85]
Cyp80G2	<i>Coptis japonica</i>	[86]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Cloned genes and characterized enzymes involved in the biosynthesis of monoterpene indole alkaloids. *, Not an alkaloidal plant.

Enzyme	Plant	Reference
Tryptophan decarboxylase (TDC)	<i>Oryza sativa</i> *, <i>Catharanthus roseus</i> , <i>Camptotheca acuminata</i> , <i>Ophiorrhiza pumila</i>	[62, 93-95]
Geraniol 10-hydroxylase (G10H or Cyp76B6)	<i>Catharanthus roseus</i>	[96]
Cyp72A1	<i>Catharanthus roseus</i>	[97]
Loganic acid <i>O</i> -methyltransferase (LAMT)	<i>Catharanthus roseus</i>	[98]
Strictosidine synthase (STR)	<i>Rauvolfia serpentina</i> , <i>Ophiorrhiza pumila</i> , <i>Catharanthus roseus</i> , <i>Rauvolfia mannii</i> , <i>Rauvolfia verticillata</i>	[89, 95, 99-101]
Strictosidine beta- <i>D</i> -glucosidase (SGD)	<i>Catharanthus roseus</i> , <i>Rauvolfia serpentina</i>	[102, 103]
Tabersonine 16-hydroxylase (T16H or Cyp71D12)	<i>Catharanthus roseus</i>	[104]
16-Hydroxytabersonine-16- <i>O</i> -methyltransferase (16OMT)	<i>Catharanthus roseus</i>	[105]
Desacetoxyvindoline-4-hydroxylase (D4H)	<i>Catharanthus roseus</i>	[106]
Deacetylvindoline 4- <i>O</i> -acetyltransferase (DAT)	<i>Catharanthus roseus</i>	[107]

Minovincinine-19- <i>O</i> - acetyltransferase (MAT)	<i>Catharanthus roseus</i>	[108]
Polyneuridine aldehyde esterase (PNAE)	<i>Rauvolfia serpentina</i>	[109]
Vinorine synthase (VS)	<i>Rauvolfia serpentina</i>	[110, 122]
Acetylajmalan esterase (AAE)	<i>Rauvolfia serpentina</i>	[87]
Raucaffricine- <i>O</i> -beta- <i>D</i> - glucosidase	<i>Rauvolfia serpentina</i>	[111]
Perakine reductase (PR)	<i>Rauvolfia serpentina</i>	[112]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Cloned and characterized enzymes involved in the biosynthesis of ergot alkaloids

Enzyme	Fungus	Reference
4-Dimethylallyltryptophan synthase	<i>Claviceps purpurea</i> , <i>Neotyphodium sp.</i> , <i>Aspergillus fumigatus</i> , <i>Malbranchea aurantiaca</i>	[116-119]
7-Dimethylallyltryptophan synthase	<i>Aspergillus fumigatus</i>	[120]
4-Dimethylallyltryptophan <i>N</i> -methyltransferase	<i>Aspergillus fumigatus</i>	[121]
Brevianamide F prenyltransferase	<i>Aspergillus fumigatus</i>	[113]

Table 4. Cloned and characterized enzymes involved in the biosynthesis of purine alkaloids

Enzyme	Plant	Reference
Xanthosine 7- <i>N</i> -methyltransferase (XMT) or 7-methylxanthosine synthase (XRS)	<i>Coffea arabica</i>	[126, 127]
Caffeine synthase (CS)	<i>Camellia sinensis</i> , <i>Coffea arabica</i>	[124, 126, 128]
7-Methylxanthine methyltransferase (MXMT)	<i>Coffea arabica</i>	[127, 129]
Dimethylxanthosine methyltransferase (DXMT)	<i>Coffea arabica</i>	[127]

Table 5. Cloned and characterized enzymes involved in the biosynthesis of paclitaxel

Enzyme	Plant	Reference
Taxa-4(5), 11(12)-diene synthase	<i>Taxus brevifolia</i> , <i>T. chinensis</i> , <i>T. media</i>	[131-133]
Geranylgeranyl diphosphate synthase	<i>Taxus canadensis</i>	[134]
Taxane 2 α -O-benzoyltransferase	<i>Taxus cuspidata</i>	[135]
10-Deacetylbaccatin III-10-O-acetyltransferase	<i>Taxus cuspidata</i> , <i>T. media</i>	[136, 137]
Taxa-4(20),11(12)-dien-5 α -ol-O-acetyl transferase	<i>Taxus sp.</i>	[138]
Taxane 13 α -hydroxylase	<i>Taxus sp.</i>	[139]
Taxane 10 β -hydroxylase	<i>Taxus sp.</i> , <i>T. media</i>	[140, 141]
Taxoid 14 β -hydroxylase	<i>Taxus sp.</i>	[142]
Taxoid 2 α -hydroxylase	<i>Taxus sp.</i>	[143]
Taxoid 7 β -hydroxylase	<i>Taxus sp.</i>	[144]
Taxadiene 5 α -hydroxylase	<i>Taxus sp.</i>	[145]
C-13 phenylpropanoid side chain-CoA acetyl transferase	<i>Taxus sp.</i>	[146]
C13-side-chain <i>N</i> -benzoyl transferase	<i>Taxus sp.</i>	[147]
Phenylalanine ammonium mutase	<i>Taxus cuspidata</i> , <i>T. chinensis</i>	[148, 149]
Taxadiene 5 α -ol-O-acetyltransferase	<i>Taxus sp.</i>	[150]

Table 6. Cloned and characterized enzymes involved in the biosynthesis of tropane alkaloids and nicotine

Enzyme	Plant	Reference
Ornithine decarboxylase (ODC)	<i>Datura stramonium</i> , <i>Nicotiana glutinosa</i> , <i>N. tabacum</i> , <i>Capsicum annuum</i>	[152-155]
Putrescine <i>N</i> -methyltransferase (PMT)	<i>Nicotiana tabacum</i> , <i>Atropa belladonna</i> , <i>Hyoscyamus niger</i> , <i>Anisodus tanguticus</i> , <i>Solanum spec.</i> , <i>Calystegia sepium</i> , <i>Datura spec.</i> , <i>Physalis divaricarpa</i> , <i>Anisodus acutangulus</i>	[156-161]
<i>N</i> -methylputrescine oxidase (MPO)	<i>Nicotiana tabacum</i>	[162, 163]
Tropinone reductase I (TRI)	<i>Hyoscyamus niger</i> , <i>Datura stramonium</i> , <i>Solanum tuberosum</i>	[164-166]
Tropinone reductase II (TRII)	<i>Hyoscyamus niger</i> , <i>Datura stramonium</i> , <i>Solanum tuberosum</i>	[164, 165, 167]
Cyp80F1	<i>Hyoscyamus niger</i>	[168]
Hyoscyamine 6 β -hydroxylase (H6H)	<i>Hyoscyamus niger</i> , <i>Atropa belladonna</i> , <i>Anisodus tanguticus</i> , <i>A. acutangulus</i> , <i>Brugmansia candida</i> , <i>Atropa beatica</i>	[169-174]
Nicotine <i>N</i> -demethylase (NND)	<i>Nicotiana tabacum</i>	[175]
Arginine decarboxylase (ADC)	<i>Nicotiana tabacum</i>	[176]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 7. Cloned and characterized enzymes involved in the biosynthesis of pyrrolizidine alkaloids

Enzyme	Plant	Reference
Homospermidine synthase (HSS)	Boraginaceae (<i>Heliotropium</i> , <i>Cynoglossum</i> , <i>Symphytum</i>), Asteraceae (<i>Eupatorium</i> , <i>Senecio</i> , <i>Petasites</i>), Convolvulaceae (<i>Ipomoea hederifolia</i>), Solanaceae (<i>Nicotiana tabacum</i>), Fabaceae (<i>Crotalaria</i>), Orchidaceae (<i>Phalaenopsis</i>)	[178]

Table 8. Cloned and characterized enzymes involved in the biosynthesis of acridone alkaloids

Enzyme	Plant	Reference
Acridone synthase	<i>Ruta graveolens</i> , <i>Huperzia serrata</i>	[181, 182]
Anthranilate <i>N</i> -methyltransferase	<i>Ruta graveolens</i>	[183]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 9. Cloned and characterized enzymes involved in the biosynthesis of betalains

Enzyme	Plant	Reference
DOPA 4,5-dioxygenase	<i>Amanita muscaria</i> , <i>Portulacca</i> <i>grandiflora</i> , <i>P.</i> <i>americana</i> ,	[184-187]
Polyphenol oxidase	<i>Phytolacca</i> <i>americana</i>	[188]
Betanidin 5- <i>O</i> - glucosyltransferase	<i>Dorotheanthus</i> <i>bellidiformis</i>	[189]

Table 10. Expression of tryptophan decarboxylase and strictosidine synthase; metabolic engineering of the monoterpene indole alkaloid biosynthesis

Enzyme	Source	Transgenic plant	Goal/remark	Ref.
Tryptophan decarboxylase	<i>Catharanthus roseus</i>	<i>Brassica napus</i>	Reduction of tryptophan derived glucosinolates	[196]
	<i>Catharanthus roseus</i>	<i>Nicotina tabacum</i>	Investigation of influence of subcellular localization on enzyme activity and tryptamine accumulation	[197]
	<i>Camptotheca acuminata</i>	<i>Populus sp.</i>	Study of tryptamine accumulation and effect on insect pests	[198]
	<i>Catharanthus roseus</i>	<i>Catharanthus roseus</i>	Stimulation of tryptamine production but no change of MIA content	[199]
Tryptophan decarboxylase + strictosidine synthase	<i>Catharanthus roseus</i>	<i>Nicotina tabacum</i>	Plasmid construct with both genes	[200]
	<i>Catharanthus roseus</i>	<i>Cinchona officinalis</i>	Functional expression of TDC and STR; enhanced levels of strictosidine and Cinchona alkaloids in hairy root cultures	[201]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<i>Catharanthus roseus</i>	<i>Catharanthus roseus</i>	Over-expression of TDC and STR with stimulation of MIA synthesis	[202, 203]
	<i>Catharanthus roseus</i>	<i>Nicotina tabacum</i>	Functional expression of TDC and STR in tobacco cell cultures and strictosidine production after feeding od secologanin	[204]

1
2
3 **((Biographical Material))**
4
5

6
7 **Holger Schäfer** (Dr. rer. nat.) studied biology at the University of Darmstadt and obtained his
8
9 doctorate in 2000 at the University of Heidelberg. He is working as a research assistant at the
10
11 Institute of Pharmacy and Molecular Biotechnology, Heidelberg, Germany. His research
12
13 interests are in molecular biology with a focus on the recombinant production of proteins and
14
15 natural products.
16

17
18 **Michael Wink** (Dr. rer. nat.) is a professor of Pharmaceutical Biology at Heidelberg
19
20 University and director at the Institute of Pharmacy and Molecular Biotechnology,
21
22 Heidelberg, Germany. He received his research training in biology and chemistry at the
23
24 Universities of Bonn, Braunschweig and Munich. In 1988 he became a professor at the
25
26 University of Mainz and in 1989 at Heidelberg University. His research interests range from
27
28 plant natural products (identification, biochemistry, pharmacology, biotechnology), medicinal
29
30 and toxic plants to evolution (phylogeny and phylogeography). He is author of more than 15
31
32 books or monographs and author/coauthor of more than 450 refereed publications.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

morphine

taxol

Figure 2. Overview of biosynthetic pathways of major groups of alkaloids
271x183mm (300 x 300 DPI)

Figure 3. Biosynthesis of benzylisoquinoline alkaloids
A. Pathway from tyrosine to S-reticuline; B. pathway from reticuline to to protoberberine alkaloids;
C. pathway from reticuline to morphinan alkaloids, D. pathway from scoulerine to benzophenanthridine alkaloids.

175x252mm (300 x 300 DPI)

Figure 3. Biosynthesis of benzyloquinoline alkaloids

A. Pathway from tyrosine to S-reticuline; B. pathway from reticuline to protoberberine alkaloids; C. pathway from reticuline to morphinan alkaloids, D. pathway from scoulerine to benzophenanthridine alkaloids.

265x169mm (300 x 300 DPI)

Figure 3. Biosynthesis of benzylisoquinoline alkaloids
A. Pathway from tyrosine to S-reticuline; B. pathway from reticuline to to protoberberine alkaloids;
C. pathway from reticuline to morphinan alkaloids, D. pathway from scoulerine to benzophenanthridine alkaloids.

255x160mm (300 x 300 DPI)

Figure 3. Biosynthesis of benzyloquinoline alkaloids

A. Pathway from tyrosine to S-reticuline; B. pathway from reticuline to protoberberine alkaloids; C. pathway from reticuline to morphinan alkaloids, D. pathway from scoulerine to benzophenanthridine alkaloids.

265x156mm (300 x 300 DPI)

Figure 4. Biosynthesis of monoterpene indole alkaloids
A. Pathway from tryptophan to ajmalicine; B. pathway from strictosidine to ajmalin C. pathway from strictosidine to vinblastine

267x177mm (300 x 300 DPI)

Figure 4. Biosynthesis of monoterpene indole alkaloids
A. Pathway from tryptophan to ajmalicine; B. pathway from strictosidine to ajmalin C. pathway from strictosidine to vinblastine

265x159mm (300 x 300 DPI)

Figure 4. Biosynthesis of monoterpene indole alkaloids
A. Pathway from tryptophan to ajmalicine; B. pathway from strictosidine to ajmalin C. pathway from strictosidine to vinblastine

261x175mm (300 x 300 DPI)

Figure 5. Biosynthesis of paclitaxel
266x177mm (300 x 300 DPI)

Figure 6. Biosynthetic pathway from ornithine to tropane alkaloids
178x261mm (300 x 300 DPI)

784x1078mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

914x1371mm (72 x 72 DPI)