The effect of the calcium source, level and the particle size on calcium retention, eggshell quality and the overall calcium requirement in laying hens

Martina Lichovnikova

► To cite this version:

Martina Lichovnikova. The effect of the calcium source, level and the particle size on calcium retention, eggshell quality and the overall calcium requirement in laying hens. *British Poultry Science*, 2007, 48 (01), pp.71-75. 10.1080/00071660601148203 . hal-00545309

HAL Id: hal-00545309

<https://hal.science/hal-00545309>

Submitted on 10 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**The effect of the calcium source, level and the particle size
on calcium retention, eggshell quality and the overall
calcium requirement in laying hens**

Journal:	<i>British Poultry Science</i>
Manuscript ID:	CBPS-2006-246.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	20-Oct-2006
Complete List of Authors:	Lichovnikova, Martina; Mendel University of Agriculture and Forestry
Keywords:	Digestion, Egg shell, Laying hens

CBPS_2006_246
edited M G MacLeod, November 2006

1
2
3
4 1
5
6 2
7
8
9 3
10
11 4
12
13 5
14
15
16 6
17
18 7
19
20
21 8
22
23
24 9
25
26 10
27
28 11
29
30
31 12
32
33 13
34
35
36 14
37
38 15
39
40 16
41
42
43 17
44
45 18
46
47 19
48
49
50 20
51
52 21
53
54 22
55
56
57 23
58
59 24
60

**The effect of dietary calcium source, concentration and particle size
on calcium retention, eggshell quality and overall calcium
requirement in laying hens**

M. LICHNIKOVA
*Department of Animal Breeding, Mendel University of Agriculture and Forestry in
Brno, Zemedelska 1, 613 00 Brno, Czech Republic*

RUNNING TITLE: CALCIUM SOURCE AND SHELL QUALITY

Telephone: ++420 54513 3229
Email: lichov@mendelu.cz

Accepted for publication 16th November 2006

Abstract 1. Four different sources of calcium in isonitrogenous and isoenergetic diets were fed to laying hens for two weeks when they were 56 and 57 weeks old. The calcium source blends were as follows: 29% fine limestone (LF) + 71% large limestone (LG), 32% fine limestone (LF) + 68% eggshell, 32% fine limestone (LF) + 68% oyster shell, 50% fine limestone + 50% large limestone (LG). The contents of these blends of calcium in the diets were as follows: 103.3 g/kg, 93.3 g/kg, 93.3 g/kg and 93.3 g/kg respectively.

2. The coefficients of calcium retention were significantly higher in 50:50 LF:LG (0.578) and 32:68 LF:eggshell (0.576). The midnight feeding significantly improved the coefficient of calcium retention in all mixtures except 50:50 LF:LG.

3. In the mixture 29:71 LF:LG and 32:68 LF:oyster shell, there was significantly greater eggshell quality, eggshell weight, eggshell thickness and eggshell strength. Midnight feeding had no significant effect on eggshell quality.

4. In the ration with oyster shells, 96.5% of the retained calcium was deposited in the eggshell, but in ration 32:68 LF:eggshells and 50:50 LF:LG the utilisation was only 73.9% and 78.6% respectively.

5. To ensure good quality eggshells in the last third of production, the recommendation for calcium is 4.1 g/kg (900 g/kg dry matter, feed intake 110 g/d). As a source of calcium in this stage of production, a feed mixture containing two-thirds large particles should be used (limestone grit or oyster shell).

INTRODUCTION

The quality of the eggshells has a major influence on the economics of egg production. Damaged or broken shells account for 6-8% of all the eggs laid (Bain, 1997) and the highest incidence of cracked eggs occurs mainly in the last third of the laying period (after 53 weeks of age). The source and particle size of mineral

50 nutrients, mainly calcium, play an important role in maintaining eggshell quality. It
51 is known that hens given ground limestone as a single source of calcium had lower
52 eggshell quality than hens fed blends of particulate and ground limestone (Guinotte
53 and Nys, 1991; Richter *et al.*, 1999; Pavlovski *et al.*, 2003; Koreleski and
54 Swiatkiewicz, 2004).

55 Rao *et al.* (1992) reported that limestone solubility in laying hens improves if
56 retention time is prolonged in the gizzard, which means feeding a minimum particle
57 size of 1.0 mm. A minimum particle size less than 1.0 mm did not sustain retention
58 in the gizzard. Scheideler (1998) reported significantly greater specific gravity
59 ($P<0.05$) of the eggs from the hens on the diets, which included large particle size
60 calcium (fine and large limestone 50:50 or 75:25 limestone:oyster shell in the diets)
61 compared with hens fed with diets containing 100% fine limestone or 100% ground
62 eggshell. Scheideler (2004), on the basis of her results, recommends that laying hens
63 be fed at least 25% of their calcium from a large particle calcium source. Also,
64 Ahmad and Balander (2003) reported higher egg specific gravity with the partial
65 replacement of limestone (50%) with oyster shell as the calcium source. However,
66 not enough data exists at this time to justify increasing the large particle portion to
67 more than 50% of the ration. Also the requirement of calcium may not be the same,
68 as reported in earlier studies or NRC (1994), to meet the demand for relatively high
69 production in the last third of the laying period. The aim of the present study was to
70 evaluate the effects of a higher proportion of the large particle calcium (taken from
71 different sources) and higher calcium concentration on the calcium retention and the
72 eggshell quality in the last third of the laying period.

73 MATERIALS AND METHODS

1
2
3
4 74 A total of 24 laying hens were divided into 4 groups and they were kept in individual
5
6 75 balance cages. They were all fed on the same basic diet (corn 407 g/kg%, wheat 340
7
8 76 g/kg, soybean meal 138 g/kg). The 4 diets, however, varied in the source of calcium
9
10
11 77 as follows: 29% fine limestone (LF) + 71% large limestone (LG: 1 – 2 mm x 1 mm),
12
13 78 32% fine limestone and 68% eggshell (2 - 5 mm x 0.4 mm in the first digestion trial
14
15 79 and 1-2 mm x 0.4 mm x 0.8 mm in the second digestion trial), 32% fine limestone
16
17 80 and 68% oyster shell (2 - 5 mm, Oyta Oystershells no. 1), 50% fine limestone and
18
19 81 50% large limestone (1 - 2 mm). The contents of these sources of calcium in the diets
20
21 82 were as follows: 103.3 g/kg, 93.3 g/kg, 93.3 g/kg and 93.3 g/kg respectively;
22
23 83 consequently, the content of calcium was higher in the diet 29:68 LF:LG. The fine
24
25 84 limestone and the large limestone were from the same source. The compositions of
26
27 85 the diets are shown in Table 1. After a 5-d adjustment period, when the hens were
28
29 86 fed *ad libitum*, two digestion trials were done. Each digestion trial took 5 d and
30
31 87 between digestion trials there was a 2-d pause. In both trials the same hens received
32
33 88 the same diets. The results in each group were calculated as average of 12 samples (2
34
35 89 x 6). The hens were 56 and 57 weeks of age. Light was provided for 16 h per d from
36
37 90 0400 h to 2000 h. During the first digestion trial the hens were given 120 g of the
38
39 91 mixtures at 0830 h. In the second digestion trial, a midnight feeding was added and
40
41 92 the hens were fed 90 g of the mixtures during the light period and 30 g during the
42
43 93 dark period. At the end of each digestion trial the remainder of the diets in the
44
45 94 feeders were weighed and average daily food intake was calculated for each hen.
46
47 95 During the dark period the light was turned on for 1 h from 2400 h to 0100 h. Water
48
49 96 was provided *ad libitum* throughout the day. Droppings were collected every 24 h
50
51 97 and dried at 65 °C each day after collection. The retention of calcium, energy,
52
53 98 nitrogen, ash, phosphorus and fat were estimated using the indicator method.
54
55
56
57
58
59
60

99 Insoluble ash in 3 M HCl was used as an indicator. The content of nitrogen was
100 determined according to Kjeldahl. The content of fat was determined according to
101 Soxhlet. Ash was determined after combustion at 550 °C. The content of calcium and
102 phosphorus were determined by spectrophotometry. The energy was determined in
103 an automatic bomb calorimeter PARR 1281 (Parr Instrument Company, Illinois,
104 USA). An acid correction was not done. All the eggs were collected and weighed.
105 The strength of the eggshells (N) was measured manually by destructive methods.
106 Eggs were compressed between two parallel plates by a steadily increasing load until
107 failure resulted. The force was recorded throughout each test and the strength of the
108 eggshell was given in terms of the force at failure. The force was measured vertically
109 to the axis. Also the thickness (the average of both ends and at the middle) of
110 eggshell was evaluated and dry eggshells were weighed.

Table 1 near here

111 On the basis of the weight of the eggshells, the calcium content in the
112 eggshells and the calcium retention, the proportions of retained calcium deposited in
113 the eggshells were determined in each group. The following equations were used:

114 $Ca_p = 100 * Ca_a / Ca_r$

115 $Ca_p = \text{proportion of retained Ca deposited into the eggshells (\%)}$

116 $Ca_a = \text{amount of calcium deposited to the eggshells (g/hen/d)}$

117 $Ca_a = Wt_e \times Ca_e / 100$

118 $Wt_e = \text{weight of dry eggshells (g/hen.d)}$

119 $Ca_e = \text{calcium content in the eggshells (\%)}$

120 $Ca_r = \text{calcium retention (g/hen.d)}$

121 $Ca_r = Ca_i \times Ca_c$

122 $Ca_i = \text{calcium intake (g/hen.d)}$

123 $Ca_c = \text{coefficient of calcium retention}$

124 Data were analysed throughout by one-way analysis of variance (ANOVA)
125 using the software package *Unistat 5.1* (UNISTAT Ltd, England). Tukey-HSD
126 (eggshell quality), and Least Significant Difference (the coefficients of nutrient
127 retention) were used as the *post hoc* test for all possible pair-wise comparisons within
128 groups.

129 The *in vitro* solubility of the limestone used in the study was determined by
130 the method described by Zhang and Coon (1997). A 2.0 g limestone sample was
131 poured into a 400 ml beaker containing 200 ml of 0.2 N HCl solution that was
132 warmed at 42 °C until the temperature of the solution became constant in a water
133 bath oscillation at 80 Hz. After allowing 10 min for reaction, the undissolved
134 limestone was filtered onto a preweighed filter paper and weighted after drying in a
135 60 °C oven for 20 h. The *in vitro* solubility of limestone was expressed as the
136 percentage weight loss.

137 RESULTS AND DISCUSSION

138 The coefficients of the retention of the dry matter, fat, energy, calcium, nitrogen, ash
139 and phosphorus are shown in Table 2. The coefficients of the calcium retention were
140 significantly higher ($P=0.0271$) in 50:50 LF:LG and LF:Eggshell in comparison with
141 29:71 LF:LG and LF:Oystershell. Also, Scheideler (1998) found that the highest
142 retention of calcium was from eggshells. The midnight feeding significantly
143 improved the retention of calcium ($P<0.01$) in all of the groups except 50:50 LF:LG
144 (Table 4). The highest retention of calcium and the highest retention of all others
145 nutrients (energy, nitrogen, ash and phosphorus) were found in the rations with
146 eggshells.

Tables 2,3,4 near here

147 The weight of the eggs and the quality of the eggshells is shown in Table 3.
148 The weight of the eggs was significantly lower ($P=0.0422$) in the 29:71 LF:LG. In

1
2
3
4 149 this group the amount of calcium was the highest. The retention of energy was
5
6 150 significantly lower ($P<0.05$). However, Rao *et al.* (2003), Chandramoni *et al.*
7
8 151 (1998), and Keshavarz and Nakajima (1993) did not find any indications of calcium
9
10 152 contents greater than 32.5 g/kg (max. 55.0 Ca g/kg) of the egg weight. The midnight
11
12 153 feeding significantly increased the weight of the eggs ($P<0.05$) except in the 50:50
13
14 154 LF:LG group (Table 4). The midnight feeding did not have a significant effect on the
15
16 155 eggshell quality. The eggshell weight ($P<0.05$), thickness ($P<0.001$) and strength
17
18 156 ($P=0.001$) were significantly higher in the group with oyster shell and 29:71 LF:LG
19
20 157 in comparison with the diets containing eggshell and 50:50 LF:LG. The same
21
22 158 improvement of eggshell strength was reported by Richter *et al.* (1999) when the
23
24 159 hens were given limestone with a particle size of 0.5 - 2.0 mm, or a mixture of one-
25
26 160 third finely ground limestone and two-thirds oyster shell. The beneficial effects of
27
28 161 oyster shell on eggshell quality were consistent with the reports of Keshavarz and
29
30 162 Nakajima (1993). Although also in the group 32:68 LF:eggshell the greater the
31
32 163 amount of large particles the worse was the quality of the eggshell (eggshell weight,
33
34 164 eggshell weight ratio and eggshell thickness). This may have been due to the low
35
36 165 solubility of the eggshells. The *in vitro* solubility was as follows: fine limestone
37
38 166 85.0%, eggshells 14.0%, oyster shell 44.0% and large limestone 49.5%. Cheng and
39
40 167 Coon (1990) reported that the eggshell quality (and bone status) were more closely
41
42 168 related with limestone *in vitro* solubility than particle size. The researchers indicated
43
44 169 a potential difference in calcium retention for layers when two calcium sources of the
45
46 170 same particle size with different *in vitro* solubility were compared. On the other
47
48 171 hand, Rao and Roland (1990) reported that *in vivo* calcium solubilisation in laying
49
50 172 hens, for the particle size tested, was not influenced by *in vitro* limestone solubility.
51
52
53
54
55
56
57
58
59
60

The source and the particle size of the calcium probably have a greater effect on eggshell quality than does the calcium level. Although in the 29:71 LF:LG diet the content of calcium was about 12% higher than in the 32:68 LF:oyster shell diet, the quality of the eggshell was not improved. There was no significant difference in eggshell strength, thickness, weight ratio or eggshell weight between these groups. Neither Chandramoni *et al.* (1998) who used calcium concentrations of 32.5, 36.0 and 39.0 g/kg nor Rao *et al.* (2003), who used 32.5, 35.0, 37.5, 40.0, 42.5 and 45.0 g/kg, observed any improvement in the eggshell quality (eggshell weight, eggshell weight per unit surface area and eggshell weight, and eggshell thickness respectively). However, Chandramoni *et al.* (1998) used limestone powder and bone meal. Rao *et al.* (2003) used oyster shell (powder and grit 1 - 2 mm) but the weights of the eggs in their experiments were almost 10 g lighter than the eggs in this experiment were, and consequently the requirement of calcium was lower.

Contrary to Clunies *et al.* (1992), there was no significant difference in the total retention of calcium between the groups, which allowed higher and lower eggshell quality (thickness). Also Keshavarz and Nakajima (1993) did not find any significant difference in the total retention of calcium among concentrations of 35.0 - 55.0 g/kg in the diets. Probably the ability to utilise the calcium from the retained calcium for eggshell formation plays an important role. The utilisation of calcium for the formation of the eggshell was calculated and shown in Table 3. It was calculated on the basis of calcium retention, eggshell production (Figure 1) and the calcium content of the eggshells. The daily intake of calcium and the content of calcium in the eggshells are shown in Table 3. In the group of hens given oyster shells, 96.5 % of the retained calcium was deposited in the eggshells. In the groups given eggshells and 50:50 LF:LG the utilisation was only 73.9 and 78.6 % respectively, although the

coefficients of calcium retention were significantly higher ($P<0.05$) in these groups. There was no difference in the daily eggshell production (g) among the groups (Figure 1). This was probably due to the fact that the hens digest the calcium from the large particle oyster shell (2 - 5 mm) slowly but constantly, and it is utilised immediately for eggshell formation.

Figure 1 near here

Conclusions

Two thirds of the calcium source should be fed in the form of large particles (limestone grit or oyster shell) in the last third of the laying period to ensure good eggshell quality. Leeson *et al.* (1993) indicated that 3.4 g calcium/d is enough for brown-egg layers because they did not observe any effect of higher levels of calcium on eggshell deformation in these hens, but they used only limestone as a source of calcium in all of the rations. On the basis of the present study, eggshell quality can be improved by using suitable sources and particle sizes of calcium. Higher retention or concentration of calcium does not, in itself, mean better eggshell quality.

ACKNOWLEDGEMENTS

The author thanks Judy McPherson for the language correction and Assistant Professor Stanislav Kracmar, Dr.Sc. for the measurements of energy in the samples. The results were obtained with financial support of project 42/2005 IGA of Mendel University of Agriculture and Forestry in Brno.

REFERENCES

AHMAD, H.A. & BALANDER, R.J. (2003) Alternative Feeding Regimen of Calcium Source and Phosphorus Level for Better Eggshell Quality in Commercial Layers. *Journal of Applied Poultry Research* **12**: 509-514.

BAIN, M.M. (1997) A reinterpretation of eggshell strength. In SOLOMON, S.E. *Egg and Eggshell Quality*. London: Manson Publishing Ltd. 131-142.

- 223 CHANDRAMONI, JADHAO, S.B. & SINHA, R.P. (1998) Effect of Dietary
224 Calcium and Phosphorus Concentrations on Retention of these Nutrients by
225 Caged Layers. *British Poultry Science* **39**: 544-548.
- 226 CHENG, T.K. & COON, C.N. (1990) Effect of Ca Source, Particle Size, Limestone
227 Solubility *in vitro*, and Ca Intake Level on Layer Bone Status and Performance.
228 *Poultry Science* **69**: 2228-2230.
- 229 CLUNIES, M., PARKS, D. & LEESON, S. (1992) Calcium and Phosphorus
230 Metabolism and Eggshell Thickness in Laying Hens Producing Thick or Thin
231 Shells. *Poultry Science* **71**: 490-498.
- 232 GUINOTTE, F. & NYS, Y. (1991) Effects of Particle Size and Origin of Calcium
233 Sources on Eggshell Quality and Bone Mineralization in Egg Laying Hens.
234 *Poultry Science* **70**: 583-592.
- 235 KESHA VARZ, K. & NAKAJIMA, S. (1993) Re-Evaluation of Calcium and
236 Phosphorus REquirements of Laying Hens for Optimum Performance and
237 Eggshell Quality. *Poultry Science* **72**: 144-153.
- 238 KORELESKI, J. & SWIATKIEWICZ, S. (2004) Calcium from Limestone Meal and
239 Grit in Laying Hen Diets – Effect on Performance, Eggshell and Bone Quality.
240 *Journal of Animal and Feed Sciences* **13**: 635-645.
- 241 LEESON, S., SUMMERS, D. & CASTON, L. (1993) Response of Brown-Egg
242 Strain Laywrs to Dietary Calcium or Phosphorus. *Poultry Science* **72**: 1510-1514.
- 243 PAVLOVSKI, Z., VITOROVIC, D., LUKIC, M. & SPASOJEVIC, I. (2003)
244 Improving Eggshell Quality by Replacement of Pulverised Limestone by
245 Granular Limestone in the Hen Diet. *Acta Veterinaria-Bograd* **53**: 35-40.
- 246 RAO, K.S. & ROLAND, D.A. (1990) *In Vivo* Limestone Solubilization in
247 Commercial Leghorns – Role of Dietary Calcium Level, Limestone Particle Size,

248 *In Vitro* Limestone solubility Rate, and the Calcium Status of the Hen. *Poultry*
249 *Science* **69**: 2170-2176.

250 RAO, K.S., ROLAND, D.A., ADAMS, J.L. & DURBORAW, W.M. (1992)
251 Improved Limestone Retention in the Gizzard of Commercial Leghorn Hens.
252 *Journal of Applied Poultry Research* **1**: 6-10.

253 RAO, S.V.R., PNADA, A.K., RAJU, M.V.L.N., SUNDER, G.S. & PRAHARAJ,
254 N.K. (2003) Requirement of Calcium for Commercial Broilers and White
255 Leghorn Layers at Low Dietary Phosphorus Levels. *Animal Feed Science and*
256 *Technology* **106**: 199-208.

257 RICHTER, G., KIESSLING, G., OCHRIMENKO, W.I. & LUDKE, H. (1999)
258 Influence of Particle Size and Calcium Source on Limestone Solubility *in vitro*,
259 Performance and Eggshell Quality in Laying Hens. *Archiv für Geflügelkunde* **63**:
260 208-213.

261 SCHEIDELER, S.E. (1998) Eggshell calcium effects on egg quality and Ca
262 digestibility in first- or third-cycle laying hens. *Journal of Applied Poultry*
263 *Research* **7**: 69-74.

264 SCHEIDELER, S.E. (2004) Calcium Source, Particle Size and Calcium Level for
265 Laying Hens. In *Proceedings of the Midwest Poultry Federation Convention*. St.
266 Paul, MN, MN: Midwest Poultry Federation.

267 ZANG, B. & COON C.N. (1997) The Relationship of Calcium Intake, Source, Size,
268 Solubility *In Vitro* and *In Vivo*, and Gizzard Limestone Retention in Laying
269 Hens. *Poultry Science* **76**: 1702-1706.

Table 1. *The composition of the diets*

Composition (g/kg)	29:71 LF:LG	32:68 LF:eggshell	32:68 LF:oyster shell	50:50 LF:LG
Maize		407		
Wheat		340		
Soybean meal		138		
Monocalcium phosphate		9.5		
Methionine		1.25		
L-lysine		3.0		
Salt		3.0		
Premix		5.0		
Limestone Fine	30	30	30	46.65
Lime Grit	73.3*	-	-	46.65
Eggshells	-	63.3	-	-
Oyster shells	-	-	63.3	-
Calcium (g/kg)	44.6	39.7	39.8	40.8
Total phosphorus (g/kg)	5.7	5.2	5.4	5.2

* The content of wheat was 330 g/kg in this diet.

Table 2. The coefficients of nutrient retention

	29:71 LF:LG	32:68 LF:eggshell	32:68 LF:oyster shell	50:50 LF:LG	P-values
	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE	
Retention of nutrients					
Dry matter	0.714 ± 0.0139 ^a	0.784 ± 0.0136 ^b	0.748 ± 0.0173	0.751 ± 0.0121	<0.05
Energy	0.758 ± 0.0135 ^a	0.818 ± 0.0136 ^b	0.786 ± 0.0168	0.786 ± 0.0123	<0.05
Nitrogen	0.367 ± 0.0353 ^a	0.541 ± 0.0290 ^b	0.465 ± 0.0441	0.455 ± 0.0290	<0.05
Ash	0.488 ± 0.0134 ^b	0.577 ± 0.0134 ^a	0.518 ± 0.0165 ^b	0.515 ± 0.0131 ^b	<0.001
Calcium - Ca _c	0.506 ± 0.0142 ^a	0.576 ± 0.0177 ^b	0.515 ± 0.0334 ^a	0.578 ± 0.0123 ^b	<0.05
Phosphorus	0.219 ± 0.0464 ^a	0.415 ± 0.0420 ^b	0.353 ± 0.0388 ^b	0.293 ± 0.0484	<0.05
Fat	0.683 ± 0.0247	0.735 ± 0.0231	0.712 ± 0.0271	0.677 ± 0.0217	NS

SE - standard error.

NS - non-significant.

^{a,b,c}Values in a row not sharing a common superscript are significantly different at P<0.05.

Table 3. *Eggshell quality and calcium requirements*

	29:71 LF:LG	32:68 LF:Eggshell	32:68 LF:Oyster shell	50:50 LF:LG	<i>P</i> -values
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	
Egg production (eggs)	52	49	46	53	-
Food intake (g/d)	109.3	108.5	104.9	109.3	-
Ca intake - Ca_i (g/d)	4.9 $\pm$ 0.24	4.3 $\pm$ 0.20	4.2 $\pm$ 0.30	4.5 $\pm$ 0.13	NS
Weight of hens (kg)	1.63 $\pm$ 0.074	1.66 $\pm$ 0.057	1.76 $\pm$ 0.075	1.65 $\pm$ 0.091	NS
Egg weight (g)	61.7 $\pm$ 0.45 ^a	63.7 $\pm$ 0.66 ^b	64.0 $\pm$ 0.59 ^b	63.9 $\pm$ 0.78 ^b	<0.05
Eggshell weight (g)	6.63 $\pm$ 0.058 ^b	6.24 $\pm$ 0.086 ^a	6.69 $\pm$ 0.110 ^b	6.40 $\pm$ 0.130	<0.01
Eggshell weight ratio (%)	10.8 $\pm$ 0.087 ^b	9.81 $\pm$ 0.108 ^{ac}	10.4 $\pm$ 0.126 ^{bc}	10.0 $\pm$ 0.154 ^c	<0.001
Eggshell thickness (mm)	0.432 $\pm$ 0.0034 ^b	0.397 $\pm$ 0.0045 ^a	0.428 $\pm$ 0.0050 ^b	0.406 $\pm$ 0.0067 ^a	<0.001
Eggshell strength (N)	38.1 $\pm$ 1.01 ^b	35.7 $\pm$ 0.97	39.1 $\pm$ 1.14 ^b	32.6 $\pm$ 1.65 ^a	<0.01
Ca in eggshell - Ca_e (g/kg)	359.7 $\pm$ 1.75	359.1 $\pm$ 1.30	359.8 $\pm$ 1.26	359.4 $\pm$ 1.26	NS
Ca eggshell : Ca retention - Ca_p (%)	84.1 $\pm$ 4.18	73.9 $\pm$ 8.04	96.5 $\pm$ 10.0	78.6 $\pm$ 3.97	NS

SE - standard error.

NS - non-significant.

^{a,b,c}Values in a row not sharing a common superscript are significantly different at *P*<0.05.

Table 4. *The effect of midnight feeding on the coefficients of calcium retention and egg weight*

	29:71 LF:LG	32:68 LF:Eggshell	32:68 LF:Oyster shell	50:50 LF:LG
	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE
Retention of calcium				
Daily feeding	0.471 ± 0.0161	0.525 ± 0.0163	0.418 ± 0.0324	0.564 ± 0.0136
Midnight feeding	0.540 ± 0.0124	0.628 ± 0.0061	0.613 ± 0.0747	0.591 ± 0.0203
<i>P</i> -values	<0.01	<0.001	<0.001	NS
Weight of eggs				
Daily feeding	60.8 ± 0.62	61.7 ± 0.81	62.3 ± 0.77	62.8 ± 1.02
Midnight feeding	62.6 ± 0.60	65.4 ± 0.83	65.8 ± 0.82	64.5 ± 1.12
<i>P</i> -values	<0.05	<0.01	<0.01	NS

SE - standard error.
NS - non-significant.