Characterisation of European varieties of triticale with special emphasis on the ability of plant phytase to improve phytate P availability to chickens

Yves Nys

► To cite this version:

Yves Nys. Characterisation of European varieties of triticale with special emphasis on the ability of plant phytase to improve phytate P availability to chickens. *British Poultry Science*, 2007, 48 (06), pp.678-689. 10.1080/00071660701691292 . hal-00545322

HAL Id: hal-00545322

<https://hal.science/hal-00545322>

Submitted on 10 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Characterisation of European varieties of triticale with special emphasis on the ability of plant phytase to improve phytate P availability to chickens

Journal:	<i>British Poultry Science</i>
Manuscript ID:	CBPS-2007-148
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	15-Jun-2007
Complete List of Authors:	Nys, Yves; INRA, Recherches Avicoles
Keywords:	chicken, triticale, phytase, phosphorus, viscosity

Characterisation of European varieties of triticale with special emphasis on the ability of plant phytase to improve phytate phosphorus availability to chickens

C. JONDREVILLE, C. GENTHON¹, A. BOUGUENNEC², B. CARRE¹, Y. NYS^{1*}

Institut National de la Recherche Agronomique INRA, UMR1079 Systèmes d'élevage Nutrition Animale et Humaine, Agrocampus, F-35590 Saint-Gilles, ¹INRA, UR83 Recherches Avicoles, F-37380 Nouzilly, ²INRA UMR1095 Amélioration et Santé des Plantes, Domaine de Crouelle, UBP, F-63000 Clermont-Ferrand, France

RUNNING TITLE: TRITICALE FOR CHICKENS

To whom correspondence should be addressed.
Fax: +33 (0) 247427778
Tel: +33 (0) 247427700
E-mail: nys@tours.inra.fr

Accepted for publication 14th June 2007

Abstract 1. A total of 30 varieties and selection lines of triticale grown under similar conditions were characterised. Thousand grain weight, specific weight, Hagberg falling number and N were 50.2 ± 5.0 g, 72.4 ± 2.1 kg/hl, 96 ± 48 s and 16.1 ± 0.11 g/kg, respectively.

2. Mean phosphorus (P) concentration was 2.86 ± 0.31 g/kg, of which 77% was of phytic origin. Mean phytase activity was 1018 ± 319 PU/kg. A genotypic effect on phytase activity was detected amongst 5 varieties studied out of 30. Potential and real applied viscosities were positively correlated and mean values were 3.53 ± 0.66 and 2.15 ± 0.31 ml/g, respectively.

3. The efficacy of plant phytase in improving P availability was assessed in chickens up to 3 weeks of age. Growth performance and bone ash concentration were compared in birds given either a maize (450 g/kg) and soybean meal (230 g/kg) phosphorus deficient diet containing 3.5 g P/kg, this basal diet supplemented with 1 or 2 g P/kg as monocalcium phosphate (MCP) or triticale (450 g/kg) and soybean meal (230 g/kg) diets containing 3.2 to 3.8 g P/kg with no MCP. To achieve graded levels of phytase activity, 4 varieties of triticale, intact or in which phytase was denaturated by heat treatment, were used. Estimated metabolisable energy, protein, amino acids and calcium concentrations were similar in all diets.

4. Phytase activity in the triticale-based diets ranged between 135 and 1390 PU/kg. Growth performance and bone ash were responsive to plant phytase and to MCP. Non-linear models of these responses were adjusted with the best fit for bone ash parameters. 250, 500 and 1000 PU of plant phytase were estimated to be equivalent to 0.46, 0.67 and 0.81 g P as MCP, respectively.

INTRODUCTION

Triticale is a hybrid of wheat and rye that has been proposed as an alternative cereal in animal feeding because of its potential combination of wheat feeding characteristics and rye winter

1
2
3 1 hardiness and disease resistance (Gatel *et al.*, 1985; Vieira *et al.*, 1995). Several grain
4
5 2 constituents play a role in optimum utilisation of this cereal by poultry. Among them, both
6
7 3 soluble non-starch-polysaccharides (NSP), mainly arabinoxylans present in the albumen, and
8
9 4 available phosphorus (P) are of great impact with regard to the nutritive value of the cereal
10
11 5 and to the alimentary strategies to control environmental pollution problems (Carré *et al.*,
12
13 6 1994; Barrier-Guillot *et al.*, 1996b) and are worth consideration in varietal selection
14
15 7 programmes. Improvements in nutritional value of triticale for broilers will increase the
16
17 8 economical interest in replacing wheat by triticale in broilers (Korver *et al.*, 2004). Protein
18
19 9 concentration and its amino acid profile as well as metabolisable energy are key contributors to
20
21 10 feeding value. High contents of arabinoxylans should be avoided, because this might increase
22
23 11 feed viscosity, reduce digestibility of various components and induce over-consumption of
24
25 12 water by birds, which results in aqueous excreta (Carré *et al.*, 2002). In contrast, high P
26
27 13 availability reduces the requirement for supplementing diets with mineral P leading to lower
28
29 14 excretion of this element by the animals.

30
31
32
33
34
35
36 15 The effect of arabinoxylans on viscosity can be assessed by means of real applied
37
38 16 viscosity (RAV) and potential applied viscosity (PAV) which refer to viscosity measured
39
40 17 when endogenous xylanases are allowed to act and are inactivated, respectively (Carré *et al.*,
41
42 18 1994; Carré, 2002). In wheat, these parameters are factors of variation in metabolisable
43
44 19 energy (Carré *et al.*, 2002). Bouguennec *et al.* (2000) reported PAV values from 1.6 to 5.1 for
45
46 20 49 varieties of triticale from the official French catalogue and highlighted that PAV is mainly
47
48 21 under genetic control, even if it may also be affected by environmental conditions. Similarly
49
50 22 in wheat, RAV and PAV are dependent on the genotype, although RAV, which depends on an
51
52 23 enzymatic activity, is affected by an environment x genotype interaction (Oury *et al.*, 1998).

53
54
55
56
57 24 In wheat, variability in P availability for poultry is mainly related to phytase activity
58
59 25 but not to phytic P concentration (Barrier-Guillot *et al.*, 1996b). Moreover, phytase activity

1 displays a significant genotypic effect in wheat, although it is also influenced by a genotype x
2 environment interaction (Barrier-Guillot *et al.*, 1996a; Oury *et al.*, 1998). Triticale has been
3 reported to display a phytase activity intermediate between those of wheat and rye (770, 460
4 and 5350 phytase units (PU)/kg, respectively according to INRA-AFZ, 2004) but the
5 available studies (Eeckhout and De Paepe, 1994; Nys *et al.*, 1996; Skiba *et al.*, 2004) do not
6 provide information about the actual genotypic variability. Besides, the effectiveness of plant
7 phytase to improve P availability in chickens has mainly been studied in wheat (Barrier-
8 Guillot *et al.*, 1996b; Potkansky, 2000; Paik, 2003; Juanpere *et al.*, 2004), with little
9 information available for triticale. Almost no information about the potential sparing effect of
10 triticale on the need for mineral P supplementation is available.

11 The aim of the present study was to evaluate the genotypic variability, in triticale, of
12 several characteristics including viscosity and phytase activity. In addition, the efficacy of
13 plant phytase contained in triticale to reduce the required mineral P supplementation in diets
14 for chickens was evaluated.

15 MATERIALS AND METHODS

16 **Triticale varieties and lines**

17 A total of 30 varieties and selection lines of triticale originating from different areas in Europe
18 and covering a wide range of genotypic variability were selected. They were grown at the
19 INRA research station in Clermont-Ferrand under a single crop management system adapted
20 to the local conditions. The 30 batches were each collected from a single plot (Table 1). In
21 addition, 5e varieties (DI34-2, Trimaran, Aubrac, Capitale and Calao) were chosen for the
22 wide range in phytase activity they displayed. They were each grown on 4 - 6 plots and were
23 used to assess the genotypic effect on phytase activity. Four out of these 5 latter varieties
24 (DI34-2, Trimaran, Aubrac and Calao) were used to assess response of chickens to graded

1 levels of plant phytase. For this latter study, a mixture in equal proportions of the 4 to 6
2 available sub-batches of each variety was constituted.

Table 1 near here

3 **Broiler experiment**

4 The experiment was conducted under the animal research guidelines of the French Ministry of
5 Agriculture. From hatching till 5 d of age, 220 male Ross white chicks were placed in plastic-
6 coated cages of two or three birds each and fed on a standard diet covering all nutrient
7 requirements (INRA, 1989). At d 5, chicks were individually weighed after an overnight fast
8 and the 192 chicks closest to the mean weight of 75.7 ± 1.78 g were placed in 96 plastic-
9 coated cages, with two birds of similar initial live weight in each. For the subsequent 16-d
10 experiment, chicks were fed on one of 12 experimental diets (8 cages per diet). The initial
11 room temperature of 32°C was gradually decreased to 30 and 28°C when the chicks
12 reached the ages of 2 and 14 d, respectively. During the first 2 d, birds were kept under 24 h
13 light a d and 23 h light per d thereafter. Birds had free access to water throughout the
14 experiment. After a 12 to 14-h fast, each chicken was weighed and killed and its right
15 tibiotarsus was collected. Feed consumption was recorded per cage for the 16-d experimental
16 period. Feed conversion ratio (FCR) was calculated as the ratio of feed intake to weight gain
17 over the 16-d experimental period.

18 Twelve diets were formulated to meet all nutrient requirements of the birds (INRA,
19 1989), except available P. Feedstuffs were ground in a hammer mill fitted with a 2.5 mm
20 screen prior to incorporation in the diet. Mixed diets were pelleted without steam addition.
21 Damage to plant phytase was not expected, as outlet temperature did not exceed 50°C. The
22 three first diets, with a phytase activity below the detection limit of 50 PU/kg, were based on
23 maize (450 g/kg) and soybean meal (230 g/kg) and contained graded levels of monocalcium
24 phosphate (MCP, 0, 1 and 2 g/kg in diets 1, 2 and 3, respectively) (Table 2). The basal diet
25 without any inorganic phosphorus contained 3,5 g P/kg and was deficient in non-phytic

phosphorus compared to the requirement for growing chickens. Diets 4 to 12, did not contain any maize but 450 g/kg of triticale. No MCP was added. In order to achieve graded levels of phytase activity, four different genotypes of triticale (DI34-2, Trimaran, Aubrac and Calao), raw or heat-treated, were introduced in the diets 4 to 7 and 8 to 11, respectively. A mixture of raw (200 g/kg) and heat-treated Calao (250 g/kg) was introduced in diet 12. Heat treatment, which aimed at reducing phytase activity, consisted in two successive heatings in a microwave oven at 600 W for 2 minutes. Measured phytase activity in the raw triticale-based diets and in the heat-treated triticale-based diets was 620 to 1390 PU/kg and 135 to 180 PU/kg, respectively. It was 645 PU/kg in the diet containing the mixture of raw and heated Calao. Maize starch, wheat straw, soy isolate, vegetable oil, calcium carbonate and DL-methionine were used to balance the diets so that estimates of metabolisable energy, protein, lysine, sulphur amino acids, phytic P and Ca concentrations were similar in maize-based diets and in triticale-based diets (INRA, 1989). Moreover, the varieties of triticale were chosen so that dietary RAV remained below 1.40 mL/g to ensure a limited effect of this parameter on dietary metabolisable energy (Maisonnier *et al.*, 2001). At the end, the 12 diets provided similar and adequate amounts of nutrients for chicks except available P (INRA, 1989).

Table 2
near here

Sampling and analyses

Analyses were performed in duplicate or in triplicate. Thousand grain weight (TGW) was assessed by passing 200 g of fresh grain through an automated Decca seed counter and adjusting weights to 15% moisture content. Specific weight (SW) was measured by means of a Nilema apparatus (Tripette et Renaud, France). The Hagberg falling number (HFN) was assessed by means of a Falling Number apparatus type 1400 (Falling Number AB, Sweden) after grinding with a Falling Number mill type KT120 (Falling Number AB, Sweden). Samples of triticale and diets were ground to pass through a 0.5 mm screen and stored at 4°C prior to the other analyses. N was determined by the Kjeldahl method according to the French

1 standard AFNOR (NFV 18-100) using a Kjelfoss apparatus (A/S N Foss Electric, Denmark).
2 Dry matter (DM) was measured by drying at 103°C until constant weight. P was analysed by
3 means of the vanadate colorimetric method according to the AFNOR method (NFV 18-106).
4 Phytic P was determined by ion-pair HPLC (Column C₁₈, Hypersyl C 18-5 µm 200 x 2 mm,
5 Interchim) after acidic extraction and anionic exchange purification according to the method
6 developed by Sandberg and Adherrine (1986) and modified by Lehrfeld (1989). Phytase
7 activity (PA) was measured colorimetrically after incubation in a sodium phytate solution
8 (Engelen *et al.*, 1994). One phytase unit (PU) is the amount of enzyme that liberates 1 µmol
9 per minute of inorganic P from 5.1 mmol/l solution of sodium phytate, at pH 5.5 and 37°C.
10 Prior to viscosity determination, extraction (pH = 4.5, temperature = 19-23°C) was performed
11 with or without pre-treatment in hot ethanol: water (80: 20). RAV and PAV refer to viscosity
12 measured when endogenous xylanases were allowed to act and were inactivated by treatment
13 of the sample, respectively. The viscosity data were divided by the viscosity of the buffer,
14 which gave relative viscosities (Vr), transformed into natural logarithm and then divided by
15 the concentration (g/mL) of the plant material in the buffer extraction volume. The results
16 “(Ln(Vr))/(g/ml)” were expressed as ml /g (Carré *et al.*, 1994; Carré, 2002).

17 Tibiotarsi were pooled per cage. They were cleaned of all soft tissues, defatted (24
18 hours in ether), dried (105°C for 12 hours) and weighed. Thereafter, they were ashed at 550°C
19 for 14 h in a muffle furnace and weighed. Ash concentration was calculated on a fat-free dry
20 matter (FF DM) basis.

21 **Statistical analyses**

22 Statistical analysis of data was performed by means of the GLM procedure of the Statistical
23 Analysis Systems software package version 8.1 (SAS, 1990). Correlation coefficients (*r*)
24 between physical and analytical characteristics of the 30 batches of triticale were calculated.
25 The effect of the variety on phytase activity was assessed by an analysis of variance using the

plot as the experimental unit, followed by a comparison of means. The chickens' responses were analysed using the cage as the experimental unit. Data were first submitted to an analysis of variance with the diet as main factor, followed by a comparison of means. Polynomial regression of the indicators of performance and of bone mineralisation was used to determine the presence of linear and quadratic effects of mineral P added in the diet and of dietary phytase activity. Effects were considered significant when $P < 0.05$.

Non-linear (NLIN procedure of SAS) functions were fitted to the response of performance and bone FF DM and ash to dietary P as MCP and phytase activity, using treatment means. Exponential models were chosen to describe the response to phytase and linear or exponential models were chosen to describe the response to mineral P, according to the results obtained by polynomial regression. The exponential model was chosen because it was extensively used to describe the effect of microbial phytase on P availability (Kornegay *et al.*, 1996; Kornegay, 2001). The model was $Y = a + b(1 - e^{-l \text{MinP}}) + c(1 - e^{-k \text{Phyt}})$ or $Y = a + b \text{MinP} + c(1 - e^{-k \text{Phyt}})$ with Y = response measurement, MinP = P added as MCP (g/kg diet), Phyt = phytase activity (PU/kg diet).

The coefficient of determination (R^2) of each equation generated was calculated as the square of the correlation coefficient between predicted and observed individual values. An equivalency value of mineral P as MCP (g) for plant phytase (PU) was calculated by setting equal the terms corresponding to mineral P and phytase activity.

RESULTS AND DISCUSSION

Physical and chemical characteristics of the 30 batches of triticale

Physical and chemical characteristics of the 30 batches of triticale are presented in Table 1. Phytase activity and especially HFN displayed the highest degree of variation, with coefficients of variation (CV) of 31 and 49%, respectively. PAV and RAV exhibited a CV of 15-20%. Phytic P displayed a CV of similar order, although with only 11 measurements

1 performed. The other parameters (TGW, N and total P) had lower amplitude of variation (CV
2 7-10%). This amplitude of variation represents genotypic variability since all the varieties
3 were grown under similar conditions.
4 Total P concentration in the 30 batches of triticale ranged between 2.06 and 3.57 g/kg with a
5 mean of 2.86 ± 0.31 g/kg. This average P concentration was slightly below the values of $3.7 \pm$
6 0.2 and 3.9 ± 0.1 g P/kg previously observed by Eeckhout and De Paepe (1994) and Skiba *et*
7 *al.* (2004) when analysing 6 and 4 varieties of triticale, respectively. Phytic P was measured
8 in only 11 out of the 30 batches and ranged between 1.50 and 2.62 g/kg, with a mean of 2.18
9 ± 0.40 g/kg. These values compare with the 2.5 ± 0.2 and 2.4 ± 0.2 g phytic P/kg reported in
10 the two aforementioned studies. As previously observed in wheat (Barrier-Guillot *et al.*,
11 1996a; Viveros *et al.*, 2000; Kim *et al.*, 2002), P and phytic P were positively correlated ($r =$
12 0.60 , $P < 0.05$) (Table 3). However, the low value of r hampers a reliable prediction of phytate
13 P from total P. In the current study, 77% of P was, on average, of phytic origin. This is quite
14 a high proportion compared with the 67 and 61% reported by Eeckhout and De Paepe (1994)
15 and Skiba *et al.* (2004), which may be a consequence of the lower total P observed in the
16 current study. P and phytic P concentrations in wheat are influenced by crop management,
17 especially N and P fertilisation (Barrier-Guillot *et al.*, 1996a; Oury *et al.*, 1998).
18 Consequently, the current results may have been influenced by the conditions under which
19 triticale was grown.

Table 3 near here

20 Phytase activity measured in the 30 batches of triticale ranged between 447 and 1843
21 PU/kg, with a mean value of 1018 ± 319 PU/kg. As expected, these values are intermediate
22 between the phytase activities reported in wheat and in rye (460 and 5350 PU/kg, according to
23 INRA-AFZ, 2004), although they are closer to those of wheat. These values are below the
24 1688 ± 227 and 1784 ± 386 PU/kg published by Eeckhout and De Paepe (1994) and Skiba *et*
25 *al.* (2004), respectively, but compare with the 1190 ± 52 PU/kg measured by Nys *et al.* (1996)

1 in 5 batches of triticale. These differences may originate from the huge inter-laboratory
2 variability in phytase activity determination mentioned by Tran and Skiba (2005), even when
3 similar analytical methodologies are implemented. Phytase activity determination might be
4 sensitive to the condition under which batches are stored prior to analysis (temperature,
5 duration of storage if the grain is ground) (Nys *et al.*, 1996) and to the way phytase is extracted
6 before analysis of the liberated inorganic phosphates (Greiner and Egli, 2003).

7 Phytase activity measured in each of the 5 varieties of triticale grown on 4 to 6 plots
8 (Table 4) was 20 to 48% higher than the value previously obtained on the single batch of the
9 same variety (Table 1). Nevertheless, the varieties were similarly ranked in the two sets of
10 data, with a coefficient of correlation of 0.95 ($P < 0.05$, $n = 5$). The origin of this systematic
11 difference is not clear but it may be ascribable to the interaction between the grinding of the
12 samples before analysis and the duration of their storage (Nys *et al.*, 1996). Phytase activity
13 differed ($P < 0.01$) between varieties. In Calao, Capitale, Aubrac and Trimaran, it exceeded by
14 112, 79, 41 and 30%, respectively, the value of 1012 PU/ kg in DI34-2. This genotypic effect
15 on phytase activity was previously reported in wheat (Barrier-Guillot *et al.*, 1996a; Kim *et al.*,
16 2002; Oury *et al.*, 1998; Tran and Skiba, 2005). However, Oury *et al.* (1998) detected a strong
17 genotype x environment interaction in the phytase activity of wheat. This interaction, which
18 may also exist in triticale, was not investigated in the current study. Phytase activity was
19 negatively correlated ($P < 0.05$) with SW and HFN, which is consistent with the fact that low
20 HFN (and possibly SW) might be linked with the stimulation of enzymatic activity in the
21 grain at germination as observed for amylase (Niziolek *et al.*, 1994). As previously reported in
22 wheat (Eeckhout et de Paepe, 1994; Barrier-Guillot *et al.*, 1996a) no relationship between
23 phytase activity and P or phytic P could be detected. In contrast, Viveros *et al.* (2000) found a
24 positive correlation between phytase activity and P in wheat; however, they were not able to
25 predict phytate P or phytase activity reliably from total P.

Table 4 near here

PAV and RAV ranged between 2.35 and 4.65 ml/g and 1.71 and 2.86 ml/g, respectively, with average values of 3.53 ± 0.66 and 2.15 ± 0.31 mL/g, respectively (Table 1). The PAV values are in the range of those reported by Bouguennec *et al.* (2000) in triticale. They are between those observed in rye (PAV = 28 mL/g, Carré *et al.*, 1994) and wheat (PAV = 2.9 mL/g, Oury *et al.*, 1998; Carré *et al.*, 2002), although they are closer to those of wheat. As previously reported (Oury *et al.*, 1998), PAV and RAV were positively correlated ($r = 0.43$, $P < 0.05$) with each other but not with any other characteristic.

Nitrogen concentration in the 30 varieties of triticale ranged between 14.1 and 18.1 g N/kg (Table 1), corresponding to a mean level of protein of 101 ± 8.8 g/kg with a range of 88 to 113 g/kg. This crude protein level is slightly lower than those observed in 8 Australian cultivars (range 101-135 g/kg; Johnson and Eason, 1988) or that analysed by Vieira *et al.* (1995) (129 g/kg) but is in agreement with levels observed by other authors: 72-110 g/kg (Zacarias *et al.*, 1982), 93 g/kg (Proodfoot and Hulan, 1988).

Triticale phosphorus availability in broilers

The microwave treatment applied to triticale was effective in reducing phytase activity, which was reduced after treatment by 85-90%, down to 142, 156, 126, 127 PU/kg in DI34-2, Trimaran, Aubrac and Calao, respectively. Phytase activity measured in diets exceeded the value expected from measurements performed on the batches of triticale, especially at low levels of phytase activity (Table 2). However, phytase activity measured in the batches of triticale and in the diets were highly correlated ($r = 0.99$, $P < 0.001$, $n = 8$). RAV in non-heated triticale-based diets was 10 to 23% lower than in heated triticale-based diets, suggesting that heat treatment reduced xylanase activity. However, it remained below 1.40 ml/g, even in heated triticale-based diets. In the current study, heat treatment was considered to have no effect on phytate concentration because phytates present in cereals are very stable to heating (Reddy *et al.*, 1989; Juanpere *et al.*, 2004).

Performance and bone characteristics of birds are presented in Table 5. Five birds fed the maize based diet without MCP (diet 1) died before the end of the experiment, explaining the removal of data from three cages. A high mortality level has previously been observed in chicks fed on maize based diets without supplemented mineral P (Kornegay *et al.*, 1996; Paik, 2003).

Table 5 near here

For all the variables studied, significant differences ($P < 0.001$) among diets were detected. Chicks given the maize-based diets without phosphate and supplemented with 2 g P as MCP displayed the lowest performance and the highest bone characteristics. The linear response of bone ash to graded level of inorganic phosphorus demonstrates that the chicks fed on negative control diet were deficient in phosphorus. Performance and bone characteristics in chickens given the raw triticales-based diets did not differ ($P > 0.05$) from those of chickens given the maize-based diet supplemented with 1 g P as MCP/kg, except for bone ash concentration in the Trimaran-based diet, which was lower. Overall, bone characteristics and final weight of chickens given the heated triticales-based diets were intermediate between those of chickens given the unsupplemented maize-based diet and the raw triticales-based diets. Except for the heated DI-34-2-based diet, feed intake in chicks given the heated triticales-based diets did not differ from that in chicks fed the unsupplemented maize-based diet, while their FCR did not differ from that in chicks fed the raw triticales-based diets. Chickens given the maize-based diet supplemented with 2 g P as MCP/kg ingested 2.1 times more feed than chickens given the maize-based diet without MCP. Feed intake increased linearly ($P < 0.001$) and quadratically ($P < 0.05$ and $P < 0.001$, respectively) with supplemental P supply and with the dietary level of phytase. By the end of the experiment, birds fed on the maize-based diet supplemented with 2 g P as MCP/kg were 2.2 times heavier than birds given the maize-based diet without added P. Final weight increased linearly ($P < 0.001$) and quadratically ($P < 0.01$ and $P < 0.001$, respectively) with P addition and with dietary phytase

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

activity (Figure). FCR decreased linearly and quadratically with added P ($P<0.01$ and $P<0.05$, respectively) and dietary phytase activity ($P<0.001$ and $P<0.01$, respectively). FCR was decreased by 30% by supplementing the maize-based diet with 1 g P as MCP, but no further improvement was achieved with 2 g P as MCP. Chickens given the heated triticale-based diets, still containing some phytase (135 to 200 PU), displayed a FCR 25% lower than chicks on the unsupplemented maize-based diet with no further improvement with raw triticale-based diets.

Improvements in weight gain, feed intake, and FCR with increased available P provision through P or phytase supplementation of low P diets have been previously reported (Broz *et al.*, 1994; Kornegay *et al.*, 1996; Sebastian *et al.*, 1996a and b; Paik, 2003). From a literature review including 298 observations from 18 literature references, Lescoat *et al.* (2005) established an exponential relationship between body weight gain of chickens slaughtered at 21 to 24 d and P intake. The two-fold increase in final weight with the addition of 2 g P as MCP in the maize-based diet fits well with the relationship established by these authors. Increased weight gain with supplemental P may have resulted not only from an increase in feed intake but also from a specific effect of P on growth. A specific effect of P is expected in the chick because of the important role of this element in the formation of the skeletal system and in body metabolism (*e.g.* nucleic acids, high-energy compounds and various enzymatic reactions), and because of the fast rate of chick growth and its low capacity of P storage (Kornegay *et al.*, 1996).

Bone FF DM followed a trend very similar to final weight ($r = 0.90$, $P<0.001$, $n = 90$) with an increase in bone FF DM by 2.9 times when 2 g of P were added to the maize based diet. Bone FF DM responded linearly to P added ($P<0.001$) and linearly ($P<0.001$) and quadratically ($P<0.01$) to phytase activity. The amplitude of the response of bone ash (g) was a 5.1-fold increase when 2 g of P were added to the maize based diet. Bone ash (g and relative

to FF DM) increased linearly with added P and with phytase activity ($P < 0.001$) and quadratically with phytase activity ($P < 0.001$). The linearity and the higher amplitude of the response of bone ash to dietary P compared to body weight gain indicates the ability of bone to incorporate P beyond the dietary supply needed to maximize weight gain (Figure). Relying on 316 observations collected in 15 literature references, Lescoat *et al.* (2005) established that bone ash concentration in chickens reached a plateau when 250-300 mg total P were ingested per d, a value that is higher than the 220 mg total P daily ingested by the chickens fed on the maize diet supplemented with 2 g P as MCP. Moreover, the increase in bone ash concentration fits well with the range of variation observed by these authors for similar dietary P supply. The improvement in bone ash in chickens fed on triticale diets indicates that plant phytase was effective in releasing P from the phytate-mineral complex. The maximum responses appeared to occur at around 600 PU plant phytase/kg diet whatever the indicator.

Figure
near here

Before equivalency values of plant phytase for P as MCP can be estimated, the question arises whether responses of performance and bone characteristics observed in the current study can be ascribed solely to the improvement in available P supply by means of P as MCP in maize-based diets and by means of increased phytase activity in triticale-based diets. In the control diet, maize was preferred to a mixture of heated triticale batches to avoid any side effect due to heating other than lowering phytase activity and because maize is the reference cereal used when evaluating any source of phosphorus in the literature. The results of the current study cannot be used to compare the relative efficiency of maize and triticale for chick performance. Nevertheless, maize- and triticale-based diets provided sufficient amounts of all nutrients to fulfill the requirement of growing chicks up to 3 weeks of age, except for P (INRA, 1989). Particularly, because all diets were balanced for Ca, there is a very high probability that the response of bone ash concentration was related to available P supply rather than to any other dietary parameter. The magnitude of variation in performance

in the current study is very large compared with the slight decrease in performance reported when maize is replaced by wheat (Singh *et al.*, 2003) or by triticale (Proudfoot and Hulan, 1988) in isocaloric, isonitrogenous, iso-Ca and iso-P diets fed to chickens. The former authors ascribe this depressed performance to the presence of arabinoxylans in wheat. In the current study, RAV in triticale-based diets was not greater than 1.40 ml/g, and, according to Maisonnier *et al.* (2001), such concentrations of RAV do not influence performance in chicks.

Non-linear models of the response of feed intake, final weight and bone characteristics to dietary mineral P and plant phytase activity as well as the derived equivalency values of P as MCP for plant phytase are presented in Table 6. Models were adjusted with a coefficient of determination of 0.77 to 0.89. All the indicators used were sensitive to plant phytase or MCP supply, but the fit was better for bone ash than for the other parameters. Moreover, the actual weight of bone ash displayed the highest R^2 for both the non-linear and polynomial adjustments. Other authors have concluded that bone ash weight is the most sensitive indicator to assess phytase efficacy in chickens (Zhang *et al.*, 2000; Hall *et al.*, 2003). Random variability in the organic matrix left after water and lipid removal may be responsible for the decreased R^2 value for ash expressed on a fat-free dry matter basis (Hall *et al.*, 2003). Based on bone ash parameters, increase in plant phytase activity from 0 to 250 PU was estimated to be equivalent to 0.46-0.47 g P as MCP. Between 250 and 500 PU, 250 PU were estimated to allow the release of an amount of P equivalent to 0.20-0.22 g of P as MCP. Between 500 and 1000 PU, an equivalency of 0.12-0.15 g P as MCP for 500 PU was obtained.

Table 6 near here

The equivalency values are not easy to compare with literature data because the efficacy of phytase has been shown to depend on several parameters including the dietary non-phytic P and Ca concentrations. These equivalency values may be relatively high because

the triticale-based diets were not supplemented with P. Kornegay *et al.* (1996) demonstrated that, probably because of an available P supply closer to the requirement, the equivalency values decrease when the dietary non-phytate P concentration increases. On the contrary, the wide Ca:P ratio in the current study, greater than 2.8:1, may have reduced the efficacy of phytase in releasing phytate P. Increasing the Ca:P ratio in poultry diets was reported to decrease the efficacy of microbial phytase (Schoner *et al.*, 1993; Qian *et al.*, 1996; Sebastian *et al.*, 1996b), with the optimum responses to phytase obtained at a Ca:P ratio as low as 1.1:1.

In accordance with our results, Paik (2003) observed that 650 PU of plant phytase achieved by the addition of wheat and wheat bran in a maize-soybean meal-based diet were equivalent to at least 1 g P as tricalcium phosphate in terms of growth performance and mineral retention in chickens up to 35 d. Similarly, an equivalency of 0.65-1 g P as MCP for 500 PU was reported by Frapin (1996) when evaluating plant phytase efficiency from various wheat varieties. At variance with our results, Juanpere *et al.* (2004) did not observe any effect of around 150 units of phytase from barley on performance and toe ash concentration in chicks. This absence of efficacy of plant phytase may be ascribable to a higher non-phytate P concentration than in the current study. From literature data, Kornegay *et al.* (2001) estimated that 500 PU of microbial phytase (3-phytase) were equivalent to 0.8 g P as MCP for 500 PU in terms of mineral retention in broilers. This higher figure may originate from the lower efficacy of plant phytase compared to microbial 3-phytase demonstrated in broilers (Frapin, 1996; Potkansky, 2000) and in pigs (Zimmermann *et al.*, 2002). The activity of plant phytase decreases rapidly when the pH decreases below the optimum pH of 5.0-5.5, whereas the 3-phytase still displays an activity of 60% of the optimum at pH 2 (Eeckhout and De Paepe, 1996). Plant phytase would be 40 to 80% less efficient in releasing P from phytates *in vivo* than 3-phytase due to the low pH in the stomach of pigs and gizzard of poultry (Frapin, 1996; Potkansky, 2000; Zimmermann *et al.*, 2002). The acidic pH of the gizzard inactivates plant

1 phytase in contrast to protected microbial phytase and consequently plant phytase activity is
2 limited to the crop (Frapin, 1996). Moreover, plant phytase may be more sensitive to the
3 presence of pepsin than 3-phytase (Rapp *et al.*, 2001).

4 The present study confirmed the usefulness of the introduction of triticale in chickens'
5 diets to reduce P emission and to limit environmental pollution, 500 PU of plant phytase
6 being equivalent to 0.66-0.69 g P as MCP. Accounting for the genotypic variability of phytase
7 activity in triticale (447 to 1843 PU/kg), a diet containing 450 g/kg triticale may display a
8 phytase activity of 201 to 829 PU/kg, provided plant phytase is not denaturated during feed
9 processing. In such a diet, P supplementation as MCP may be reduced by 0.4 to 0.8 g/kg,
10 compared to a low phytase diet, such as a maize-based diet.

11 REFERENCES

12 BARRIER-GUILLOT, B., CASADO, P., MAUPETIT, P., JONDREVILLE, C. & GATEL, F.
13 (1996a) Wheat phosphorus availability: 1 - In vitro study; factors affecting endogenous
14 phytasic activity and phytic phosphorus content *Journal of the Science of Food and*
15 *Agriculture*, **70**: 62-68.

16 BARRIER-GUILLOT, B., CASADO, P., MAUPETIT, P., JONDREVILLE, C. & GATEL, F.
17 (1996b) Wheat P availability: 2 - In vivo study in broilers and pigs; relationship with
18 endogenous phytasic activity and phytic phosphorus content in wheat. *Journal of the*
19 *Science of Food and Agriculture*, **70**: 69-74.

20 BOUGUENNEC, A., OURY, F.X. & JESTIN, L. (2000) Viscosity related to arabinoxylans in
21 triticale: genetic and environmental variation in France. *Vorträge für Pflanzenzüchtung*,
22 **49**: 161-169.

23 BROZ, J., OLDALE, P., PERRINVOLTZ, A. H., RYCHEN, G., SCHULZE, J. & SIMOES
24 NUNES, C. (1994) Effects of supplemental phytase on performance and phosphorus

- 1 utilisation in broiler chickens fed a low phosphorus diet without addition of inorganic
2 phosphates. *British Poultry Science*, **35**: 273-280.
- 3 CARRÉ, B. (2002) Carbohydrate chemistry of the feedstuffs used for poultry, in: McNAB, B.
4 & BOORMAN, N. (Eds) *Poultry feedstuffs: supply, composition and nutritive value*, pp
5 39-56 (Wallingford, UK, CAB International Publishing).
- 6 CARRE, B., GOMEZ, J., MELCION, J.P. & GIBOULOT, B. (1994) La viscosité des
7 aliments destinés à l'aviculture. Utilisation pour prédire la consommation et l'excrétion
8 d'eau. *INRA Productions Animales*, **7**: 369-379.
- 9 CARRÉ, B., IDI, A., MAISONNIER, S., MELCION, J. P., OURY, F. X., GOMEZ, J. &
10 PLUCHARD, P. (2002) Relationships between digestibilities of food components and
11 characteristics of wheat (*Triticum aestivum*) introduced as the only cereal source in a
12 broiler chicken diet. *British Poultry Science* **43**: 404-415.
- 13 EECKHOUT, W. & DE PAEPE, M. (1994) Total phosphorus, phytate-phosphorus and
14 phytase activity in plant feedstuffs. *Animal Feed Science and Technology*, **47**: 19-29.
- 15 EECKHOUT, W. & DE PAEPE, M. (1996) In vitro and in vivo comparison of microbial and
16 plant phytase, in: COELHO, M.B. & KORNEGAY, E.T. (Eds) *Phytase in Animal*
17 *Nutrition and Waste Management*, pp 237-240 (Mount Olive, NJ, BASF corporation).
- 18 ENGELN, A.J., VAN DER HEEFT, F.C., RANDSDORP, P.H.G. & SMIT, E.L.C. (1994)
19 Simple and rapid determination of phytase activity. *Journal of AOAC International*, **77**:
20 760-764.
- 21 FRAPIN, D. (1996) Valorisation du phosphore phytique végétal chez l'oiseau : intérêt et
22 mode d'action des phytases végétales et microbiennes. *Thèse de l'école nationale*
23 *supérieure agronomique de Rennes*, 134 p.

- 1
- 2
- 3 1 GATEL, F., LAVOREL, O., FEKETE, J., GROSJEAN, F. & CASTAING, J. (1985) Feeding
- 4
- 5 2 value of triticale for monogastrics: weaned piglets, growing finishing pigs and broilers, in:
- 6
- 7 3 BERNARD, M. & BERNARD, S. (Eds) *Genetics and Breeding of Triticale*, pp 659-670
- 8
- 9 4 (Paris, France, INRA).
- 10
- 11
- 12
- 13 5 GREINER, R. & EGLI, I. (2003) Determination of the activity of acidic phytase-degrading
- 14
- 15 6 enzymes in cereal seeds. *Journal of Agricultural and Food Chemistry*, **51**: 847-850.
- 16
- 17
- 18 7 HALL, L. E., SHIRLEY, R. B., BAKALLI, R. I., AGGREY, S. E., PESTI, G. M. &
- 19
- 20 8 EDWARDS, H. M. (2003) Power of two methods for the estimation of bone ash of
- 21
- 22 9 broilers. *Poultry Science*, **82**:414-418.
- 23
- 24
- 25
- 26 10 INRA (1989) L'Alimentation des Animaux Monogastriques: Porc, Lapin, Volailles (Paris,
- 27
- 28 11 France, Institut National de la Recherche Agronomique).
- 29
- 30
- 31 12 INRA-AFZ (2004) Tables of composition and nutritional value of feed materials, in:
- 32
- 33 13 SAUVANT, D., PEREZ, J.M. & TRAN, G. (Eds) 304 pp (Paris, France, Institut National de
- 34
- 35 14 la Recherche Agronomique, Association Française de Zootechnie).
- 36
- 37
- 38
- 39 15 JOHNSON, R. & EASON, P. (1988) Evaluation of triticale for use in diets for meat-type
- 40
- 41 16 chickens. *Journal of the Science of Food and Agriculture*, **42**: 95-108.
- 42
- 43
- 44 17 JUANPERE, J., PEREZ-VENDRELL, A. M. & BRUFAU, J. (2004) Effect of microbial
- 45
- 46 18 phytase on broilers fed barley-based diets in the presence or not of endogenous phytase.
- 47
- 48 19 Animal Feed Science and Technology, **115**: 265-279.
- 49
- 50
- 51
- 52 20 KIM, J. C., MULLAN, B. P., SELLE, P. H. & PLUSKE, J. R. (2002) Levels of total
- 53
- 54 21 phosphorus, phytate-phosphorus, and phytase activity in three varieties of Western
- 55
- 56 22 Australian wheats in response to growing region, growing season, and storage. *Australian*
- 57
- 58 23 *Journal of Agricultural Research*, **53**: 1361-1366.
- 59
- 60

- 1 KORNEGAY, E. T. (2001) Digestion of phosphorus and other nutrients: the role of phytases
2 and factors influencing their activity, in: Bedford, M. R. & Partridge, G. G. (Eds)
3 *Enzymes in Farm Animal Nutrition*, pp 237-271 (Wallingford, CAB International).
- 4 KORNEGAY, E. T., DENBOW, D., Yi, Z. & RAVINDRAN, V. (1996) Response of broilers
5 to graded levels of microbial phytase added to maize-soyabean-meal-based diets
6 containing three levels of non-phytate phosphorus. *British Journal of Nutrition*, **75**:839-
7 852.
- 8 KORVER, D.R., ZUIDHOF, M.J. & LAWES, K.R. (2004) Performance characteristics and
9 economic comparison of broiler chickens fed wheat- and triticale-based diets. *Poultry
10 Science* **83**: 716-725.
- 11 LEHRFELD, J. (1989) High performance liquid chromatography analysis of phytic acid on a
12 pH-stable, macroporous polymer column. *Cereal Chemistry*, **66**: 510-515.
- 13 LESCOAT, P., TRAVEL, A. & NYS, Y. (2005) Lois de réponse des volailles de chair à
14 l'apport de phosphore. *INRA Productions Animales*, **18**: 193-201.
- 15 MAISONNIER, S., GOMEZ, J., CARRE, B. (2001) Nutrient digestibility and intestinal
16 viscosities in broiler chickens fed on wheat diets, as compared to maize diets with added
17 guar gum. *British Poultry Science*, **42**: 102-110.
- 18 NIZIOLEK, S., BARTOSZEWICZ, K. & KACZKOWSKI, J. (1994) Some hydrolases and
19 their substrates occurring in initial period of germination in triticale grains of various pre-
20 harvest sprouting resistance. *Acta Physiologiae Plantarum*, **16**: 171-176.
- 21 NYS, Y., FRAPIN, D. & POINTILLART, A. (1996) Occurrence of phytase in plants, animals
22 and microorganisms, in: COELHO, M.B. & KORNEGAY, E.T. (Eds) *Phytase in Animal
23 Nutrition and Waste Management*, pp 213-236 (Mount Olive, NJ, BASF corporation).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 OURY, F.X., CARRE, B., PLUCHARD, P., BERARD, P., NYS, Y. & LECLERCQ, B.
2 (1998) Genetic variability and stability of poultry feeding related characters in wheat, in
3 relation to environmental variation. *Agronomie*, **18**: 139-150.

4 PAIK, I. (2003) Application of phytase, microbial or plant origin, to reduce P excretion in
5 poultry production. *Asian-Australasian Journal of Animal Sciences*, **16**:124-135.

6 POTKANSKY, A. (2000) The comparison of plant and microbial phytases in the feeding, in:
7 Grela (Ed) *Proceedings of the International Symposium on Phytase in Animal Nutrition*,
8 pp 21-27 (Lublin).

9 PROUDFOOT, F.G. & HULAN, H.W. (1988) Nutritive value of triticale as a feed ingredient
10 for broiler chickens. *Poultry Science*, **67**: 1743-1749.

11 QIAN, H., KORNEGAY, E. T. & DENBOW, D. M. (1996) Phosphorus equivalence of
12 microbial phytase in turkey diets as influenced by calcium to phosphorus ratios and
13 phosphorus levels. *Poultry Science*, **75**: 69-81.

14 RAPP, C., LANTZSCH, H. J. & DROCHNER, W. (2001) Hydrolysis of phytic acid by
15 intrinsic plant or supplemented microbial phytase (*Aspergillus niger*) in the stomach and
16 small intestine of minipigs fitted with re-entrant cannulas. 2. Phytase activity. *Journal of*
17 *Animal Physiology and Animal Nutrition*, **85**: 414-419.

18 REDDY, N.K., PIERSON, M.D., SATHE, S.K. & SALUNKHE, D.K., 1989. Phytates in
19 Cereals and Legumes. CRC Press, Boca Raton.

20 SANDBERG, A.S. & ADHERINNE, R. (1986) HPLC method for determination of inositol
21 tri, tetra, penta and hexaphosphates in foods and intestinal contents. *Journal of Food*
22 *Science*, **51**: 547-550.

23 SAS, 1990. Statistical Analysis Systems software package version 8.1. Statistical Analysis
24 Systems Institute, Inc., Cary, NC, USA.

- SCHONER, F. J., HOPPE, P. P., SCHWARZ, G. & WIESCHE, H. (1993) Comparison of microbial phytase and inorganic-phosphate in male chickens - the influence on performance and mineral retention at various calcium levels. *Journal of Animal Physiology and Animal Nutrition*, **69**: 235-244.
- SEBASTIAN, S., TOUCHBURN, S., CHAVEZ, E. R. & LAGUE, P. C. (1996a) The effects of supplemental microbial phytase on the performance and utilization of dietary calcium, phosphorus, copper, and zinc in broiler chickens fed corn-soybean diets. *Poultry Science*, **75**: 729-736.
- SEBASTIAN, S., TOUCHBURN, S., CHAVEZ, E. R. & LAGUE, P. C. (1996b) Efficacy of supplemental microbial phytase at different dietary calcium levels on growth performance and mineral utilization of broiler chickens. *Poultry Science*, **75**: 1516-1523.
- SINGH, P. K., KHATTA, V. K., THAKUR, R. S., DEY, S. & SANGWAN, M. L. (2003) Effect of phytase supplementation on the performance of broiler chickens fed maize and wheat based diets with different levels of non-phytate phosphorus. *Asian-Australasian Journal of Animal Sciences*, **16**: 1642-1649.
- SKIBA, F., CALLU, P., CASTAING, J., PABOEUF, F., CHAUVEL, J. & JONDREVILLE, C. (2004) Variabilité intra-matière première de la digestibilité du phosphore des céréales et du pois chez le porc en croissance. *Journées de la Recherche Porcine en France*, **36**: 9-10.
- TRAN, G. & SKIBA, F. (2005) Variabilité inter et intra matière première de la teneur en phosphore total et phytique de l'activité phytasique. *INRA Productions Animales*, **18**: 159-168.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 VIVEROS, A., CENTENO, C., BRENES, A., CANALES, R. & LOZANO, A. (2000)
2 Phytase and acid phosphatase activities in plant feedstuffs. *Journal of Agricultural and*
3 *Food Chemistry*, **48**: 4009-4013.
4 VIEIRA, S.L., PENZ, A.M., KESSLER, A.M. & CATELLAN, E.V., Jr (1995) A nutritional
5 evaluation of triticale in broiler diets. *Journal of Applied Poultry Research*, **4**: 352-355.
6 ZACARIAS, I., YANEZ, E., ESCOBAR, M., HEWSTONE, C. & WULF, H. (1982)
7 Chemical and nutritional evaluation of triticale (*Secale* sp.) cultivated in Chili. *Archivos*
8 *Latinoamericanos de Nutrition*, **32**: 713-724.
9 ZHANG, Z. B., KORNEGAY, E. T., RADCLIFFE, J. S., DENBOW, D. M., VEIT, H. P. &
10 LARSEN, C. T. (2000) Comparison of genetically engineered microbial and plant phytase
11 for young broilers. *Poultry Science*, **79**: 709-717.
12 ZIMMERMANN, B., LANTZSCH, H. J., MOSENTHIN, R., BIESALSKI, H. K. &
13 DROCHNER, W. (2002) Comparative evaluation of the efficacy of cereal and microbial
14 phytases in growing pigs fed diets with marginal phosphorus supply. *Journal of the*
15 *Science of Food and Agriculture*, **82**: 1298-1304.

Table 1. *Physical and chemical characteristics¹ of 30 batches of triticale*

Variety ²	Origin	TGW	SW	HFN	DM	N	P	Phytic P	PA	PAV	RAV
		g	kg/hl	s	g/kg	g/kg	g/kg	g/kg	PU/kg	mL/g	mL/g
Trimaran	France	45.0	70.7	76	865	15,9	2.63	1.53	1023	2.35	1.79
A	France	47.0	69.8	67	871	16,1	3.11		987	2.89	1.84
Capitale	France	46.5	74.8	62	869	15,7	2.83	2.12	1226	3.42	1.76
Vision	Germany	57.5	75.5	109	874	16,1	3.13		974	3.56	2.28
B	Germany	48.5	72.3	84	856	16,1	2.75		842	3.92	2.01
Galtjo	Netherlands	55.0	75.2	260	857	17,1	3.22		728	3.84	1.97
Ego	Netherlands	53.5	73.9	158	857	15,0	3.11	2.32	1287	4.57	1.94
C	Sweden	45.5	73.0	70	853	14,6	2.83		881	3.50	1.83
D	Slovakia	53.0	72.2	64	862	17,2	2.75		1098	3.22	1.88
Colina	Romania	58.5	71.0	62	858	16,1	2.83	2.62	1346	2.44	2.07
E	Romania	54.5	74.5	75	859	17,8	3.57		1067	4.65	2.86
F	Poland	47.5	73.9	224	859	18,1	3.29	2.54	691	4.61	2.36
G	Poland	56.0	73.2	68	853	16,6	2.91		993	3.74	2.20
Binova	Germany	41.5	70.8	146	850	14,9	2.98		722	4.24	2.28
H	Germany	54.0	72.5	92	858	14,9	3.21		970	2.57	2.01
I	Poland	48.0	71.4	65	858	16,6	2.36		807	3.20	2.19
J	Poland	54.0	73.4	68	867	16,9	3.13		731	3.12	2.78
K	Poland	46.0	73.6	97	869	17,8	2.96	2.58	447	3.29	2.07
L	Poland	54.5	73.4	67	868	14,4	2.74		848	4.43	2.40
M	Slovakia	50.0	69.3	114	870	17,7	2.73		880	3.20	2.35
Aubrac	France	40.5	73.2	92	874	15,4	2.50	1.50	1103	3.86	2.32
Calao	France	41.5	68.1	62	858	16,1	2.83	1.95	1700	3.97	2.70
DI34-2	France	54.0	71.5	103	861	16,5	2.93	2.21	768	2.95	2.45
N	France	43.0	66.7	66	865	17,5	2.53	2.07	1843	3.52	2.06
O	France	49.0	74.4	94	859	15,4	2.06		1604	3.12	1.71
P	France	55.5	73.0	64	855	16,4	2.89		1064	2.87	2.01
Q	France	49.5	74.6	119	871	14,6	2.54		803	4.47	2.63
R	France	52.0	75.3	130	873	16,2	2.65	2.54	638	2.72	2.01
S	France	53.0	71.3	65	864	14,1	2.62		1101	4.12	1.98
T	France	52.0	71.3	68	873	16,4	3.21		1360	3.51	1.90
Mean		50.2	72.4	96	863	16,1	2.86	2.18	1018	3.53	2.15
SD		5.0	2.1	48	7	1,09	0.31	0.40	319	0.66	0.31
Min		40.5	66.7	62	850	14,1	2.06	1.50	447	2.35	1.71
Max		58.5	75.5	260	874	18,1	3.57	2.62	1843	4.65	2.86

¹TGW, thousand grain weight; SW, specific weight; HFN, Hagberg falling number; DM, dry matter; PA, phytase

activity; PAV, potential applied viscosity; RAV, real applied viscosity.

² cultivars or selection lines.

Table 2. *Composition and analytical characteristics of the experimental diets*

Diet	1 to 3	4 to 12 ¹		1 to 3	4 to 12 ¹
Ingredients	(g/kg diet as fed)		Analytical characteristics	(/ kg diet)	
Maize	450	0	Metabolisable energy, MJ ⁴	13.2	13.3
Soybean meal	230	230	Protein (N x 6.25), g ⁴	216	216 to 219
Soy isolate	70	60	Crude fibre, g ⁴	42 to 40	36
Triticale	0	450	Lysine ⁴	12.0	12.0
Maize starch	110	125	Methionine + Cystine ⁴	9.0	9.0
Wheat straw ²	59 to 54	39	Ca, g ⁴	10.6	10.6
Vegetable oil	45	60	P, g ⁵	3.5 to 5.7	3.2 to 3.8
Calcium carbonate ²	25 to 21	25	Phytic P, g ⁴	2.2	2.0 to 2.3
Monocalcium phosphate ²	0.0 to 8.8	0.0	Phytase activity, PU ⁵	<50	135 to 1390
DL-methionine	2.0	2.0	RAV, mL/g ⁵	0.63 to 0.80	0.96 to 1.40
Sodium chloride	3.0	3.0			
Minerals and vitamins premix ³	6.0	6.0			

¹ Without heat treatment, variety DI34-2, Trimaran, Aubrac and Calao in diets 4, 5, 6 and 7, respectively; heat-treated, variety DI34-2, Trimaran, Aubrac and Calao in diets 8, 9, 10 and 11, respectively; variety Calao, 200 g/kg without heat treatment and 250 g/kg heat-treated in diet 12. Heat treatment, which aimed at reducing phytase activity, consisted in two successive heating in a microwave oven at 600 W for 2 minutes.

² Wheat straw: 59, 56 and 54; calcium carbonate, 25, 23 and 21; monocalcium phosphate, 0.0, 4.4 and 8.8 g/kg in diets 1, 2 and 3, respectively.

³ Vitamin-trace mineral mix that provided the following per kg diet: vitamin A (retinyl acetate), 10000 IU; vitamin D3 (cholecalciferol), 4000 IU; vitamin E (DL-alpha-tocopherol), 80 mg; vitamin K3 (menadione), 4 mg; vitamin B1 (thiamin), 4 mg; vitamin B2 (riboflavin), 6.4 mg; vitamin B3 (PP, niacin), 80 mg; vitamin B5 (Ca pantothenate), 20 mg; vitamin B6 (pyridoxine), 5.6 mg; vitamin B8 (biotin, H), 0.2 mg; vitamin B9 (folic acid), 2.4 mg; vitamin B12 (cyanocobalamin), 0.016 mg; choline, 440 mg, Fe (FeSO₄), 40 mg ; Cu (CuSO₄), 16 mg; Mn (MnO), 64 mg; Zn (ZnSO₄), 72 mg; Co (CoSO₄), 0.5 mg; I (Ca(IO₃)₂), 1.6 mg; Se (Na₂SeO₃), 0.16 mg.

⁴ Estimated from INRA (1989) and the analysed N and phytic P concentrations in the four varieties of triticale.

⁵ Analysed according to the methods described in the materials and methods section: P, 3.5, 4.5, 5.7, 3.8, 3.2, 3.6, 3.6, 3.7, 3.3, 3.3, 3.7 and 3.8 g/kg; phytase activity, < 50, < 50, < 50, 620, 875, 920, 1390, 180, 200, 180, 135 and 645 PU/kg; RAV, Real Applied Viscosity, 0.76, 0.63, 0.80, 1.04, 0.96, 1.08, 1.03, 1.16, 1.13, 1.40, 1.22 and 1.02 ml/g in diets 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12, respectively.

Table 3. Correlation coefficient (r)¹ between physical and chemical characteristics² of the 30 batches of
triticales

	TGW	SW	HFN	DM	N	P	Phytic P ³	PA	PAV
SW	0.42 (*)								
HFN	0.04	0.38 (*)							
DM	-0.05	0.09	-0.14						
N	0.05	-0.11	0.19	0.05					
P	0.35	0.20	0.31	-0.11	0.32				
Phytic P ³	0.66 (*)	0.34	0.40	-0.27	0.45	0.60(*)			
PA	-0.15	-0.46 (*)	-0.38 (*)	-0.05	-0.10	-0.27	-0.36		
PAV	0.06	0.00	0.03	0.05	0.21	0.29	-0.03	-0.18	
RAV	-0.13	0.21	0.35	-0.13	-0.12	0.23	0.03	-0.03	0.43 (*)

¹*, $P < 0.05$.

²TGW, thousand grain weight; SW, specific weight; HFN, Hagberg falling number; DM, dry matter; PA, phytase activity; PAV, potential applied viscosity; RAV, real applied viscosity.

³ n = 11.

Table 4. *Phytase activity in 5 varieties of triticale*

Variety	n	Phytase activity, PU/kg ¹
DI34-2	4	1012 ± 102 a
Aubrac	6	1320 ± 87 b
Trimaran	6	1424 ± 125 b
Capitale	4	1815 ± 126 c
Calao	6	2146 ± 145 d
<i>P</i>		<0.001
RSD ²		119

¹ values are means ± standard deviation.

² RSD, residual standard deviation.

a-d; Means within a column not followed by the same letter differ at *P*<0.05.

Table 5. *Growth performance and bone¹ characteristics in chickens given the maize-based diets supplemented with graded levels of P as monocalcium phosphate or the triticale-based diets containing graded levels of phytase²*

Diet	Plant phytase	n	Initial weight	Final weight ³	Feed intake ³	FCR ^{3, 4}	Bone DM ^{3, 4}	Bone ash ^{3, 4}	Bone ash g/kg FF DM ^{3, 4}	
	PU/kg diet		g	g	g		g	g		
Maize-based diets										
P as monocalcium phosphate (g/kg)										
1	0	< 50	5	154	411 a	594 a	2.33 a	1.00 a	0.214 a	215 a
2	1	< 50	7	151	747 d	982 c	1.65 cd	2.02 c	0.589 c	292 f
3	2	< 50	8	150	901 e	1225 d	1.63 d	2.88 d	1.090 d	379 g
Triticale-based diets										
45 raw triticale										
4	DI-34-2	620	7	154	764 d	1039 c	1.70 bc	2.07 c	0.587 c	285 ef
5	Trimaran	875	8	151	751 d	1055 c	1.76 bc	1.98 c	0.524 c	266 de
6	Aubrac	920	8	150	755 d	1032 c	1.71 bc	2.00 c	0.572 c	285 ef
7	Calao	1390	8	151	737 d	1045 c	1.79 bc	1.99 c	0.542 c	272 ef
450 g/kg heated triticale										
8	DI-34-2	180	7	151	620 c	739 b	1.77 bc	1.46 b	0.364 b	251 cd
9	Trimaran	200	8	152	536 b	646 ab	1.84 b	1.27 ab	0.301 b	237 bc
10	Aubrac	180	8	150	539 b	693 ab	1.79 bc	1.37 b	0.327 b	238 bc
11	Calao	135	8	150	567 bc	706 ab	1.68 c	1.43 b	0.330 b	231 ab
Raw (200 g/kg) and heated (250 g/kg) triticale										
12	Calao	645	8	153	712 d	959 c	1.72 bc	1.92 c	0.504 c	264 d
P				<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	
RSD ⁴				65	102	0.15	0.25	0.071	19	

¹ Right tibiotarsus.

² Values are given on the basis of one cage of two birds.

³ Linear (L) and quadratic (Q) effects of P added as monocalcium phosphate (g/kg diet) and of phytase activity in the diet (PU/kg diet):

Final weight: P added, L ($P < 0.001$), Q ($P < 0.01$); phytase activity, L ($P < 0.001$), Q ($P < 0.001$), RSD = 69, $R^2 = 0.77$.

Feed intake: P added, L ($P < 0.001$), Q ($P < 0.05$); phytase activity, L ($P < 0.001$), Q ($P < 0.001$), RSD = 104, $R^2 = 0.78$.

FCR: P added, L ($P < 0.01$), Q ($P < 0.05$); phytase activity, L ($P < 0.001$), Q ($P < 0.01$), RSD = 0.19, $R^2 = 0.23$.

Bone DM: P added, L ($P < 0.001$); phytase activity, L ($P < 0.001$), Q ($P < 0.001$), RSD = 0.25, $R^2 = 0.78$.

Bone ash (g): P added, L ($P < 0.001$); phytase activity, L ($P < 0.001$), Q ($P < 0.001$), RSD = 0.075, $R^2 = 0.90$.

Bone ash (g/kg FF DM): P added, L ($P < 0.001$); phytase activity, L ($P < 0.001$), Q ($P < 0.001$), RSD = 20, $R^2 = 0.81$.

⁴ DM, dry matter; FF DM, fat-free dry matter; FCR, feed conversion ratio; RSD, residual standard deviation.

a-g; Means within a column not followed by a common letter differ at $P < 0.05$.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 6. Adjustment of final weight, feed intake, bone¹ dry matter and bone ash to P as monocalcium phosphate (MCP, g/kg diet) and phytase activity (PU/kg diet) and equivalency values of P as MCP (g) for plant phytase (PU)

	Final weight ³ g	Feed intake ³ g	Bone FF DM ^{4,6} g	Bone ash ⁴ g	Bone ash ⁴ g/kg FF DM ⁶
Non-linear adjustment ²					
Coefficients ²					
a	409	553	0.996	0.180	211
b	621	990	0.958	0.446	83.3
c	353	571	1.09	0.393	67.1
l	0.786	0.568			
k	0.00352	0.00200	0.00281	0.00303	0.00338
R ²	0.77	0.77	0.77	0.89	0.81
RSD	67	105	0.25	0.075	20
Equivalency values ⁵					
Plant phytase (PU)	P as monocalcium phosphate (g)				
150	0.34	0.29	0.42	0.32	0.32
250	0.52	0.45	0.60	0.47	0.46
500	0.81	0.80	0.89	0.69	0.66
750	0.96	1.05	1.02	0.79	0.74
1000	1.02	1.22	1.08	0.84	0.78
1250	1.05	1.33	1.11	0.86	0.79

¹ Right tibiotarsus.

² Models were generated using treatment means, R² (coefficient of determination) and RSD (residual standard deviation) are calculated relative to individual observations.

³ The model was $Y = a + b (1 - e^{-l \text{MinP}}) + c (1 - e^{-k \text{Phyt}})$, with Y = response measurement, MinP = P added as MCP (g/kg diet), Phyt = phytase activity (PU/kg diet).

⁴ The model was $Y = a + b \text{MinP} + c (1 - e^{-k \text{Phyt}})$, with Y = response measurement, MinP = P added as MCP (g/ kg diet), Phyt = phytase activity (PU/kg diet).

⁵ Final weight, feed intake, calculated as $\text{MinP} = -1/l \ln [1 - A (1 - e^{-k \text{Phyt}})]$; Bone dry matter and bone ash, calculated as $\text{MinP} = A (1 - e^{-k \text{Phyt}})$, with $A = c/b$, MinP = P as MCP (g), Phyt = phytase activity (PU)

⁶ FF DM, fat-free dry matter.

Figure heading

Figure. *Response of final weight and bone¹ ash to P as monocalcium phosphate in maize based diets and to plant phytase in triticale based diets.*

Figure footnotes

¹ Right tibiotarsus.

Values are means $\pm$ S.E.

Parameters of the models are presented in Table 6.