Accepted Manuscript

Title: Capacity of type I and II ligands to confer to estrogen receptor alpha an appropriate conformation for the recruitment of coactivators containing a LxxLL motif – Relationship with the regulation of receptor level and ERE-dependent transcription in MCF-7 cells

Authors: Sandrine Bourgoin-Voillard, Dominique Gallo, Ioanna Laïos, Anny Cleeren, Latifa El Bali, Yves Jacquot, Denis Nonclercq, Guy Laurent, Jean-Claude Tabet, Guy Leclercq

PII: S0006-2952(09)00930-7
DOI: doi:10.1016/j.bcp.2009.10.015
Reference: BCP 10361

To appear in: *BCP*

Received date: 3-9-2009
Revised date: 19-10-2009
Accepted date: 19-10-2009

Please cite this article as: Bourgoin-Voillard S, Gallo D, Laïos I, Cleeren A, Bali LE, Jacquot Y, Nonclercq D, Laurent G, Tabet J-C, Leclercq G, Capacity of type I and II ligands to confer to estrogen receptor alpha an appropriate conformation for the recruitment of coactivators containing a LxxLL motif – Relationship with the regulation of receptor level and ERE-dependent transcription in MCF-7 cells, *Biochemical Pharmacology* (2008), doi:10.1016/j.bcp.2009.10.015

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Capacity of type I and II ligands to confer to estrogen receptor alpha an appropriate conformation for the recruitment of coactivators containing a LxxLL motif – Relationship with the regulation of receptor level and ERE-dependent transcription in MCF-7 cells

Sandrine Bourgoin-Voillard ^{a, b}, Dominique Gallo ^a, Ioanna Laïos ^a, Anny Cleeren ^a, Latifa El Bali ^a, Yves Jacquot ^c, Denis Nonclercq ^d, Guy Laurent ^d, Jean-Claude Tabet ^b and Guy Leclercq ^{a, *}

^a Laboratoire J.-C. Heuson de Cancérologie Mammaire, Université Libre de Bruxelles (U.L.B.), Institut Jules Bordet, 1, rue Héger-Bordet, B-1000 Brussels, Belgium.

^b Université Pierre et Marie Curie – Paris 6, UMR 7201, Paris, F-75 005 France

^c Université Pierre et Marie Curie – Paris 6, UMR 7203, Paris, F-75 005 France

^d Service d'Histologie et de Cytologie Expérimentale, Faculté de Médecine et de Pharmacie, Université de Mons-Hainaut, 6, avenue du Champ de Mars, B-7000 Mons, Belgium.

* Corresponding author:

Pr. Guy Leclercq,
Laboratoire J.-C. Heuson de Cancérologie Mammaire
Institut Jules Bordet
1, rue Héger-Bordet
B-1000 Brussels
Belgium
Tél : +32 2 541 37 44
Fax : +32 2 541 34 98
e-mail : lcanmamm@ulb.ac.be

Abstract

Estrogen receptor α (ER α) belongs to the superfamily of nuclear receptors and as such acts as a ligand-modulated transcription factor. Ligands elicit in ER α conformational changes leading to the recruitment of coactivators required for the transactivation of target genes *via* cognate response elements. In many cells, activated ER α also undergoes downregulation by proteolysis mediated by the ubiquitin/proteasome system. Although these various molecular processes have been well characterized, little is known as to which extent they are interrelated. In the present study, we used a panel of type I (estradiol derivatives and “linear”, non-steroidal ligands) and type II (“angular” ligands) estrogens, in order to identify possible relationships between ligand binding affinity, recruitment of LxxLL-containing coactivators, ER α downregulation in MCF-7 cells and related transactivation activity of ligand-bound ER α . For type I estrogens, there was a clear-cut relationship between ligand binding affinity, hydrophobicity around C-11 of estradiol and ability of ER α to associate with LxxLL motifs, both in cell-free condition and *in vivo* (MCF-7 cells). Moreover, LxxLL motif recruitment by ER α seemed to be a prerequisite for the downregulation of the receptor. By contrast, type II ligands, as well as estradiol derivatives bearing a bulky side chain at 11 β , had much less tendency to promote ER α -LxxLL interaction or even behaved as antagonists in this respect, in agreement with the well known partial estrogenicity/antiestrogenicity of some of these compounds. Interestingly, some type II ligands which antagonized LxxLL motif recruitment were nonetheless able to enhance ER α -mediated gene transactivation.

Keywords: Estrogen Receptor alpha; Ligands; LxxLL motif; Downregulation; Transcriptional activity.

Introduction

Estrogen receptor alpha (ER α) is a member of the nuclear receptor superfamily, the members of which function as ligand-regulated transcription factors [1]. Estrogens are essential for the development and the maintenance of the female reproductive system. Besides, ER α is also known to play a pivotal role in the etiology of hormone-dependent forms of breast cancer [2]. Hence, for the last 30 years or so there has been an intensive search for agents capable of modulating and/or inhibiting ER α -mediated cell proliferation. This has led to the identification of two main classes of ligands both containing two hydroxyl functions that contribute to their binding to the receptor (Tables I and II) [3]. Type I ligands include estrogenic steroids and diphenolic structural analogues that similarly stimulate target tissues (*trans* stilbene derivatives such as diethylstilbestrol, isoflavones such as genistein, coumestanes such as coumestrol). These planar / linear agonists differentiate from angular weak agonists (type II) for which two subsets have been described: *cis* stilbene-like and *geminal* structures. Such angular ligands can in some cases antagonize the effect of strong type I estrogens [4]. Insofar as their pharmacological profile varies among different tissues, they are currently referred to as SERMs (Selective Estrogen Receptor Modulators).

All investigated type I ligands have been reported to fit within a cleft of the hormone binding domain (HBD) in such a way that their hydroxyl groups can form hydrogen bonds with a few amino acids (for E₂, 3-OH: Glu-353 and Arg-394; 17 β -OH: His-524; Fig. 1) [5;6]. Although this property does not hold for type II ligands because of inappropriate orientation of their hydroxyl groups, interactions with Glu-353 and Arg-394 are conserved and concur to produce non covalent ligand-receptor complexes. Stability of these complexes results from additional electrostatic interactions between the second phenolic ring of these ligands and

Thr-347, a residue located in a subregion of the ligand binding domain (LBD) known to attract substituents in C-11 of estradiol (E_2) [6;7]. While hydrogen bonding between the hydroxyl of this phenolic ring and Thr-347 (Fig. 1) is of importance for $ER\alpha$ binding, hydrophobic interactions between the C-11 subregion of the steroid core and the LBD are largely dominant [8]. Such differences in binding modes between type I and II ligands have been reported to confer to $ER\alpha$ distinct conformations that should influence its ability to recruit coactivators containing a consensus LxxLL binding motif (L = Leucine, x = any amino acid) [9-13] such as those of the CBP / p300 or the SRC / p160 families known as acetyltransferase enzymes and / or adaptator proteins for the transcription machinery recruitment (see [14-16]).

Even though there is a general consensus that the ability to recruit such coactivators determines the capacity of $ER\alpha$ to induce gene transactivation, no structure / activity study substantiating this concept has been reported so far. Similarly, there are no systematic data concerning a potential relationship between coactivator recruitment (or the absence of recruitment) and the capacity of a ligand to modulate $ER\alpha$ turnover, a factor known to influence receptor-mediated gene transactivation [17-19]. This gap in our knowledge has led us to conduct such a comparative study based on a panel of type I and type II estrogens. Of note, a part of our investigations focuses on several E_2 derivatives substituted in C-3, C-11 and C-17 since, as discussed above, these three reactive positions of the steroid core are known to play a prominent role in ligand binding and to influence the receptor conformations leading to an estrogenic response. Our experimental approach for this program relied on an ELISA-based assay allowing the quantitative measurements of ligand-activated $ER\alpha$ associated with immobilized LxxLL motif-containing peptide. In previous studies, the specificity of the assay was established by competition experiments showing that a peptide

derived from the SRC-1 coactivator effectively suppresses receptor binding to LxxLL-coated plates through a complexation process [20].

Accepted Manuscript

Material and methods

Compounds

Type I ligands were purchased from Sigma (St Louis, MO) or Steraloids (Newport, RI) while type II [4;21] were obtained from Prof. R. Gust (Pharmacological Institute, Free University Berlin, Germany) ; 11 β long chain derivatives of E₂ [22] were from Prof. J.-C. Blazejewski (University of Versailles, France). For assays, stock ethanol solutions of these compounds were diluted in buffer (cell-free assays) or medium (cell culture) in order to have a final concentration of solvent below 0.1%. Routine biochemical reagents were from standard suppliers.

Cell-free assays with human ER α recombinant

Assays were conducted with a highly purified human ER α (hER α) recombinant obtained from Calbiochem (Euro-Biochem, Bierges, Belgium). Upon receipt, the protein was diluted ($\approx$ 2.5 pmol/ μ L) in 50 mM Tris-HCl pH 7.5 containing 500 mM KCl, 2 mM DTT, 1 mM EDTA, 1 mM sodium orthovanadate and 10 % glycerol. Diluted aliquots were stored at -80°C .

Relative Binding Affinity of compounds for hER α

Binding affinity of the compounds for hER α was evaluated by a solid phase competition assay based on the adsorbing property of the receptor onto hydroxylapatite (HAP) at low ionic strength [23]. For that purpose, hER α diluted in a buffered bovine serum albumin (BSA) solution (1:500 in 10mM Tris-HCl pH 8 containing 1 mg/mL BSA) was adsorbed onto HAP (Bio-Rad). After removal of unbound receptor by centrifugation, the HAP suspension was incubated during 24 hours at 4°C in the presence of 1 nM [³H]E₂ (GE

Healthcare Biosciences, Eindhoven, NL) with or without increasing amounts of the investigated compounds or unlabeled E₂ taken as a reference. Bound [³H]E₂ was then extracted with ethanol and measured by liquid scintillation counting. Relative concentrations of investigated compounds and E₂ required to inhibit the binding of [³H]E₂ by 50% yielded the Relative Binding Affinity values ($RBA = [IC_{50}]_{E_2} / [IC_{50}]_X \times 100$). Assays were performed twice, each time in duplicate. In each competition curve, variations between bound [³H]E₂ values were extremely low (SD = 3 %), suggesting a very high reproducibility of the data.

Ability of compounds to confer to hER α a conformation appropriate for recruitment of a LxxLL-containing peptide.

The ability of hER α to associate with a capture LxxLL containing peptide was assessed with an ELISA-based assay (Elisa NR peptide ER α -chemiluminescent assay from Active Motif, Rixensart, Belgium). hER α (5 ng in the diluent buffer) was incubated on ice with increasing concentrations of the investigated compounds (from 10 pM to 1 μ M) for 30 min. These samples were then added to LxxLL peptide-coated multiwell plates and further incubated for 1 hour at 20°C; controls were run in parallel without any compound. Binding of hER α to the wells was finally assessed by successive exposures to an anti-ER α primary antibody, an HRP-conjugated secondary antibody and a chemiluminescent development reagent according to manufacturer's instructions. Light intensity was measured using an Infinite® 200 microplate reader (Tecan, Mechelen, Belgium). Assays were performed in triplicate and the recruitment capacities (RCs) values expressed as percentages $\pm$ SD of the optimal value established with 1 μ M E₂.

Assays on MCF-7 cells

Cell culture

MCF-7 and MVLN cells (MCF-7 cells stably transfected with a vitERE-tk-Luc reporter plasmid [24]) were propagated at 37 °C (5 % CO₂, humid atmosphere) in Earle's based minimal essential medium (EMEM; Invitrogen, Carlsbad, California) supplemented with Phenol Red, 2 mM L-glutamine, 100 U/mL penicillin, 100 mg/mL streptomycin (from Invitrogen) and 10 % heat-inactivated fetal bovine serum (FBS; HyClone, Logan, Utah). Experiments were conducted in Phenol Red-free EMEM containing 10% charcoal-stripped FBS (estrogen-free medium, EFM).

Ability of compounds to confer to ER α a conformation appropriate for recruitment of an LxxLL-containing peptide.

MCF-7 cells were cultured for 2 days in EFM (6-wells dishes) before treatment with increasing concentrations of investigated compounds. After treatment (usually 15 min), cell monolayers were washed twice with ice-cold PBS (40 mM Na₂HPO₄, 10 mM KH₂PO₄, 120 mM NaCl, pH 7.2) and the cells were lysed in 50 μ L ice-cold RIPA (50 mM Tris-HCl, pH 7.5, 150 mM NaCl containing 1% NP-40, 0.1% SDS, 0.5% sodium deoxycholate, 50 mM NaF, 0.1 mM orthovanadate, 0.6 mM PMSF, 0.3 mM TPCK). Lysates were diluted by adding 200 μ L of diluent buffer (reagent # 101251 of the Active Motif's ELISA kit) and clarified by fine needle aspirations followed by centrifugation. Ability of ER α from these samples (150 μ L) to associate with LxxLL motif was then assessed as described above.

Ability of compounds to bind to ER α ("whole cell assay")

The binding of test compounds to ER α in MCF-7 cells was evaluated by a [3 H]E $_2$ competitive assay, using E $_2$ as a reference compound (RBA = 100 %) [25]. For that purpose, cells were exposed for 45 min to 1 nM [3 H]E $_2$ alone (control) or in the presence of increasing amounts of unlabeled E $_2$ or investigated compounds in serum-free medium; additional cell cultures were exposed to 500-fold excess unlabeled E $_2$ for assessment of non-specific binding (NSB). Bound [3 H]E $_2$ was measured by scintillation counting after extraction and the data (total – NSB) expressed in percentages of the control value in order to establish RBA values (*i.e.* [IC $_{50}$] $_{E_2}$ / [IC $_{50}$] $_X$ X 100).

Ability of compounds to regulate ER α level

MCF-7 cells were plated in 10 cm-diameter Petri dishes (500,000 cells per dish) containing EFM. After 3 days of culture, medium was removed and cells were exposed to increasing amounts of the investigated compounds for 4 hours in a fresh medium ; control cells were maintained in culture without any compound (this short time period was selected to avoid an effect on ER α synthesis) [17]. Cells were then washed with TBS (50 mM Tris, pH 7.5, 150 mM NaCl) and lysed for 30 min at 4°C in RIPA buffer (see above). ER α levels in these lysates were finally assessed by Western blotting following a procedure described previously[17]. Briefly, ER α was detected with a mouse monoclonal antibody (F-10, Santa Cruz Biotechnology, Santa Cruz, CA). Actin, used as a loading control, was detected with mouse monoclonal antibody MAB1501R from Chemicon International (Temecula, CA). Exposure to primary antibodies was followed by incubation in presence of a horseradish peroxidase-conjugated goat anti-mouse secondary antibody (Pierce, Rockford, IL). Peroxidase activity was revealed with Supersignal West Pico Chemiluminescent substrate from Pierce. ER α and actin bands were captured with a FLA-3000 digital camera (Fuji, Tokyo, Japan). Band intensities were quantified using AIDA software (Raytest,

Straubenhardt, Germany). ER α data were normalized according to actin (ER α / actin) and expressed in percent of control (untreated cells). Ability of bisphenol (# 23) to antagonize E₂-induced downregulation of ER α was further assessed by immunofluorescence microscopy using an experimental protocol described previously [26].

Ability of compounds to enhance ER α -mediated transactivation

MVLN cells were cultured for 3 days in 6-well plates (plating density 100,000 cells/well) containing EFM. Medium was then removed and replaced by fresh medium containing increasing concentrations of the investigated compounds ; control cell cultures were run in parallel in the absence of any compound. After 24 hours of incubation, luciferase activity was assayed according to a previously described procedure [17]. Efficiency of compounds was expressed as percentages of luciferase activity induced by 10 nM E₂.

RESULTS

1. Ability of estradiol and investigated derivatives to modulate the capacity of a hER α recombinant to recruit coactivators containing LxxLL motifs

1.1. Importance of ligand binding affinity for ER α

Interaction of each ER α -ligand complex with LxxLL-coated plates is likely to depend on various factors, including in particular the binding affinity of the ligand for the LBD. In other words, the extent of LxxLL motif recruitment by ER α should be related to the relative ligand binding capacity, even though the measurement of LxxLL recruitment and of ligand binding rely on very different procedures. To validate this view, we checked whether the recruitment curve established with increasing amount of E₂ correlated with the [³H]E₂ / E₂ competition curve. As expected, both curves mirrored each other (Fig. 2) with RC₅₀ (3 nM) and IC₅₀ (2 nM) values in the same order of magnitude.

We subsequently assessed whether this property holds for E₂ derivatives whose functions / regions of the steroid core known to influence ER α binding had been modified. The next sections refer to this study.

1.2. Importance of C-3 and C-17 hydroxyl groups of estradiol

In the binding pocket of ER α , Glu-353 and Arg-394, with the participation of water molecules, interact with the 3-phenolic function of E₂ while His-524, modeled as the ϵ tautomer, attracts its 17 β -alcoholic group (Fig. 1) [5;27]. To assess the importance of these bridging properties on LxxLL coactivator recruitment, we tested three E₂ derivatives bearing modified OH group at these critical positions (i.e., -OCH₃ / -OSO₂O⁻Na⁺ for 3-OH, and -H for

17 β -OH; compounds # 2, 3 and 4, respectively) (Table I). As shown in Fig. 3a and 3b, these substitutions largely decreased the capacity of the hormone to promote ER α binding to LxxLL-coated plates, reflecting the acidic role of phenol in the complex. Interestingly, stereochemistry change of the β OH at C-17 into the α configuration (# 5) decreased by 40% ER α association with LxxLL motifs while it decreased by 70 % ligand binding to the receptor, stressing the importance of the hydrophobic steroidal core of the hormone in coactivator recruitment. Studies concerning substitutions in C-11 of E₂, reported below, support this view.

1.3. *Influence of substitutions in C-11 of estradiol*

The LBD is highly hydrophobic and therefore not well suited to accommodate E₂ derivatives in which an hydrogen in C-11 is substituted by a polar residue. In contrast, hydrophobic substituents are readily accepted if their size is not too large (see Table I) and if they stabilize ER α -ligand complexes at high temperature [8].

Figure 3c shows that the grafting of a hydroxyl in C-11 α or β of E₂ (compounds # 6 α and 6 β) modify its binding properties to the LxxLL-coated plate. While a decrease of the ability of hormone to promote high binding was observed at all concentrations for 11 α -OH E₂, this property was solely recorded at concentrations ≥ 10 nM for 11 β -OH E₂. The latter ligand was indeed more effective than E₂ at 1 nM indicating that regulation of LxxLL binding by the C-11 subregion of the steroid core is influenced by factors other than polarity, that antagonize its insertion within the LBD. In this regard, gas phase acidity study of these steroids has shown that 11 β -OH E₂ presents a higher acidity than 11 α -OH E₂ and E₂, indicating that particular electronic effects occur in 11 β -OH E₂ [28]. Polarity appeared nevertheless of prime importance for LxxLL-containing coregulators recruitment because

acetylation of the 11 β -OH function (# 7) resulted in a ligand able to induce “supraphysiological” receptor- LxxLL motif association.

According to this concept, hydrophobic substitutions in the C-11 β stereochemistry should confer to ER α a conformation favoring extensive LxxLL recruitment. In agreement with this view, we found that -CH₃ (# 8), -CH₂Cl (# 9) and -C $\equiv$ CH (# 10) substituents enhanced ligand-induced association of ER α with LxxLL motifs (Fig. 3d). Interestingly, distinct LxxLL binding profiles were recorded for these three ligands showing that the chemical nature of the substituent influence the ability of the receptor to interact with LxxLL motifs. The observation that the introduction of a double bond between C-9 and C-11 (# 11) produced a shift in ER α -LxxLL binding (as compared to the curve generated by E₂), similar to that produced by the grafting of -CH₂Cl or -C $\equiv$ CH groups, revealed that it is the whole region around C-11 which exerts an influence on LxxLL recruitment (Fig. 3f).

In contrast to these small substituents, large hydrophobic side chains grafted at 11 β (# 12, 13 and 14) totally abrogated the basal LxxLL binding capacity of ER α (Fig. 3e). Hence, such E₂ derivatives, usually not well accommodated by the LBD [8;22], must be considered as LxxLL recruitment antagonists, a property confirmed by the fact that a -C₆H₁₃ derivative - tested as a prototype - blunted the enhancing effect of E₂ on ER α -LxxLL association.

2. Ability of non-steroidal estrogens to modulate the capacity of ER α to recruit co-regulators containing LxxLL motifs

2.2. Type I ligands

Diethylstilbestrol (#15), whose binding affinity for ER α largely exceeds that of coumestrol (#16) and genistein (#17) enhanced the binding of the receptor to LxxLL-coated plates with a much higher efficiency as compared to the latter compounds (Table II and Fig. 4), suggesting a direct relationship between ligand affinity and capacity to induce receptor-LxxLL association. Hence, "linear" estrogens, the phenolic functions of which interact with Glu-353, Arg-394 and His-524 may be assimilated to steroidal estrogens in terms of coactivator recruitment.

2.2. *Type II ligands*

None of the "angular" ligands tested in this study enhanced the binding of ER α to LxxLL-coated plates, with the notable exception of a diphenolic imidazoline derivative (# 18) for which a weak ability to induce LxxLL recruitment had already been reported [29] (Table II, Fig. 5a-b). Moreover, *gem*-diphenolic ligands (# 21 - 25) antagonized ER α binding to LxxLL motifs in the presence as well as in the absence of E₂, as clearly demonstrated with bisphenol (# 23) and its analogs (Fig. 5c). This antagonism could not be ascribed to a lack of hydrogen bonding with His-524 in view of the fact that substitution of 17 β -OH of E₂ by a hydrogen atom (# 4) failed to show such a property. Hence, hydrogen interaction with Thr-347 seems directly responsible of the antagonistic property of these ligands. Thus, type II angular ligands totally distinguish themselves from type I linear ligands in terms of coactivator recruitment.

3. **Recruitment of coactivators containing an LxxLL motif by ER α in MCF-7 cells**

Exposure of MCF-7 cells to E₂ produced in ER α a rapid conformational change favoring the recruitment of LxxLL-containing coactivators, as shown by measuring in cell extracts the extent of ER α binding to LxxLL-coated plates (Fig. 6). Enhanced ER α binding,

which culminated within the first minutes of incubation, became undetectable after 4 hours. Thus, experiments aiming to assess whether observations under cell-free conditions can be extended to living cells were based on short-term (15 min) incubations of cells with the ligands. After this treatment duration, the LxxLL recruitment curve established with increasing amounts of E₂ displayed a profile quite similar to that observed for the recombinant receptor, although the RC₅₀ value was about 10 times higher (Fig. 6, Table III), probably reflecting the existence of cellular factors (*i.e.* endogenous receptor coregulators) limiting the binding of ER α to LxxLL-coated plates. The fact that this property was not recorded for the hydrophobic 11 β -substituted derivatives (# 9, 8 and 10) of E₂ (Table III) suggests that the stability of ER α -ligand complexes modulates the impact of these factors. In agreement with this view, we found that the inhibitory effect of 11 β -OH E₂ on receptor-LxxLL interactions was exacerbated in cell culture experiments, leading to an almost total suppression of the recruitment capacity of the receptor. As expected, substitutions of 3 or 17 β -hydroxyls of E₂ (# 2, 3, 4 and 5) also abrogated its recruitment ability (weak effect at 10 μ M).

ER α from MCF-7 cells treated with DES bound to the LxxLL-coated plates like the receptor from E₂-treated cells. By contrast, other non steroidal type I ligands had almost no effect on receptor-LxxLL interactions, most probably because of their low affinity for ER α in cultured cells (Table III). As expected, compounds antagonizing hER α interaction with LxxLL motifs in cell-free conditions (# 12, 23 and 25) produced a similar effect in cell extracts.

On the other hand, kinetic studies of the ER α -LxxLL peptide interaction induced by E₂ and its 11 β -CH₂Cl derivative revealed identical profiles. Thus, any ER α -ligand complexes potentially capable of recruiting LxxLL-motif coactivators progressively disappeared. This

property may reflect a competition with cellular coactivators, a strong ER α association with an insoluble target, an alteration of its LxxLL binding motif or a proteasomal degradation of the receptor (these various possibilities being non exclusive) [30;31]. On the other hand, concerning the partial estrogenicity of bisphenol (# 23), we found that this compound failed to display any ability to recruit LxxLL motifs even after 4 hours of cell treatment. Hence, the agonistic activity of this ligand [32] seems to be unrelated to a delayed recruitment of LxxLL motif-harboring coactivators.

4. Relationship between ligand-induced recruitment of coactivators containing LxxLL motif and ER α level in MCF-7 cells

Early studies from our laboratory have shown that E₂-induced depletion (downregulation) of ER α in MCF-7 cells is evident after 4 hours of hormone treatment [17]. As shown in Fig. 7a, after this duration of exposure, E₂ concentrations ranging from 0.1 nM to 1 μ M induced a similar extent of ER α decrease (~30-40 %). Thus in the case of E₂, there was no quantitative relationship between the ligand-induced downregulation of ER α and the degree of ER α association with LxxLL motifs. This apparent absence of relationship was further illustrated by the behavior of 11 β -substituted E₂ derivatives (# 9, 8 and 10) as well as the C9=C11 analog (# 11) which at 10 nM produced ER α downregulation with an efficiency similar to that of E₂, even though they were more efficient at recruiting the LxxLL motif (Fig. 7b and Table III) (note that at higher concentrations of these compounds did not produce more pronounced downregulation). However, the low sensitivity of Western blotting (see SD of the data) could preclude the detection of small ER α level changes which might occur during the 15-min time period used for the assessment of LxxLL binding, obscuring thereby a potential relationship between recruitment of LxxLL-containing coactivators and ER α .

downregulation. Indeed, because of the limited sensitivity of Western blotting, downregulation of ER α can only be evidenced after several hours of drug exposure [17]. Nevertheless, such a relationship may exist in view of the fact that, within the exception of 11 α -OH E₂, compounds provoking an extensive LxxLL recruitment downregulated ER α with a higher efficiency than compounds which were less active in this respect (Table III). This view is supported by the additional finding that a 100 fold excess of bisphenol, which impedes the recruitment of LxxLL-containing peptides, (# 23) abrogated E₂-induced ER α downregulation (immunofluorescence : see Fig. 8 ; Western blotting : residual ER α level 43 % $\rightarrow$ 108 %). Note also in this context that the 11 β -C₆H₁₃ derivative of E₂ (# 12), which similarly antagonized this recruitment, was able to stabilize the receptor when added at concentrations $\geq$ 10 mM, producing an upregulation similar to that described for tamoxifen [17].

Unexpectedly, 11 β -OH E₂ downregulated ER α with an efficiency as strong as E₂. The lack of capacity of this compound to promote an ER α conformation appropriate for the recruitment of LxxLL coregulators, as well as its limited binding to the receptor suggest a mechanism of receptor downregulation different from that induced by other steroids.

Previous studies have revealed that cycloheximide (CHX) at 50 μ M abrogates E₂-induced ER α degradation by the ubiquitin-proteasome system [33-35]. Our investigations extended this finding to all ligands able to induce LxxLL-motif recruitment (data not shown). Downregulation induced by the E₂ derivative C₆F₁₃-substituted in 11 β (# 14), unable to enhance the ER α -LxxLL peptide interaction, was also sensitive to CHX (% residual ER α level -/+ CHX : 43/95), excluding any major implication of LxxLL-coactivator recruitment in the action of CHX.

5. Relationship between ligand-induced recruitment of coactivators containing LxxLL motif and ERE-dependent transcription in MCF-7 cells stably transfected with a vitERE-tk-Luc plasmid (MVLN cells)

Steroidal and nonsteroidal ligands able to enhance the binding of ER α to the LxxLL-coated plate increased transcription of the luciferase gene (reporter gene) with an efficiency closely related to their ability to down regulate the receptor (see Table III and representative compounds in Fig. 9). Hence, as already reported [18;19], proteasomal degradation of ER α appeared as an important step in ER α -mediated transcription induced by compounds sharing some structural homology with E₂. We believe, therefore, that physiological ER α -induced gene transactivation might be relevant to a process involving LxxLL motif-coactivator recruitment followed by the proteasomal degradation of the receptor.

Paradoxically, 11 β C₆F₁₃-E₂ (# 14), which failed to modify the binding of ER α to the LxxLL-coated plate, and compounds that antagonized this binding (# 12, 23, 25) increased luciferase expression. This indicates that recruitment of LxxLL-containing coactivators is not absolutely required for ERE-dependent transcription (Table III, Fig. 9). ER α downregulation seems of higher importance in this regard since enhancement of luciferase expression was recorded at ligand concentrations which decreased receptor level. The additional observation that 11 α OH-E₂, which down regulated ER α by a mechanism most probably independent of LxxLL-motif recruitment, increased luciferase expression supports this view.

At concentrations which failed to decrease ER α level, 11 β C₆H₁₃-E₂ (# 12) lost its capacity to enhance transcription, while bisphenol (# 23) remained able to induce transcriptional activity (Fig. 9), in agreement with the ability of these compounds to either up

regulate or stabilize the receptor. The fact that bisphenol was reported to antagonize E₂-induced ERE-dependent transcription [4] clearly indicates that this compound interferes with the normal action of the hormone, most probably by abrogating the capacity of the hormone to confer a conformation appropriate for the recruitment of LxxLL- motif containing coregulators.

DISCUSSION

The present study evaluates the ability of various ER α ligands to induce the recruitment of LxxLL-containing coactivators by the receptor and examines a possible relationship between this recruitment, ER α level and ER α -mediated gene transactivation. Our data clearly show that both steroidal and planar / linear type I estrogens enhance LxxLL motif recruitment with an efficiency related to their binding affinity. By contrast, angular type II estrogens have much less tendency to promote receptor binding to LxxLL motifs or even act as antagonists in this respect. This is in agreement with the usual classification of these nonsteroidal compounds as partial estrogens/antiestrogens, as opposed to pure antagonists which exclusively abrogate the recruitment of LxxLL-containing coactivators [36].

The insertion of E₂ (and probably of most type I estrogens) into the LBD of ER α provokes a rapid shift of its helix H12 which seals the ligand binding cavity [5]. In this new conformation, H12 covers the binding pocket and plays a major role in stabilizing ligand-receptor complexes. In contrast, type II estrogens behave differently inasmuch as they maintain the pocket in an “open conformation”, this resulting in less stable complexes. The latter conformation being generated by ligands belonging to various chemical families, one may reasonably postulate that it is associated with a large panel of H12 helix orientations explaining the agonistic / antagonistic profile of these compounds in terms of LxxLL-coactivator recruitment.

Hydrogen bridges involved in E₂ binding (3-OH and 17 β -OH), which contribute to the displacement of H12 [5], are found here to be essential for optimal LxxLL recruitment. We show in this study that small hydrophobic substituents in C-11 β , as well as the introduction of

a double bond between C-9 and C-11, enhance the recruitment of LxxLL motifs. This observation suggests that hydrophobic interactions between the LBD and substituents around the C-11 position of the ligand strengthen the affinity of the receptor toward LxxLL coactivators. Polar residues, as well as long hydrophobic side chains not readily accommodated by the LBD largely limit this promoting effect. However, the 11 β -OH E₂ shows an unexpected behavior that could be relevant to electronic effects [28]. In fact, the nature of the interactions between ligand substituents and the LBD most likely influence the affinity of ER α for coactivators since (-CH₃) *versus* (-CH₂Cl and -C $\equiv$ CH) gave distinct LxxLL binding profiles. Hence, steric hindrance and/or electronic delocalization may also play a role. In this context, it should be stressed that C-11 β hydrophobic substituents often increase E₂ binding to ER α at 25° C [8]. This property, most probably due to a higher plasticity of the LBD at this temperature, was associated with an increase (-CH₃, -CH₂Cl, -C $\equiv$ CH) or a decrease (-C₆H₁₃) of LxxLL recruitment capacity, stressing its relevance to the agonistic as well as antagonistic character of a ligand.

Stereochemistry change of β OH at C-17 of E₂ significantly decreases its binding affinity for ER α . Our data reveal that such a change would not markedly affect the ability of the hormone to recruit LxxLL-containing coregulators, in agreement with the known estrogenicity of 17 α -E₂. On the contrary, α/β configuration of a hydroxyl at C-11 strongly affects the recruitment of a LxxLL motif in a way closely related to the decrease of binding affinity.

According to our data, Glu-353/Arg-394 and His-524, which contribute to the binding of type I ligands through hydrogen bonds, appear to be similarly implicated in LxxLL-coactivator recruitment. This property does not hold for Thr-347, even though this residue

participates to the interaction of type II ligands with the LBD. Indeed, bisphenol displays an evident antagonistic activity on LxxLL motif recruitment. Thus, H-bonds between the aforementioned residues and the phenol / alcohol groups of these two classes of ligands govern the coactivator recruitment capacity of the receptor. It should be noted in this context that the spatial orientation of these hydroxyl groups is also of importance, as revealed by the distinct LxxLL recruitment capacity of steroidal estrogens and the type I estrogens (e.g., genistein and coumestrol).

Influence of substitutions in position 11β of E_2 on LxxLL-motif recruitment could not similarly be explained because of the lack of information concerning the interactions of such substituents with specific residues of the LBD. While their insertion in the subpocket containing Thr-347 may be advocated, a “ 7α subpocket” could not be excluded in view of the fact that E_2 may rotate around a $3OH-17\beta OH$ axis [37]. This 7α subpocket, essentially hydrophobic, is known to accept alkyl substituents which generate ligands with RBA values overpassing that of E_2 [38]. Hence, this mode of insertion within the LBD appears particularly well suited for our $-CH_3$, $-CH_2Cl$ and $-C\equiv CH$ derivatives of E_2 , in contrast to the Thr-347 insertion mode which is more appropriate for ligand with aromatic substituents [38]. Hydroxyl groups being hardly accepted by this 7α subpocket, $11\beta-OH$ would be rejected away from the latter with a concomitant decrease of the capacity of $ER\alpha$ to recruit LxxLL-motifs.

The acquisition of an $ER\alpha$ conformation appropriate for LxxLL motif recruitment seems to be necessary for its proteasomal degradation in MCF-7 cells. This property does not hold for ERE-dependent gene transactivation, since bisphenol, which antagonizes LxxLL motif recruitment, enhances the transcription of reporter genes at concentrations which

upregulate the receptor. The fact that this ligand suppresses the stimulatory effect of E₂ on LxxLL motif recruitment by ER α suggests that its ability to enhance transactivation in estrogen-free conditions results from a mechanism involving a distinct class of coactivators. Another possibility would be the induction of intermolecular interactions *via* the expression of LxxLL-like motifs, which would generate a competitive inhibition in our LxxLL binding assay. To our knowledge, this model, which explains the antagonistic property of SERMs (*i.e.* the LLEML motif of H12 that fits between H3 and H5 [39]), has never been proposed in the context of type II agonists probably because of the lack of any description of auto-activating mechanism of ER α .

In agreement with previous observations, CHX was found to systematically abrogate the proteasomal degradation of ER α induced by the ligands which enhanced ERE-dependent transcription in MCF-7 cells, even the 11 β -C₆F₁₃ derivative of E₂ which failed to promote LxxLL-motif recruitment. Hence, the absence of CHX effect on the downregulation of ER α induced by the pure antiestrogen RU 58668 [34;35] which contains a functionalized side chain in C-11 β like this C₆F₁₃ derivative, would be related to the antiestrogenic activity of this compound. This interpretation similarly holds for the ER α downregulation induced by ICI 182,780, which is also insensitive to the action of CHX [35]. All these considerations indicate that newly synthesized protein(s) operate in concert with the activated receptor to induce its degradation, in a process independent of LxxLL-containing coactivator recruitment.

In MCF-7 cells, E₂-induced LxxLL-coactivator recruitment was fast and transient. Although the underlying mechanism has not been investigated as yet, this observation is consistent with the existence of repeated cyclic interactions between ER α and cognate partners during target gene transactivation, a concept developed by several authors

[18;19;30;40;41]. Whether the membrane / cytoplasmic receptor pool, which participates to early estrogenic responses [31, 42, 43], is implicated in this dynamic LxxLL-coactivator recruitment is unknown. Investigating such a possibility seems to us of primary importance. Use of ligands that selectively interact with this pool [44-47] might be an adequate approach to initiate such a study.

In conclusion, our study shows that ER α -mediated transcription induced by type I estrogens depends upon LxxLL-coactivator recruitment. In contrast, type II estrogens are largely less dependent of this recruitment, a fact which may explain their mixed estrogenic/antiestrogenic properties. Moreover, same hydrophilic and hydrophobic interactions appear to govern the stability of receptor-ligand complexes, the attraction of LxxLL motifs and associated biological responses.

ACKNOWLEDGMENTS

This work was supported by grants from the Belgian Fund for Medical Scientific Research (n° 3.4501.09), the "Fonds Lambeau-Marteaux", the "Fonds Jean-Claude Heuson" for breast cancer research and from "Les amis de l'Institut Bordet". We also acknowledge the French Ministry of Research, CNRS and UPMC for financial supports. Dominique Gallo is recipient of an award from the "Fondation David et Alice Van Buuren". Guy Laurent is Senior Research Associate of the National Fund for Scientific Research (Belgium). We are grateful to Xavier Rousseau from the "Institut Meurice" for its active contribution in this work.

REFERENCES

- [1] Tsai MJ, O'Malley BW. Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu Rev Biochem* 1994; 63: 451-486.
- [2] McGuire WL. Hormone receptors: their role in predicting prognosis and response to endocrine therapy. *Semin Oncol* 1978; 5: 428-433.
- [3] Jordan VC, Schafer JM, Levenson AS, Liu H, Pease KM, Simons LA, Zapf JW. Molecular classification of estrogens. *Cancer Res* 2001; 61: 6619-6623.
- [4] Lubczyk V, Bachmann H, Gust R. Investigations on estrogen receptor binding. The estrogenic, antiestrogenic, and cytotoxic properties of C2-alkyl-substituted 1,1-bis(4-hydroxyphenyl)-2-phenylethenes. *J Med Chem* 2002; 45: 5358-5364.
- [5] Brzozowski AM, Pike AC, Dauter Z, Hubbard RE, Bonn T, Engstrom O, Ohman L, Greene GL, Gustafsson JA, Carlquist M. Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature* 1997; 389: 753-758.
- [6] Kekenyes-Huskey PM, Muegge I, von RM, Gust R, Knapp EW. A molecular docking study of estrogenically active compounds with 1,2-diarylethane and 1,2-diarylethene pharmacophores. *Bioorg Med Chem* 2004; 12: 6527-6537.
- [7] Bai Z, Gust R. Breast cancer, estrogen receptor and ligands. *Arch Pharm (Weinheim)* 2009; 342: 133-149.
- [8] Anstead GM, Carlson KE, Katzenellenbogen JA. The estradiol pharmacophore: ligand structure-estrogen receptor binding affinity relationships and a model for the receptor binding site. *Steroids* 1997; 62: 268-303.
- [9] Heery DM, Kalkhoven E, Hoare S, Parker MG. A signature motif in transcriptional co-activators mediates binding to nuclear receptors. *Nature* 1997; 387: 733-736.
- [10] Iannone MA, Simmons CA, Kadwell SH, Svoboda DL, Vanderwall DE, Deng SJ, Consler TG, Shearin J, Gray JG, Pearce KH. Correlation between in vitro peptide binding profiles and cellular activities for estrogen receptor-modulating compounds. *Mol Endocrinol* 2004; 18: 1064-1081.
- [11] Ozers MS, Ervin KM, Steffen CL, Fronczak JA, Lebakken CS, Carnahan KA, Lowery RG, Burke TJ. Analysis of ligand-dependent recruitment of coactivator peptides to estrogen receptor using fluorescence polarization. *Mol Endocrinol* 2005; 19: 25-34.
- [12] Savkur RS, Burris TP. The coactivator LXXLL nuclear receptor recognition motif. *J Pept Res* 2004; 63: 207-212.
- [13] Tamrazi A, Carlson KE, Rodriguez AL, Katzenellenbogen JA. Coactivator proteins as determinants of estrogen receptor structure and function: spectroscopic evidence for a novel coactivator-stabilized receptor conformation. *Mol Endocrinol* 2005; 19: 1516-1528.

- [14] Klinge CM. Estrogen receptor interaction with co-activators and co-repressors. *Steroids* 2000; 65: 227-251.
- [15] McKenna NJ, Lanz RB, O'Malley BW. Nuclear receptor coregulators: cellular and molecular biology. *Endocr Rev* 1999; 20: 321-344.
- [16] Rosenfeld MG, Lunyak VV, Glass CK. Sensors and signals: a coactivator/corepressor/epigenetic code for integrating signal-dependent programs of transcriptional response. *Genes Dev* 2006; 20: 1405-1428.
- [17] Laios I, Journe F, Nonclercq D, Vidal DS, Toillon RA, Laurent G, Leclercq G. Role of the proteasome in the regulation of estrogen receptor alpha turnover and function in MCF-7 breast carcinoma cells. *J Steroid Biochem Mol Biol* 2005; 94: 347-359.
- [18] Lonard DM, Nawaz Z, Smith CL, O'Malley BW. The 26S proteasome is required for estrogen receptor-alpha and coactivator turnover and for efficient estrogen receptor-alpha transactivation. *Mol Cell* 2000; 5: 939-948.
- [19] Reid G, Hubner MR, Metivier R, Brand H, Denger S, Manu D, Beaudouin J, Ellenberg J, Gannon F. Cyclic, proteasome-mediated turnover of unliganded and liganded ERalpha on responsive promoters is an integral feature of estrogen signaling. *Mol Cell* 2003; 11: 695-707.
- [20] Gallo D, Jacquot Y, Cleeren A, Jacquemotte F, Laios I, Laurent G, Leclercq G. Molecular basis of agonistic activity of ER alpha 7p, a synthetic peptide corresponding to a sequence located at the N-terminal part of the estrogen receptor alpha ligand binding domain. *Letters in Drug Design & Discovery* 2007; 4: 346-355.
- [21] von RM, Schlenk M, Gust R. Effects of C2-alkylation, N-alkylation, and N,N'-dialkylation on the stability and estrogen receptor interaction of (4R,5S)/(4S,5R)-4,5-bis(4-hydroxyphenyl)-2-imidazolines. *J Med Chem* 2004; 47: 915-927.
- [22] Agouridas V, Blazejewski JC, Cleeren A, Laios I, Leclercq G, Magnier E. Fluorous tolerance of the estrogen receptor alpha as probed by 11-polyfluoroalkylestradiol derivatives. *Steroids* 2008; 73: 320-327.
- [23] Maaroufi Y, Leclercq G. Importance of A/B and C domains of the estrogen receptor for its adsorption to hydroxylapatite. *J Steroid Biochem Mol Biol* 1994; 48: 155-163.
- [24] Pons M, Gagne D, Nicolas JC, Mehtali M. A new cellular model of response to estrogens: a bioluminescent test to characterize (anti) estrogen molecules. *Biotechniques* 1990; 9: 450-459.
- [25] Stoessel S, Leclercq G. Competitive binding assay for estrogen receptor in monolayer culture: measure of receptor activation potency. *J Steroid Biochem* 1986; 25: 677-682.
- [26] Jacquot Y, Laios I, Cleeren A, Nonclercq D, Bermont L, Refouvelet B, Boubekur K, Xicluna A, Leclercq G, Laurent G. Synthesis, structure, and estrogenic activity of 4-

- amino-3-(2-methylbenzyl)coumarins on human breast carcinoma cells. *Bioorg Med Chem* 2007; 15: 2269-2282.
- [27] Celik L, Lund JD, Schiott B. Conformational dynamics of the estrogen receptor alpha: molecular dynamics simulations of the influence of binding site structure on protein dynamics. *Biochemistry* 2007; 46: 1743-1758.
- [28] Bourgoin-Voillard S, Zins EL, Fournier F, Jacquot Y, Afonso C, Pèpe C, Leclercq G, Tabet JC. Stereochemical effects during [M-H]⁻ dissociations of epimeric 11,17 β -estratriols and distant electronical effects of substituents at C₍₁₁₎ position on gas phase acidity. *J Am Soc Mass Spectrom* 2009; doi:10.1016/j.jasms.2009.08.017.
- [29] Laios I, Cleeren A, Leclercq G, Nonclercq D, Laurent G, Schlenk M, Wellner A, Gust R. Effects of (R,S)/(S,R)-4,5-bis(2-chloro-4-hydroxyphenyl)-2-imidazolines and (R,S)/(S,R)-2,3-bis(2-chloro-4-hydroxyphenyl)piperazines on estrogen receptor alpha level and transcriptional activity in MCF-7 cells. *Biochem Pharmacol* 2007; 74: 1029-1038.
- [30] Alarid ET. Lives and times of nuclear receptors. *Mol Endocrinol* 2006; 20: 1972-1981.
- [31] Leclercq G, Lacroix M, Laios I, Laurent G. Estrogen receptor alpha: impact of ligands on intracellular shuttling and turnover rate in breast cancer cells. *Curr Cancer Drug Targets* 2006; 6: 39-64.
- [32] Legros N, Leclercq G, McCague R. Effect of estrogenic and antiestrogenic triphenylethylene derivatives on progesterone and estrogen receptors levels of MCF-7 cells. *Biochem Pharmacol* 1991; 42: 1837-1841.
- [33] El Khissiin A., Cleeren A, Borrás M, Leclercq G. Protein synthesis is not implicated in the ligand-dependent activation of the estrogen receptor in MCF-7 cells. *J Steroid Biochem Mol Biol* 1997; 62: 269-276.
- [34] El Khissiin A, Leclercq G. Implication of proteasome in estrogen receptor degradation. *FEBS Lett* 1999; 448: 160-166.
- [35] Marsaud V, Gougelet A, Maillard S, Renoir JM. Various phosphorylation pathways, depending on agonist and antagonist binding to endogenous estrogen receptor alpha (ERalpha), differentially affect ERalpha extractability, proteasome-mediated stability, and transcriptional activity in human breast cancer cells. *Mol Endocrinol* 2003; 17: 2013-2027.
- [36] Agouridas V, Magnier E, Blazejewski JC, Laios I, Cleeren A, Nonclercq D, Laurent G, Leclercq G. Effect of fluorination on the pharmacological profile of 11beta isomers of fulvestrant in breast carcinoma cells. *J Med Chem* 2009; 52: 883-887.
- [37] Tedesco R, Katzenellenbogen JA, Napolitano E. 7 alpha,11 beta-disubstituted estrogens: Probes for the shape of the ligand binding pocket in the estrogen receptor. *Bioorganic & Medicinal Chemistry Letters* 1997; 7: 2919-2924.

- [38] Seo JW, Comninos JS, Chi DY, Kim DW, Carlson KE, Katzenellenbogen JA. Fluorine-substituted cyclofenil derivatives as estrogen receptor ligands: synthesis and structure-affinity relationship study of potential positron emission tomography agents for imaging estrogen receptors in breast cancer. *J Med Chem* 2006; 49: 2496-2511.
- [39] Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL. The structural basis of estrogen receptor/coactivator recognition and the antagonism of this interaction by tamoxifen. *Cell* 1998; 95: 927-937.
- [40] Fowler AM, Alarid ET. Dynamic control of nuclear receptor transcription. *Sci STKE* 2004; e51.
- [41] Metivier R, Penot G, Hubner MR, Reid G, Brand H, Kos M, Gannon F. Estrogen receptor- α directs ordered, cyclical, and combinatorial recruitment of cofactors on a natural target promoter. *Cell* 2003; 115: 751-763.
- [42] Hammes SR, Levin ER. Extranuclear steroid receptors: nature and actions. *Endocr Rev* 2007; 28: 726-741.
- [43] Kampa M, Pelekanou V, Castanas E. Membrane-initiated steroid action in breast and prostate cancer. *Steroids* 2008; 73: 953-960.
- [44] Seo HS, Leclercq G. Evaluation of potential implication of membrane estrogen binding sites on ERE-dependent transcriptional activity and intracellular estrogen receptor- α regulation in MCF-7 breast cancer cells. *J Steroid Biochem Mol Biol* 2002; 80: 109-123.
- [45] Taguchi Y, Koslowski M, Bodenner DL. Binding of estrogen receptor with estrogen conjugated to bovine serum albumin (BSA). *Nucl Recept* 2004; 2: 5.
- [46] Wessler S, Otto C, Wilck N, Stangl V, Fritzemeier KH. Identification of estrogen receptor ligands leading to activation of non-genomic signaling pathways while exhibiting only weak transcriptional activity. *J Steroid Biochem Mol Biol* 2006; 98: 25-35.
- [47] Harrington WR, Kim SH, Funk CC, Madak-Erdogan Z, Schiff R, Katzenellenbogen JA, Katzenellenbogen BS. Estrogen dendrimer conjugates that preferentially activate extranuclear, nongenomic versus genomic pathways of estrogen action. *Mol Endocrinol* 2006; 20: 491-502.

FIGURE LEGENDS

Figure 1. Stabilizing interactions in (a) the type I estrogen/ER α complex (b) the type II estrogen/ER α complex.

Figure 2. Relationship between the capacity of E₂ to compete with [³H]E₂ for binding to purified hER α recombinant and the binding of the receptor to the LxxLL-coated plate. hER α recombinant was incubated with increasing amounts of E₂ and processed for each assay as described in the methods section.

Figure 3. Influence of substitution onto E₂ on its capacity to enhance the binding of hER α recombinant to the LxxLL-coated plate. hER α recombinant was incubated with increasing amounts of compounds and processed for LxxLL binding assay. Hydroxyl substitutions: Panel on the left (a) refers to 3-OH, panel on the right (b) to the 17 β -OH. Substitutions at C-11: Panels refer to the linkage of (c) a polar group, (d) an hydrophobic group or (e) a long side chain. Panel (f) refers to the introduction of a double bond in C₉-C₁₁ onto the steroid.

Figure 4. Capacity of non steroidal type I estrogens to enhance the binding of hER α recombinant to the LxxLL-coated plate. hER α recombinant was incubated with increasing amounts of compounds and processed for LxxLL binding assay.

Figure 5. Capacity of angular type II ligands to modulate the binding of hER α recombinant to the LxxLL-coated plate. hER α recombinant was incubated with increasing amounts of compounds and processed for LxxLL binding assay. Upper part: The panel on the left (a) refers to “*cis* diphenolic” compounds, the panel on the right (b) to “*gem* diphenolic” compounds. The figures evidence that former compounds are agonists while the latter are antagonists. Lower part (c): Strong antagonists (# 21 - 23) at 100 nM abrogate the promoting effect of 1nM E₂ while weak antagonists (# 24, 25) display a moderate inhibition, agonists (# 18 - 20) are ineffective.

Figure 6. Capacity of E₂ to modulate the LxxLL motif recruitment in MCF-7 cells. (a) Kinetics of recruitment: Cells were incubated with 0.1 μ M E₂ for various periods (from 15 min to 4 hours; insert: for 5 to 60 min). (b) Dose-effect study: Cells were treated for 15 min with increasing concentrations of E₂ (from 1 nM to 10 μ M). Values are expressed as percentage of untreated cells (control). Figure refers to an experiment performed (a) twice and (b) five times (identical results).

Figure 7. Effect of E₂ and C-11 β derivatives with strong LxxLL recruitment potency on ER α level in MCF-7 cells. Western blotting data established with extracts from cells incubated for 4 hours with the compounds were normalized according to actin and expressed in percentage of ER α level in control, untreated cells. Panel (a) refers to increasing concentrations of E₂ and (b) to C-11 β derivatives at 10 nM.

Figure 8. Demonstration by immunofluorescence that bisphenol suppresses the down regulation of ER α induced by E₂. MCF-7 cells were exposed to drugs for a period of 6 hours and processed for immunofluorescence microscopy as described in Methods. (a): no drug, (b): 1 nM E₂, (c): 100 nM bisphenol, (d): E₂ and bisphenol in combination. ER α was revealed by using polyclonal antibody HC-20 raised against the receptor. Texas Red labeling.

Figure 9. Influence of (a) steroidal and (b) nonsteroidal ligands on ERE-dependent transcription. MVLN cells were incubated for 24 hours with increasing amounts of a set of representative ligands varying in their ability to modulate the binding of ER α in our LxxLL binding assay. (— : agonists; - - - : no effect; - - - : antagonists). Luciferase activity was then assayed in cell extracts as described in methods.

Figure 2

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Table I

Relative efficiency of steroidal ligands to associate with the E₂ binding site and induce the recruitment of coregulators containing a LxxLL motif.

N°	Structure	E ₂ binding (RBA)*	LxxLL recruitment (RRC)*		
17β estradiol derivatives

	R₃	R₁₇	R₁₁		
1 (E ₂)	OH	OH	H	100	100
2	O-CH ₃	OH	H	3	10
3	O- SO ₂ -O ⁻ Na ⁺	OH	H	<<0.1	0
4	OH	H	H	10	<<0.01
5	OH	OH (α)	H	30	60
6α	OH	OH	OH (α)	0.3	1
6β	OH	OH	OH (β)	10	4000**
7	OH	OH	O-CO-CH ₃	30	800
8	OH	OH	CH ₃	100	1300
9	OH	OH	CH ₂ -Cl	100	600
10	OH	OH	C≡CH	30	400
11	OH	OH	C ₍₉₎ = C ₍₁₁₎	10	200
12	OH	OH		10	Antagonist
13	OH	OH		10	Antagonist
14	OH	OH		3	No effect

*Ratio of concentrations of E₂ and investigated compounds to produce 50 % of maximal effect. Value are expressed in percentages of E₂ efficiency to compete with [³H]E₂ for ERα binding (RBA) or confer a conformation to ERα appropriate for binding to a plate coated with a peptide containing a LxxLL motif (RRC).

**Value most probably overestimated due to the low optimal RC value (see Figure 6).

Table II

Relative efficiency of nonsteroidal ligands to associate with the E₂ binding site and induce the recruitment of coregulators containing a LxxLL motif.

N°	Structure	E ₂ binding (RBA)*	LxxLL recruitment (RRC)*	
1 (E ₂)		100	100	
a. Non steroidal “linear” estrogens (Type I)				
15		50	200	
16		1	1	
17		1	0.1	
b. Angular diphenolic estrogens (Type II)				
18		<0.01	0.1	
19		<0.01	<<0.1	
20		<0.01	<<0.1	
<hr/>				
	R			
21		-H	10	Antagonist
22		-CH ₃	10	Antagonist
23		-CH ₂ CH ₃ (bisphenol)	10	Antagonist
24		-CH ₂ CH ₂ CH ₃	3	Antagonist
25			5	Antagonist

*Ratio of concentrations of E₂ and investigated compounds to produce 50 % of maximal effect. Value are expressed in percentage of E₂ efficiency to compete with [³H]E₂ for ER α binding (RBA) or confer a conformation to ER α appropriate for binding to a plate coated with a peptide containing a LxxLL motif (RRC).

Table III

Relationship between ligand-induced recruitment of coregulators containing a LxxLL motif, ER α downregulation and ERE-dependent transcription

Ligand	RBA ^a	LxxLL-recruitment ^b	ER α downregulation ^c	ERE-dependent transcription ^d
<u>Steroidal</u>				
		<i>Minimal concentration (Log C) producing ~50 % change</i>		
1 (E₂)	100	-7	-10	-11
9	100	-10 / -9	-10	-12 / -11
8	100	-10 / -9	-10	-11
10	100	-9	-10	-11
11	40	-9 / -8	-9	-10
6 β	4	> -5	-10	-11
14	1	no effect	-8	-9 / -8
12	10	antagonism	upregulation at -8	weak increase at -10 / no effect at -8
<u>Non steroidal</u>				
15	80	-7	-10	-11 / -10
18	<0.01	-6	-9	-9 / -8
16	3	no effect	-8	-9 / -8
17	1	no effect	-7	-7
23	2	antagonism	weak decrease at -9 / no effect at -7	-9
25	0.1	antagonism	-6	-7

^a Binding affinity for ER α ("whole cell assay") established by measuring the capacity of a ligand to compete with [³H]E₂ for uptake in MCF-7 cells.

^b Ability of ligands to confer an ER α conformation for a LxxLL motif containing coactivators in MCF-7 cells. NR peptide assay performed on cell extracts from cells preincubated for 15 min with the ligands.

^c Downregulation established by Western blotting from MCF-7 cells treated with ligands for 4 hours.

^d Ligand-induced luciferase induction in MVLN cells.