Fish-farm impact on metazoan meiofauna in the Mediterranean Sea: analysis of regional vs. habitat effects

Simone Mirto, Silvia Bianchelli, Cristina Gambi, Maja Krzelj, Antonio Pusceddu, Mariaspina Scopa, Marianne Holmer, Roberto Danovaro

► To cite this version:

Simone Mirto, Silvia Bianchelli, Cristina Gambi, Maja Krzelj, Antonio Pusceddu, et al.. Fish-farm impact on metazoan meiofauna in the Mediterranean Sea: analysis of regional vs. habitat effects. *Marine Environmental Research*, 2009, 69 (1), pp.38. 10.1016/j.marenvres.2009.07.005 . hal-00547650

HAL Id: hal-00547650

<https://hal.science/hal-00547650>

Submitted on 17 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Fish-farm impact on metazoan meiofauna in the Mediterranean Sea: analysis of regional vs. habitat effects

Simone Mirto, Silvia Bianchelli, Cristina Gambi, Maja Krzelj, Antonio Pusceddu, Mariaspina Scopa, Marianne Holmer, Roberto Danovaro

PII: S0141-1136(09)00104-4
DOI: [10.1016/j.marenvres.2009.07.005](https://doi.org/10.1016/j.marenvres.2009.07.005)
Reference: MERE 3361

To appear in: *Marine Environmental Research*

Received Date: 2 February 2009
Revised Date: 24 July 2009
Accepted Date: 28 July 2009

Please cite this article as: Mirto, S., Bianchelli, S., Gambi, C., Krzelj, M., Pusceddu, A., Scopa, M., Holmer, M., Danovaro, R., Fish-farm impact on metazoan meiofauna in the Mediterranean Sea: analysis of regional vs. habitat effects, *Marine Environmental Research* (2009), doi: [10.1016/j.marenvres.2009.07.005](https://doi.org/10.1016/j.marenvres.2009.07.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Fish-farm impact on metazoan meiofauna in the Mediterranean Sea:
analysis of regional vs. habitat effects**

Simone Mirto^{a,b}, Silvia Bianchelli^a, Cristina Gambi^a, Maja Krzelj^{a,c}, Antonio Pusceddu^a,
Mariaspina Scopa^a, Marianne Holmer^d, Roberto Danovaro^{a*}

^a *Department of Marine Sciences, Polytechnic University of Marche, Via Brecce Bianche, 60131 Ancona, Italy*

^b *Institute for the Marine Coastal Environment, National Council of Research, Spianata S. Raineri, 86 - 98122 Messina, Italy*

^c *Center of Marine Studies, University of Split, Livanjska 5/III, 21000 Split, Croatia*

^d *Institute of Biology, University of Southern Denmark, Campusvej 55, DK5230, Odense M., Denmark*

Submitted to
Marine Environmental Research

*** Corresponding author:**

Prof. Roberto Danovaro
Dipartimento Scienze del Mare,
Università Politecnica delle Marche,
Via Brecce Bianche, 60131 Ancona, Italy
Tel: +390712204654
Fax: +390712204560
r.danovaro@univpm.it

ABSTRACT:

The worldwide exponential growth of off-shore mariculture is raising severe concerns about the impacts of this industry on marine habitats and their biodiversity. We investigated the metazoan meiofaunal response to fish farm impact in four regions of the Mediterranean Sea. Meiofaunal assemblages were investigated in two habitats (seagrass meadows of *Posidonia oceanica* and non-vegetated soft bottoms) comparing sites receiving faeces and uneaten food pellets from fish farms to control sites. We report here that, consistently across different regions, the meiofaunal abundance typically responded positively to fish-farm effluents. Biodeposition caused also significant changes in assemblage structure and the reduction in the richness of higher meiofaunal taxa, but the multivariate analysis of variance revealed that the effects were region- and habitat-specific. In non-vegetated systems, three of the four regions investigated displayed significant effects of the fish farms on richness of meiofaunal taxa. In vegetated habitats, meiofauna did not respond to biodeposition (except in one region), suggesting that seagrass meadows can mask the effects of fish-farm effluents on benthic biodiversity. We conclude that different indicators of fish-farm impact are needed in vegetated and non-vegetated benthic systems.

Key-words: fish-farm impact, meiofauna, soft sediments, *Posidonia oceanica*, Mediterranean Sea.

1. Introduction

Eutrophication is recognized as one of the most important emerging problems of the coastal oceans and, during the past four decades, has increased exponentially in intensity, geographical area and environmental consequences (Cloern, 2001). Eutrophication is typically related to the increase of nutrient and organic matter loads, which cause a progressive decrease of dissolved oxygen concentrations (Dell'Anno et al., 2008). The accumulation of large amounts of organic matter in coastal sediments, associated with eutrophication processes, also induce significant changes in the composition of sediment organic matter, which can have putative effects on the benthos (see Pusceddu et al., 2009 for a review).

Aquaculture is a fast-growing industry which, through the release of organic and inorganic N and P contributes to the progressive eutrophication of coastal areas (see Holmer et al., 2008 and citations therein). Aquaculture activities are now relevant at local and regional scales, and recent estimates indicate that, in Mediterranean coastal areas, the release of nutrients from fish farming contributes for up to 7 and 10% of N and P total discharge, respectively (Pitta et al., 1999).

Aquaculture installations can produce relevant shifts of the whole natural environment (Boyra et al., 2004; Machias et al., 2004), threatening the environmental quality of coastal zones (Gowen and Bradbury, 1987) and generating conflicts between aquaculture and the conservation of marine habitats, including the protection of large benthic primary producers, such as the seagrass *Posidonia oceanica*.

The organic enrichment of the sediments immediately beneath the sea cages is a direct result of the sedimentation of particulate waste products from the fish farm (Hargrave et al., 1997; Karakassis et al., 1998; Holmer et al., 2008) and a decreasing concentration with the increasing distance from the point source is typically observed (Hargrave et al., 1993; Mazzola

et al., 1999; Pusceddu et al., 2007; Holmer et al., 2007). The continuous flow of faeces and food pellets from fish cages alters the quantity and the biochemical composition of sediment organic matter, but with potentially different effects in different regions and/or habitats.

In fact, previous studies, carried out in non-vegetated and seagrass habitats of four Mediterranean regions (Cyrus, Greece, Italy and Spain), have shown that the response of the benthic biochemistry to aquaculture biodeposition is idiosyncratic, and significant increase in the organic load can be detected only in those regions characterized by biopolymeric C contents typically, 2 mg C g⁻¹ (Pusceddu et al., 2007).

To identify the changes induced by aquaculture activities on benthic ecosystems, we investigated meiofaunal assemblages in different regions of the Mediterranean Sea along a longitudinal gradient of trophic conditions (in terms of primary productivity, Pusceddu et al., 2007). Metazoan meiofauna, for their ecological importance in the benthic ecosystems and the lack of larval dispersion, are considered a sensitive tool for investigating structural and functional changes of natural and anthropogenically-impacted ecosystems (see Vincx and Heip, 1987 for a review).

Here we tested, using a hierarchical sampling strategy under uni- and multi-variate contexts, the hypothesis that the presence of fish farms influences significantly the meiofaunal assemblages in terms of abundance, community structure and diversity (i.e. richness and evenness of taxa) and that different patterns can be expected in the four regions according to their different trophic backgrounds.

2. Materials and Methods

2.1 Study site and sampling strategy

To cover different environmental conditions characterizing Mediterranean coastal zones along an east-to-west longitudinal transect (ca. 3,500 km wide), four regions were selected: Akrotiri Bay (Cyprus; 34° 39' N, 34° 04' E; July, 2002), Sounion Bay (southern Greece; 37°39' N, 24°01' E; July, 2003), Pachino Bay (Italy; 36°43' N, 15°05'E; September, 2002), and Gulf of Alicante (Spain; 38° 24' N, 0° 24' W; September, 2003) (Figure 1). All of these regions, located at similar latitudes and depths were selected on the basis of the presence of the fish farms, which have been previously characterized in terms of their main environmental features (Table 1). In each of the sampling regions, the effects of the fish farm on the meiofauna were investigated in two different habitats: meadows of the seagrass *Posidonia oceanica*, and soft non- vegetated bottoms. In each region, a preliminary survey was carried out to ascertain the presence of both the seagrass and the soft substrates, and to characterize the environmental settings of the areas in terms of mean depth and temperature, bottom currents, sediment type and porosity, and chlorophyll-a and inorganic nutrient concentrations in the water column (Karakassis et al., 2005). In each habitat, the impact was quantified by contrasting the fish-farm sites with control sites characterized by relatively pristine conditions and by environmental features comparable to those found beneath the cages (Table 1). Controls were located upstream of the main currents, and at least 1,000 m from the fish farms. It has been demonstrated that this distance avoids any potential effect of fish farms (Holmer et al., 2008) and allows maintaining the spatial variability in background (control) conditions as low as possible.

At each site, three replicates were collected randomly from the central area of each fish-farm site (i.e. beneath the cages) and in each control site, by means of manual corers (diameter 3.7 cm, 10.7 cm² surface area, down to a depth of 10 cm) operated by SCUBA divers.

2.2 Sediment biochemistry

Sediment protein, carbohydrate and lipid contents were determined spectrophotometrically and concentrations were converted into carbon equivalents using 0.40 and 0.49 and 0.75 g C g⁻¹ conversion factors, respectively, and their sum reported as biopolymeric C (Pusceddu et al., 2009). Data relative to the quantity and biochemical composition (i.e. protein, carbohydrate and lipid contents) of organic matter in the sediments under scrutiny have been detailed elsewhere (Pusceddu et al., 2007). In this study, we focused our attention on the biopolymeric C concentrations and on the values of the protein to carbohydrate ratio as descriptors of the quantity and quality of sediment organic matter, respectively (Pusceddu et al., 2009).

2.3 Meiofaunal analyses

Each sediment sample was fixed with 4% buffered formaldehyde (in filtered seawater solution) and was sieved through 1000 µm sieve (to retain macrobenthos and macroalgae) and 32 µm sieve (to retain smaller meiofauna) in laboratory. The sample fraction retained by a 32 µm mesh net was added to Ludox HS 40 (density arranged to 1.18 g cm⁻³), for density centrifugation extraction (10 min, 800 × g, for 3 times) from the sediment (Heip et al., 1985). All metazoan animals were counted and classified per taxon under a stereomicroscope using Delfuss cuvettes, after staining with Rose Bengal (0.5 g L⁻¹). Community evenness was calculated as the Pielou's index using the different meiofaunal taxa as entries, by means of the PRIMER software (Clarke and Gorley, 2003).

2.4 Statistical analyses

Differences between control and impact sediments were assessed for all of the investigated variables using a three-way analysis of variance (ANOVA). The analysis treated the factor region

(R, 4 levels) as random, habitat (H, 2 levels) as fixed and crossed with R, and impact (I, 2 levels) as fixed and orthogonal to R and H. When significant differences ($p < 0.05$) were observed, a post-hoc Student-Newman-Kuels' test (SNK) was also performed. ANOVA and SNK tests were carried out using the GMAV software (University of Sidney).

Distance-based permutational multivariate analyses of variance (PERMANOVA, Anderson, 2001) were used to quantify in the two benthic habitats the effects of the fish farms on i) the organic matter quantity and quality and ii) the meiofaunal assemblages. The analysis treated the factor region (R, 4 levels) as random, habitat (H, 2 levels) as fixed and crossed with R, and impact (I, 2 levels) as fixed and orthogonal to R and H. For organic matter, the data set included 48 observations on concentrations of biopolymeric C and the values of the protein to carbohydrate ratio. For meiofauna, the data set included 48 observations and the abundance of all the taxa. Since the interaction term $R \times H \times I$ was seen to be significant either for organic matter and meiofauna, pairwise comparisons, which also used 499 random permutations to obtain P-values, were also carried out to ascertain differences between control and impact sediments in all regions, separately. The P-values were calculated using 499 Monte Carlo draws from the asymptotic permutation distribution (Anderson and Robinson, 2003). PERMANOVA tests were carried out (either in the uni- and multivariate contexts) also removing the effects of the main environmental variables (i.e., water depth, current speed, sediment type, porosity and water content, biopolymeric C and protein to carbohydrate ratio) as covariates.

In order to assess whether and how much environmental variables explained changes in meiofaunal abundance, richness and evenness of taxa, a non-parametric multivariate multiple regression analysis that was based on Euclidean distances was carried out using the routine DISTLM forward (Mc Ardle and Anderson, 2001). The forward selection of the predictor variables was carried out with tests by permutation. P values were obtained using 4,999

permutations of raw data for the marginal tests (tests of individual variables), while for all of the conditional tests, the routine uses 4,999 permutations of residuals under a reduced model.

3. Results

Three-way univariate ANOVA revealed that biopolymeric C concentrations and values of the protein to carbohydrate ratio displayed a significant Region (R) $\times$ Habitat (H) $\times$ Impact (I) effect (Table 2), with differences between impacted and control sites varying between the two habitats and the four geographical areas. In particular, the SNK tests revealed that biopolymeric C concentrations in farm-impacted sediments increased consistently in vegetated sediments in Cyprus, in both habitats in Greece and in non-vegetated sediments in Italy (Figure 2A), whereas values of the protein to carbohydrate ratio increased in impact sites in non-vegetated sediments in Cyprus, in both habitats in Italy and in non-vegetated sediments in Spain and decreased in impact sediments of both habitats in Greece (Figure 2B).

A general increase of meiofaunal abundance was observed in impacted sediments (Figure 3A) and the ANOVA revealed significant R $\times$ H $\times$ I variations in total meiofaunal abundance (Table 3). Moreover, the SNK tests indicated that total meiofaunal abundance increased significantly in impact sites in vegetated sediments in Cyprus and non-vegetated sediments in Italy, whereas no significant differences between impact and control sediments were observed in all other regions and habitats (Figure 3A).

Abundance of all of the meiofaunal taxa in the sediments are reported in Table 4. ANOVA revealed significant R $\times$ H $\times$ I effects on the abundance of all of the meiofaunal taxa with differences between impact and control sites differing in extent and direction in different regions and habitats and for the different taxa (Table 5). Significant interaction of the R $\times$ H $\times$ I factors, indicating idiosyncratic responses to fish-farm impact in different regions and habitats,

was observed for polychaetes, gastrotrichs, bivalves, cumaceans and tanaidaceans. The SNK tests (data not shown) revealed increasing nematode abundance beneath the cages in vegetated sediments in Cyprus and in both habitats in Greece. Copepods increased in both fish-impacted habitats in Cyprus. The abundance of ostracods varied between impacted and control sites only in Spain (in both habitats), whereas oligochaetes, kinorhynchs and amphipods did not display any significant difference between impact and control sediments in any of the investigated region and habitat. Turbellarians' abundance exhibited significant differences only between regions.

The PERMANOVA test applied on the meiofaunal community composition showed a significant $R \times H \times I$ effect, indicating that the differences between control and fish-farm impacted sites varied between the two habitats and the four regions (Table 6, Figure 3B-C). The analysis carried out separately for all of the four regions and the pairwise comparison tests revealed that differences between impact and control sites in the meiofaunal community composition were significant only in non-vegetated sediments either in Cyprus and Greece.

At all sites nematodes and copepods were the most dominant taxa, followed by polychaetes, ostracods, turbellarians, oligochaetes, gastrotrichs and all other taxa. The richness of meiofaunal taxa decreased significantly in impacted sites in all non-vegetated sediments, whereas no significant differences between impact and control sites were observed in seagrass sediments (Table 7 and Figure 4A). On the other hand, significant differences in community evenness were observed between regions and between impact and control sites only in non-vegetated sediments in Cyprus (Table 7, Figure 4B).

4. Discussion

4.1 Impacts of fish-farm effluents on meiofaunal abundance

Ecosystem alterations induced by different typologies of disturbance, including human impacts, generate variable shifts in benthic species, assemblages and community composition. Fish farming, by modifying the whole attributes of the benthic environment beneath the cages, produces relevant modifications in the abundance, biomass, species composition and evenness of meio- and macrofauna (e.g. Mirto et al., 2002; Kalantzi and Karakassis, 2006).

The responses of metazoan meiofauna to various ecosystem alterations are clearly detected at the highest taxonomic level, with a resolution similar to that provided by the analysis of lower taxonomic levels (Kennedy and Jacoby, 1999; Warwick 1988), also under the influence of fish farms (Sutherland et al. 2007).

In this study, we analysed the potential effects of fish farm effluents on the abundance and community composition of meiofauna, by comparing, for the first time, two different habitats in four different regions with different background trophic conditions. The general outcome from the available literature is that fish-farm effluents typically alter meiofaunal abundance, diversity, biomass, and species composition (Duplisea and Hargrave, 1996; Kennedy and Jacoby, 1999; Mirto et al., 2000; -2002; La Rosa et al., 2001; Sutherland et al., 2007).

However, changes associated with the presence of fish-farm effluents are often not consistent, as meiofaunal abundance may either increase or decrease beneath the fish cages, depending on the site or the farm characteristics.

In the present study, the idiosyncratic meiofaunal responses to fish farm impact was confirmed. Although total meiofaunal abundance was generally higher in the fish-farm than in the control sediments, such changes were not always statistically significant. This result is apparently in contrast to previous investigations reporting a decrease of meiofaunal abundance in

non-vegetated systems subjected to bio-deposition from fish cages (Mirto et al., 2002; Sutherland et al., 2007). The general positive response of meiofaunal abundance to fish-farm biodeposition could be related to the relatively limited organic enrichment in the sediments beneath the cages, as reported in sediments subjected to mussel farm bio-deposition, where the organic inputs to the bottom were similarly low (Danovaro et al., 2004).

The results of the PERMANOVA, carried out in the univariate context, revealed that differences in total meiofaunal abundance between impact and control sites were due to the effects of the environmental covariates (Table 8). The effect of organic matter enrichment in fish-farm sediments was significant only in oligotrophic regions (i.e., Cyprus and Greece). The multiple regression analysis, indeed, revealed that, among the different environmental variables, only the current speed and the biopolymeric C concentrations explained significant proportions (15% and 9%, respectively) of the total meiofaunal abundance (Table 9). This result supports the expectation that changes in meiofaunal abundance in fish-farm sediments are linked to the organic matter inputs released by the fish farm, as well as to the local hydrodynamism.

The results of this study suggest that the impact of aquaculture effluents on the meiofaunal abundance is both region- and habitat-specific and also highlight that the direction and intensity of the response of total meiofaunal abundance to aquaculture effluents depend on the environmental conditions.

4.2 Impacts of fish-farm effluents on meiofaunal communities

The results presented here provide evidence that, in all investigated regions, consistent significant effects of fish-farming effluents on meiofaunal biodiversity can only be detected in non-vegetated sediments. In all soft bottoms beneath fish farms the richness of higher taxa decreased significantly when compared to control sites (from 13-16 in all control sediments to a

minimum of 9 taxa in non-vegetated impact sediments in Spain), whereas no significant differences were observed in terms of taxa evenness. The taxa that disappeared beneath the cages changed amongst regions, but always included the rare taxa (i.e., taxa representing <1% of the total meiofaunal abundance; Figure 3C) of the control sites. Meiofaunal taxa disappeared in fish-farm sediments included kinorhynchs (only in non-vegetated sediments in Spain), cumaceans (in 37.5% of the total cases, i.e., in non-vegetated sediments in Cyprus, Italy and Spain), isopods (37.5%, in non vegetated sediments in Cyprus and Greece and in vegetated sediments in Spain), thanaidaceans (37.5% in non-vegetated sediments in Cyprus and both habitats in Greece), amphipods (50%, in non-vegetated sediments in Greece, Italy and Spain and in vegetated sediments in Italy), and tardigrades (25% of the total cases: in non-vegetated sediments in Cyprus and Spain). These results do not allow identifying specific taxa as indicators of biodeposition impact.

The analysis of meiofaunal assemblage attributes (meiofaunal abundance, community composition, richness of taxa and community evenness) within seagrass sediments revealed that all of these variables responded idiosyncratically to the presence of fish farming activities. The lack of a clear and consistent meiofaunal response to the fish-farm biodeposition in seagrass sediments could have several possible explanations. *Posidonia oceanica* is a seagrass endemic of the Mediterranean Sea, which covering ca 50.000 km², plays a key ecological role for a wide range of organisms and assemblages, preserving their biodiversity and protecting the integrity of the coastal areas (Hemminga and Duarte, 2000). The effects of fish-farm biodeposition on seagrass meadows can be difficult to detect due to the fact that the presence of the seagrass masks the changes in organic matter concentrations and composition (Pusceddu et al., 2007). However, the presence of a large number of filter feeders and detritus feeders associated with the seagrass bed can also act as a buffer for the organic enrichment, by consuming a large fraction of the

biodeposits (Balata et al., 2008). Moreover, the presence of a large assemblage of prokaryotes, protozoa and meiofauna within the seagrass sediments makes it difficult to detect the presence of significant changes in these components (Danovaro 1996, Bongiorni et al., 2005). *Posidonia oceanica* is known to be highly sensitive to anthropogenic impacts, and it has been demonstrated that, in the long term (i.e. years to decades), the increased sedimentation of waste particles and the accumulation of organic matter lead to the deterioration of the seagrass system (Pergent-Martini et al., 2006). In the regions investigated, the seagrass systems displayed a clear impact and several indicators of deterioration of the *Posidonia* meadows were reported (Holmer et al., 2008). However, the effects of biodeposition on meiofaunal variables were not evident, indicating that in seagrass systems the search for meiofaunal indicators of impact is unnecessary as it provides, if any, a slow or late response to the fish-farm impact.

4.3 Conclusions and perspectives

In this study we showed that fish-farm biodeposition in the Mediterranean Sea can provoke changes in meiofaunal abundance, community structure and the biodiversity at high taxonomic levels. However, the hierarchical sampling conducted here demonstrated that the meiofaunal responses varied among regions and habitats. In all non-vegetated sediments the richness of meiofaunal taxa decreased in response to fish-farm biodeposition, and therefore independently from the background environmental conditions. Conversely, in seagrass sediments, the idiosyncratic response did not allow any clear indicator of impacts from fish-farm bio-deposition to be identified. The results of this study highlight the importance of using different indicators of fish-farm impact in vegetated and non-vegetated systems.

Acknowledgements

The authors thank the EU project MedVeg (n° EU Quality of Life & Management of Living Resources Programme, Contract No. Q5RS-2001-02456.).

References

- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26, 32–46.
- Anderson, M.J., Robinson, R., 2003. Generalised discriminant analysis based on distances. *Australian and New Zealand Journal of Statistics* 45, 301–318.
- Balata, D., Bertocci, I., Piazzzi, L., Nesti, U. 2008. Comparison between epiphyte assemblages of leaves and rhizomes of the seagrass *Posidonia oceanica* subjected to different levels of anthropogenic eutrophication. *Estuarine Coastal and Shelf Science* 79, 533–540.
- Bongiorni, L., Mirto, S., Pusceddu, A., Danovaro, R., 2005. Response of benthic protozoa and thraustochytrid protists to fish farm impact in seagrass (*Posidonia oceanica*) and soft-bottom sediments. *Microbial Ecology* 50, 268–276.
- Boyra, A., Sanchez-Jerez, P., Tuya, F., Espino, F., Haroun R. 2004. Attraction of wild coastal fishes to an Atlantic subtropical cage fish farms, Gran Canaria, Canary Islands. *Environmental Biology of Fishes* 70, 393–401.
- Clarke, K.R., Gorley, R.N., 2003. *PRIMER 5* (Plymouth Routines in Multivariate Ecological Research). Plymouth Marine Laboratory, Plymouth, United Kingdom.
- Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. *Marine Ecology Progress Series* 210, 223–253.
- Danovaro, R., 1996. Detritus-Bacteria-Meiofauna Interactions in a seagrass bed (*Posidonia oceanica*) of the NW Mediterranean. *Marine Biology* 127, 1–13.

- 329 Danovaro, R., Gambi, C., Luna, G.M., Mirto, S., 2004. Sustainable impact of mussel farming in
330 the Adriatic Sea (Mediterranean Sea): evidence from biochemical, microbial and meiofaunal
331 indicators. *Marine Pollution Bulletin* 49, 325-333.
- 332 Dell'Anno, A., Pusceddu, A., Langone, L., Danovaro, R. 2008. Early diagenesis of organic
333 matter in coastal sediments influenced by riverine inputs. *Chemistry and Ecology* 24, 75-85.
- 334 Duplisea, D.E., Hargrave, B.T., 1996. Response of meiobenthic size-structure, biomass and
335 respiration to sediment organic enrichment. *Hydrobiologia* 339, 161-170.
- 336 Gowen, R.J., Bradbury, N.B., 1987. The ecological impact of salmonid farming in coastal waters:
337 a review. *Oceanography and Marine Biology: An Annual Review* 25, 563– 575.
- 338 Hargrave, B.T., Duplisea, D.E., Pfeifer, E., Wildish, D.J., 1993. Seasonal changes in benthic
339 fluxes of dissolved oxygen and ammonium associated with marine cultured Atlantic salmon.
340 *Marine Ecology Progress Series* 96, 249-257.
- 341 Hargrave, B.T., Phillips, G.A., Doucette, L.I., White, M.J., Milligan, T.G., Wildish, D.J.,
342 Cranston, R.E., 1997. Assessing benthic impacts of organic enrichment from marine
343 aquaculture. *Water Air and Soil Pollution* 99, 641-650.
- 344 Heip, C., Vincx, M., Vranken, G. 1985. The ecology of marine nematodes. *Oceanography Marine*
345 *Biology Annual Review* 23, 399-489.
- 346 Hemminga, M.A., Duarte, C.M., 2000. *Seagrass Ecology*, Cambridge University Press,
347 Cambridge.
- 348 Holmer, M., Marbà, N., Diaz-Almela, E., Duarte, C.M., Tsapakis, M., Danovaro, R., 2007.
349 Sedimentation of organic matter from fish farms in oligotrophic Mediterranean assessed
350 through bulk and stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) *Aquaculture* 262, 268-280.
- 351 Holmer, M., Frederiksen, M., Pusceddu, A., Danovaro, R., Mirto, S., Perez, M., Marbà, N.,
352 Duarte, C.M., Diaz-Almela, E., Tsapakis, M., Karakassis, Y., 2008. Effects of fish-farm

- 1
2
3 353 waste on *Posidonia oceanica* meadows: synthesis and provision of management tools.
4
5
6 354 Marine Pollution Bulletin 56, 1618-1629.
7
8 355 Kalantzi, I., Karakassis, I., 2006. Benthic impacts of fish farming: Meta-analysis of community
9
10 356 and geochemical data. Marine Pollution Bulletin 52, 484-493.
11
12
13 357 Karakassis, I., Tsapakis, M., Hatziyanni, E., 1998. Seasonal variability in sediments profiles
14
15 358 beneath fish farm cages in the Mediterranean. Marine Ecology Progress Series 162, 243-252.
16
17 359 Karakassis, I., Pitta, P., Krom, M.D., 2005. Contribution of fish farming to the nutrient loading of
18
19
20 360 the Mediterranean. Scientia Marina 69, 313-321.
21
22 361 Kennedy, A.D., Jacoby, C.A., 1999. Biological indicators of marine environmental health:
23
24
25 362 meiofauna - a neglected benthic component? Environmental Monitoring Assessment 54, 47-
26
27 363 68.
28
29 364 La Rosa, T., Mirto, S., Mazzola, A., Danovaro, R., 2001. Differential responses of benthic
30
31
32 365 microbes and meiofauna to fish-farm disturbance in coastal sediments. Environmental
33
34 366 Pollution 112, 427-434.
35
36 367 Machias, A., Karakassis, I., Labropoulou M., Somarakis, S., Papadopoulou K.N.,
37
38
39 368 Papaconstantinou C., 2004. Changes in wild fish assemblages after the establishment of a
40
41 369 fish farming zone in an oligotrophic marine ecosystem. Estuarine Coastal and Shelf Science
42
43
44 370 60, 771-779.
45
46 371 Mazzola, A., Mirto, S., Danovaro, R., 1999. Initial fish-farm impact on meiofaunal assemblages
47
48
49 372 in coastal sediments of the Western Mediterranean. Marine Pollution Bulletin 38, 1126-
50
51 373 1133.
52
53 374 McArdle, B.H., Anderson, M.J., 2001. Fitting multivariate models to community data: A
54
55
56 375 comment on distance-based redundancy analysis. Ecology 82, 290-297.
57
58 376 Mirto, S., La Rosa, T., Danovaro, R., Mazzola, A., 2000. Microbial and meiofaunal response to

- intensive mussel-farm biodeposition in coastal sediments of the Western Mediterranean. Marine Pollution Bulletin 40, 244-252.
- Mirto, S., La Rosa, T., Gambi, C., Danovaro, R., Mazzola, A., 2002. Nematode community response to fish-farm impact in the western Mediterranean. Environmental Pollution 116, 203-214.
- Pergent-Martini, C., Boudouresque, C.-F., Pasqualini, V., Pergent, G. 2006. Impact of fish farming facilities on *Posidonia oceanica* meadows: a review. Marine Ecology and Evolutionary Perspective 27, 310-319.
- Pitta, P., Karakassis, I., Tsapakis, M., Zivanovic, S., 1999. Natural vs. mariculture induced variability in nutrients and plankton in Eastern Mediterranean. Hydrobiologia 391, 181-194.
- Pusceddu, A., Fraschetti, S., Mirto, S., Holmer, M., Danovaro, R., 2007. Effects of intensive mariculture on sediment biochemistry. Ecological Applications 17, 1366-1378.
- Pusceddu, A., Dell'Anno, A., Fabiano, M., Danovaro, R., 2009. Quantity and bioavailability of sediment organic matter as signatures of benthic trophic status. Marine Ecology Progress Series 375, 41-52.
- Sutherland, T.F., Levings, C.D., Petersen, S.A., Poon, P., Piercey, B., 2007. The use of meiofauna as an indicator of benthic organic enrichment associated with salmonid aquaculture. Marine Pollution Bulletin 54, 1249-1261.
- Vincx, M., Heip, C., 1987. The use of meiofauna in pollution monitoring studies. A review. ICES Journal of Marine Science 16, 50-67.
- Warwick, R. M., 1988. The level of taxonomic discrimination required to detect pollution effects on marine benthic communities. Marine Pollution Bulletin 19, 259-268.

Figure Captions

Figure 1. Sites of the four fish-farms investigated in Cyprus, Italy, Greece and Spain in the Mediterranean Sea.

Figure 2. Biopolymeric C concentrations (A) and values of the protein to carbohydrate ratio (B) in the investigated sediments. According to the SNK tests (at $p < 0.05$), + = increase; - = decrease; ns = not significant. Data are extracted from Pusceddu et al. (2007).

Figure 3. Total meiofaunal abundance (A), meiofaunal community composition (B) and the relative abundance of rare taxa (C) in Cyprus, Greece, Italy and Spain in non-vegetated and vegetated habitats, in control (C) and impact (I) sediments.

Figure 4. Taxa richness (A) and community evenness (B) in Cyprus, Greece, Italy and Spain in non-vegetated and vegetated habitats, in control (C) and impact (I) sediments. The statistical significance of the differences among bars (SNK test: + = increase $p < 0.05$, - = decrease $p < 0.05$, ns = not significant) are also reported.

Table 1 - Environmental characteristics reported in all sampling sites (distance of the sampling sites from the fish farm, water depth, current speed, sediment type, porosity, water content and grain size).

Region	Habitat	Site	Distance from fish farm (m)	Water Depth (m)	Current speed (cm sec ⁻¹)	Sediment type Silt/Clay (%)	Sediment Porosity (%)	Sediment Water Content (%)	Grain Size (Mean Diameter, mm)
Cyprus	non-vegetated	Impact	0-20	30	20-40	80.7	1.2	42.9	0.01
	vegetated	Control	4000 westward	30	20-40	75.9	1.2	44.9	0.02
	Vegetated	Impact	300 northward	20	20-40	68.4	1.2	46.4	0.02
		Control	1000 eastward	20	20-40	2.5	1.1	36.5	0.10
Italy	non-vegetated	Impact	0	22	20	5.7	1.2	41.6	0.47
	vegetated	Control	3200 north-east	22	20	3.9	0.8	13.5	0.84
	Vegetated	Impact	5	22	20	4.6	1.2	46.9	0.45
		Control	1000 northward	22	20	4.8	1.1	33.7	0.90
Greece	non-vegetated	Impact	0	16	6.3	6.7	1.1	29.5	0.38
	vegetated	Control	800 westward	16	6.3	2.2	1.6	18	1.89
	Vegetated	Impact	5	16	6.3	20.2	1.0	27.3	0.33
		Control	1000 southward	15	6.3	13.1	1.1	34.4	0.76
Spain	non-vegetated	Impact	0	29	4.7	7.7	1.1	31.7	0.32
	vegetated	Control	1000 westward	29	4.7	15.6	1.2	42.3	0.43
	Vegetated	Impact	10	29	4.7	16.2	1.1	32.5	0.40
		Control	1000 southward	29	4.7	15.3	1.1	32.4	0.9

Table 2 - Results of ANOVA testing for differences between impact and control sites in quantity and quality of sediment organic matter in different habitats and regions. R = region; H = habitat; I = impact; SS = sum of squares; DF = degrees of freedom; MS = means square; F = F value; P = probability level: *** = $P < 0.001$; ns = not significant.

	Source	SS	DF	MS	F	P
Biopolymeric C	R	38.74	3	12.91	28.63	***
	H	3.54	1	3.54	0.73	ns
	I	4.40	1	4.39	0.53	ns
	R × H	14.53	3	4.84	10.74	***
	R × I	25.01	3	8.34	18.48	***
	H × I	3.99	1	3.99	1.00	ns
	R × H × I	11.99	3	3.99	8.86	***
	Residuals	14.43	32	0.45		
	Total	116.64	47			
Protein to carbohydrate ratio	R	12.83	3	4.28	285.87	***
	H	1.723	1	1.72	2.65	ns
	I	0.05	1	0.05	0.01	ns
	R × H	1.95	3	0.65	43.42	***
	R × I	10.66	3	3.55	237.45	***
	H × I	1.23	1	1.23	5.12	ns
	R × H × I	0.72	3	0.24	16.04	***
	Residuals	0.48	32	0.02		
	Total	29.64	47			

Table 3 - Results of ANOVA testing for differences between impact and control sites in total meiofaunal abundance in different habitats and regions. R = region; H = habitat; I = impact; SS = sum of squares; DF = degrees of freedom; MS = means square; F = F value; P = probability level: *** = $P < 0.001$; ns = not significant.

Source	DF	SS	MS	F	P
R	3	27255106	9085035	11.7511	**
H	1	909986	909986	0.2431	ns
I	1	13917504	13917504	4.4563	ns
R × H	3	11228668	3742889	4.8413	**
R × I	3	9369245	3123082	4.0396	*
H × I	1	347142.6	347142.6	0.1521	ns
R × H × I	3	6848726	2282909	2.9528	*
Residuals	32	24739857	773120.5		
Total	47	94616234			

Table 4 - Meiofaunal taxa abundance (ind 10 cm⁻²) in fish-farm impacted and control sites across the four regions and the two habitats. Reported are average value $\pm$ standard deviation.

Site	Location	Site	Nematoda	Copepoda	Polychaeta	Ostracoda	Turbellaria	Oligochaeta	Gastrotricha	Bivalvia
Cyprus	Non-vegetated	Control	1903 ± 499	192 ± 43	64 ± 14	3 ± 0	9 ± 8	9 ± 6	2 ± 0	1 ± 0
		Impact	1751 ± 365	438 ± 56	69 ± 5	0 ± 0	4 ± 2	53 ± 20	20 ± 17	1 ± 0
	Vegetated	Control	898 ± 157	186 ± 26	52 ± 0	3 ± 2	1 ± 1	8 ± 6	2 ± 2	2 ± 0
		Impact	1919 ± 145	312 ± 65	90 ± 2	5 ± 2	2 ± 2	7 ± 7	9 ± 2	1 ± 0
Greece	Non-vegetated	Control	813 ± 248	595 ± 253	37 ± 13	27 ± 17	16 ± 3	20 ± 6	61 ± 4	1 ± 1
		Impact	3406 ± 1798	757 ± 147	549 ± 217	45 ± 47	23 ± 6	155 ± 73	344 ± 26	1 ± 1
	Vegetated	Control	583 ± 262	582 ± 258	87 ± 53	15 ± 5	7 ± 2	13 ± 12	43 ± 8	86 ± 40
		Impact	1218 ± 201	784 ± 809	87 ± 103	19 ± 27	18 ± 6	35 ± 16	224 ± 80	0 ± 0
Italy	Non-vegetated	Control	367 ± 69	409 ± 48	52 ± 18	14 ± 3	1 ± 0	2 ± 0	5 ± 1	0 ± 0
		Impact	291 ± 7	394 ± 88	65 ± 13	16 ± 0	5 ± 0	6 ± 4	13 ± 4	1 ± 1
	Vegetated	Control	723 ± 95	284 ± 145	69 ± 42	16 ± 10	3 ± 1	5 ± 2	12 ± 8	0 ± 0
		Impact	938 ± 239	255 ± 39	77 ± 36	21 ± 9	2 ± 0	6 ± 1	28 ± 11	0 ± 0
Spain	Non-vegetated	Control	1467 ± 521	539 ± 63	144 ± 32	6 ± 6	3 ± 4	24 ± 3	150 ± 131	0 ± 0
		Impact	2342 ± 737	305 ± 107	222 ± 108	1 ± 2	1 ± 1	72 ± 52	10 ± 12	0 ± 0
	Vegetated	Control	2084 ± 639	488 ± 186	129 ± 65	7 ± 2	2 ± 3	252 ± 102	11 ± 10	0 ± 0
		Impact	3381 ± 406	432 ± 420	125 ± 60	37 ± 12	4 ± 5	33 ± 27	41 ± 47	0 ± 0
			Kynorhyncha	Tardigrada	Cumacea	Amphipoda	Isopoda	Tanaidacea	Acarina	Nemertina
Cyprus	Non-vegetated	Control	11 ± 3	0 ± 0	0 ± 0	0 ± 1	0 ± 0	1 ± 0	0 ± 0	20 ± 7
		Impact	9 ± 9	0 ± 0	0 ± 0	0 ± 1	0 ± 0	0 ± 0	0 ± 0	5 ± 3
	Vegetated	Control	1 ± 0	0 ± 0	1 ± 0	1 ± 1	1 ± 0	0 ± 0	0 ± 0	4 ± 4
		Impact	3 ± 0	2 ± 2	0 ± 0	1 ± 1	1 ± 0	2 ± 0	1 ± 1	7 ± 7
Greece	Non-vegetated	Control	19 ± 10	9 ± 9	0 ± 0	2 ± 1	0 ± 0	1 ± 1	2 ± 2	46 ± 34
		Impact	22 ± 7	2 ± 2	0 ± 0	0 ± 1	0 ± 0	0 ± 0	0 ± 0	292 ± 160
	Vegetated	Control	15 ± 5	12 ± 9	44 ± 17	0 ± 1	2 ± 0	5 ± 3	0 ± 0	0 ± 0
		Impact	14 ± 5	20 ± 15	2 ± 1	0 ± 1	1 ± 0	0 ± 0	4 ± 3	62 ± 46
Italy	Non-vegetated	Control	2 ± 2	36 ± 4	0 ± 0	1 ± 1	1 ± 0	7 ± 3	7 ± 0	1 ± 1
		Impact	0 ± 0	0 ± 0	0 ± 0	0 ± 1	0 ± 0	0 ± 0	0 ± 0	1 ± 0
	Vegetated	Control	7 ± 1	27 ± 12	1 ± 1	0 ± 1	1 ± 0	3 ± 2	3 ± 2	5 ± 4
		Impact	0 ± 0	14 ± 8	0 ± 0	0 ± 1	0 ± 0	0 ± 0	1 ± 0	5 ± 1
Spain	Non-vegetated	Control	9 ± 7	26 ± 19	1 ± 1	2 ± 1	1 ± 0	1 ± 0	5 ± 3	4 ± 2
		Impact	0 ± 0	0 ± 0	0 ± 0	0 ± 1	0 ± 0	2 ±	0 ± 0	77 ± 56
	Vegetated	Control	8 ± 0	44 ± 26	0 ± 0	2 ± 1	1 ± 0	2 ± 0	1 ± 1	4 ± 2
		Impact	8 ± 0	66 ± 57	0 ± 0	1 ± 1	0 ± 0	2 ± 0	3 ± 3	30 ± 23

Table 5 - Results of ANOVA testing for differences between impact and control sites in abundance of the meiofaunal taxa in different habitats and regions. R = region; H = habitat; I = impact; SS = sum of squares; DF = degrees of freedom; MS = means square; F = F value; P = probability level: *** = $P < 0.001$; ** = $P < 0.01$; * = $P < 0.05$; ns = not significant.

	Source	SS	DF	MS	F	P		Source	SS	DF	MS	F
Nematodes	R	13.15	3	4.38	38.98	***	Turbellarians	R	18.88	3	6.29	12.80
	H	0.06	1	0.06	0.04	ns		H	2.28	1	2.28	3.60
	I	2.66	1	2.66	4.47	ns		I	0.41	1	0.41	1.90
	R × H	4.34	3	1.45	12.86	***		R × H	1.86	3	0.62	1.20
	R × I	1.78	3	0.59	5.29	**		R × I	0.65	3	0.22	0.40
	H × I	0.10	1	0.10	0.36	ns		H × I	0.03	1	0.03	0.00
	R × H × I	0.82	3	0.27	2.42	ns		R × H × I	1.96	3	0.65	1.30
	Residuals	3.60	32	0.11				Residuals	15.66	32	0.49	
	Total	26.50	47					Total	41.73	47		
Copepods	R	1116932.28	3	372310.76	6.23	**	Oligochaetes	R	59266.22	3	19755.41	1.80
	H	17542.98	1	17542.98	1.01	ns		H	56.14	1	56.14	0.00
	I	30403.35	1	30403.35	0.38	ns		I	223.39	1	223.39	0.00
	R × H	52331.00	3	17443.67	0.29	ns		R × H	40553.05	3	13517.68	1.20
	R × I	237589.60	3	79196.53	1.33	ns		R × I	41567.71	3	13855.90	1.20
	H × I	1332.47	1	1332.47	0.11	ns		H × I	34558.65	1	34558.65	3.40
	R × H × I	35031.09	3	11677.03	0.20	ns		R × H × I	29968.09	3	9989.36	0.90
	Residuals	1912625.14	32	59769.54				Residuals	349658.16	32	10926.82	
	Total	3403787.91	47					Total	555851.41	47		
Polychaetes	R	170.08	3	56.69	8.01	***	Gastrotrichs	R	49.57	3	16.52	19.50
	H	39.79	1	39.79	1.39	ns		H	0.66	1	0.66	0.50
	I	96.90	1	96.90	2.63	ns		I	5.48	1	5.48	1.80
	R × H	85.71	3	28.57	4.04	*		R × H	3.67	3	1.22	1.40
	R × I	110.65	3	36.88	5.21	**		R × I	9.05	3	3.02	3.50
	H × I	66.82	1	66.82	1.12	ns		H × I	1.80	1	1.80	0.40
	R × H × I	179.26	3	59.75	8.44	***		R × H × I	12.30	3	4.10	4.80
	Residuals	226.52	32	7.08				Residuals	27.00	32	0.84	
	Total	975.73	47					Total	109.54	47		
Ostracods	R	24.69	3	8.23	16.96	***	Nemertins	R	33.06	3	11.02	23.50
	H	1.53	1	1.53	0.60	ns		H	5.29	1	5.29	0.80
	I	0.01	1	0.01	0.04	ns		I	14.88	1	14.88	2.00
	R × H	7.58	3	2.53	5.21	**		R × H	19.28	3	6.43	13.70
	R × I	0.40	3	0.13	0.28	ns		R × I	22.38	3	7.46	15.90
	H × I	1.75	1	1.75	1.39	ns		H × I	1.25	1	1.25	0.90
	R × H × I	3.77	3	1.26	2.59	ns		R × H × I	4.16	3	1.39	2.90
	Residuals	15.53	32	0.49				Residuals	15.00	32	0.47	
	Total	55.26	47					Total	115.29	47		

	Source	SS	DF	MS	F	P		Source	SS	DF	MS	F	P
Acarines	R	39.94	3	13.31	6.11	***	Isopods	R	1.12	3	0.37	1.77	ns
	H	0.76	1	0.76	0.16	ns		H	2.99	1	2.99	3.12	ns
	I	11.47	1	11.47	0.68	ns		I	3.89	1	3.89	3.91	ns
	R × H	14.28	3	4.76	2.18	ns		R × H	2.87	3	0.96	4.57	**
	R × I	50.90	3	16.97	7.78	**		R × I	2.99	3	1.00	4.75	**
	H × I	50.51	1	50.51	11.65	*		H × I	0.04	1	0.04	0.65	ns
	R × H × I	13.01	3	4.34	1.99	ns		R × H × I	0.17	3	0.06	0.26	ns
	Residuals	69.78	32	2.18				Residuals	6.70	32	0.21		
	Total	250.64	47					Total	20.76	47			
Amphipods	R	4.06	3	1.35	2.78	ns	Kinorhynchs	R	1566.82	3	522.27	20.47	***
	H	0.04	1	0.04	0.03	ns		H	40.43	1	40.43	0.42	ns
	I	4.25	1	4.25	8.04	ns		I	39.30	1	39.30	1.42	ns
	R × H	4.27	3	1.42	2.92	*		R × H	285.49	3	95.16	3.73	*
	R × I	1.59	3	0.53	1.09	ns		R × I	83.07	3	27.69	1.09	ns
	H × I	3.03	1	3.03	3.46	ns		H × I	2.49	1	2.49	0.08	ns
	R × H × I	2.62	3	0.87	1.79	ns		R × H × I	97.20	3	32.40	1.27	ns
	Residuals	15.58	32	0.49				Residuals	816.26	32	25.51		
	Total	35.44	47					Total	2931.04	47			
Bivalves	R	15.59	3	5.20	79.49	***	Tanaidaceans	R	1.94	3	0.65	5.74	**
	H	2.58	1	2.58	1.16	ns		H	0.10	1	0.10	0.25	ns
	I	2.37	1	2.37	0.72	ns		I	2.28	1	2.28	1.57	ns
	R × H	6.67	3	2.22	34.01	***		R × H	1.27	3	0.42	3.76	*
	R × I	9.89	3	3.30	50.41	***		R × I	4.35	3	1.45	12.85	***
	H × I	3.50	1	3.50	1.22	ns		H × I	0.05	1	0.05	0.07	ns
	R × H × I	8.61	3	2.87	43.89	***		R × H × I	2.02	3	0.67	5.96	***
	Residuals	2.09	32	0.07				Residuals	3.61	32	0.11		
	Total	51.30	47					Total	15.64	47			
Cumaceans	R	9.05	3	3.02	63.63	***	Tardigrades	R	41.73	3	13.91	27.05	***
	H	5.28	1	5.28	1.25	ns		H	18.12	1	18.12	11.49	*
	I	2.94	1	2.94	2.79	ns		I	9.57	1	9.57	2.81	ns
	R × H	12.65	3	4.22	88.95	***		R × H	4.73	3	1.58	3.07	*
	R × I	3.17	3	1.06	22.28	***		R × I	10.23	3	3.41	6.63	**
	H × I	0.91	1	0.91	0.54	ns		H × I	12.22	1	12.22	12.82	*
	R × H × I	5.05	3	1.68	35.48	***		R × H × I	2.86	3	0.95	1.85	ns
	Residuals	1.52	32	0.05				Residuals	16.46	32	0.51		
	Total	40.57	47					Total	115.91	47			

Table 6 - Results of PERMANOVA testing for differences in the whole set of meiofaunal taxa. The analysis was done on the distance matrix calculated using Bray-Curtis similarities on untransformed data. Each term was tested using 4,999 random permutations of the appropriate units (** = $P < 0.01$; * = $P < 0.05$; ns = not significant). DF = degrees of freedom, MS = mean square, F = F-value; P (MC) = Monte Carlo probability level. Reported are also the results of the pairwise comparisons testing differences between impact and control sites in the two habitats separately for the four regions (* $P < 0.02$). R = region; H = habitat; I = impact

PERMANOVA outputs

	DF	MS	F	P(MC)
R	3	5835.49	19.49	***
H	1	1111.62	0.76	ns
I	1	3614.80	3.56	ns
R × H	3	1455.63	4.86	**
R × I	3	1016.26	3.39	**
H × I	1	418.29	0.51	ns
R × H × I	3	818.36	2.73	**
Residuals	32	299.37		
Total	47			

Pairwise comparisons: Impact vs. Control sites

Habitat	Region	P
Cyprus	Non-vegetated	**
	Vegetated	ns
Greece	Non-vegetated	**
	Vegetated	ns
Italy	Non-vegetated	ns
	Vegetated	ns
Spain	Non-vegetated	ns
	Vegetated	ns

Table 7 - Results of ANOVA testing for differences between impact and control sites in richness and evenness of meiofaunal taxa in different habitats and regions. R = region; H = habitat; I = impact; SS = sum of squares; DF = degrees of freedom; MS = means square; F = F value; P = probability level: *** = $P < 0.001$; ns = not significant.

	Source	SS	DF	MS	F	P
Number of taxa	R	47.5	3	15.8	126.67	***
	H	24.1	1	24.1	9.32	ns
	I	48.0	1	48.0	32.00	*
	R × H	7.8	3	2.6	20.67	***
	R × I	4.5	3	1.5	12.00	***
	H × I	30.1	1	30.1	15.70	*
	R × H × I	5.8	3	1.9	15.33	***
	Residuals	4	32	0.1		
	Total	171.7	47			
Community Evenness	R	0.327	3	0.109	27.31	***
	H	0.018	1	0.018	8.59	ns
	I	0.003	1	0.003	0.53	ns
	R × H	0.006	3	0.002	0.53	ns
	R × I	0.015	3	0.005	1.28	ns
	H × I	0.027	1	0.028	4.48	ns
	R × H × I	0.018	3	0.006	1.54	ns
	Residuals	0.128	32	0.004		
	Total	0.543	47			

Table 8 - Results of PERMANOVA testing for differences between impact and control sites in total meiofaunal abundance in different habitats and regions, after the removal of the covariate effects. R = region; H = habitat; I = impact; SS = sum of squares; DF = degrees of freedom; MS = means square; F = F value; P = probability level: *** = $P < 0.001$; ns = not significant. Covariates included: biopolymeric C contents, protein to carbohydrate values, water depth, sediment porosity and water content and current.

Source	DF	SS	MS	F	P
Covariates	6	7.395	1.232	7.45	**
R	3	0.406	0.135	1.28	ns
H	1	0.082	0.081	0.09	ns
I	1	1.128	1.127	-0.78	ns
R × H	3	489.232	163.077	-38.78	ns
R × I	3	-12.148	-4.049	2.22	ns
H × I	1	1.851	1.851	-0.35	ns
R × H × I	3	0.609	0.203	0.61	ns
Residual	26	8.521	0.327		
Total	47	13.398			

Table 9 – Results of the multiple regression analysis of meiofaunal abundance (square root transformed), taxa richness (untransformed) and community evenness over the environmental variables. Prop = proportional; Cu, = cumulative.

	Variable	pseudo-F	P	Explained variance (%)	
				Prop	Cum
Meiofaunal abundance	Current	8.17	0.012	15.08	15.08
	Biopolymeric C	5.13	0.032	8.7	23.77
	Protein to carbohydrate ratio	2.64	ns	4.31	28.09
	Water depth	0.13	ns	0.21	28.30
	Sediment porosity	0.07	ns	0.11	28.41
	Sediment water content	0.01	ns	0.01	28.42
Taxa richness	Water depth	11.19	0.004	19.56	19.56
	Current	9.07	0.014	13.49	33.05
	Protein to carbohydrate ratio	12.81	0.002	15.09	48.14
	Sediment porosity	0.58	ns	0.69	48.84
	Biopolymeric C	0.30	ns	0.36	49.20
	Sediment water content	0.87	ns	1.05	50.25
Community evenness	Current	19.71	0.002	29.99	29.99
	Water depth	14.85	0.002	17.37	47.37
	Biopolymeric C	1.15	ns	1.34	48.71
	Sediment porosity	0.90	ns	1.06	49.76
	Protein to carbohydrate ratio	0.85	ns	0.99	50.76
	Sediment water content	0.07	ns	0.09	50.84

Figure 1

Figure 2

Figure 3

Figure 4