**[Pb₂I₃(18-crown-6)₂][SnI₅] and CdI₂(18-crown-6) 2I₂:
Two Layered Iodine Networks with Crown-ether
Coordinated Pb²⁺ and Cd²⁺**

Michael Wolff, Claus Feldmann

► **To cite this version:**

Michael Wolff, Claus Feldmann. [Pb₂I₃(18-crown-6)₂][SnI₅] and CdI₂(18-crown-6) 2I₂: Two Layered Iodine Networks with Crown-ether Coordinated Pb²⁺ and Cd²⁺. Journal of Inorganic and General Chemistry / Zeitschrift für anorganische und allgemeine Chemie, 2010, 636 (9-10), pp.1787. 10.1002/zaac.201000072 . hal-00552458

HAL Id: hal-00552458

<https://hal.science/hal-00552458>

Submitted on 6 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Zeitschrift für Anorganische und
Allgemeine Chemie

**[Pb₂I₃(18-crown-6)₂][SnI₅] and CdI₂(18-crown-6) · 2I₂:
Two Layered Iodine Networks with Crown-ether
Coordinated Pb²⁺ and Cd²⁺**

Journal:	<i>Zeitschrift für Anorganische und Allgemeine Chemie</i>
Manuscript ID:	zaac.201000072.R1
Wiley - Manuscript type:	Article
Date Submitted by the Author:	15-Mar-2010
Complete List of Authors:	Wolff, Michael; Karlsruhe Institut of Technology (KIT), Institut für Anorganische Chemie Feldmann, Claus; Karlsruhe Institut of Technology (KIT), Institut für Anorganische Chemie
Keywords:	metal iodide, crystal structure, ionic liquid, iodine, synthesis

[Pb₂I₃(18-crown-6)₂][SnI₅] and CdI₂(18-crown-6) · 2I₂: Two Layered Iodine Networks with Crown-ether Coordinated Pb²⁺ and Cd²⁺

Dedicated to Professor Bernd Harbrecht on the occasion of his 60th birthday

Michael Wolff and Claus Feldmann*

Institut für Anorganische Chemie, Karlsruhe Institute of Technology (KIT), Germany

Prof. Dr. C. Feldmann

Institut für Anorganische Chemie, Karlsruher Institut für Technologie (KIT)

Engesserstraße 15, D-76131 Karlsruhe, Germany

Tel: +49-721-6082855

Fax: +49-721-6084892

E-mail: claus.feldmann@kit.edu

Abstract. [Pb₂I₃(18-crown-6)₂][SnI₅] (**1**) and CdI₂(18-crown-6) · 2I₂ (**2**) were prepared by reaction of PbI₂, SnI₄ and 18-crown-6 as well as CdI₂, I₂ and 18-crown-6. All reactions were performed in the ionic liquid [NMe(*n*-Bu)₃][N(Tf)₂], which serves as polar but aprotic liquid-phase providing a fast diffusion of the reactants at moderate temperatures (≤100 °C). **1** exhibits V-shaped [Pb₂I₃(18-crown-6)₂]⁺ cations and trigonal-bipyramidal [SnI₅][−] anions. In CdI₂(18-crown-6) · 2I₂ (**2**), the building unit CdI₂(18-crown-6) formally contains linear CdI₂ molecules that are interlinked by molecular iodine via long-ranging I–I contacts.

Keywords: Iodometallate; Crown-ether complex; ionic liquid; lead iodide; cadmium iodide.

Table of content text: $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ with a V-shaped $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2]^+$ cation and $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$, formally containing a linear CdI_2 molecule.

Introduction

In comparison to standard solvents used in inorganic synthesis – ranging from liquid ammonia, water, alcohols to molten alkali metal halogenides – up-coming ionic liquids (IL's) exhibit a very unusual portfolio of properties [1]. Namely, this is a wide liquid range (-50 to $+400$ °C), an excellent thermal stability (up to 400 °C), a wide electrochemical window (-4 eV to $+4$ eV) and a low vapor pressure. Since the composition nowadays can be selected from more than 100 different cations and anions, the physical and chemical properties of IL's (e.g. polarity, miscibility with other solvents, solubility of salts) can be adjusted and fine-tuned to an extent, which can be hardly realized based on standard solvents [2,3]. Comparably high costs and a dramatical drop in quality and properties when impurified with even low traces of water or impurities – e.g. stemming from the synthesis or thermal decomposition of the IL – can be regarded as major current drawbacks.

Due to there exceptional features, IL's have attracted considerable interest in recent years and especially have gained significant impact on organic synthesis [4] as well as on the synthesis of functional nanomaterials [5,6]. On the contrary, exploration towards a beneficial use of IL's in inorganic synthesis has just begun and already lead to promising results, for instance, fascinating germanium clathrates [7] or the chemistry of non-coordinating anions [8]. As part of our studies, IL's are used as polar, but aprotic liquids that guarantee a sufficient diffusion at comparably low temperatures (<100 °C) [9]. In the following we aim at the synthesis of novel iodometallates in ionic liquids. Our focus is especially on halogen-rich compounds and novel structural building units [10,11]. In general, iodometallates have already gained interest due to complex supramolecular networks and unexpected semiconductor-type properties [12–15]. With $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ and $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ in the following a first compound in the system Pb(II)–Sn(IV)–I as well as a compound that formally contains I_2 -interlinked CdI_2 molecules are presented.

Experimental Section

All experimental and analytical studies were performed under argon or vacuum (i.e. glove-boxes or Schlenk techniques). The commercially available starting materials (SnI_4 : ≥ 99.999 %, Sigma-Aldrich; PbI_2 : ≥ 99.999 %, Sigma-Aldrich; CdI_2 : ≥ 99.999 %, Sigma-Aldrich; 18-crown-6: ≥ 99 %, Acros) were used subsequent to overnight drying in vacuum. Iodine (≥ 99.5 %, Sigma-Aldrich) was purified by double sublimation. The ionic liquid (IL), $[\text{NMe}(n\text{-Bu})_3][\text{N}(\text{Tf})_2]$ ($\text{Tf} = \text{SO}_2\text{CF}_3$), was synthesized as described elsewhere [16].

Synthesis of $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ (I): 135.9 mg ($2.17 \cdot 10^{-4}$ mol) of SnI_4 , 200 mg ($4.34 \cdot 10^{-4}$ mol) of PbI_2 and 104.8 mg ($4.34 \cdot 10^{-4}$ mol) of 18-crown-6 were added to 1 mL of $[\text{NMe}(n\text{-Bu})_3][\text{N}(\text{Tf})_2]$ at 100 °C. Dark-red, plate-shaped, moisture-sensitive crystals were obtained after cooling

to room temperature with at 5 °C/h. These crystals were separated from the IL by filtration through a glass filter and by washing three times with 4 mL of dried tetrahydrofuran. Finally, the crystals were dried in vacuum and obtained with a yield of about 90 %.

Synthesis of $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ (2): 100 mg ($2.73 \cdot 10^{-4}$ mol) of CdI_2 , 72.2 mg ($2.73 \cdot 10^{-4}$ mol) of 18-crown-6 and 69.3 mg ($2.73 \cdot 10^{-4}$ mol) of I_2 were added to 1 mL of $[\text{NMe}(n\text{-Bu})_3][\text{N}(\text{Tf})_2]$ at 80 °C. **2** crystallized as dark-red cubes within 14 days. The crystals were separated from the IL by filtration through a glass filter and by washing several times with 5 mL of dried diethyl ether. Finally, $\text{CdI}_2(\text{I}_2)_2 \cdot (18\text{-crown-6})$ was dried in vacuum and obtained with a yield of about 50 %. Surprisingly, the crystals turned out to be stable in air.

Data collection and refinement: Single crystal X-ray diffraction analysis was carried out with an IPDS II diffractometer (STOE, Darmstadt) using Mo- K_α radiation ($\lambda = 0.71073 \text{ \AA}$, graphite monochromator). A suited crystal was selected for both title compounds directly from the mother liquor and fixed onto a glass filament begird in perfluorated oil (Kel-F). Data collection was performed at 200 K. Subsequent to data reduction via X-RED [17], space group determination was carried out on the basis of the reflection conditions via XPREP. Structure solution and refinement were performed with SHELXTL [18]. The results are listed in Table 1. All non-hydrogen atoms were refined with anisotropic parameters of displacement. The position of hydrogen could not be fixed via Fourier refinement and was therefore modeled based on idealized C–H bonds. X-SHAPE was used to apply a numerical absorption correction based on crystal-shape optimization for the data of compound (**1**) [19]. Illustrations of the crystal structures were prepared via DIAMOND [20]. More details concerning the crystallographic data can be requested from the Cambridge Crystallographic Data Center [deposit@ccdc.cam.ac.uk] with the deposition numbers CCDC ?????? and ??????.

Powder diffraction pattern were obtained with a STADI-P diffractometer (Stoe, Darmstadt) using germanium monochromatized Cu- $\text{K}_{\alpha 1}$ radiation. The samples were filled in a glass capillary and measured at ambient temperature. Powder diffraction pattern (not pictured) were used to confirm the setting of the unit cells as well as the purity of the title compounds.

Energy-dispersive X-ray analysis (EDX): EDX was carried out using an AMETEC EDAX mounted on a Zeiss SEM Supra 35 VP scanning electron microscope. Crystals of $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ were fixed on conductive carbon pads on aluminum sample holders. In contrast $[\text{Sn}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ [11] the presence of lead was clearly visible in the title compound **1**.

Results and Discussion

$[Pb_2I_3(18\text{-crown-}6)_2][SnI_5]$ (**1**)

By reaction of PbI_2 , SnI_4 and 18-crown-6 in the ionic liquid $[NMe(n\text{-}Bu)_3][N(Tf)_2]$, moisture-sensitive dark-red crystals with the composition $[Pb_2I_3(18\text{-crown-}6)_2][SnI_5]$ were obtained. According to X-ray structure analysis based on single crystals, the title compound crystallizes monoclinically with space group $P2_1/n$ (Table 1). The unit cell displays $[SnI_5]^-$ anions and $[Pb_2I_3(18\text{-crown-}6)_2]^+$ -cations as the constituents of **1** (Figure 1).

The $[SnI_5]^-$ anion exhibits only a slight deviation from a trigonal-bipyramidal symmetry (Figure 1). This view is confirmed by angles $I_{eq}\text{--}Sn\text{--}I_{eq}$ of $120\pm 2^\circ$, and $I_{ax}\text{--}Sn\text{--}I_{eq}$ of $90\pm 1^\circ$. The Sn–I bond lengths with 271.7 to 285.4 pm are close to those of SnI_4 (263 pm), but differ significantly from SnI_2 (300–325 pm) [21,22]. Taking the widened coordination from 4 (SnI_4) to 5 (SnI_5^-) into account, the bond lengths clearly point to the presence of Sn^{4+} . Such a trigonal-bipyramidal coordination, in general, is much less common for iodostannates(IV) than the well-known octahedral coordination, most often leading to merged building units [23]. Beside $[SnI_5]^-$, which has been first observed in the mixed-valent iodostannate $[Sn(II)I_3(18\text{-crown-}6)_2][Sn(IV)I_5]$ [11], only $[SnI_6]^{2-}$ has been described as an isolated building unit so far [24].

The novel **dinuclear** cation $[Pb_2I_3(18\text{-crown-}6)_2]^+$ consists of two symmetry equivalent crown-ether molecules, each of them entrapping Pb^{2+} . Similar to $[Sn_2I_3(18\text{-crown-}6)]^+$ [11], each lead cation is coordinated not only by 18-crown-6, but also to a terminal iodine atom as well as to a joint bridging iodide (Figure 2). This connectivity results in a V-shaped $[Pb_2I_3(18\text{-crown-}6)]^+$ building unit that exhibits angles of 164.2° (I–Pb–I) and 142.8° (Pb–I–Pb). Noteworthy, the terminal iodide with 292.7 pm shows a significantly shorter Pb–I distance than the bridging I $^-$ (350.4 pm) (Table 2). Consequently, the bonding situation can be also described as two $[PbI(18\text{-crown-}6)]^+$ units linked by a single iodide anion. A cation $[PbI(18\text{-crown-}6)]^+$ is known from $Pb(18\text{-crown-}6)I_2$ [36] and exhibits a similar Pb $^{2+}$ –I distance (290 pm) as observed here (293 pm). Pb^{2+} is positioned exactly inside the plane of the crown ether molecule, but slightly deflected from its center (Figure 3). This finding results in a shortened (266.8(1) pm), a lengthened (285.1(1) pm) and four largely equivalent (270.9(1)–274.5(1) pm) Pb–O distances. All relevant Pb–O and Pb–I distances are summarized and compared to literature data in Table 2. In comparison to $[Sn_2I_3(18\text{-crown-}6)]^+$, Pb^{2+} is less deflected from the center of the crown ether than Sn^{2+} . This finding is in accordance with the increased radius as well as with a less pronounced lone-pair effect [25,26]. All C–O and C–C bond lengths of 18-crown-6 match with literature data [27].

Although a description of **1** based on ionic constituents $[Pb_2I_3(18\text{-crown-}6)_2]^+$ and $[SnI_5]^-$ is reasonable, long-ranging I–I interactions can not be disregarded (Figure 4). Attractive bonding with distances below the doubled van-der-Waals radius of iodine (i.e. 420 pm, [28]) are observed between I1 and I3 (402.1

pm) as well as between I2 and I4 (406.4 pm). Only I5 remains without any significant I–I contact. In sum, long-ranging I–I contacts interconnect $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2]^+$ and $[\text{SnI}_5]^-$ to infinite helices along [001] and [100] to form a two-dimensional iodometallate network. Finally it should be mentioned that $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ first contains Pb^{2+} , Sn^{4+} and I^- in combination. In general, the system Pb–Sn–I is with limited examples namely $[(\text{NH}_3\text{CH}_3)\text{Pb}(\text{II})_{0.17}\text{Sn}(\text{II})_{0.83}]\text{I}_3$ and PbSnI_4 [29,30].

CdI₂(18-crown-6) · 2I₂ (2)

$\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ was prepared by reaction of one equivalent of CdI_2 and 18-crown-6 with two equivalents of I_2 in the ionic liquid $[\text{NMe}(n\text{-Bu})_3][\text{N}(\text{Tf})_2]$. Dark reddish, cube-shaped and air-stable single crystals were crystallized within two weeks. According to X-ray structure analysis based on single crystals, **2** crystallizes monoclinically with space group *C2/c* (Table 1). The unit cell displays $\text{CdI}_2(18\text{-crown-6})$ units as well as molecular iodine (Figure 5).

Cd^{2+} is positioned almost centrally inside the crown ether with Cd–O distances ranging from 265.7(1) to 280.8(1) pm. All C–O and C–C distances match with the expectation [27]. In addition to 18-crown-6, cadmium is further coordinated by two iodine atoms, which are themselves in contact to two I_2 molecules (Figure 6). Here, the question arises whether the observed structural building unit is indeed related to a molecular situation such as $[\text{Cd}(18\text{-crown-6})\text{I}_2] \cdot 2\text{I}_2$ or whether a description $[\text{Cd}(18\text{-crown-6})]^{2+}[\text{I}_3^-]_2$ with the well-known triiodide I_3^- is more appropriate. According to the bond lengths (Figure 6), I2–I3 (273.3(1) pm) corresponds very well to solid I_2 (I–I: 272 pm, [31]), whereas I1–I2 (321.9(1) pm) is considerably elongated. With regard to the angles, I1–Cd–I1 (180 °) turned out to be exactly linear, whereas I1–I2–I3 (172.9(1) °) clearly deviates from linearity. Altogether, these findings are in contrast to what was reported for various crystal structures addressing I_3^- as a polyiodide anion. The latter in fact possesses an almost linear geometry and I–I bond lengths ranging from 285 to 295 pm [28]. The above argumentation is also consistent with the related compound $\text{CdI}_2(\text{benzo-18-crown-6}) \cdot \text{I}_2$ [32]. Herein, the $\text{CdI}_2(\text{benzo-18-crown-6})$ building unit is in contact to I_2 , too. Bond lengths (I1–I2: 314.8 pm, I2–I3: 275.3 pm) and angles (I1–I2–I3: 170.9 °) are very similar to the title compound **2**. Compared to $\text{CdI}_2(\text{benzo-18-crown-6}) \cdot \text{I}_2$, however, two significant differences are evident for $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ (**2**): First, the I–Cd–I angle in **2** matches exactly 180 ° due to inversion symmetry at the Cd^{2+} site whereas $\text{CdI}_2(\text{benzo-18-crown-6}) \cdot \text{I}_2$ shows an angulated conformation (I1–Cd–I1: 167.0 °). Second, the CdI_2 molecule in **2** is connected to two I_2 molecules, in contrast to only a single I_2 molecule in $\text{CdI}_2(\text{benzo-18-crown-6}) \cdot \text{I}_2$. The latter can be most probably ascribed to the volume and steric demand of benzo-18-crown-6, which obviously prevents addition of a second iodine molecule.

Surprisingly, the Cd–I1 distance in $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ with 269.3(1) pm is significantly shorter as compared to bulk- CdI_2 (Cd–I: 299 pm, [33]). In contrast, the observed value matches very well with

what is discussed for CdI_2 molecules in the gas phase (260 pm, [34,35]). Taking both aspects – the I–I as well as the Cd–I distances – into account, the structural building unit $\text{CdI}_2(18\text{-crown-6})$ in sum can be described as a linear CdI_2 molecule with Cd^{2+} additionally coordinated by the crown ether and each iodide atom weakly bound to I_2 molecules. $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ altogether exhibits the highest iodine content of all cadmium compounds known by now.

Taking all long-ranging I–I contacts with distances below the doubled van-der-Waals radius (420 pm, [28]) into account, additional attractive interactions (I2–I3 with 399.5(1) pm) become obvious (Figure 6). Accordingly, $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ forms an infinite two-dimensional network via I_2 molecules. Formally, this network is constituted of distorted Cd_2I_{14} -rings that form zigzagged layers along [110] (Figure 7). Even though no hexagonal close-packing of iodide is observed here, the structure is nevertheless related to the CdI_2 -type of structure. Based on a 2x2 super cell, CdI_2 as well as **2** show stacked $\cdots\text{I}-\text{Cd}-\text{I}-\text{I}-\text{Cd}-\text{I}\cdots$ layers, in general. However, the stacking sequence I(A)–Cd(1)–I(B)–I(A)–Cd(2)–I(B) in CdI_2 is different from I(A)–Cd(1)–I(A)–I(B)–Cd(2)–I(B) in **2**.

In summary, two novel layered iodide networks have been gained with $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ and $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ via ionic liquid based reaction. Although we can naturally not exclude other preparative methods as access to the title compounds, a sufficient diffusion and reactivity of the starting materials in the IL at moderate temperature (<100 °C) is obvious. Concretely, thermal decomposition of metal iodides, temperature-driven loss of iodine as well as undesired redox processes due to elemental iodine and metal iodides can be avoided.

Acknowledgments

The authors are grateful to the Center of Nanostructured Materials (CFN) of the Deutsche Forschungsgemeinschaft (DFG) at the Karlsruhe Institute of Technology (KIT) for financial support.

Table 1. Crystallographic data of the title compounds.

	[Pb ₂ I ₃ (18-crown-6) ₂][SnI ₅] (1)	CdI ₂ (18-crown-6) · 2I ₂ (2)
Formula	C ₂₄ H ₄₈ O ₁₂ Pb ₂ SnI ₈	C ₁₂ H ₂₄ O ₆ CdI ₆
crystal system / space group	monoclinic / <i>P</i> 2/ <i>n</i> (no. 13)	monoclinic / <i>C</i> 2/ <i>c</i> (no. 15)
cell parameters	<i>a</i> = 12.375(3) Å <i>b</i> = 7.866(2) Å <i>c</i> = 24.395(5) Å <i>β</i> = 91.99(3)° <i>V</i> = 2373.2(9) Å ³	<i>a</i> = 12.591(3) Å <i>b</i> = 10.137(2) Å <i>c</i> = 20.637(4) Å <i>β</i> = 95.47(3)° <i>V</i> = 2621.9(9) Å ³
formula units per cell	<i>Z</i> = 2	<i>Z</i> = 4
calculated density	$\rho_{calc} = 2.91 \text{ g cm}^{-3}$	$\rho_{calc} = 2.88 \text{ g cm}^{-3}$
temperature	<i>T</i> = 200 K	<i>T</i> = 200 K
measurement device	imaging plate diffraction system (IPDS II, STOE)	imaging plate diffraction system (IPDS II, STOE)
radiation	graphite-monochromated Mo K _α (<i>λ</i> = 0.71073 Å)	graphite-monochromated Mo-K _α (<i>λ</i> = 0.71073 Å)
measurement limits	3.3° ≤ 2 <i>θ</i> ≤ 58.6° −16 ≤ <i>h</i> ≤ 16, −10 ≤ <i>k</i> ≤ 10, −31 ≤ <i>l</i> ≤ 33	4.0° ≤ 2 <i>θ</i> ≤ 58.4° −17 ≤ <i>h</i> ≤ 17, −13 ≤ <i>k</i> ≤ 13, −28 ≤ <i>l</i> ≤ 26
absorption correction	numerical	numerical
absorption coefficient	<i>T</i> _{min} = 0.165; <i>T</i> _{max} = 0.223 <i>μ</i> = 12.81	<i>T</i> _{min} = 0.032; <i>T</i> _{max} = 0.085 <i>μ</i> = 7.94
no. of reflections	23184 (6351 unique) measured	11619 (3536 unique)
data averaging	<i>R</i> _{int} = 0.037	<i>R</i> _{int} = 0.044
structure refinement	full matrix least-squares on <i>F</i> _o ² anisotropic displacement parameters for all non-hydrogen atoms	full-matrix least-squares on <i>F</i> _o ² anisotropic displacement parameters for all non-hydrogen atoms
no. of parameters	216	117
residual electron density	−3.45 to 2.35 e Å ^{−3}	−1.54 to 2.03 e Å ^{−3}
figures of merit	<i>R</i> 1 = 0.049 (all data) <i>R</i> 1 (3834 <i>F</i> _o ² > 4σ(<i>F</i> _o ²)) = 0.045 <i>wR</i> 2 = 0.131 <i>Goof</i> = 1.091	<i>R</i> 1 = 0.060 (all data) <i>R</i> 1 (2893 <i>F</i> _o ² > 4σ(<i>F</i> _o ²)) = 0.047 <i>wR</i> 2 = 0.137 <i>Goof</i> = 1.086

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Selected bond lengths (pm) in $[M_nI_m(18\text{-crown-6})]^+$ complexes (M = Pb, Sn).

	$[Pb_2I_3(18\text{-crown-6})_2]^+$	$[PbI(18\text{-crown-6})]^+$ [36]	$[Sn_2I_3(18\text{-crown-6})]^+$ [11]
M–O	266.8 – 285.1	271.5 – 278.5	261.2 – 293.7
M–I1	292.7	290.1	287.2
M–I2	350.4	–	355.0

Figure 1. Unit cell of $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ (**1**) (thermal ellipsoids with 50 % probability of finding).

Figure 2. $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2]^+$ -cation in $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ (**1**) (thermal ellipsoids with 50 % probability of finding).

Figure 3. Coordination of Pb^{2+} by 18-crown-6 in **1** (bond lengths in pm; thermal ellipsoids with 50 % probability of finding).

Figure 4. The two-dimensional iodometalate network in **1** interlinking the cation $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2]^+$ as well as the anion $[\text{SnI}_5]^-$ via long-ranging I–I contacts, (18-crown-6 not shown for clarity).

Figure 5. Unit cell of $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ (**2**) (thermal ellipsoids with 50 % probability of finding).

Figure 6. Connectivity and selected bond length in $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ (**2**) (in pm; thermal ellipsoids with 50 % probability of finding).

Figure 7. The two-dimensional network of $[\text{CdI}_2(18\text{-crown-6})]^{2+}$ and molecular I_2 in **2** along $[110]$ (thermal ellipsoids with 50 % probability of finding).

References

-
- [1] P. Wasserscheid, T. Welton (Eds.), *Ionic Liquids in Synthesis*; Wiley-VCH, Weinheim 2002.
- [2] J. M. Slattery, C. Daguenet, P. J. Dyson, T. J. S. Schubert, I. Krossing, *Angew. Chem.* **2007**, *119*, 5480; *Angew. Chem. Int. Ed.* **2007**, *46*, 5384.
- [3] M. Armand, F. Endres, D. R. MacFarlane, H. Ohno, B. Scrosati, *Nature Mater.* **2009**, *8*, 621.
- [4] J. Dupont, R. F. de Souza, P. A. Z. Suarez, *Chem. Rev.* **2002**, *102*, 3667.
- [5] M. Antonietti, D. Kuang, B. Smarsly, Y. Zhou, *Angew. Chem.* **2004**, *116*, 5096; *Angew. Chem. Int. Ed.* **2004**, *43*, 4988.
- [6] H. Goesmann, C. Feldmann, *Angew. Chem.* **2010**, in press; *Angew. Chem. Int. Ed.* **2010**, in press.
- [7] A. M. Guloy, R. Ramlau, Z. Tang, W. Schnelle, M. Baitinger, Y. Grin, *Nature* **2006**, *443*, 320.
- [8] I. Krossing, I. Raabe, *Angew. Chem.* **2004**, *116*, 2116; *Angew. Chem. Int. Ed.* **2004**, *43*, 2066.
- [9] G. Bühler, C. Feldmann, *Angew. Chem.* **2006**, *118*, 4982; *Angew. Chem. Int. Ed.* **2006**, *45*, 4864.
- [10] A. Okrut, C. Feldmann, *Inorg. Chem.* **2008**, *47*, 3084.
- [11] M. Wolff, T. Harmening, R. Pöttgen, C. Feldmann, *Inorg. Chem.* **2009**, *48*, 3153.
- [12] C. Lode, H. Krautscheid, *Z. anorg. allg. Chem.* **2007**, *633*, 1691.
- [13] H. Li, Z. Chen, L. Cheng, J. Liu, X. Chen, J. Li, *J. Crystal Growth* **2008**, *8*, 4355.
- [14] C. R. Kagan, D. B. Mitzi, C. D. Dimitrakopoulos, *Science* **1999**, *286*, 945.
- [15] J. C. Papaioannou, V. G. Charalampopoulos, P. Xynogalas, K. Viras, *J. Phys. Chem. Solids* **2006**, *67*, 1379.
- [16] D. R. MacFarlane, P. Meakin, J. Sun, N. Amini, N. Forsyth, *J. Phys. Chem. B* **1999**, *103*, 4164.
- [17] X-RED, *Data Reduction Program*, Version 1.14, Stoe & Cie GmbH, Darmstadt **1999**.
- [18] SHELXTL, *Structure Solution and Refinement Package*, Version 5.1, Bruker AXS, Karlsruhe 1998.
- [19] X-SHAPE, *Crystal Optimisation for Numerical Absorption Correction*, Version 1.06, Stoe & Cie GmbH, Darmstadt 1999.
- [20] DIAMOND, *Visuelles Informationssystem für Kristallstrukturen*, Version 3.0d, Crystal Impact GbR, Bonn 2005.
- [21] R. G. Dickinson, *J. Am. Chem. Soc.* **1923**, *45*, 958.
- [22] R. A. Howie, W. Moser, I. C. Treventa, *Acta Crystallogr.* **1972**, *28*, 2965.
- [23] C. Lode, H. Krautscheid, *Z. anorg. allg. Chem.* **2001**, *627*, 1454.
- [24] J. Guan, Z. Tang, A. M. Guloy, *Chem. Commun.* **2005**, 48.
- [25] R. D. Rogers, A. H. Bond, *Inorg. Chim. Acta* **1992**, *192*, 163.
- [26] R. P. Power, *Chem. Rev.* **1999**, *99*, 3463.

-
- [27] J. D. Dunitz, P. Seiler, *Acta Crystallogr.* **1974**, 30, 2739.
- [28] A. F. Holleman, E. Wiberg, *Lehrbuch der Anorganischen Chemie*; Walter de Gruyter: Berlin-New York 2007.
- [29] D. Weber, *Z. Naturforsch. B* **1979**, 34, 939.
- [30] T. A. Kuhn, *Thin Solid Films* **1998**, 340, 292.
- [31] F. van Bolhuis, P. B. Koster, T. Migchelsen, *Acta Crystallogr.* **1967**, 23, 90.
- [32] R. Striebinger, C. Walbaum, I. Pantenburg, G. Meyer, *Z. anorg. allg. Chem.* **2008**, 634, 2743.
- [33] G. K Chadha, *Z. Kristallogr.* **1974**, 139, 147.
- [34] P. W. Allen, L. E. Sutton, *Acta Crystallogr.* **1950**, 3, 46.
- [35] O. Hassel, L. C. Stromme, *Z. Phys. Chem. B.* **1938**, 38, 466.
- [36] K. F Tebbe, M. E. Essawi, S. A. E. Khalik, *Z. Naturforsch. B* **1995**, 50, 1429.

Figure 1. Unit cell of [Pb₂I₃(18-crown-6)₂][SnI₅] (1) (thermal ellipsoids with 50 % probability of finding).

Figure 2. [Pb₂I₃(18-crown-6)₂]⁺-cation in [Pb₂I₃(18-crown-6)₂][SnI₅] (1) (thermal ellipsoids with 50 % probability of finding).

Figure 3. Coordination of Pb²⁺ by 18-crown-6 in 1 (bond lengths in pm; thermal ellipsoids with 50 % probability of finding).

Figure 4. The two-dimensional iodometalate network in **1** interlinking the cation $[Pb_2I_3(18\text{-crown-}6)_2]^+$ as well as the anion $[SnI_5]^-$ via long-ranging I-I contacts, (18-crown-6 not shown for clarity).

Figure 5. Unit cell of $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$ (2) (thermal ellipsoids with 50 % probability of finding).

Figure 6. Connectivity and selected bond length in CdI₂(18-crown-6) · 2H₂O (2) (in pm; thermal ellipsoids with 50 % probability of finding).

Figure 7. The two-dimensional network of $[CdI_2(18\text{-crown-}6)]^{2+}$ and molecular I_2 in 2 along $[110]$ (thermal ellipsoids with 50 % probability of finding).

Table of contents text: $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2][\text{SnI}_5]$ with a V-shaped $[\text{Pb}_2\text{I}_3(18\text{-crown-6})_2]^+$ cation and $\text{CdI}_2(18\text{-crown-6}) \cdot 2\text{I}_2$, formally containing a linear CdI_2 molecule.