Morbidity and Mortality in Craniopharyngioma Patients after Surgery

Rachel K Crowley, Ole Petter Hamnvik, Eoin P O'Sullivan, Lucyann Behan, Diarmuid Smith, Amar Agha, Christopher J Thompson

► To cite this version:

Rachel K Crowley, Ole Petter Hamnvik, Eoin P O'Sullivan, Lucyann Behan, Diarmuid Smith, et al.. Morbidity and Mortality in Craniopharyngioma Patients after Surgery. *Clinical Endocrinology*, 2010, 10.1111/j.1365-2265.2010.03838.x . hal-00552614

HAL Id: hal-00552614

<https://hal.science/hal-00552614>

Submitted on 6 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Morbidity and Mortality in Craniopharyngioma Patients after Surgery

Journal:	<i>Clinical Endocrinology</i>
Manuscript ID:	CEN-2009-000833.R2
Manuscript Type/Office:	1 Original Article - UK/Europe
Date Submitted by the Author:	04-May-2010
Complete List of Authors:	Crowley, Rachel; Beaumont Hospital, Academic Department of Endocrinology Hamnvik, Ole Petter; Beaumont Hospital, Academic Department of Endocrinology O'Sullivan, Eoin P; Beaumont Hospital, Academic Department of Endocrinology Behan, LucyAnn; Beaumont Hospital, Academic Department of Endocrinology Smith, Diarmuid; Beaumont Hospital, Academic Department of Endocrinology Agha, Amar; Beaumont Hospital, Academic Department of Endocrinology Thompson, Christopher J; Beaumont Hospital, Academic Department of Endocrinology
Key Words:	craniopharyngioma, hypopituitarism, mortality, obesity

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Morbidity and Mortality in Craniopharyngioma Patients after Surgery

**Short title: Morbidity and Mortality in
Craniopharyngioma**

**RK Crowley, OP Hamnvik, EP O’Sullivan, LA
Behan, D Smith, A Agha, CJ Thompson**

**Academic Department of Endocrinology, Beaumont
Hospital, Dublin 9, Ireland**

Correspondence: RK Crowley, Beaumont Hospital, Dublin 9, Ireland.

rachelcrowley@hotmail.com

Keywords: craniopharyngioma, hypopituitarism, mortality, obesity

SUMMARY

Objective: Craniopharyngioma (CP) is a benign tumour of the suprasellar region that is associated with increased morbidity and mortality in comparison to other causes of hypopituitarism. We aimed to establish the rate and causes of mortality and morbidity in patients with CP who attended our centre.

Design: We performed a retrospective case note audit of CP patients who were managed by our service. We established the Standardised Mortality Ratio for CP patients. We compared obesity prevalence with two other hypopituitary groups who are managed by our service.

Patients: We identified 70 patients with craniopharyngioma, 97% of whom had undergone surgery and 42% radiotherapy. We compared the prevalence of obesity with that of 89 patients with hypopituitarism secondary to surgery for non-functioning pituitary adenoma, and 29 patients with post-traumatic hypopituitarism.

Measurements: Standardised Mortality ratio for CP patients was 8.75 (95% CI of 5.4-13.3); SMR for women was 10.51 (95% CI 5.04-19.3) and 7.55 (95% CI 3.77-13.52) for men. The rates of GH, gonadotrophin, ACTH and TSH deficiencies were 91%, 93.5%, 92% and 86%, respectively. The rate of diabetes insipidus was 81%; 7.1% had adipsic diabetes insipidus. Dyslipidaemia was present in 46.9% and diabetes mellitus in 11.5%. Obesity affected 66% of CP patients, 47% of non-functioning adenoma

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

patients and 31% of those with post-traumatic hypopituitarism ($p < 0.001$).

Conclusions: Patients with craniopharyngioma suffer from high rates of mortality and morbidity. The underlying causes for mortality and for obesity in this population remain poorly-understood.

Word count: 242

For Peer Review

INTRODUCTION

Craniopharyngioma (CP) is a benign tumour of the suprasellar region which has been associated with increased morbidity and mortality in comparison to other causes of hypopituitarism¹⁻⁴. The increased mortality has been attributed to both cardiovascular and respiratory disease, but the reported susceptibility to cardiorespiratory disease is poorly understood. The reported morbidity associated with craniopharyngioma includes high rates of hypopituitarism, hypothalamic complications and impaired quality of life⁵.

The aim of this study was to establish mortality rates in CP patients treated at our institution and to calculate the Standardised Mortality Ratio (SMR). We also wished to document the spectrum of morbidity, and the potential contribution of co-morbidities, such as obesity, hypertension, dyslipidaemia and diabetes mellitus, to the increased mortality in CP. We compared the prevalence of obesity in our CP population to that of another patient cohort with tumoural hypopituitarism (non-functioning pituitary adenoma) and to that of a hypopituitary group of similar age (post-traumatic hypopituitarism), who were managed in the same centre.

METHODS

Patients were identified from the hospital pituitary database, which collates details from all patients treated between 1980 and 2008. 70 patients (39 male) with craniopharyngioma were identified from the database; 21 patients had died at the time of analysis.

Mortality

Patients who died while under follow-up in our hospital had cause of death identified from hospital casenotes and local post-mortem information. For patients who were lost to follow-up, or who were followed up in other institutions, the family practitioner was contacted in order to identify patients who had died.

Cause of death was established from death certificates from the Office of the Registrar General and from post mortem data. Mortality rates for the Irish population were obtained from the Central Statistics Office of Ireland. The background Irish mortality rate per 1000 population for each age group and gender was expressed as a decimal and converted into a cumulative mortality by summation of previous yearly mortality.

The expected mortality by age and gender of each craniopharyngioma patient was calculated:

Background cumulative mortality for year of death - Background cumulative mortality for the year of diagnosis with craniopharyngioma

1
2
3
4
5
6 The craniopharyngioma Standardised Mortality Ratio (SMR) was defined
7
8 as:

9
10 Number of deaths observed / number of expected deaths in an age- and
11
12 gender-matched population
13
14
15
16

17 Irish data were available from the years 1950-2004; thus the background
18
19 mortality in 2005 to 2008 was assumed to equal that of 2004. Poisson
20
21 probability distribution was used to calculate the 95% confidence
22
23 interval (CI) for the SMR.
24
25
26
27

28 29 Morbidity 30 31 32 33

34 Case note review was performed for collection of morbidity data and
35
36 details were recorded in a spreadsheet (Windows Excel). Biographical
37
38 details included gender, date of birth and age at diagnosis and death.
39
40 Case management details included use of surgery and radiotherapy,
41
42 surgical approach, recurrence of tumour, pituitary hormone replacement
43
44 therapy and dosage and prescription of statin, anti-hypertensive and
45
46 anti-epileptic therapy. Prevalence of anterior pituitary hormone
47
48 deficiency and diabetes insipidus was recorded.
49
50
51
52

53
54
55 The most recent measurement of fasting blood glucose and lipid profile,
56
57 and blood pressure, height, weight and BMI measurements from the
58
59 most recent outpatient office measurement were noted. Historical
60

diagnoses of diabetes mellitus, hypertension or dyslipidaemia were also recorded. The diagnosis of diabetes mellitus was based on standard WHO criteria of positive OGTT results, two fasting blood glucose concentrations > 7.0 mmol/l or a random blood glucose of > 11.1 mmol/l when accompanied by osmotic symptoms. Hypertension was diagnosed by three office readings of > 140 / 90mmHg. Dyslipidaemia was defined as total cholesterol > 5mmol/l or LDL cholesterol > 2.6 mmol/l.

GH deficiency (GHD) was defined as a peak GH response of less than 3 µg/l in response to insulin tolerance test (ITT) or glucagon stimulation test (GST), as defined by locally derived normative data ⁶. ACTH deficiency was defined as a peak cortisol less than 500 nmol/l on ITT or 450 nmol/l on GST, as previously defined ⁶. Gonadotrophin deficiency in pre-menopausal women was defined as amenorrhea for more than 6 months in the presence of low oestradiol and normal or low gonadotrophins. In post-menopausal women gonadotrophin deficiency was defined as gonadotrophins of pre-menopausal range. In men gonadotrophin deficiency was defined as symptomatic low testosterone, and inappropriately low gonadotrophins. TSH deficiency was defined as low T4 in the presence of low or normal TSH. Diabetes insipidus was diagnosed in the post-operative period by the presence of polyuria and hypernatraemia, according to the criteria of Seckl and Dunger ⁷ and confirmed with water deprivation test six weeks after discharge. Adipsia was diagnosed by use of an infusion of hypertonic saline (855 mmol/l)

with measurement of thirst as previously described⁸. Prevalence of other morbidity such as obesity, hydrocephalus, seizures, visual loss and venous thrombosis was recorded from the case notes.

Body mass index (BMI) data were compared to that of other hypopituitary groups; 89 patients with hypopituitarism after surgery for non-functioning pituitary adenoma (NFPA) and 29 patients with hypopituitarism after traumatic brain injury (TBI), also identified from the pituitary database.

Statistics

Mean and standard deviation were used to describe data that were normally distributed; otherwise the median and range was reported. BMI data were compared with Wilcoxon's test and subanalysed with Tukey's HSD. Significance of subanalysis findings was determined by post hoc t test with correction for multiple comparisons. JMP 5 (SAS Institute, California, USA) was used for statistical analysis. The study was approved by the local Ethics Committee.

RESULTS

Mortality

Median follow-up of craniopharyngioma patients from time of diagnosis was 8 years (range 1-50 years). Twenty-four patients (34%) were diagnosed with CP in childhood. Twenty-one CP patient deaths were observed between 1988 and 2007. The expected mortality in this population was 2.4, therefore SMR was 8.75 with a 95% CI of 5.4-13.3. SMR for women (10.51, 95% CI 5.04-19.3) was higher than for men (7.55, 95% CI 3.77-13.52), but this was not statistically significant (RR 1.388, CI 0.55-3.51) ⁹. Causes of death are reported in Table 1.

Morbidity

Age at diagnosis was distributed in a bimodal pattern (Fig 1). Sixty-eight patients underwent surgical intervention; 2 patients were diagnosed with CP on radiological appearance alone and did not undergo surgery because of preservation of visual fields. Eight patients underwent surgery by a trans-sphenoidal approach, 5 of which required a later craniotomy. An average of 60 pituitary resections were performed in the centre each year; there were eight neurosurgeons in the centre, 1 of whom was a specialist pituitary surgeon. Three successive pituitary surgeons were responsible for the service over the time period studied.

Radiotherapy details were not available for 6 patients (4 deceased, 1 attended follow-up elsewhere, 1 failed to follow up). Of the remaining 64 patients, 27 (42%) underwent radiotherapy; 16 after recurrence, 1 as prophylaxis with early recurrence and 10 of these patients received prophylactic radiotherapy without recurrence. Records were incomplete in 3 deceased patients; of the remaining 67, 31 patients (46.2%) suffered from tumour recurrence.

Growth hormone deficiency (GHD) was diagnosed in 53 of 58 patients (91%) for whom the GH axis was tested; of the remaining 12 patients, 8 died prior to 1997 when assessment for adult GHD became routine in our unit and 4 did not follow-up with our service. Of those 5 patients considered to have intact GH secretion, 2 did not undergo surgery and the remaining 3 were considered to be GH replete because of a normal IGF1 result. Of the patients identified with GHD, 62% received GH replacement. GH was not replaced because of tumour mass or recurrence in 8% of GHD patients, discontinued after perceived lack of benefit in 16%, discontinued after the development of diabetes in 5.4%, or not commenced because of patient age (over 70 years) in 5.4%. GH replacement therapy was monitored by serial measurement of IGF1 and titrated to maintain IGF1 in the upper half of the normal reference range. Median GH dose was 0.35 mg with a range of 0.2 – 1.8 mg. Analysis of mortality in GHD patients who received replacement compared with those who did not, was not significant (χ^2 , $p = 0.2$).

ACTH status was not documented in 4 patients; 3 were deceased and 1 attended another institution. Of those for whom data was available, 61 patients (92%) had ACTH deficiency. 2 of the patients with intact ACTH secretion had not undergone surgery, 2 failed to follow-up with our service and 1 was managed elsewhere. All patients diagnosed with ACTH deficiency received replacement therapy with hydrocortisone or prednisolone. Median daily dose of steroid was equivalent to 20 mg of hydrocortisone, range 10 – 30mg, where 5mg prednisolone was considered equivalent to 20mg of hydrocortisone.

Gonadotrophin status was not assessed in 1 paediatric patient, not documented for 5 deceased patients and unknown in 2 patients who did not follow-up in our institution. Of those for whom data was available, 58 patients (93.5%) were deficient. Gonadotrophin deficiency was treated with sex hormone replacement in all men and in women who were younger than 51 years of age at the time of diagnosis.

Thyrotrophin (TSH) status was not documented for 1 patient who attended another institution and for 4 deceased patients. Of the remaining patients, 56 (86%) were TSH deficient. All patients with TSH deficiency received replacement therapy with thyroxine. Median daily dose was 112 µg of thyroxine, range 50 – 250 µg.

Diabetes insipidus (DI) was documented in 56 patients (81%), who all received replacement desmopressin (dDAVP) therapy. Median dDAVP

dosage was twice per day, with a range of nil to 6 doses per day.

Patients were advised to miss one dose at the weekend in order to avoid fluid overload. Four patients (7.1%) had adipsic diabetes insipidus.

Lipid profile and blood pressure were recorded after appropriate hormone replacement therapy. Median total cholesterol was 5.0 mmol/l, with a range of 3.3 – 8.2 mmol/l (target reference range <5 mmol/l). Median LDL cholesterol was 3.2 mmol/l, with a range of 1.1 – 5.2 mmol/l (target < 2.6 mmol/l), HDL was 1.1 mmol/l (range 0.6 – 2.8 mmol/l, target > 1 mmol/l) and triglyceride was 2.2 mmol/l (range 0.6 – 6.7 mmol/l, target <1.9 mmol/l). Twenty-three patients of the 49 survivors (46.9%) had total cholesterol of more than 5 mmol/l, of whom only 11 were known to be on a statin. There was no difference between CP patients who died and those who continued to attend the outpatients in any lipid parameter.

Five of the surviving 49 patients had a blood pressure of equal to or greater than 140/90 mmHg, 2 of whom were treated with antihypertensives. One other patient had achieved normal blood pressure with an ACE inhibitor. Ten patients of historically normal blood pressure were followed up elsewhere. Current data therefore suggests that 6/39 (15.3%) have hypertension.

Six patients out of 52 for whom results were available were had diabetes mellitus (11.5%); 1 was diagnosed with Type 1 diabetes in childhood

prior to the diagnosis of CP. Median fasting glucose was 4.8 mmol/l (range 2.3 – 11 mmol/l).

Median BMI in CP patients was 31.8 kg/m² (range 20.3 – 61.2 kg/m²). 66% of CP patients had a BMI in the obese range (greater than 30 kg/m²) and 26% were overweight (BMI 25 – 29.9 kg/m²). The CP patients were compared to two other hypopituitary cohorts identified from the local database: non-functioning adenoma patients (NFPA; median BMI 29.8, 47% obese) and those with post-traumatic hypopituitarism (PTHP; median BMI 26.6, 31% obese). There was a significant difference in BMI across the three groups (Figure 2 and Table 2, $p < 0.001$, Wilcoxon). Subanalysis revealed that there was no difference between the BMI of CP and NFPA patients (Tukey HSD) but there was a difference between both CP and NFPA patients compared to PTHP patients. Student t test with correction for multiple comparisons revealed no significant difference between CP and NFPA BMI, but a significant difference between CP and PTHP ($p < 0.001$) and between NFPA and PTHP ($p < 0.05$). There was also a significant difference in age between the groups (Table 2).

Hydrocephalus was documented radiologically in 40.3% of patients, and was more prevalent in patients who were deceased at the time of analysis than in live patients (chi square, $p < 0.0001$). Seizures affected 49% of patients and were associated with hydrocephalus (chi square, p

1
2
3 < 0.0001). Seventy-seven percent of patients had documented visual
4
5 field loss.
6
7

8
9
10 Seven patients had a history of development of either a deep venous
11
12 thrombus (DVT) or a pulmonary embolus (PE). Post-mortem studies
13
14 revealed that PE was the cause of death in 2 / 21 deceased patients
15
16
17 (Table 1).
18
19

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

DISCUSSION

Craniopharyngioma (CP) is a tumour with a higher mortality than pituitary adenomata that is not explained by endocrine morbidity alone. Our management of craniopharyngioma is comparable to that in other tertiary centres, thus our findings are representative of the wider experience with CP. The elevated SMR for CP in our centre (8.75) is comparable to that of published international studies. In an analysis of the West Midlands database reported by Tomlinson et al, the SMR was 9.28 in craniopharyngioma patients ², an excess mortality which was mainly attributed to respiratory and cardiovascular disease. A single-centre Swedish study, which analysed mortality in 60 CP patients, and which documented an elevated SMR of 5.5 ³, reported that 7 of 27 deaths were related to pneumonia or PE. Our single-centre study also documents an excess of deaths due to respiratory disease. There is a consistent reporting of excess respiratory and vascular disease in the available literature. Because this was a retrospective study, we were forced to depend on the accuracy of death certificate reports, the majority of which were completed in hospitals other than our own. In Ireland, physicians are legally required to report cause rather than mechanism of death. Knowledge of the mechanism of death, such as respiratory failure, would have been of greater clinical value in assessment of mortality. For instance, our clinical suspicion is that some of the deaths were attributable to respiratory failure due to documented severe obstructive sleep apnoea.

1
2
3 The authors of both the UK and the Swedish studies suggested that
4
5 untreated GH deficiency contributed to the excess mortality of
6
7 hypopituitary patients. There are confounding factors in their studies
8
9 that should be considered, however, when analysing whether their data
10
11 fully substantiates this suggestion. Both series included patients
12
13 diagnosed prior to the use of dynamic testing for GHD in adults, which
14
15 questions the accuracy of diagnosis of GH deficiency in some patients.
16
17 In addition, early post-operative deaths, prior to testing for GH
18
19 deficiency may have led to under-estimation of the rate of GHD. We
20
21 have a policy of offering GH replacement to all eligible GHD adults with
22
23 craniopharyngioma, which differs slightly from historical practice in the
24
25 West Midlands centre; our data did not show any survival benefit of GH
26
27 treatment compared with those who were not treated. Larger studies,
28
29 with multicentre cooperation would probably be required to answer this
30
31 question.
32
33
34
35
36
37
38
39
40

41 Female patients with hypopituitarism due to any cause have a higher
42
43 mortality than their male counterparts ^{2, 3, 10}. Our data suggested a trend
44
45 in this direction, though the relative risk of death for women compared
46
47 to men did not reach statistical significance. This is likely to reflect the
48
49 small absolute numbers of deaths involved in our series. It has been
50
51 speculated that sub-optimal sex steroid replacement in women may
52
53 contribute to this reversal of gender mortality when compared to the
54
55 background population, and the Swedish group have suggested that the
56
57 pre-diagnosis clinical history of oestrogen deficiency should be
58
59
60

1
2
3 included in the total time of oestrogen deficiency in order to give some
4
5 indication of increased cardiovascular risk ^{10, 11}. We would observe,
6
7 however, that one of our patients, who subsequently died of respiratory
8
9 failure, developed multiple pulmonary emboli on oestrogen replacement,
10
11 during an episode of hypernatraemic dehydration due to adipsic
12
13 diabetes insipidus. Hormonal treatment is not always straightforward,
14
15 therefore, in this group of patients.
16
17
18
19

20
21
22 The excess mortality in CP patients may be related to hypothalamic
23
24 damage and its complications, such as adipsic DI ¹² and obesity ^{2, 10, 13,}
25
26 ¹⁴. A recent, single centre study in Holland reported an elevated SMR of
27
28 2.88 in 54 CP patients, with higher rates of obesity and diabetes than the
29
30 background Dutch population ¹⁰. Obesity is a recognised complication
31
32 of CP and also of the effects of treatment of CP, with surgery and
33
34 radiotherapy ¹⁵⁻¹⁷. We report a very high prevalence of CP obesity (66%)
35
36 in our CP cohort compared to the prevalence in Irish people of 18% ¹⁸.
37
38 We expected a high rate of obesity in our CP cohort, but were surprised
39
40 by the high incidence of obesity in our NFPA cohort, which was not
41
42 statistically different to that in CP. The NFPA group was significantly
43
44 older than the CP cohort, and as age is an independent associate of
45
46 obesity, we compared the rates of obesity in CP to age-matched patients
47
48 with hypopituitarism secondary to brain trauma; our data showed a
49
50 considerable excess of obesity in the groups with tumoural
51
52 hypopituitarism. Patients with PTHP appear to suffer from significantly
53
54 less obesity than patients with tumoural hypopituitarism, but they have
55
56
57
58
59
60

1
2
3 a much shorter duration of follow-up because PTHP has been
4
5 recognised relatively recently. The excess of obesity in the younger CP
6
7 group is clinically very relevant, as obesity in a younger population
8
9 predicts a higher risk of future morbidity and mortality. Obesity has
10
11 been suggested, for instance, to contribute to cardiovascular mortality
12
13 in CP ¹⁰. We have reported deaths due to sleep apnoea in CP patients ¹²,
14
15 and the possible contribution of obesity-related sleep apnoea to
16
17 respiratory mortality in CP patients deserves formal study.
18
19
20
21
22
23

24
25 Neurosurgical intervention in CP is often performed before full pre-
26
27 operative endocrine assessment ³. Thus we cannot comment on the
28
29 relative effects of pre-operative CP tumour bulk and surgical
30
31 intervention on pituitary function. The 3 patients who were assessed
32
33 before the adoption in 1997 of formal dynamic testing for GH deficiency,
34
35 and who were considered to have adequate GH secretion on the basis of
36
37 a normal IGF1 result, would probably have failed GH stimulation testing,
38
39 given that they had evidence of deficiency of all other pituitary
40
41 hormones ¹⁹. The true rate of GHD in our CP cohort is therefore more
42
43 likely to be 96%. GH, gonadotrophin and TSH deficiencies in our cohort
44
45 were comparable to international reports ^{10, 20}. ACTH deficiency has
46
47 been reported to range from 55 to 88% in a review of the literature by
48
49 Karavitaki, but our prevalence figures were a little higher at 92%.
50
51
52
53
54
55
56

57
58 Craniotomy is a frequently used surgical approach for management of
59
60 craniopharyngioma, particularly in cases of large tumour size and

1
2
3 suprasellar location ^{15, 20, 21}. Our use of the craniotomy approach is at
4
5 the higher end of international practice and might have contributed to
6
7 the high rates of hypopituitarism and hypothalamic complications
8
9 reported in our cohort. Gross total resection is rarely achieved in CP
10
11 patients, and where surgical clearance is attempted, imaging studies
12
13 can reveal a remnant in 18 – 26% of cases ²⁰. Thus a more conservative
14
15 surgical debulking procedure followed by irradiation of the remnant is
16
17 currently favoured in the hope that this approach might be less
18
19 destructive to hypothalamic and pituitary function. In our centre,
20
21 radiotherapy was used both for prevention of recurrence and as a
22
23 therapeutic intervention for recurrence. In spite of our local practice to
24
25 attempt gross total resection, recurrence rate (46%) was comparable to
26
27 previous studies ²⁰.
28
29
30
31
32
33

34
35
36 Craniopharyngioma patients are at increased cardiac and
37
38 cerebrovascular risk. Our cardiovascular risk assessment policy
39
40 identified significant dyslipidaemia. The low rate of statin prescription
41
42 probably reflects the youth of the CP population. A recent analysis of
43
44 Irish cardiovascular mortality since the 1980s attributed 30% of
45
46 improvement in survival to a 4.6% decrease in the population total
47
48 cholesterol ²², which suggests that our Irish CP cohort could benefit
49
50 from more aggressive management of dyslipidaemia. Hypertension
51
52 does not appear to be as prevalent a cardiovascular risk factor in the CP
53
54 group, but the rate of diabetes (11.5%) is much higher than that of the
55
56 background Irish population (4.8%) ²³. The majority of measurements of
57
58
59
60

fasting glucose were recorded at baseline of pituitary stimulation testing, prior to hormone replacement. The true rate of diabetes mellitus may be higher because GH and glucocorticoids are known to affect glucose metabolism ^{24, 25}, hence measurement of fasting glucose after hormone replacement may be a more appropriate evaluation. Given the spectrum of cardiorespiratory disease in CP, it may be valuable to adopt a formal screening programme for vascular risk factors such as diabetes mellitus, hypertension and hyperlipidaemia, and a more aggressive treatment schedule.

Frequent visual field loss and hydrocephalus reflected the large size of the tumours. The prevalence of seizures in our patient cohort was closely correlated with hydrocephalus and was higher than the rate of seizure in Oxford CP patients, which reported a 26% probability of seizure over 20 years of follow-up ¹⁶. Hydrocephalus was more prevalent in patients who subsequently died, but not in patients diagnosed in childhood; this data from our study is a novel finding which deserves further study ²⁰.

CP patients are recognised to suffer high rates of DI ⁸. DI patients are prone to the development of dehydration, volume contraction and hypernatraemia if they omit dDAVP replacement therapy or have limited access to fluid. Hypernatraemic dehydration increases the risk of thrombosis; this complication is particularly common in patients with adipsic DI, in whom we have described an increased frequency of

thrombotic disease¹². We recommend treatment of patients with adipsic DI with low molecular weight heparin prophylaxis during hypernatraemic episodes¹² and this intervention for DI patients with intact thirst might have averted two of the deaths from pulmonary embolus in this cohort. Adipsic DI is a well recognised, but not a common complication of CP; in a series of 13 patients with ADI, 4 had a diagnosis of CP, which reflected the destructive effect of tumour bulk and surgery on the hypothalamus^{12, 26}. As adipsic DI is recognised to confer increased likelihood of premature death, and one of the young deaths in this cohort was attributable to complications of adipsic DI, we feel that this is a significant marker for increased morbidity and mortality in CP.

CONCLUSION

We have shown that mortality is increased in an Irish CP population and that women are more affected than male patients. In our cohort increased mortality was not related to GH deficiency, or replacement, but was associated with a diagnosis of hydrocephalus, a probable marker of tumour size. There was a high rate of respiratory mortality, comparable to that previously published. The cause for increased mortality in CP patients is likely to be multifactorial, and includes excess hypothalamic morbidity, the high rate of obesity, hypopituitarism and adipsic diabetes insipidus. We did not find evidence for inappropriate hormone replacement therapy as a cause of morbidity or mortality in this study.

Strategies aimed at reducing mortality in CP are therefore only likely to be successful if they include a multifactorial approach, including preservation of hypothalamic structures at surgery, aggressive management of obesity and cardiovascular parameters, and management aimed at reducing the dangerous effects of sleep disturbance and adipsic diabetes insipidus.

Our study results confirm the high morbidity and mortality associated with a diagnosis of craniopharyngioma.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

The authors wish to acknowledge the assistance of G O'Connor,
actuary, for calculation of Standardised Mortality Ratio.

Dr Rachel Crowley is the beneficiary of an unrestricted educational
grant from Novo Nordisk.

For Peer Review

References

- 1 Dekkers, O.M., Biermasz, N.R., Smit, J.W., Groot, L.E., Roelfsema, F., Romijn, J.A. & Pereira, A.M. (2006) Quality of life in treated adult craniopharyngioma patients. *Eur J Endocrinol* 154, 483-489.
- 2 Tomlinson, J.W., Holden, N., Hills, R.K., Wheatley, K., Clayton, R.N., Bates, A.S., Sheppard, M.C. & Stewart, P.M. (2001) Association between premature mortality and hypopituitarism. West Midlands Prospective Hypopituitary Study Group. *Lancet* 357, 425-431.
- 3 Bulow, B., Attewell, R., Hagmar, L., Malmstrom, P., Nordstrom, C.H. & Erfurth, E.M. (1998) Postoperative prognosis in craniopharyngioma with respect to cardiovascular mortality, survival, and tumor recurrence. *J Clin Endocrinol Metab* 83, 3897-3904.
- 4 DeVile, C.J., Grant, D.B., Hayward, R.D. & Stanhope, R. (1996) Growth and endocrine sequelae of craniopharyngioma. *Arch Dis Child* 75, 108-114.
- 5 Poretti, A., Grotzer, M.A., Ribi, K., Schonle, E. & Boltshauser, E. (2004) Outcome of craniopharyngioma in children: long-term complications and quality of life. *Dev Med Child Neurol* 46, 220-229.
- 6 Agha, A., Rogers, B., Mylotte, D., Taleb, F., Tormey, W., Phillips, J. & Thompson, C.J. (2004) Neuroendocrine dysfunction in the acute phase of traumatic brain injury. *Clinical Endocrinology* 60, 584-591.
- 7 Seckl, J. & Dunger, D. (1989) Postoperative diabetes insipidus. *BMJ* 298, 2-3.
- 8 Smith, D., Finucane, F., Phillips, J., Baylis, P.H., Finucane, J., Tormey, W. & Thompson, C.J. (2004) Abnormal regulation of thirst and

vasopressin secretion following surgery for craniopharyngioma. *Clin Endocrinol (Oxf)* 61, 273-279.

Altman, D.G. & Bland, J.M. (2003) Interaction revisited: the difference between two estimates. *British Medical Journal* 326, 219.

Pereira, A.M., Schmid, E.M., Schutte, P.J., Voormolen, J.H., Biermasz, N.R., van Thiel, S.W., Corssmit, E.P., Smit, J.W., Roelfsema, F. & Romijn, J.A. (2005) High prevalence of long-term cardiovascular, neurological and psychosocial morbidity after treatment for craniopharyngioma. *Clin Endocrinol (Oxf)* 62, 197-204.

Erfurth, E.M., Bulow, B., Svahn-Tapper, G., Norrving, B., Odh, K., Mikoczy, Z., Bjork, J. & Hagmar, L. (2002) Risk factors for cerebrovascular deaths in patients operated and irradiated for pituitary tumours. *J Clin Endocrinol Metab* 87, 4892-4899.

Crowley, R.K., Sherlock, M., Agha, A., Smith, D. & Thompson, C.J. (2007) Clinical insights into adipsic diabetes insipidus: a large case series. *Clin Endocrinol (Oxf)* 66, 475-482.

Daousi, C., Dunn, A.J., Foy, P.M., MacFarlane, I.A. & Pinkney, J.H. (2005) Endocrine and neuroanatomic features associated with weight gain and obesity in adult patients with hypothalamic damage. *Am J Med* 118, 45-50.

de Vile, C.J., Grant, D.B., Hayward, R.D., Kendall, B.E., Neville, B.G. & Stanhope, R. (1996) Obesity in childhood craniopharyngioma: relation to post-operative hypothalamic damage shown by magnetic resonance imaging. *J Clin Endocrinol Metab* 81, 2734-2737.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 15 Verhelst, J., Kendall-Taylor, P., Erfurth, E.M., Price, D.A., Geffner, M., Koltowska-Haggstrom, M., Jonsson, P.J., Wilton, P. & Abs, R. (2005) Baseline characteristics and response to 2 years of growth hormone (GH) replacement of hypopituitary patients with GH deficiency due to adult-onset craniopharyngioma in comparison with patients with nonfunctioning pituitary adenoma: data from KIMS (Pfizer International Metabolic Database). *J Clin Endocrinol Metab* 90, 4636-4643.
- 16 Karavitaki, N., Brufani, C., Warner, J.T., Adams, C.B., Richards, P., Ansorge, O., Shine, B., Turner, H.E. & Wass, J.A. (2005) Craniopharyngiomas in children and adults: systematic analysis of 121 cases with long-term follow-up. *Clin Endocrinol (Oxf)* 62, 397-409.
- 17 Pinkney, J., Wilding, J., Williams, G. & MacFarlane, I. (2002) Hypothalamic obesity in humans: what do we know and what can be done? *Obes Rev* 3, 27-34.
- 18 (2005) Obesity: the Policy Challenges - The report of the National Taskforce on Obesity. Irish Department of Health and Children.
- 19 Toogood, A.A., Beardwell, C.G. & Shalet, S.M. (1994) The severity of growth hormone deficiency in adults with pituitary disease is related to the degree of hypopituitarism. *Clin Endocrinol (Oxf)* 41, 511-516.
- 20 Karavitaki, N., Cudlip, S., Adams, C.B. & Wass, J.A. (2006) Craniopharyngiomas. *Endocr Rev* 27, 371-397.
- 21 Kendall-Taylor, P., Jonsson, P.J., Abs, R., Erfurth, E.M., Koltowska-Haggstrom, M., Price, D.A. & Verhelst, J. (2005) The clinical, metabolic and endocrine features and the quality of life in adults with

childhood-onset craniopharyngioma compared with adult-onset craniopharyngioma. *Eur J Endocrinol* 152, 557-567.

22 Bennett, K., Kabir, Z., Unal, B., Shelley, E., Critchley, J., Perry, I., Feely, J. & Capewell, S. (2006) Explaining the recent decrease in coronary heart disease mortality rates in Ireland, 1985-2000. *J Epidemiol Community Health* 60, 322-327.

23 The Institute of Public Health in Ireland (2006) Making Diabetes Count, a systematic approach to estimating population prevalence on the island of Ireland in 2005. The Institute of Public Health in Ireland, Belfast.

24 Gathercole, L.L., Bujalska, I.J., Stewart, P.M. & Tomlinson, J.W. (2007) Glucocorticoid modulation of insulin signalling in human subcutaneous adipose tissue. *Journal of Clinical Endocrinology and Metabolism* 92, 4332-4339.

25 Jorgensen, J.O., Moller, L., Krag, M., Billestrup, N. & Christiansen, J.S. (2007) Effects of growth hormone and fat metabolism in human subjects. *Endocrinol Metab Clin North Am* 36, 75-87.

26 Smith, D., McKenna, K., Moore, K., Tormey, W., Finucane, J., Phillips, J., Baylis, P. & Thompson, C.J. (2002) Baroregulation of vasopressin release in adipsic diabetes insipidus. *J Clin Endocrinol Metab* 87, 4564-4568.

TABLES

<u>Patient</u>	<u>Gender</u>	<u>Age at Diagnosis CP</u>	<u>Age at death</u>	<u>Cause of Death</u>
1	F	64	71	Myocardial infarction
2	F	48	51	CP
3	F	74	81	Respiratory arrest
4	M	47	49	Respiratory arrest
5	M	63	68	Not reported
6	M	26	61	Epilepsy
7	M	34	38	Metastatic Lung Carcinoma
8	F	52	58	Small cell lung carcinoma
9	F	30	32	Pulmonary Embolus
10	F	66	68	Respiratory Tract Infection
11	M	70	82	Cerebral Thrombus
12	M	36	45	Aspiration Pneumonia
13	M	43	43	Cerebral Infarct
14	F	44	46	CP
15	F	39	40	CP
16	M	3	8	Pneumonia
17	F	16	23	Increased Intracranial pressure
18	F	13	13	Pulmonary Embolus
19	M	7	19	Glioma
20	M	48	53	Left ventricular failure
21	M	11	20	Head trauma secondary to seizure

Table 1. Mortality data of 21 craniopharyngioma patients. F = female, M = male, CP = craniopharyngioma. Cause of death reported as stated on death certificate (not located for patient 5).

<u>Diagnosis</u>	<u>n</u>	<u>Median Age</u> <u>years</u>	<u>Median BMI</u> <u>kg/m²</u>	<u>% BMI > 30</u> <u>kg/m²</u>
CP	70	27.5	31.7	66
NFPA	89	52	29.8	47
PTHP	29	41	26.6	31
p value		< 0.001	< 0.001	

Table 2. Comparison of CP patients to NFPA and PTHP groups. Age at diagnosis is different for NFPA group from both CP and PTHP group. Difference between CP and PTHP group is borderline non-significant (p = 0.052). BMI is different in PTHP group from both CP and NFPA.

Figures

Figure 1. Distribution of CP patients by age at diagnosis. Line shows bimodal distribution

Figure 2. Comparison of BMI in Hypopituitary patients by underlying cause of hypopituitarism. CP = craniopharyngioma (n = 70), NFPA = non-functioning adenoma (89), PTHP = post-traumatic hypopituitarism (n = 29).