Study of lead phytoavailability for atmospheric industrial micronic and sub-micronic particles in relation with lead speciation

Gaëlle Uzu, Sophie Sobanska, Yassine Aliouane, Philippe Pradere, Camille Dumat

► To cite this version:

Gaëlle Uzu, Sophie Sobanska, Yassine Aliouane, Philippe Pradere, Camille Dumat. Study of lead phytoavailability for atmospheric industrial micronic and sub-micronic particles in relation with lead speciation. *Environmental Pollution*, 2009, 157 (4), pp.1178-1185. 10.1016/j.envpol.2008.09.053 . hal-00559688

HAL Id: hal-00559688

<https://hal.science/hal-00559688>

Submitted on 26 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Study of lead phytoavailability for atmospheric industrial micronic and submicronic particles in relation with lead speciation.

G. Uzu^a, S. Sobanska^b, Y. Aliouane^a, P. Pradere^c and C. Dumat^{a*}

^a-EcoLab UMR 5245 CNRS-INPT-UPS, ENSAT BP 32607 Auzeville Tolosane, 31326 Castanet Tolosan, France.

^b-LASIR UMR 8516 - Université des Sciences et Technologies de Lille - Bâtiment C5 - 59655 Villeneuve d'Ascq Cedex, France.

^c- Chemical Metal Treatment Company, STCM, 30-32 chemin de Fondreyre, 31200 Toulouse, France.

* Corresponding author: Camille Dumat. Tel.: 335 62 19 39 03; fax: 335 62 19 39 01.

E-mail address: camille.dumat@ensat.fr.

gaelle.uzu@ensat.fr; Sophie.Sobanska@univ-lille1.fr; p.pradere@stc-metaux.com;

Capsule

The soil-lettuce lead transfer from atmospheric industrial sub-micronic and micronic particles depends on particle size.

Abstract

Particles from channelled emissions of a battery recycling facility were size-segregated and investigated to correlate their speciation and morphology with their transfer towards lettuce. Microculture experiments carried out with various calcareous soils spiked with micronic and submicronic particles ($1650 \pm 20 \text{ mgPb.kg}^{-1}$) highlighted a greater transfer in soils mixed with the finest particles. According to XRD and Raman spectroscopy results, the two fractions presented differences in the amount of minor lead compounds

like carbonates, but their speciation was quite similar, in decreasing order of abundance: PbS, PbSO₄, PbSO₄.PbO, α -PbO and Pb⁰. Morphology investigations revealed that PM_{2.5} (i.e. Particulate Matter 2.5 composed of particles suspended in air with aerodynamic diameters of 2.5 micrometers or less) contained many Pb nano balls and nano crystals which could influence lead availability. The soil-plant transfer of lead was mainly influenced by size and was very well estimated by 0.01M CaCl₂ extraction.

Keywords: lead, PM_{2.5} and PM₁₀, soil-lettuce transfer, phytoavailability.

1. Introduction

Due to its numerous past and present uses and high persistence, lead is a major environmental contaminant (Chen et al., 2006). Potentially toxic for living organisms even at low concentrations, lead constitutes a risk for humans who can absorb it in various ways (Canfield et al., 2003). In the context of contaminated gardens, elevated lead intake by humans can be due to the consumption of crop plants grown on soils with relatively high plant-available metal concentrations, ingestion of contaminated soil, either accidentally or intentionally (pica), inhalation of soil particles and drinking water with high soluble concentrations of metals (Alexander et al., 2006). The total quantities of lead emitted in the environment by industries have decreased sharply in recent decades (Glorennec et al., 2007) and are strictly controlled in Europe nowadays. Lead was recently classified as a substance of very high concern in the European REACH law (Regulation EC 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals). However, particles enriched with lead are still generated especially by

lead-recycling plants (Batonneau et al., 2004; Ohmsen, 2001) and constitute the main source of lead pollution for soils (Miquel, 2001; Donisa et al., 2000).

According to Zhang et al. (2005), emitted particles present a large granulometric spectrum in the atmosphere, but during the last decade the proportion of fine particle matter (PM) increased with the use of more effective filters in industry. Indeed PM₁₀ are target species of the World Health Organization (WHO, 2001) and the European Union Framework Directive on ambient air quality assessment (EU, 1999), due to their adverse effects on the environment and human health. While micrometric and sub-micrometric fractions contribute very little to ambient particle mass, they may occur in substantial number concentrations. Most of the studies dealing with the characterization of metal-enriched particles in the ambient air provide information on quantitative measurements for PM₁₀ fractions (EU directives 96/62 and 99/30) and very few on the sub-micronic fraction (Lazaridis et al., 2002). The lack of knowledge regarding metal speciation in the industrial particles results mainly from a lack of analytical tools, both sensitive and specific to the size of the particles.

These fine particles are highly reactive due to their high specific area and can be transported over long distances in the troposphere (Barrie, 1992). They could therefore present a greater impact on the biosphere than coarse particles (Fernandez Espinosa, 2005). Ruby et al. (1992) concluded that the bioaccessibility of lead rises strongly in particles under 2.4 µm size. But, the phytoavailability of lead in industrial particles as a function of their size and speciation have not been studied yet. In comparison with zinc, lead generally shows a relatively low mobility in soils (Dumat et al., 2006). It can however migrate through the soil with dissolved organic matter (Cecchi et al., 2008) or be mobilized by certain plants (Arshad et al., 2008). Moreover, carried from the air to

the soils as fine particles, lead could be released more easily in soil solution (Komarnicki, 2005).

We therefore focused our study on the links between soil-plant transfer of lead, size and speciation of particles emitted by a lead recycling plant, currently the main source of atmospheric emissions (Cecchi et al., 2008). The objectives were the following: (i) the elemental and molecular characterisation of micrometer and nanometer sized lead-rich particles and (ii) to study the influence of particle characteristics on lead soil-plant transfer.

The physico-chemical characterization of industrial PM₁₀ and PM_{2.5} particles collected in the plant was investigated using both bulk and micro analysis techniques: (i) MEB-EDS to determine the morphology and chemistry on the scale of a particle (Laskin et al., 2006; Choel et al., 2005); (ii) Raman microspectrometry to study particle speciation (Batonneau et al., 2004; 2006; Falgayrac et al., 2006; Sobanska et al., 2006). The transfer of lead from particles to the lettuce *Lactuca sativa* a widely grown garden vegetable was investigated in the laboratory: two different uncontaminated calcareous soils were spiked with PM₁₀ and PM_{2.5} for soil-plant experiments with a biotest device that enabled careful study of rhizosphere and roots in addition to the transfer to the shoots (Chaignon and Hinsinger, 2003). The study was finally completed by CaCl₂ extraction experiments carried out according to Houba et al. (1996) to estimate lead phyto-availability.

The hypothesis tested throughout all these experiments was that particle characteristics have a significant influence on lead soil-plant transfer and translocation.

2. Materials and methods

2.1. Particle sampling and size separation.

A secondary lead smelter which currently recycles batteries was chosen as a representative example of the smelter metal industry to develop a methodology aimed at the risk assessment of industrial lead particles. The plant of the Chemical Metal Treatment Company (STCM) is located in the urban area of Toulouse (43°38'12" N, 01°25'34" E). According to the French authorities (DRIRE, 2007), 328 kg of Total Suspended Particles (TSP) including 31 kg of lead were emitted by this factory in 2007.

Three sources of particles corresponding to the three work units involving different steps in the process are identified in the plant: (i) the battery grinding unit where battery components are separated under wet conditions (ii) the smelter where lead pastes are processed in rotary furnaces at 1200°C and finally (iii) the refinery where lead is purified from unwanted metals or enriched. The same process has been used for thirty years in the plant. This study focuses on channeled emissions only generated by the furnace unit, considering the transfer towards the biosphere. The particle collection, performed three months after complete cleaning of the three work units (March, 12, 2007), is therefore representative of that emission period. Our aim was to characterize the reactivity of the particles in relation with their previously determined speciation and size, not to follow one specific parameter over time.

1kg of particles was collected in polyethylene bags from the air-sleeve filters of the furnace, then sealed and transferred in sealed opaque containers for transport to the laboratory. Samples were passed through a 2-mm stainless steel AFNOR sieve. Process dust was stored in a cool (4°C) dark place with Merck desiccant.

Because 80 % of emitted particles were smaller than 10 µm, PM_{2.5} and PM₁₀ were size-segregated by artificial resuspension in a Teflon bag (Batonneau et al., 2004; Young et al. 2002) and collected by impaction onto a PM_{10/2.5} Dekati inertial impactor. The cascade impactor consists of two successive stages with aerodynamic cut-off

diameters of 10 and 2.5 μm when it operates at 10 L/min airflow. PM_{10} present an aerodynamic diameter between 2.5 and 10 μm , whereas the $\text{PM}_{2.5}$ stage collects particles $< 2.5\mu\text{m}$.

2.2. Characterization of particles

Elemental total contents of the two size fractions were determined by ICP-OES (IRIS Intrepid II XDL) after heated digestion with standard acid (HNO_3 , HCl and HF , Suprapur, Merk) in a PTFE vessel. Levels of C, H, S and O were determined after burning in an elemental analyser with coulometric-catharometric detection and IR.

X-ray diffraction patterns of PM_{10} and $\text{PM}_{2.5}$ were recorded on an INEL diffractometer equipped with a curved detector CPS 120 and $\text{Co (K}\alpha\text{)}$ radiation allowing a $120^\circ/2\theta$ detection. Crystallized compounds were identified by comparison with the diffraction patterns of the JCPDS database. The relative abundance of each crystallized phase detected in samples was qualitatively estimated by using the relative intensity of the strongest X-ray pattern peaks of each considered phase. It should be noticed that only crystallized phases with an abundance superior to 5 % in weight can be detected by XRD.

Particles were characterized using complementary microscopy and imaging techniques to determine elementary and molecular composition, size, morphology and heterogeneity of the individual lead-rich particles.

X-ray elemental mapping and secondary images were obtained using an Environmental Scanning Electron Microscope (ESEM) working in high-vacuum mode. The Quanta 200 FEI instrument was equipped with an energy-dispersive X-ray (EDX) detector. ESEM was operated at 10-25 kV. Particles were dispersed on carbon

substrates and analysed without further preparation. Because of the ESEM configuration, light element detection (C, N, and O) was ambiguous.

Raman microspectrometric measurements were carried out with a LabRAM confocal spectrometer (Jobin Yvon, Horiba Gr, France). The spot size of the laser focused by a 100×objective (numerical aperture (NA) = 0.90) was estimated to be $1\text{ }\mu\text{m}^2$ in size. A liquid nitrogen-cooled CCD (Jobin-Yvon, 2048 pixels $\times$ 512 pixels) was used for detection. The Raman backscattering was excited at 632.8 nm provided by an internal, air-cooled, linearly polarized helium-neon laser. The laser power delivered to the sample was 8 mW. The microscope stage was XY-motorized and computer-controlled for point by point scanning with 0.1 μm resolution, 1 μm reproducibility and 90 mm $\times$ 60 mm spatial range. The glass plate with the impacted particles was mounted on the microscope stage without any further preparation. Data acquisition consisted in recording many spectra in point-by-point scanning mode with 1 μm as a minimum step, one accumulation and 30 s spectrum acquisition time. For identification of Raman spectra the experimental spectra were compared to reference spectra using Spectral Library Search ID 301 software (Thermo Galactic).

2.3. Measure of soil-plant lead transfer performing microculture experiments

Lettuce was chosen because it is a common vegetable widely grown for human consumption and has recently been used by several authors for metal transfer studies (Khan et al., 2007; Waisberg et al., 2004; Alexander et al., 2006). Moreover, lettuce is often used around plants to estimate atmospheric pollution.

The biotest presented on figure 1 was first described by Niebes et al. (1993) and then adapted by Chaignon and Hinsinger (2003). A small PVC cylinder (25 mm inner diameter) was closed by a polyamide net (900 μm mesh) inserted into a larger cylinder,

itself closed by a finer polyamide mesh (30 μm , Fyltis/Nytel, Sefar filtration). A space of 3 mm was left between the net and the finer mesh, where the roots could develop as a mat.

Commercial lettuce seeds, “Batavia blonde dorée” cultivar, were surface sterilized with 0.9% CaClO for 15 min and rinsed with deionised water. Three seeds were sown in the container on the surface of the coarser mesh.

Lettuces were first grown hydroponically for 21 days to obtain a large flat mass of roots that fully covered the fine mesh.

The devices were placed on top of troughs containing an aerated complete nutrient solution with the macroelements: 5 mM KNO_3 , 5 mM $\text{Ca}(\text{NO}_3)_2$, 2 mM KH_2PO_4 and 1.5 mM MgSO_4 and oligoelements: 9.11 μM MnSO_4 , 1.53 μM ZnSO_4 , 0.235 μM CuSO_4 , 24.05 μM H_3BO_3 , 0.1 μM Na_2MoO_4 and 268.6 μM Fe/EDTA . The height of the nutrient solution was adjusted daily to keep the fine mesh wet during the whole pre-culture period. The experiment was conducted in a growth chamber (temperature 24 ± 0.5 $^\circ\text{C}$ /18 $\pm$ 0.5 $^\circ\text{C}$ day/night cycles; photoperiod 16 h under daylight fluorescent lamps providing 400 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ [Philips 600W, Eindhoven, Netherlands] and 8 h darkness; relative humidity 70%).

Then, the second step involved an eight-day soil-plant contact period: five replicate plants for each treatment were transferred onto the various soil samples.

Top soils of two calcic cambisol profiles (FAO, 1998) produced by quaternary alluvial deposits, were sampled. These two calcareous soils (noted soil-1 and soil-2) with various physicochemical properties (table 1) were chosen because that kind of soil happens to be observed around several battery recycling plants. Moreover, if the lettuce species could favor lead transfer, the choice of calcareous soils could reduce it. The soils differ by texture, soil cultivation, organic matter (OM) and CEC: soil-1 was

uncultivated loamy-sandy clay with 2% of OM and soil-2 was sandy-clayey loam under sunflowers with 1% of OM and a higher CEC.

These two soils were spiked with $PM_{2.5}$ and PM_{10} up to $[Pb] = 1650 \pm 20 \text{ mg.kg}^{-1}$. 60 mg of particles were added per box containing 10 g of soil and were agitated for a day. Soils were turned over every fifteen days for four months. This long period was chosen to allow time for natural equilibration of the various sorption mechanisms in the soil (Alexander et al., 2006). The relatively high total lead soil concentration was chosen to be representative of a real soil pollution situation observed for atmospheric fallout from an STCM plant (Cecchi et al., 2008) and we wanted to be sure that enough lead was available for the plant in soil solution. Surveys of garden soils in several countries have shown wide ranges of concentrations of heavy metals. For example, Culbard et al. (1988) found concentrations of up to $14,100 \text{ mg Pb kg}^{-1}$. Some soils were left without crops, to determine the pH possible variation due to soil solution influence. During the growth, pH was measured daily and nutrient solution level was maintained constant. pH of nutrient solution was 5.5 ± 0.3 during the hydroponic and soil contact phases.

This biotest device presents two main advantages: (i) the roots are physically separated from the soil which enables total recovery of the shoots, roots and soil; (ii) the thickness of the soil layer used enables it to be considered it as rhizosphere soil and provides enough rhizosphere material to evaluate root-induced changes in metal speciation.

Roots and aerial parts were collected and analysed separately after the soil plant contact period. Biomass was determined before oven-drying at 80°C for 48H. Lead bound to the outer root cell walls, called $[Pb]_{\text{adsorbed}}$, was determined according to the method of acidic desorption as described by Ferrand et al. (2006): roots were shaken end over end with 40mL of 0.001M HCl for 3 min, and then 360 μL of 1 M HCl was

added to yield a final concentration of 0.01 M HCl. After shaking for another 5 min, the suspension was filtered through ashless paper. Then, plant roots and aerial parts were mineralized separately in a 1:1 mixture of HNO₃ and H₂O₂ at 80°C for 4h. After filtration, the major and trace element concentrations were determined with an IRIS Intrepid II XDL ICP-OES. The concentration in roots after mineralization is considered as uptaken lead ([Pb]_{roots}). The accuracy of the acidic digestion and analytical procedures was verified using the reference material Virginia tobacco leaves, CTA-VTL-2, ICHTJ.

2.4. Estimation of lead availability by CaCl₂ extraction experiments

For the determination of the phytoavailable fraction, according to Menzies et al. (2007) neutral salts extractants provide the most useful indication. The 0.01 M CaCl₂ extraction procedure gives a good indication of lead phytoavailability (Meers et al., 2007). The CaCl₂ procedure, first described by Houba, et al. (1996), was performed on the two contaminated soils and particle samples (PM₁₀ and PM_{2.5}). 30 mL of 0.01 M CaCl₂ solution was mixed with 3 g of soil (1:10 solid solution ratio) in 50ml polypropylene centrifugation tubes placed on a shaker table (Heidolph promax 1020) at 50 oscillations/min for 2h, 20°C. After extraction, the tubes were centrifuged at 10,000 g for 30 min (Avanti 30 centrifuge Beckman). The supernatant liquid was then filtered (0.22µm), acidified to 2% with distilled HNO₃ (15M, suprapur 99.9%) and stored at 4°C before analysis. That CaCl₂ extraction procedure was performed on particles, reference soils, and the two spiked soils before and after culture. Extracted metal concentrations were determined by an IRIS Intrepid II XDL/ Thermo Electron Corporation model ICP-AES Calibration used reference materials (ion 915 from National Water Research Institute, Canada).

2.5. Statistical data treatment

The plant absorption and chemical extraction data obtained were subjected to analysis of variance (ANOVA) with one factor, using the software Statistica, Edition'98 (StatSoft Inc., Tulsa, OK, USA). For each bioassay, mean values with different letters represent a significant difference ($p < 0.05$) as measured by the LSD Fisher test. Letters are reported on the figures and tables.

3. Results

3.1. PM₁₀ and PM_{2.5} characterization:

Elemental concentrations in particles are shown in table 2. All results are given as the mean of the three replicates for each sample (PM₁₀ and PM_{2.5}) and standard deviations never exceed 7%. No significant difference except for Fe in the total elemental concentrations was observed in relation with the size of the particle. Major elements found in the samples were, by mass: Pb (27%), O (15%) and S (7.5%) for both fractions. High levels of Na (3-4%) were due to the industrial recycling process where Na is used to lower the melting point of Pb. Several other metals: Cd (2.5%), Zn (0.5%), Fe (0.1-0.4%) and Sb (0.1%) are also present. The remaining elements to complete to 100% are expected to be chloride and other trace metals.

XRD patterns of PM₁₀ and PM_{2.5} provide identification of major crystallized species in bulk samples. The results are presented in table 3. Particles are mainly composed of metallic sulfides, sulfates, oxides and perchlorates. A significant amount of Na₂SO₄ was found in PM₁₀ samples. Regardless of the size, the major phases identified were the same and one more Fe species was detected for PM_{2.5}.

The ESEM-EDX analysis provided the morphology and elemental composition of individual particles in the two size fractions. PM₁₀ samples exhibited both particles with a size range between 2 and 10 μm and large aggregates composed of many micron-sized particles without specific shapes (figure 2). The main elements detected by EDX in these aggregates were, by order of importance, Pb, S, Cl, Sn, Na, and Fe. The chemical complexity of aggregates did not allow the detection of minor elements. Elemental mapping recorded on PM₁₀ showed the chemical heterogeneity of aggregates. PM_{2.5} samples are composed of fine aggregates of few submicronic particles exhibiting characteristic features i.e. needles (< 100 nm), nano crystals (< 500 nm), cubes (~300 nm) and balls (~500 nm). Elemental analysis of particles showed that needles mainly contain Pb and S, nano crystals are composed of Pb, S and Cl while Na and S are detected in cubic particles. Compared to the PM₁₀ sample, numerous Pb only-rich nanoballs were observed in PM_{2.5} (figure 2).

Raman microspectrometry combines the analytical capability of Raman spectroscopy to distinguish *in situ* a wide range of chemical substances in aerosols with the spatial resolution of a confocal optical microscope (~1 μm^3) which enables investigation of individual particles. The scanning of a large area of sample (300 μm^2) by automated analyses enables the analysis of a huge number of particles. The data treatment of all the spectra recorded in this area leads to the major and minor component identification. $x\text{PbO} \cdot \text{PbSO}_4$ ($x = 1, 2, 3$), PbSO_4 , $\alpha\text{-PbO}$, PbCO_3 , Na_2SO_4 , and ZnSO_4 were identified in both PM₁₀ and PM_{2.5} samples as major species. The laser-damaging effect generates complete and irreversible oxidation of PbS to PbSO_4 as described previously (Batonneau et al., 2000). The Raman mapping of species within particles exhibits a large heterogeneity of particle composition. A typical Raman image of PM_{2.5} particles is shown in figure 3. A large number of particles are composed of

lead sulphate and are agglomerated with other compounds such as sodium sulphate or carbonates in both PM₁₀ and PM_{2.5} size fractions. The results suggest that no significant differences in lead speciation are observed in accordance with the particle size. However, for the minor lead compounds (like carbonates) not exactly quantified by XRD, differences in percentages between the two size fractions could exist.

3.2. Assessment of the available fraction with CaCl₂ extraction

Figure 4A and 4B shows respectively total lead concentrations extracted by CaCl₂ and percentages of extracted lead with CaCl₂ (relative, with respect to total lead concentrations in different substrates) for various samples: the single particles, soil-1 and spiked soil-1 before and after an eight-day exposure of *Lactuca sativa*.

Due to high levels of lead in particles (330,000 mg.kg⁻¹), maximum lead quantities were extracted from PM₁₀ and PM_{2.5}. But, in comparison with total lead contents, lead in uncultivated spiked soils was 0.2% exchangeable, while PM_{10-2.5} presented a 0.01% exchangeability. Reference soils were under ICP-OES limit detection. After culture, a significantly greater lead quantity was extracted by CaCl₂ for the soil spiked with the finest PM_{2.5} fraction than in PM₁₀. A 1 unit pH decrease was measured after plant-soil contact while the pH values of the control soils (left without crops but with nutrient solution influence) remained constant. Concerning CaCl₂ extraction, no significant differences were observed between the two particle sizes in the PM_{10-2.5} and spiked soil-1 before culture; The same trends described above for soil-1 were also observed for soil-2.

3.3. Influence of lead exposure on plant growth.

Fresh biomass data are shown in figure 5. After eight days of soil-plant exposure, biomasses for aerial parts grown on spiked soils were slightly lower than in the respective unspiked soils. No significant biomass differences were observed among the two types of spiking ($PM_{10-2.5}$). In controls, fresh weights were approximately 18.5 ± 2.5 g for aerial parts and 3.7 ± 0.7 g for roots while in spiked soils, weights reached 12 ± 4 g and 2.5 ± 1 g respectively. This slight influence of lead on plant biomass could be due to lead toxicity (Sharma & Dubey, 2005) or water status (Parys et al., 1998).

3.4. Absorption and adsorption of lead in the lettuce

Figure 6 presents results of lead transfer from soil to the various compartments of the plant (roots and shoots) and the distribution between adsorption (noted $[Pb]_{adsorbed}$) and absorption for roots ($[Pb]_{roots}$) respectively described in the materials and method section as lead only adsorbed at the root surface and lead truly taken up by the plant. Measurements on the reference samples (Tobacco leaf VTL-2) validated the assay: the concentration found was 21.4 ± 1.1 mg.kg⁻¹, for a certified value of 22.1 ± 1.2 mg.kg⁻¹. Several trends were observed dealing with the influence of the particle type, soil type and lead location. For both soils, a significant increase of lead adsorption on roots, absorption and translocation throughout the shoots was observed, when the finest particles were added in comparison with the PM_{10} . Global uptakes ($[Pb]_{adsorbed} + [Pb]_{roots} + [Pb]_{shoots}$) were greater for soil 1 than for soil 2. The main lead fraction was adsorbed on root membranes where concentrations were up to 985 ± 147 .kg⁻¹ for PM_{10} contaminated soils and up to 1281 ± 195 mg.kg⁻¹ for $PM_{2.5}$ contaminated soils. The soils spiked with PM_{10} allowed a lead uptake of 230 ± 24 mg.kg⁻¹ by roots and a translocation to aerial parts of 7.7 ± 2.7 mg.kg⁻¹. While in the $PM_{2.5}$ spiked soils, root and shoot concentrations reached 275 ± 40 mg.kg⁻¹ and 12.19 ± 3 mg.kg⁻¹ respectively.

4. Discussion

(1) Influence of particle size on soil-plant transfer and lead translocation.

Whatever the soil, for a given total lead concentration ($1650 \text{ mg Pb.kg}^{-1}$ soil), higher lead soil-plant transfer and translocation were observed for the finest particles. Roots exposed to $\text{PM}_{2.5}$ spiked soils allowed a 20% greater lead uptake and a 30% increase in adsorbed lead. Shoots presented a 60% increase in translocated lead in PM_{10} spiked soils. For the first time in industrial particles, the transfer of lead to the soil solution and its translocation throughout the plant is reported to increase as the particle size decreases.

Douay et al. (2006), measuring lead concentrations in lettuces in urban gardens ($1572 \text{ mg Pb.kg}^{-1}$) observed a shoot concentration of 5 mg Pb.kg^{-1} DW. The soil to plant metal transfer measured in this study (for one week of lead exposure) was relatively high in comparison with previously reported data in publications dealing with vegetables (BAPPET, 2007), probably because of particle size, but also due to numerous factors that influence the soil-plant transfer of metals (Kabata-Pendias, 2004; Dumat et al., 2001 & 2006; Costa & Morel, 1993; Oliver et al., 1994).

Root transfer factors (TF-roots) and shoot transfer factors (TF-shoots) were calculated as ratios between $[\text{Pb}]_{\text{roots}}$, $[\text{Pb}]_{\text{shoots}}$ with $[\text{Pb}]_{\text{soil}}$. TFr ranged from 0.10 to 0.17 and TFs were between 0.005 and 0.007 (table 4). These relatively low values illustrate the known low mobility and availability of lead and are comparable to Khan et al.'s results (2008) reporting 0.15 for TFr and 0.07 for TFs. As transfer factors for vegetables decrease with increasing levels of lead in the soil (Zheng et al., 2007; Wang et al., 2006), we therefore expected a greater influence of particle size on transfer with a lower total lead concentration in soil.

According to Sobanska et al. (1999), lead smelter emissions contain PbS, PbSO₄, PbSO₄.PbO, Pb, α-PbO compounds and numerous nano balls were observed for PM_{2.5}. Dumat et al. (2001) and Cecchi et al. (2008) working on soils polluted by lead produced by industrial atmospheric fallout from smelters, concluded that lead chemical speciation strongly influences its bioavailability. The sequence of solubility constants from the CHESS data base for the lead is the following: PbCO₃ > PbSO₄ > PbO > Pb⁰. In water media and equilibrium conditions, lead carbonate and sulphate will be much more labile than PbO or Pb⁰ species. However, according to Birkefeld et al. (2006 & 2007) which used an in-situ method to study dissolution and phase transformation of lead particles from a smelter in different soils, the sequence of solubility is strongly dependent on soil characteristics (like texture, pH, lime amount...). They observed that PbO was rapidly covered by lead-hydroxy carbonates (hydrocerussite) in some soils while in other it was relatively stable. Moreover, according to the general review of Ruby et al. (1992), release of lead in the soil solution depends on particle size, speciation and soil geochemistry. In soil solution, ligands like fulvic acids or low weight organic acids excreted by roots can displace the equilibrium (Ferrand et al., 2006). Finally, as no significant difference among speciation forms was noticed between PM₁₀ and PM_{2.5}, all changes observed in reactivity were attributed to size differences.

Whatever the particle size, greater lead absorption by lettuce was observed for soil-1 than for soil-2. In comparison with soil-2, soil-1 has more clay, its CEC is higher and its level of lime is lower (see Table 1). According to Twining et al. (2004), we could expect that lead transfer would be higher for soil-2. However, several hypotheses can explain the results observed: (i) the higher amount of carbonates present in the soil-2 could reduce lead absorption (Birkefeld et al., 2006 & 2007); (ii) the complex influence of soil organic matter on the transfer of metals (Yin et al., 2002; Inaba &

Takenaka, 2005); (iii) for the total lead concentration studied, the relatively high quantity of lead could be available in the soil solution (due to high solubility of the fine particles) reducing the influence of soil characteristics.

Total lead in the roots ($[Pb]_{\text{roots}} + [Pb]_{\text{adsorbed}}$) reached 1500 mgPb/ kg dry weight. Lead in roots was predominantly adsorbed onto the cell walls, 80 % of the total lead in roots, and depends on the nature of the particles used for exposure: the uptake was greater for PM_{2.5} than PM₁₀. According to Seregin et al. (2004), Pb²⁺ binds to the carboxy groups at the root surface reducing the translocation rate of lead (Pendergrass et al., 2006; Piechalak et al., 2002). To illustrate this phenomenon, the translocation factors or shoot/root ratios calculated indicate the ability of plants to transport metals from the roots toward the aerial parts (Ferrand et al., 2006): they ranged between 0.03 and 0.04 (table 4). But considering the lower biomasses in roots than in shoots, we can also reason with lead quantities. Whatever the type of soil or spiking, the total lead quantity in shoots ($Pb_{\text{shoots}} \times \text{dry weight}$) between 0.003 and 0.007 mg Pb translocated which represents up to 33% (PM_{2.5}) of the total lead uptake. This percentage is not negligible with respect to risks concerning consumption of vegetables.

(2) Behaviour of lead in the rhizosphere and assessment of lead availability

Relatively high lead transfer was observed for the lettuce cultivated on the two alkaline spiked calcareous soils. PM_{10-2.5} can therefore release lead in particular in the rhizosphere of *Lactuca sativa*. The lead availability estimated by the CaCl₂ procedure was greater for soils spiked with the finest particles and for both soils it increased after soil-plant contact: $[Pb]_{\text{CaCl}_2}$ in spiked soil with PM_{2.5} > $[Pb]_{\text{CaCl}_2}$ in spiked soil with PM₁₀. Under the root activity influence, the CaCl₂ lead extracted from polluted soils increased and a one-unit pH decrease in soil was measured. Lin et al. (2004) and Kidd

and Monterroso (2005) also observed that exchangeable lead was much higher in the rhizosphere than in the bulk soil. Producing exudates, plants can modify metal speciation and behaviour in the rhizosphere (Lin et al., 2004; Laperche, 1997; Welch, 1995). This phenomenon has been particularly observed for calcareous soil by Chaignon and Hinsinger (2003). As pH influences metal solubility and transfer (Wang et al., 2006), the rhizosphere acidification could have displaced the equilibrium towards bicarbonates, which are less stable than carbonates (Sauvé et al., 1998). An effect of soil on particle solubility was also observed: particles present a CaCl_2 exchangeability ten times lower than in spiked soil. Quantities extracted for particles were only up to 1.5% of the solution extraction. Mixed with soil for four months, particle solubility could therefore have changed.

In order to estimate the transfer of lead from polluted soils towards lettuce, relationships were sought between: lead mobilised by CaCl_2 extraction performed on soils before culture and lead concentrations in lettuce (shoots and roots). Equations were obtained from 6 parameters (2 soils; uncontaminated, spiked with PM_{10} or $\text{PM}_{2.5}$), and every condition was studied by 5 replicates finally involving 30 observations. Significant correlations were observed between lead concentrations in shoots (Equation 1 below, with $r^2 = 0.8$) or in roots (Equation 2 below, with $r^2 = 0.9$) and lead extracted by CaCl_2 .

Equation 1: $[\text{Pb}]_{\text{shoots}} = 2.11 \times [\text{Pb}]_{\text{CaCl}_2} + 2.13 \quad r^2 = 0.797 \quad p < 0.01 \quad n = 30$

Equation 2: $[\text{Pb}]_{\text{roots}} = 35.02 \times [\text{Pb}]_{\text{CaCl}_2} + 4.24 \quad r^2 = 0.915 \quad p < 0.005 \quad n = 30$

Both for PM_{10} and $\text{PM}_{2.5}$, the CaCl_2 chemical extraction was therefore a good indicator of soil-plant transfer. As previously shown by Pueyo et al. (2004), the 0.01 mol l^{-1} CaCl_2 extraction procedure seems to be a suitable method for the determination of phytoavailable lead. This test simplifies the matrix and could avoid *in vivo* tests.

Conclusions and perspectives

A significant size influence was found for soil-plant lead transfer and translocation throughout the lettuce when micronic and nanometric industrial particles were compared: roots exposed to PM_{2.5} spiked soils allowed a 20% greater lead uptake and a 30% increase in adsorbed lead. Shoots presented a 60% increase in translocated lead in PM₁₀ spiked soils. Our results highlight that source characteristics strongly influence metal transfer: total metal soil concentration is insufficient to estimate the risk induced by soils polluted by metals. Lead speciation was quite similar in PM₁₀ and PM_{2.5} particles i.e. PbS, PbO.PbSO₄, PbO, PbCO₃ and Pb⁰, were predominant however differences could exist for minor lead components. Moreover, the high amount of Pb-rich nano balls and nano crystals much more dispersed for PM_{2.5} could influence the lead transfer in the rhizosphere acidified by plant root activity. Lead concentrations in the edible part of lettuces were very well estimated by 0.01M CaCl₂ extraction.

In order to check the generality of the size influence of industrial particles on soil-plant transfer, further experiments could be performed on other vegetables and various soils testing aging effects. Moreover, the study of other metals and metalloids measured in the particles will be performed.

Acknowledgements

ADEME, the French Agency for Environment and Energy as well as the STCM, are gratefully acknowledged for their financial support and technical help. This research project was supported by the National CNRS CYTRIX-EC2CO program. We thank Dr J. Silvestre from ENSAT for cropping advice, P. Recourt from *Laboratoire Geosystèmes*, UMR CNRS 8157, University of Lille for ESEM-EDX measurements and

J. Laureyns from LASIR UMR CNRS 8516, for Raman Microspectrometry assistance.
Finally, Dr P. Winterton from UPS is acknowledged for English review.

References

Alexander, P. D., Alloway, B. J. & Dourado, A. M., 2006. Genotypic variations in the accumulation of Cd, Cu, Pb and Zn exhibited by six commonly grown vegetables. *Environmental Pollution* 144(3), 736-745.

Arshad, M., Silvestre, J., Pinelli, E., Kallerhoff, J., Kaemmerer, M., Shahid, M., Pradere, P. & Dumat, C., 2008. A field study of lead phytoextraction by various scented *Pelargonium* cultivars. *Chemosphere* 71, 2187–2192.

Bahemuka, T. E. & Mubofu, E.B., 1999. Heavy metals in edible green vegetables grown along the sites of the Sinza and Msimbazi rivers in Dar es Salaam, Tanzania. *Food Chemistry* 66(1), 63-66.

BAPPET –Database on ETM in vegetables. ADEME, INERIS, CNAM, INP, ENSAT,ISA.2008.<http://www.sitespollues.ecologie.gouv.fr/DocumentsDiagnostics.asp#BAPPET>.

Barrie, L. A., 1992. Scavenging Ratios: Black Magic or Useful Scientific Tool?, in S. E. Schwartz and W. G. N. Slinn (eds), *Precipitation Scavenging and Atmosphere-Surface Exchange*, Volume1, Hemisphere Publishing Corporation, Washington.

Batonneau Y., Brémard, C., Gengembre, L., Laureyns, J., Le Maguer, A., Le Maguer, D., Perdrix, E., and Sobanska S., 2004. Speciation of PM₁₀ sources of airborne nonferrous metals within the 3-km zone of lead/zinc smelters. *Environ. Sci. Technol.* 38, 5281-5289.

Batonneau, Y., Sobanska, S., Laureyns, J., Bremard, C., 2006. Confocal microprobe raman imaging of urban tropospheric aerosol particles. *Environ. Sci. Technol.* 40, 1300-1306.

Batonneau, Y., Brémard, C., Laureyns, J. Merlin, J.C., 2000. Microscopic and imaging Raman scattering study of PbS and its photo-oxidation products. *Journal of Raman Spectroscopy* 31, 1113-1119.

Birkefeld, A., Schulin, R., Nowack, B., 2007. In situ transformations of fine lead oxide particles indifferent soils. *Environ. Pollut.* 145: 554-561.

Birkefeld, A., Schulin, R., Nowack, B., 2006. In situ investigation of dissolution of heavy metal containing mineral particles in an acidic forest soil. *Geochim. Cosmochim. Acta.* 70: 2726-2736.

Canfield, R.L., Henderson, C.R., Jr., Cory-Slechta, D.A., Cox, C., Jusko, T.A and Lanphear, B.P., 2003. Intellectual impairment in children with blood lead concentrations below 10 µg per deciliter. *N. Engl. J. Med.* 348, 1517-1526.

Cecchi, M., Dumat, C., Alric, A., Felix-Faure, B., Pradere, P., Guiresse, M., 2008. Multi-metal contamination of a calcic cambisol by fallout from a lead-recycling plant. *Geoderma* 144, 287-298.

Chaignon, V., Hinsinger, P., 2003. A biotest for evaluating copper bioavailability to plants in a contaminated soil. *J Environ Qual* 32, 824-833.

Chen, Y., Wang, C., Wang, Z., 2005. Residues and source identification of persistent organic pollutants in farmland soils irrigated by effluents from biological treatment plants. *Environ. Int.* 31, 778–783.

Choel, M.; Deboudt, K.; Osan, J.; Flament, P.; Van Grieken, R., 2005. quantitative determination of low-Z elements in single atmospheric particles on boron

substrates by automated scanning electron microscopy-energy-dispersive X-ray Spectrometry Anal. Chem. 77(17), 5686-5692.

Choël, M., Deboudt, K., Flament, P., Lecornet, G., Perdrix, E., Sobanska, S., 2006. Fast evolution of tropospheric Pb- and Zn-rich particles in the vicinity of a lead smelter. Atmospheric Environment 40, 4439-4449.

Costa, G., Morel, J.L., 1993. Cadmium uptake by *Lupinus albus* (L.): cadmium excretion, a possible mechanism of cadmium tolerance. Journal of plant nutrition (USA) 16(10), 1921-1929.

Culbard, E.B., Thornton, I., Watt, J., Wheatley, M., Moorcroft, S., Thompson, M., 1988. Metal contamination in British suburban dusts and soils. Journal of Environmental Quality 17, 226-234.

Donisa, C., Mocanu, R., Steinnes, E. And Vasu, A., 2000. Heavy Metal Pollution by Atmospheric Transport in Natural Soils from the Northern Part of Eastern Carpathians. Water, Air, and Soil Pollution 120(3-4), 347-358.

Douay, F., Pruvot, C., Mazzuca, M., Howsam, M., Fourier, H., and De Saint Mahieu, A.L., 2005. Cadmium, lead and zinc concentrations in soil and vegetables from kitchen gardens in urban and highly-contaminated areas of northern France: evaluation of the risk of population exposure, ConSoil 2005. Proceedings of the 9th International FZK/TNO Conference on Soil–Water Systems. Bordeaux, France (2005), 667–676.

DRIRE, Direction régionale pour l'Industrie, la Recherche et l'Environnement, <http://www.midi-pyrenees.drire.gouv.fr/>

Dumat, C., Chiquet, A., Gooddy, D., Aubry, E., Morin, G., Juillot, F., Benedetti, M.F., 2001. Metal ion geochemistry in smelter impacted soils and soil solutions. Bulletin De La Societe Geologique De France 172, 539-548.

Dumat, C., Quenea, K., Bermond, A., Toinen, S., Benedetti, M.F., 2006. Study of the trace metal ion influence on the turnover of soil organic matter in cultivated contaminated soils. *Environmental Pollution* 142, 521-529.

Ebert, M., Inerle-Hof, M. and Weinbruch, S., 2002. Environmental scanning electron microscopy as a new technique to determine the hygroscopic behaviour of individual aerosol particles. *Atmospheric Environment* 36, 5909–5916.

European Commission, 96/62/CE directive (September 27, 1996) for the quality of ambient air; directive 22/04/99 setting maximum levels for lead in ambient air; REACH directive Regulation (EC) No 1907/2006, Directive 67/548/EEC.

Falgayrac, G., Sobanska, S., Laureyns, J., Brémard, C., 2006. Heterogeneous chemistry between PbSO_4 and calcite microparticles using Raman microimaging. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 64, 1095-1101.

FAO, 1998. World reference base for soil research. *World Soil resources Report*, v. 84. FAO, Rome.

Fernandez Espinosa, A.J., Rossini Oliva, S., 2006. The composition and relationships between trace element levels in inhalable atmospheric particles (PM_{10}) and in leaves of *Nerium oleander* L. and *Lantana camara* L. *Chemosphere* 62, 1665-1672.

Ferrand, E., Dumat, C., Leclerc-Cessac, E., Benedetti, M.F., 2006. Phytoavailability of zirconium in relation to its initial added form and soil characteristics. *Plant and Soil* 287, 313-325.

Glorennec, P., Bemrah, N., Tard, A., Robin, A., Le Bot, B., Bard, D., 2007. Probabilistic modeling of young children's overall lead exposure in France: Integrated approach for various exposure media. *Environment International* 33, 937-945.

Houba, V.J.G., Lexmond, T.M., Novozamsky, I., van der Lee, J.J., 1996. State of the art and future developments in soil analysis for bioavailability assessment. *Science of The Total Environment* 178, 21-28.

Inaba, S., Takenaka, C., 2005. Effects of dissolved organic matter on toxicity and bioavailability of copper for lettuce sprouts. *Environment International*, 31, 603-608.

Kabata-Pendias, A., 2004. Soil-plant transfer of trace elements an environmental issue. *Geoderma* 122, 143-149.

Khan, S., Aijun, L., Zhang, S.Z., Hu, Q.H., Zhu, Y.G., 2008. Accumulation of polycyclic aromatic hydrocarbons and heavy metals in lettuce grown in the soils contaminated with long-term wastewater irrigation. *Journal of Hazardous Materials* 152, 506-515.

Kidd, P.S., Monterroso, C., 2005. Metal extraction by *Alyssum serpyllifolium* ssp. *lusitanicum* on mine-spoil soils from Spain. *Science of The Total Environment* 336, 1-11.

Komarnicki, G.J.K., 2005. Lead and cadmium in indoor air and the urban environment. *Environmental Pollution* 136, 47-61.

Laperche, V., Logan, T.J., Gaddam, P., Traina, S.J., 1997. Effect of apatite amendments on plant uptake of lead from contaminated soil. *Environ. Sci. Technol.* 31, 2745-2753.

Laskin, A., Cowin, J.P., Iedema, M.J., 2006. Analysis of individual environmental particles using modern methods of electron microscopy and X-ray microanalysis. *Journal of Electron Spectroscopy and Related Phenomena* 150, 260-274.

Lazaridis M., Semb A., Larssen S., Hjellbrekke A.G., Hov O., Hanssen J.E., Schaugh J., Torseth K., 2002. Long-range transport of aerosol particles and precursors -

determining the relationship between emissions and observed concentration levels. The science of the total environment 285, 209-235.

Lin, Q., Chen, Y.X., He, Y.F., Tian, G.M., 2004. Root-induced changes of lead availability in the rhizosphere of *Oryza sativa* L. Agriculture, Ecosystems & Environment 104, 605-613.

Meers, E., Samson, R., Tack, F.M.G., Ruttens, A., Vandegehuchte, M., Vangronsveld, J., Verloo, M.G., 2007. Phytoavailability assessment of heavy metals in soils by single extractions and accumulation by *Phaseolus vulgaris*. Environmental and Experimental Botany 60, 385-396.

Menzies, N.W., Donn, M.J., Kopittke, P.M., 2007. Evaluation of extractants for estimation of the phytoavailable trace metals in soils. Environmental Pollution 145, 121-130.

Miquel, G., 2001. Les effets des métaux lourds sur l'environnement et la santé. Rapport parlementaire européen.

Ohmsen, G. S., 2001. Characterization of fugitive material within a primary lead smelter. J. Air Waste Manage. Assoc. 51, 1443-1451.

Oliver, D.P., Hannam, R., Tiller, K.G., Wilhelm, N.S., Merry, R.H. and Cozens, G.D., 1994. The effects of zinc fertilization on cadmium concentration in wheat grain. J. Environ. Qual. 23, 705-711.

Parys, E., Romanowska, E., Siedlecka, M., and Poskuta, J., 1998. The effect of lead on photosynthesis and respiration in detached leaves and in mesophyll protoplasts of *Pisum sativum*. Acta Physiologiae Plantarum 20(3), 313-322.

Pendergrass, A., Butcher, D.J., 2006. Uptake of lead and arsenic in food plants grown in contaminated soil from Barber Orchard, NC. Microchemical Journal 83, 14-16.

- Piechalak, A., Tomaszewska, B., Baralkiewicz, D., Malecka, A., 2002. Accumulation and detoxification of lead ions in legumes. *Phytochemistry* 60, 153-162.
- Pueyo, M., Lopez-Sanchez, J.F., Rauret, G., 2004. Assessment of CaCl_2 , NaNO_3 and NH_4NO_3 extraction procedures for the study of Cd, Cu, Pb and Zn extractability in contaminated soils. *Analytica Chimica Acta* 504, 217-226.
- Ruby, M.V., Davis, A., Kempton, J.H., Drexler, J.W., Bergstrom, P.D., 1992. Lead Bioavailability: Dissolution Kinetics under Simulated Gastric Conditions. *Environ. Sci. Technol.* 26, 1242-1248.
- Ruby, M.V., Schoof, R., Brattin, W., Goldade, M., Post, G., Harnois, M., Mosby, D.E., Casteel, S.W., Berti, W., Carpenter, M., Edwards, D., Cragin, D., Chappell, W., 1999. Advances in evaluating the oral bioavailability of inorganics in soil for use in human health risk assessment. *Environ. Sci. Technol.* 33, 3697-3705.
- Sauvé, S., Dumestre, A., McBride, M., Hendershot, W., 1998. Derivation of soil quality criteria using predicted chemical speciation Pb^{2+} and Cu^{2+} . *Environmental Toxicology and Chemistry* 17(8), 1481-1489.
- Seregin, I.V., Kozhevnikova, A.D., 2004. Strontium transport, distribution, and toxic effects on maize seedling growth. *Russian Journal of Plant Physiology* 51, 215-221.
- Sharma, P., and Dubey, R.S., 2005. Lead toxicity in plants. *Brazilian Journal of Plant Physiology* 17, 35-52.
- Sobanska, S., Ricq, N., Laboudigue, A., Guillermo, R., Bremard, C., Laureyns, J., Merlin, J.C., Wignacourt, J.P., 1999. Microchemical investigations of dust emitted by a lead smelter. *Environ. Sci. Technol.* 33, 1334-1339.

Sobanska S., Pauwels B., Maenhaut W. and Adams F., 2002. Single particle characterisation and sources of tropospheric aerosols in the Negev desert (Israel). *Journal de Physique IV* 12(Pr10), 161-183.

Sobanska, S., Falgayrac, G., Laureyns, J., Bremard, C., 2006. Chemistry at level of individual aerosol particle using multivariate curve resolution of confocal Raman image. *Spectrochim Acta A Mol Biomol Spectrosc.* 64, pp. 1102–1109.

Twining, J.R., Payne, T.E., Itakura T., 2004. Soil-water distribution coefficients and plant transfer factors for ^{134}Cs , ^{85}Sr and ^{65}Zn under field conditions in tropical Australia. *J Environ Radioac* 71, 71-87.

Waisberg, M., Black, W.D., Waisberg, C.M., Hale, B., 2004. The effect of pH, time and dietary source of cadmium on the bioaccessibility and adsorption of cadmium to/from lettuce (*Lactuca sativa* L. cv. *Ostinata*). *Food and Chemical Toxicology* 42, 835-842.

Wang, G., Su, M.Y., Chen, Y.H., Lin, F.F., Luo, D., Gao, S.F., 2006. Transfer characteristics of cadmium and lead from soil to the edible parts of six vegetable species in southeastern China. *Environmental Pollution* 144, 127-135.

Welch, R.M., 1995. Micronutrient nutrition of plants. *Critical Reviews in Plant Sciences* 14, 49-82.

WHO, 1987. Air quality guidelines for Europe. WHO Regional Publications, European Series No. 23.

Yin, Y., Impellitteri, C.A., You, S.J., Allen, H.E., 2002. The importance of organic matter distribution and exact soil : solution ratio on the desorption of heavy metals from soils. *The Science of the Total Environment* 287, 107-119.

Young, T.M., Heeraman, D.A., Sirin, G., Ashbaugh, L.L., 2002. Resuspension of soil as a source of airborne lead near industrial facilities and highways. *Environ. Sci. Technol.* 36, 2484-2490.

Zhang Z., C. Kleinstreuer, J.F. Donohue, C.S. Kim., 2005. Comparison of micro- and nano-size particle depositions in a human upper airway model. *J Aerosol Science* 36, 211–233.

Zheng, N., Wang, Q., Zheng, D., 2007. Health risk of Hg, Pb, Cd, Zn, and Cu to the inhabitants around Huludao Zinc Plant in China via consumption of vegetables. *Science of The Total Environment* 383, 81-89.

Figure legends**Figure 1**

Microculture device.

Figure 2

ESEM images of PM_{2.5} (A) and PM₁₀ (B).

Figure 3

Raman image of PM_{2.5} particles after a smoothing procedure. Blue 4PbO.PbSO₄, pink PbO.PbSO₄, red Na₂SO₄ and green substrate.

Figure 4

Total contents (A) and relative contents (B) of lead extracted with CaCl₂ on particles (white), spiked soils with lead particles of different size before (grey) and after (black) a seven-days exposure of *Lactuca sativa*, n=5.

Figure 5

Plant biomass data for aerial parts (black) and roots (grey). The error bars indicate the standard deviation while different letters show significant differences (p<0.05) as measured by an LSD fisher test (one-factor ANOVA).

Figure 6

Lead concentrations in different plant compartments: A: adsorbed onto roots, B: root uptake and C: translocated in shoots.

Table 1

Physico-chemical characteristics of the two soils used for micro-culture experiments.

	unit	Soil-1	Soil-2
pH		8.3	8.4
CEC	cmol+/kg	19.9	8.9
Clay	(g.kg ⁻¹)	405	195
Silt		261	390
Sand		334	415
C/N		9.15	8.9
MO	(g.kg ⁻¹)	18.8	12.5
Total carbonates	(g.kg ⁻¹)	36.8	98
P₂O₅ Joret Hebert	(g.kg ⁻¹)	0.28	0.215
Cd (HF extraction)	(mg.kg ⁻¹)	0.22	0.175
Pb (HF)		35.5	21
Zn (HF)		101	48
Cu (HF)		28.5	31

Table 2

Total elemental analysis of particles.

Sample	Contents (mg.kg ⁻¹) ± 7%												
	Al	As	Cd	Cu	Fe	O	Na	Ni	Pb	Sb	Zn	S	C
PM_{2.5}	10	867	25243	480	1395	149300	31691	41	272834	1266	5194	78100	12100
PM₁₀	52	841	23139	617	4010	150000	39697	97	267353	1361	5372	74300	11500

Table 3

Determination of main chemical speciation of lead in the PM₁₀ and PM_{2.5} particles by X-Ray Powder Diffraction Analysis ^a

Sample	Cristallized compound	Estimated % of phases
PM ₁₀	PbS ,	55
	PbO.PbSO₄ , PbSO₄ , Pb(ClO ₄) ₂ , Pb,	25
	Na₂SO₄	10
	CdS, Cd(ClO ₄) ₂ , ZnO, ZnSO ₄	10
PM _{2.5}	PbS ,	55
	PbO.PbSO₄ , PbSO₄ , Pb(ClO ₄) ₂ , Pb,	25
	Na₂SO₄	10
	CdS, Cd(ClO ₄) ₂ , ZnO, ZnSO ₄ , FeS ₂	10

^a The major compounds are written in bold

Table 4

Transfer factors in roots (TFr) and shoots (TFs) and translocation ratios.

Soil	TFs	TFr	Translocation
S1 + PM₁₀	0.005	0.14	0.03
S1 + PM_{2.5}	0.007	0.17	0.04
S1	0.081	0.26	0.31
S2 + PM₁₀	0.004	0.10	0.04
S2 + PM_{2.5}	0.005	0.15	0.03
S2	0.01	0.44	0.18

Figure 1

Figure 2

(A)

(B)

Figure 3

Figure 4
(A)

(B)

Figure 5

Figure 6

(A)

(B)

(C)